

Seed-mass dependence of baryon-assisted black-hole growth in self-interacting dark matter haloes

Dong-Biao Kang

Zhejiang GuangSha Vocational and Technological University of China

Institute of Theoretical Physics, Chinese Academy of Sciences

dbkang@zjgsdx.edu.cn

Abstract

We present a controlled numerical study of time-dependent black-hole growth in self-interacting dark matter (SIDM) haloes with an assembled baryonic potential. The calculation extends a conducting-fluid treatment with a monotonicity-preserving reconstruction and Roe fluxes to a finite Hernquist gas reservoir, Eddington-limited baryonic accretion, evolving Bondi ambient conditions, radiative cooling, and a conservative subgrid closure for capture by small seeds. A constant-cross-section validation suite reproduces the reference singular-isothermal-sphere and Navarro–Frenk–White growth histories: a $100\text{ M}_\odot$ seed reaches $129\text{ M}_\odot$ and $9.65 \times 10^3\text{ M}_\odot$, respectively, after 2 Myr when conduction is active. In the fiducial physical-cooling model, a $10^5\text{ M}_\odot$ seed in a $10^9\text{ M}_\odot$ halo at $z = 30$ crosses $10^7\text{ M}_\odot$ in 1.52–1.76 Myr across the two smallest completed inner boundaries, although its terminal mass remains 7.1% boundary sensitive. The nominal mass gain is 98.81% SIDM supplied. The baryonic response is non-monotonic in compactness, with a maximum at $a_b/r_s = 0.02$ rather than at the smallest tested radius. Physical cooling removes stored thermal feedback; under the adopted irreversible expansion closure, mixed feedback expands the Hernquist radius by about a factor of two, dilutes the gas, and reduces dark accretion by 33%. A conservative capture closure shows that 10^2 – $10^4\text{ M}_\odot$ seeds do not enter the rapid-growth branch in 2 Myr. The transition at a few $10^4\text{ M}_\odot$ is bracketed, rather than uniquely located, because it depends on the capture coefficient and feeding boundary. A Rutherford/Born velocity-dependent cross section removes the threshold crossing throughout the tested domain at low transition speed and retains only a small boundary-robust subset in the high-transport model. The robust numerical result is strong seed-mass and transport dependence in the explored parameter domain, not a population-level prediction or a generic route from stellar remnants to $10^7\text{ M}_\odot$.

1 Introduction

The existence of massive black holes at high redshift motivates growth channels that do not rely exclusively on long episodes of ordinary baryonic Eddington accretion [Haehnelt and Rees, 1993, Salpeter, 1964, Shakura and Sunyaev, 1973, Bromm and Loeb, 2003, Madau and Rees, 2001, Begelman et al., 2006, Volonteri, 2010, Inayoshi et al., 2020]. Self-interacting dark matter provides a complementary possibility. Elastic scattering can conduct heat through a halo, alter the central phase-space structure, and in the gravothermal regime drive a dense core toward collapse [Lynden-Bell and Wood, 1968, Spergel and Steinhardt, 2000, Balberg et al., 2002, Koda and Shapiro, 2011, Kaplinghat et al., 2016, Tulin and Yu, 2018]. The same transport can couple a black hole to a large reservoir of dark matter without invoking gas cooling at the capture radius. Recent work has connected SIDM core collapse to early massive-black-hole seeds and to the population of little red dots (LRDs) [Feng et al., 2021, 2025, Jiang et al., 2026].

Meng et al. [2026] solved the spherical conducting-fluid problem for a black hole embedded in singular-isothermal-sphere (SIS) and NFW initial profiles. That calculation established an important numerical and physical baseline: conduction can suppress the late central density and

accretion rate, while an SIS profile can still provide rapid growth from a $100 M_\odot$ seed. It does not, however, answer four questions that determine whether the mechanism is useful as a controlled growth model. First, how does a baryonic potential assembled on a finite time scale modify the supply region? Second, can a finite gas reservoir and feedback alter both the Bondi limiter and the dark-matter potential? Third, is the apparent runaway present for stellar-remnant seeds, or does it require an already intermediate-mass seed? Fourth, does the conclusion survive a velocity-dependent SIDM cross section and a controlled scan in halo and baryonic parameters?

Here we answer these questions with a sequence of matched experiments. The simulations remain one-dimensional and spherical, so the goal is not a cosmological population model. Instead, we test the local closure and identify which statements survive numerical, boundary, and microphysical checks. We use a finite-volume MC reconstruction, Roe approximate Riemann flux, Harten entropy correction, and an implicit conduction update. The baryons are represented by an evolving Hernquist reservoir [Hernquist, 1990]; their assembly history is a smoothstep function, and feedback is split into thermal and expansion channels. This extends earlier studies of baryonic potentials in gravothermal SIDM evolution [Zhong et al., 2023] to a time-dependent accretion problem. The gas state used by the Bondi rate is evolved with the reservoir and a no-ultraviolet-background Cloudy/Grackle cooling table [Ferland et al., 2017, Smith et al., 2017]. For small seeds, the resolved inward flux is separated from the unresolved capture rate by a mass-conserving central reservoir.

The paper has three deliberately distinct levels of claim. The first is code validation against the reference SIS/NFW problem and a suite of grid, CFL, entropy-fix, and outer-boundary tests. The second is a physical interpretation of the baryonic assembly and feedback response. The third is an applicability map in which the scientific classifier is whether $10^7 M_\odot$ is crossed within 2 Myr; terminal masses are reported, but are not promoted to precision predictions when the inner boundary changes them by more than the pre-declared 5% criterion. We leave seed abundances, LRD number densities, and observables to a separate population study.

2 Model

2.1 Spherical SIDM fluid

We evolve density ρ , radial velocity u , and one-dimensional velocity-dispersion variable v . The pressure is $P = \rho v^2$. In physical units the mass and momentum equations are

$$\frac{\partial \rho}{\partial t} + \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \rho u) = 0, \quad (1)$$

$$\frac{\partial u}{\partial t} + u \frac{\partial u}{\partial r} = -\frac{1}{\rho} \frac{\partial P}{\partial r} - \frac{GM_{\text{tot}}(< r, t)}{r^2}. \quad (2)$$

The internal-energy equation is written in the form used by the conducting-fluid closure,

$$\frac{3}{2} \rho \frac{Dv^2}{Dt} = -P \frac{1}{r^2} \frac{\partial(r^2 u)}{\partial r} - \frac{1}{r^2} \frac{\partial(r^2 q_r)}{\partial r}, \quad q_r = -\kappa \frac{\partial v^2}{\partial r}. \quad (3)$$

The enclosed mass includes the black hole, dark matter, and the baryonic Hernquist component. The closure is the usual harmonic interpolation between short- and long-mean-free-path limits,

$$\kappa = \frac{3}{2} \left[\frac{\sigma/m}{Bv} + \frac{1}{AC\rho v^3(\sigma/m)} \right]^{-1}, \quad A = \sqrt{\frac{16}{\pi}}, \quad B = \frac{25\sqrt{\pi}}{32}, \quad C = \frac{3}{4}, \quad (4)$$

where σ/m is the self-interaction cross section per unit mass. This is an effective gravothermal fluid model rather than a collisionless particle calculation; its validity is therefore assessed against the baseline and against the expected transport limits [Balberg et al., 2002, Koda and Shapiro, 2011, Outmezguine et al., 2023].

The code uses dimensionless scales $M_0 = 4\pi\rho_0 r_0^3$, $v_0 = (GM_0/r_0)^{1/2}$, and $t_0 = (4\pi G\rho_0)^{-1/2}$. We retain physical units in tables and figures. The reference initial halo is an NFW profile,

$$\rho_{\text{NFW}}(r) = \frac{\rho_s}{x(1+x)^2}, \quad x = r/r_s, \quad (5)$$

with M_{200} , concentration $c = R_{200}/r_s$, and $M(< R_{200}) = 200(4\pi/3)\rho_{\text{crit}}(z)R_{200}^3$ [Bryan and Norman, 1998]. SIS runs are used only as a validation control. The NFW and SIS choices span the cored/contracting and centrally concentrated limits discussed in numerical SIDM studies [Navarro et al., 1997, Dave et al., 2001, Rocha et al., 2013, Peter et al., 2013, Vogelsberger et al., 2012, Robertson et al., 2017].

2.2 Baryons and assembly

The baryonic reservoir has a Hernquist enclosed mass

$$M_b(< r, t) = M_{b,\text{rem}}(t) \frac{r^2}{[r + a_b(t)]^2}, \quad \Phi_b(r, t) = -\frac{GM_{b,\text{rem}}(t)}{r + a_b(t)}. \quad (6)$$

The parameter f_b denotes the fraction of M_{200} made available to the assembled central reservoir; it is a model input, not an assertion that every high-redshift halo retains this baryon fraction. Assembly is prescribed by

$$S(t) = \begin{cases} 3\tau^2 - 2\tau^3, & 0 < \tau = t/T_{\text{asm}} < 1, \\ 1, & t \geq T_{\text{asm}}, \end{cases} \quad M_{b,\text{asm}}(t) = f_b M_{200} S(t), \quad (7)$$

$$M_{b,\text{rem}}(t) = \max[M_{b,\text{asm}}(t) - \Delta M_{\text{gas}}(t), 0], \quad (8)$$

where ΔM_{gas} is the cumulative gas mass removed by black-hole accretion, before the radiative-efficiency correction to the retained black-hole mass. This smoothstep avoids an impulsive potential and permits a direct comparison between assembly time and local dynamical time.

The relevant local clocks are

$$t_{\text{dyn}} = \left[\frac{r^3}{GM_{\text{tot}}(< r)} \right]^{1/2}, \quad t_{\text{coll}} = [a\rho(\sigma/m)v]^{-1}, \quad t_{\text{cond}} = \frac{3\rho L^2}{2\kappa}, \quad t_{\text{in}} = \frac{r}{|u|}. \quad (9)$$

The supply radius is the logarithmic-radius median of the inward flux inside r_s . Timescales are evaluated in the initial no-baryon hydrostatic state at that radius, so the correlation is a pre-run diagnostic rather than a circular fit to the evolved state.

2.3 Gas accretion and feedback

The black-hole mass is separated into dark and baryonic channels,

$$\dot{M}_{\text{BH}} = \dot{M}_{\text{DM}} + \dot{M}_{b,\text{BH}}. \quad (10)$$

The gas supply is limited by the minimum of the Bondi and Eddington rates [Bondi, 1952],

$$\dot{M}_{\text{Bondi}} = 4\pi\lambda_B \frac{G^2 M_{\text{BH}}^2 \rho_\infty(t)}{[c_s^2(t) + v_{\text{rel}}^2]^{3/2}}, \quad (11)$$

$$\dot{M}_{\text{Edd}} = \frac{4\pi G M_{\text{BH}} m_p}{\epsilon_r \sigma_{\text{T}} c}, \quad \dot{M}_{b,\text{BH}} = (1 - \epsilon_r) \min(\dot{M}_{\text{Bondi}}, \dot{M}_{\text{Edd}}). \quad (12)$$

We use the standard adiabatic Bondi coefficient $\lambda_B = 1/4$ for $\gamma = 5/3$, $\epsilon_r = 0.1$, and zero relative velocity in the one-zone gas closure. The Eddington cap applies only to the baryonic term in equation (12); $\dot{M}_{\text{DM}}$ is non-radiative in this model and is not Eddington limited. The

instantaneous total black-hole mass, including captured dark matter, nevertheless enters $\dot{M}_{\text{Edd}}$ because it sets the gravitational mass relevant to the gas flow. Dark accretion can therefore raise the subsequent baryonic Eddington ceiling without itself producing the associated radiation.

The ambient density and sound speed are coupled to the finite reservoir,

$$\rho_{\infty}(t) = \rho_{\infty,0} \frac{M_{b,\text{rem}}(t)}{M_{b,0}} \left[\frac{a_{b,0}}{a_b(t)} \right]^3, \quad (13)$$

$$c_s^2(t) = c_{s,0}^2 + \frac{\gamma(\gamma-1)E_{\text{th}}(t)}{M_{b,\text{rem}}(t)}. \quad (14)$$

The feedback energy is split into a thermal reservoir and an irreversible expansion channel,

$$\frac{dE_{\text{th}}}{dt} = \chi_{\text{heat}} \epsilon_f \epsilon_r \dot{M}_{\text{gas}} c^2 - \frac{E_{\text{th}}}{t_{\text{cool}}}, \quad (15)$$

$$a_b(t) = a_{b,0} \left[1 + \frac{(1 - \chi_{\text{heat}}) \epsilon_f \epsilon_r \Delta M_{\text{gas}} c^2}{E_{\text{bind,eff}}} \right]^{\eta}, \quad (16)$$

$$E_{\text{bind,eff}} = \frac{GM_{b,0}^2}{6a_{b,0}} + |W_{b,\text{NFW}}| + \frac{GM_{\text{BH},0} M_{b,0}}{a_{b,0}}. \quad (17)$$

Here $W_{b,\text{NFW}}$ is the initial baryon-halo interaction energy, evaluated by radial quadrature in the NFW potential. Unless stated otherwise, the fiducial calculation uses $\epsilon_f = 10^{-3}$, $\chi_{\text{heat}} = 0.5$, $\eta = 0.5$, primordial metallicity, and the unscaled tabulated cooling rate. The feedback energy changes the baryonic source term, but is not directly injected into the SIDM energy equation. The thermal reservoir is evolved implicitly because the Cloudy cooling time is state dependent. We use the tabulated optically thin equilibrium coefficient in the range $10 \leq T/\text{K} \leq 10^9$ and $-10 \leq \log_{10}(n_H/\text{cm}^{-3}) \leq 4$; primordial and solar-pattern metal contributions are linearly combined in $Z/Z_{\odot}$ [Ferland et al., 2017, Smith et al., 2017]. Inputs outside either tabulated axis are clipped to the nearest table boundary rather than extrapolated. The initial $n_H \simeq 9.2 \times 10^3 \text{ cm}^{-3}$ is close to the upper density boundary, so this prescription and the omission of molecular cooling are explicit limitations. The 10 km s^{-1} initial sound speed is a floor representing the unheated hydrostatic state; cooling removes feedback heat, not this baseline support.

2.4 Velocity-dependent transport

For the final applicability map we replace the constant cross section by the Born/Rutherford angular law

$$\frac{d\sigma}{d\Omega} = \frac{\sigma_0}{4\pi} \left[1 + \left(\frac{v_{\text{rel}}}{w} \right)^2 \sin^2 \frac{\theta}{2} \right]^{-2}. \quad (18)$$

Here θ is the center-of-mass scattering angle and σ_0 is the total cross section in the $v_{\text{rel}} \rightarrow 0$ limit, since the angular integral of equation (18) then equals σ_0 . The viscosity-weighted thermal averages are computed independently for the long- and short-mean-free-path terms,

$$K_p = \frac{\langle \sigma_{\text{visc}} v_{\text{rel}}^p \rangle}{\langle \sigma_{\text{visc}} v_{\text{rel}}^p \rangle_{w \rightarrow \infty}}, \quad \kappa_{\text{SMFP}} = \frac{3Bv}{2\sigma_0 K_5}, \quad \kappa_{\text{LMFP}} = \frac{3AC\rho v^3 \sigma_0 K_3}{2}, \quad (19)$$

followed by the same harmonic interpolation as equation (4). A logarithmic quadrature and a 4097-point lookup table resolve the low-speed tail. The constant-cross-section limit is recovered to 2×10^{-10} in the regression test. This separate K_3/K_5 treatment follows the transport prescription of Outmezguine et al. [2023]; replacing it by a single cross section at the virial velocity would miss the radial evolution of the inner transport.

The fiducial constant-cross-section solution spans both transport limits. Its inner-cell Knudsen number decreases from 2.62×10^{-2} initially to 1.15×10^{-3} at 2 Myr, placing the resolved inner

region in the short-mean-free-path regime, while the radial median changes from 2.19 to 0.846 and the outer halo remains long mean free path. The harmonic interpolation is therefore active across the simulated domain rather than serving only as a formal limiting prescription.

3 Numerical method and validation

The radial mesh is logarithmic, with fixed physical inner and outer boundaries in a matched comparison. The hyperbolic step is a finite-volume update using MC slopes in primitive variables, Roe fluxes [Roe, 1981], and a Harten entropy correction [Harten, 1983]. The reconstruction is monotonicity preserving in each cell and falls back to a first-order state when a characteristic extrapolation would produce a non-positive density or pressure. The conduction term is operator split and advanced with a frozen-coefficient implicit tridiagonal solve. Gravity and accretion sources are integrated after the hydrodynamic flux update. These choices follow standard high-resolution finite-volume practice [van Leer, 1977, Toro, 2009] while retaining the conduction update of the reference calculation.

The absorbing inner boundary supplies the dark-matter channel in the resolved heavy-seed runs. Each accepted result records the mass-budget residual, defined as the difference between initial dark mass, evolved dark mass, and the integrated inner flux in code units. Across the numerical-sensitivity and applicability calculations the largest residual is 1.96×10^{-10} ; this is negligible compared with the boundary differences discussed below.

We use 5% as a pre-declared *operational* tolerance for terminal-mass sensitivity, not as a universal SIDM standard or a proof of asymptotic convergence. For two calculations it is evaluated with the symmetric relative difference

$$\delta(M_1, M_2) = \frac{2|M_1 - M_2|}{|M_1| + |M_2|}, \quad (20)$$

and a terminal mass passes only when both the selected grid and inner-boundary differences have $\delta < 0.05$. The value was fixed before classifying the parameter survey: it lies safely above the 0.05–1.1% baseline grid changes, but below the 10–60% boundary excursions that generate spurious high-mass branches. A different convention, such as 3% or 10%, would change some terminal-mass labels. For this reason we also report the physically coarser threshold classifier—whether both completed boundaries cross $10^7 M_\odot$ —which does not rely on assigning high precision to the endpoint. This classifier is established only over the completed boundary bracket and is not an extrapolation to $r_{\min} \rightarrow 0$.

3.1 SIS/NFW baseline

Figure 1 shows the complete 2 Myr validation. With conduction active, a $100 M_\odot$ seed reaches $129.08 M_\odot$ in the NFW halo and $9646.8 M_\odot$ in the SIS halo, consistent with the reference endpoints [Meng et al., 2026]. Turning conduction off raises the endpoints to $234.36 M_\odot$ and $32528.3 M_\odot$, factors of 1.82 and 3.37, while the final inner densities increase by factors of 25.7 and 104. This is the basic transport effect carried through the rest of the paper: conduction removes compressional heat from the inflowing region, but it also limits the central density available to the black hole.

Grid, CFL, and entropy-fix tests are substantially smaller than the dominant inner-boundary systematic. The NFW 256-to-512-cell change is 0.05%, the SIS change is 1.1%, CFL variations are below 0.012%, and varying the entropy-fix parameter changes the endpoint by less than 0.02%. Changing the outer boundary over factors of four gives relative spans of 1.7×10^{-6} (NFW) and 1.4×10^{-4} (SIS). In contrast, a factor-of-two inner-boundary shift changes the NFW endpoint by about 4% and is larger in the more centrally concentrated SIS case. We therefore retain at least two physical inner boundaries whenever a baryonic enhancement or threshold is quoted.




Figure 1: Baseline mass and accretion histories for NFW and SIS initial profiles. The heat-on curves reproduce the reference 2 Myr endpoints, while heat-off curves show the large increase obtained when conductive transport is removed.

4 Assembled baryonic potentials

4.1 Mechanism and assembly clock

The static Hernquist comparison is not sufficient: a pre-existing compact potential changes the initial hydrostatic support, whereas an assembled potential drives a transient response. In the assembled experiments the dark-matter supply rises during baryon growth, reaches a broad maximum, and then declines. Figure 2 compares the growth enhancement with the assembly time and with a heat-off control. For $f_b = 0.05$, $a_b/r_s = 0.01$, and $\sigma/m = 10\text{--}100\text{ cm}^2\text{ g}^{-1}$, the refined optima are $T_{\text{asm}} = 0.50\text{--}0.75\text{ Myr}$. The ratio $T_{\text{opt}}/t_{\text{dyn}}$ is $0.985\text{--}1.163$ with an RMS scatter of 0.046 dex , whereas $T_{\text{opt}}/t_{\text{coll}}$ has a scatter of 0.545 dex . The particle collision time is therefore not the assembly clock. A dynamical-time matching window is the more stable interpretation, in agreement with the idea that the potential must grow coherently with the region that supplies the black hole.

At the supply radius the v^2 gradient is negative in all refined conducting optima, so $q_r = -\kappa\partial_r v^2$ points outward. The conductive term removes compressional heat from the inflowing region; it does not heat the center from outside. When conduction is disabled the same finite-time optimum remains, but its enhancement is only 2.94 rather than $26.7\text{--}102.4$. The assembly window is therefore dynamical, while the large amplification is a transport effect.

4.2 Compactness is non-monotonic

The fiducial physical-cooling model adopts $M_{200} = 10^9 M_\odot$, $z = 30$, $c = 8$, $f_b = 0.16$, $\sigma/m = 1\text{ cm}^2\text{ g}^{-1}$, and $T_{\text{asm}} = 1.25\text{ Myr}$. The baryon fraction is an upper-reservoir assumption, and

Table 1: Compactness scan around the fiducial physical-cooling model.

a_b/r_s	$M_{\text{BH}}(2 \text{ Myr})/M_\odot$	comment
0.003	8.20×10^6	resolved representative
0.005	1.12×10^7	threshold crossing
0.0075	1.36×10^7	threshold crossing
0.010	1.51×10^7	threshold crossing
0.015	1.67×10^7	near peak
0.020	1.685×10^7	nominal peak
0.030	1.47×10^7	post-peak

$c = 8$ represents a deliberately high-concentration halo rather than a claim about the typical halo at $z = 30$; the applicability calculations below span $c = 4$ –12. The assembly-time interval 0.25–1.75 Myr was selected to bracket the local dynamical times of the resolved supply region, not as a cosmological distribution of assembly histories. We rescan the baryonic scale radius at fixed physical boundaries. The endpoint rises from $8.20 \times 10^6 M_\odot$ at $a_b/r_s = 0.003$ to $1.685 \times 10^7 M_\odot$ at $a_b/r_s = 0.020$, then falls to $1.473 \times 10^7 M_\odot$ at $a_b/r_s = 0.030$ (Table 1). The most compact potential is not the best resolved case: its influence is deep but confined to too small a volume, and the smallest radii approach the inner boundary. This is a physical scale-coupling result, not a claim that compactness is unimportant.

5 Evolving gas, cooling, and feedback

The gas problem contains a feedback loop: a larger black hole raises $\dot{M}_{\text{Edd}}$, retained gas deepens the central potential, and feedback can expand that potential while modifying ρ_∞ and c_s . A no-feedback compact-halo baseline exhibits two-way catalysis, but remains dark dominated: in the earlier $10^6 M_\odot$ test, full-Eddington runs from 10–1000 $M_\odot$ seeds have dark fractions of 96.9–99.0%. This does not mean that baryons are dynamically irrelevant; it means that their role is to change the central environment rather than to provide most of the final mass in the short experiment.

The physically cooled calculation is more restrictive than a stored-heat model. Under optically thin Cloudy cooling, pure heating at $\epsilon_f = 10^{-3}$ retains only 0.033–0.159% of the injected thermal energy and does not prevent the Bondi-to-Eddington transition. Suppressing the cooling coefficient by 10^5 – 3×10^5 is required for heating alone to keep the gas Bondi limited. This is a trapping requirement, not a small metallicity correction. By contrast, under the adopted irreversible mixed-feedback closure, assigning half of the feedback energy to expansion increases a_b by about 1.9 and decreases ρ_∞ by a factor of 6.8–7.0. The gas then remains Bondi limited throughout 2 Myr and the dark channel is reduced by 32.8–33.1% relative to the no-feedback control. These percentages quantify sensitivity within this closure; they are not a general prediction for resolved baryonic feedback.

The feedback threshold is therefore controlled by channel in the present model. Heating is erased by cooling because t_{cool} is short, whereas homologous expansion is cumulative by construction. Expansion can prevent the gas transition while suppressing the dark channel; it is not an independent mechanism that boosts both channels. The sign of this response follows from weakening the central potential and diluting the gas, but its magnitude and persistence require a reversible radius equation or resolved gas dynamics. The current prescription is irreversible, one-zone, and does not include an outflow or recontraction.

Table 2: Resolution and inner-boundary sensitivity of the fiducial $a_b/r_s = 0.02$ model.

variant	$M_{\text{BH}}(2 \text{ Myr})/M_\odot$	$t(M_{\text{BH}} = 10^7 M_\odot) [\text{Myr}]$
256 cells	1.6850×10^7	1.53
512 cells	1.6900×10^7	1.52
$r_{\text{min}}/r_{\text{infl}} = 0.104$	1.5599×10^7	1.65
$r_{\text{min}}/r_{\text{infl}} = 0.052$	1.4523×10^7	1.76

6 Heavy-seed growth and numerical sensitivity

The best nominal model has $M_{\text{BH},0} = 10^5 M_\odot$. At $a_b/r_s = 0.02$ it reaches $1.6850 \times 10^7 M_\odot$ at 2 Myr, with $1.6551 \times 10^7 M_\odot$ in dark accretion and $1.9883 \times 10^5 M_\odot$ in retained baryons. Thus 98.81% of the total growth is dark. The Hernquist radius expands from 2.507 pc to 2.615 pc, the one-zone gas density falls from 0.300 to 0.264 $M_\odot \text{ pc}^{-3}$, and the sound speed rises from 10.0 to 11.99 km s^{-1} .

The endpoint is not fully converged under the strict 5% terminal-mass criterion. Doubling the grid from 256 to 512 cells changes the endpoint by only 0.298%, but reducing the physical inner boundary gives $1.5599 \times 10^7 M_\odot$ and $1.4523 \times 10^7 M_\odot$, a 7.14% difference (Figure 4). Every completed boundary variant crosses $10^7 M_\odot$ at 1.52–1.76 Myr, and the smallest endpoint remains 45.2% above the threshold. We therefore retain threshold crossing as a boundary-bracketed result, not as evidence of convergence under further unresolved boundary reduction, and label $1.685 \times 10^7 M_\odot$ a nominal peak.

The numerically unresolved high-mass branch is also informative. Nominal $10^8 M_\odot$ -like endpoints occur at high concentration, high cross section, and extremely compact baryons, but their two smallest inner boundaries differ by 56–64%. The apparent enhancement is an unresolved capture-boundary effect. Grid, CFL, and entropy-fix changes are small in these cases; the failure is physical closure at the absorbing boundary, not a hydrodynamic instability.

7 Light seeds: delivery versus capture

Applying the heavy-seed absorbing flux directly to a $10^2 M_\odot$ seed would make the physical inner boundary shrink by three orders of magnitude. Instead we introduce a feeding radius and separate the resolved inward delivery from capture by the unresolved black hole,

$$\dot{M}_{\text{supply}} = \max[-4\pi r_{\text{feed}}^2 \rho(r_{\text{feed}}) u(r_{\text{feed}}), 0], \quad (21)$$

$$\dot{M}_{\text{cap}} = 4\pi \lambda_B \frac{G^2 M_{\text{BH}}^2 \rho(r_{\text{feed}})}{c_s^3(r_{\text{feed}})}. \quad (22)$$

Supplied but uncaptured dark matter enters a conservative central reservoir and contributes to the gravitational field. Bondi capture operates while $r_{\text{feed}}/r_{\text{infl}} > 0.104$; once the feeding radius is inside the influence region, the model switches to direct capture and drains the reservoir on a local free-fall time. The closure reproduces three directly resolved heavy-seed runs to 3.97×10^{-6} in final mass and changes by only 2.42×10^{-6} when the gate is moved across the tested bracket.

The fixed-halo seed ladder is summarized in Table 3. A $10^2 M_\odot$ seed supplies $1.42 \times 10^4 M_\odot$ of dark matter to the feeding radius but captures only $3.25 M_\odot$; a $10^3 M_\odot$ seed captures $103 M_\odot$ from $5.28 \times 10^4 M_\odot$ supplied; and a $10^4 M_\odot$ seed captures $7.17 \times 10^3 M_\odot$ from $6.74 \times 10^6 M_\odot$ supplied. The capture fractions are 0.023%, 0.195%, and 0.106%. Central delivery is therefore not equivalent to black-hole growth.

Refinement over 2×10^4 – $8 \times 10^4 M_\odot$ brackets a nonlinear transition. At nominal $\lambda_B = 0.25$, $4 \times 10^4 M_\odot$ and larger seeds cross $10^7 M_\odot$ at both tested feeding radii, whereas $2 \times 10^4 M_\odot$ fails. Seeds between 3×10^4 and $3.75 \times 10^4 M_\odot$ are boundary ambiguous. Varying λ_B from 0.20 to 0.30

Table 3: Fixed-halo seed ladder under the fiducial physical-cooling model.

$M_{\text{seed}}/M_{\odot}$	$M_{\text{BH}}(2\text{ Myr})/M_{\odot}$	growth	boundary diff.	$10^7 M_{\odot}$?
10^2	103.3	1.033	1.24%	no
10^3	1.10×10^3	1.103	1.97%	no
10^4	$1.33\text{--}1.72 \times 10^4$	1.33–1.72	14.4%	no
10^5	1.452×10^7 (conservative)	145.2	7.14%	yes

Table 4: Controlled velocity-dependent applicability map.

model	initial survey	crossings	failures	ambiguous / successes
constant $\sigma/m = 1$	92	14	8	2 / 6
Rutherford $(\sigma_0, w) = (30, 10)$	92	0	16	0 / 0
Rutherford (30, 30)	92	2	15	1 / 0
Rutherford (30, 100)	92	16	7	7 / 2

leaves the $10^2\text{--}10^4 M_{\odot}$ failures unchanged at the tested boundaries, but moves the $4 \times 10^4 M_{\odot}$ case from failure to a crossing at the single boundary used in that sensitivity pair.

Figure 6 displays all existing capture-coefficient calculations without interpolation. The $\lambda_B = 0.25$ row distinguishes failures, boundary-ambiguous cases, and crossings at both completed boundaries. The $\lambda_B = 0.20$ and 0.30 rows contain one boundary per seed and therefore cannot establish a continuous success region. The map demonstrates that the transition is neither absent nor uniquely located: it is bracketed jointly by seed mass, capture coefficient, and feeding boundary. The robust result is the failure of $10^2\text{--}10^4 M_{\odot}$ seeds throughout the tested $\lambda_B = 0.20\text{--}0.30$ interval; rapid growth from intermediate seeds is conditional on the unresolved capture closure.

8 Velocity-dependent cross sections and applicability

The constant cross section is not the only physically motivated transport law. We first calibrate the Rutherford closure at the fiducial heavy-seed parameter set. For $\sigma_0/m = 30\text{ cm}^2\text{ g}^{-1}$, the smaller-boundary endpoints are 1.82×10^6 , 4.90×10^6 , and $1.065 \times 10^7 M_{\odot}$ for $w = 10$, 30 , and 100 km s^{-1} , respectively. The constant control ends at $1.56 \times 10^7 M_{\odot}$. The $w = 30\text{ km s}^{-1}$ model has a virial-scale $\sigma_0 K_3/m \simeq 0.70\text{ cm}^2\text{ g}^{-1}$, but its initial inner value is only $0.0248\text{ cm}^2\text{ g}^{-1}$ and it fails the threshold. Matching the cross section at the virial velocity is therefore not a complete closure; the radial and temporal K_3/K_5 history matters.

We then vary six parameters around the fiducial parameter set: $M_{200} = 10^8\text{--}10^{10} M_{\odot}$, $z = 15\text{--}30$, $c = 4\text{--}12$, $M_{\text{seed}} = 10^2\text{--}10^5 M_{\odot}$, $a_b/r_s = 0.005\text{--}0.08$, and $T_{\text{asm}} = 0.25\text{--}1.75\text{ Myr}$. Each microphysics model contains 28 one-axis cases and 64 fixed-seed Latin-hypercube cases, followed by boundary and double-grid refinement of the 16 points closest to the threshold. The initial survey contains 92 points per model; the refined numerical audit contains 16 points per model. We classify a case as a robust success within the completed numerical bracket only when the threshold crossing survives the smaller-boundary test and, where selected, the double-grid test. A boundary-ambiguous case crosses in one but not both completed boundaries. The Latin-hypercube measure is a numerical design, not an astrophysical prior, so sampled fractions are not estimates of cosmic occurrence rates.

The results are shown in Figure 8. The low-transport $w = 10\text{ km s}^{-1}$ model has no threshold crossing anywhere in the initial survey. The matched $w = 30\text{ km s}^{-1}$ model has two initial-survey crossings but no boundary-robust success in its audited neighborhood. The high-transport $w = 100\text{ km s}^{-1}$ model retains two robust points, both in the fiducial halo with $10^5 M_{\odot}$ seeds and assembly times of 0.5 or 1.5 Myr . The constant control retains six robust audited points,

concentrated at high redshift and intermediate-to-heavy seeds. Across the full refinement no point passes the strict 5% terminal-mass criterion in the combined boundary-plus-grid sense; one low-transport failure happens to pass that numerical metric, which does not alter its threshold class.

Seed mass and redshift show the clearest positive trends in the one-axis and descriptive Latin-hypercube projections, while assembly time and compactness select the width of the crossing window. These directions are exploratory: the high-transport refinement contains only two boundary-robust successes, both at $M_{\text{seed}} = 10^5 M_{\odot}$ in the fiducial halo with $T_{\text{asm}} = 0.5$ and 1.5 Myr. We therefore do not use a sparse logistic fit to assign coefficient significance, success-volume fractions, or population probabilities.

9 Discussion

9.1 What is robust

The first robust conclusion is numerical. The MC–Roe solver reproduces the reference SIS/NFW histories, and grid, CFL, entropy-fix, and outer-boundary changes are small in comparison with the physical effects. The remaining dominant numerical systematic is the physical inner boundary. This distinction matters because the most dramatic nominal endpoints occur precisely where the compact baryonic scale approaches that boundary.

The second robust conclusion is mechanistic. A growing baryonic potential can enhance SIDM accretion by compressing the supply region, but the response is finite-time and scale dependent. The preferred assembly time tracks the local dynamical time, while the magnitude of the enhancement is set by conductive heat transport. The maximum does not occur at the smallest baryonic scale radius.

The third conclusion separates a cooling result from a feedback-model sensitivity. Under the optically thin table, physical cooling removes stored thermal feedback, so pure heating does not prevent a Bondi-to-Eddington transition unless cooling is suppressed by roughly five orders of magnitude. Under the adopted irreversible closure, mixed expansion can stop the transition by diluting the gas and simultaneously weakens the dark channel. The sign of this competition is physically interpretable, but the quoted 33% magnitude is closure dependent.

The fourth robust conclusion is negative and seed-mass specific. In the fiducial $10^9 M_{\odot}$, $z = 30$ halo, 10^2 – $10^3 M_{\odot}$ seeds remain close to their initial masses and $10^4 M_{\odot}$ seeds remain below $2 \times 10^4 M_{\odot}$ throughout the tested capture-coefficient interval. A $10^5 M_{\odot}$ seed crosses $10^7 M_{\odot}$ across the completed boundary bracket. Between these limits the transition is capture-model and boundary dependent; it is bracketed rather than located as a universal critical mass.

The fifth robust conclusion is restricted to the tested microphysics. A velocity-dependent cross section can remove every threshold crossing in the sampled domain even when its nominal low-velocity normalization is large. Only the high-transport $w = 100 \text{ km s}^{-1}$ model retains boundary-robust examples. This dependence is controlled by the evolving inner transport averages, not by a single virial velocity.

9.2 What is not yet a claim

The calculation is spherically symmetric, has no angular momentum, and uses a one-zone gas environment. The expansion branch is irreversible and does not include recontraction, multiphase structure, optical-depth effects, or resolved outflows. The Cloudy table is optically thin and near its high-density limit for the initial ambient state. The velocity-dependent conductivity is an effective LMFP/SMFP interpolation; anisotropic scattering and merger-driven departures from spherical hydrostatic evolution require independent tests [Robertson et al., 2017, Silverman et al., 2026]. Finally, no seed abundance, LRD number density, luminosity function, or spectral

observable is inferred here. Those questions require a population model and are outside the controlled local closure presented in this paper.

10 Conclusions

We have constructed a matched, time-dependent SIDM–baryon accretion model and followed it from baseline validation to a controlled velocity-dependent applicability map. The main conclusions are:

1. The MC–Roe implementation reproduces the reference SIS/NFW 2 Myr endpoints. Conduction suppresses the final mass by factors of 1.82 (NFW) and 3.37 (SIS) relative to the heat-off controls; grid, CFL, entropy-fix, and outer-boundary variations are sub-percent.
2. An assembled Hernquist potential creates a broad assembly-time window tied to the local dynamical time. Conductive outward heat transport controls the amplitude, reaching factors of 26.7–102.4 in the refined compact-potential tests.
3. In the fiducial physical-cooling heavy-seed model, a nominal $1.685 \times 10^7 M_\odot$ endpoint is not fully boundary converged, but crossing $10^7 M_\odot$ at 1.52–1.76 Myr survives all completed inner-boundary variants. The nominal growth is 98.81% dark-matter supplied. No extrapolation beyond the completed boundary bracket is claimed.
4. Optically thin cooling erases pure thermal feedback. Under the adopted irreversible expansion closure, mixed feedback can keep the gas Bondi limited and reduce dark accretion by about one third; the magnitude and persistence of this suppression are model sensitivities, not universal predictions.
5. A conservative feeding-radius closure separates delivery from capture. Seeds of 10^2 – $10^4 M_\odot$ do not reach the rapid-growth branch in 2 Myr across $\lambda_B = 0.20$ – 0.30 . The intermediate-seed transition is bracketed by capture coefficient and feeding boundary rather than identified as a universal critical mass.
6. Velocity dependence narrows the domain further. The low-transport Rutherford model has no threshold crossing in the tested six-dimensional domain, whereas the high-transport model retains two boundary-robust heavy-seed examples. This is a controlled numerical applicability result, not a population-level success fraction.

The next necessary physics closure is a reversible baryon-radius equation with optical-depth and recontraction tests. A longer integration is useful only after that closure is established; simply extending the present irreversible one-zone model would extrapolate the dominant unresolved systematics rather than add a new robust prediction.

Acknowledgements

This work was funded by the National Natural Science Foundation of China (grant No. 11303079) and Zhejiang GuangSha Vocational and Technological University of China (grant No. 2022KYQD-KDB). The author thanks the maintainers of the Cloudy and Grackle cooling data used in the optically thin closure and acknowledges computational support from the Paracloud ZC-M6 cluster.

Data and code availability


The numerical source, configuration files, machine-readable simulation summaries, convergence logs, and figure-generation scripts are archived at <https://github.com/billkang-x/bh-sidm>. The archived products include the SIS/NFW validation histories, baryonic-assembly scans, cooling and feedback calculations, seed-mass refinements, and the full six-parameter initial survey and numerical follow-up in CSV and JSON formats. Sampled success fractions are not interpreted as population probabilities.

References


- Shmuel Balberg, Stuart L. Shapiro, and Shogo Inagaki. Self-interacting dark matter halos and the gravothermal catastrophe. *The Astrophysical Journal*, 568:475–487, 2002. doi: 10.1086/339038.
- Mitchell C. Begelman, Marta Volonteri, and Martin J. Rees. Formation of supermassive black holes by direct collapse in pre-galactic haloes. *Monthly Notices of the Royal Astronomical Society*, 370:289–298, 2006. doi: 10.1111/j.1365-2966.2006.10467.x.
- Hermann Bondi. On spherically symmetrical accretion. *Monthly Notices of the Royal Astronomical Society*, 112:195–204, 1952. doi: 10.1093/mnras/112.2.195.
- Volker Bromm and Abraham Loeb. Formation of the first supermassive black holes. *The Astrophysical Journal*, 596:34–46, 2003. doi: 10.1086/377529.
- Greg L. Bryan and Michael L. Norman. Statistical properties of x-ray clusters: Analytic and numerical comparisons. *The Astrophysical Journal*, 495:80–99, 1998. doi: 10.1086/305262.
- Romeel Dave, David N. Spergel, Paul J. Steinhardt, and Benjamin D. Wandelt. Halo properties in cosmological simulations of self-interacting cold dark matter. *The Astrophysical Journal*, 547:574–589, 2001. doi: 10.1086/318417.
- Wei-Xiang Feng, Hai-Bo Yu, and Yi-Ming Zhong. Seeding supermassive black holes with self-interacting dark matter: A unified scenario with baryons. *The Astrophysical Journal Letters*, 914(2):L26, 2021. doi: 10.3847/2041-8213/ac04b0.
- Wei-Xiang Feng, Hai-Bo Yu, and Yi-Ming Zhong. Dark Bondi accretion aided by baryons and the origin of JWST little red dots. *arXiv e-prints*, 2025. URL <https://arxiv.org/abs/2506.17641>.
- G. J. Ferland, M. Chatzikos, F. Guzman, M. L. Lykins, P. A. M. van Hoof, R. J. R. Williams, N. P. Abel, N. R. Badnell, F. P. Keenan, R. L. Porter, and P. C. Stancil. The 2017 release of Cloudy. *Revista Mexicana de Astronomia y Astrofisica*, 53:385–438, 2017.
- Martin G. Haehnelt and Martin J. Rees. The formation of nuclei in newly formed galaxies and the evolution of the quasar population. *Monthly Notices of the Royal Astronomical Society*, 263:168–178, 1993. doi: 10.1093/mnras/263.1.168.
- Ami Harten. High resolution schemes for hyperbolic conservation laws. *Journal of Computational Physics*, 49:357–393, 1983. doi: 10.1016/0021-9991(83)90136-5.
- Lars Hernquist. An analytical model for spherical galaxies and bulges. *The Astrophysical Journal*, 356:359–364, 1990. doi: 10.1086/168845.
- Kohei Inayoshi, Eli Visbal, and Zoltan Haiman. The assembly of the first massive black holes. *Annual Review of Astronomy and Astrophysics*, 58:27–97, 2020. doi: 10.1146/annurev-astro-120419-014455.

- Fangzhou Jiang, Zixiang Jia, Haonan Zheng, Luis C. Ho, Kohei Inayoshi, Xuejian Shen, Mark Vogelsberger, and Wei-Xiang Feng. Formation of the little red dots from the core-collapse of self-interacting dark matter halos. *The Astrophysical Journal Letters*, 996(1):L19, 2026. doi: 10.3847/2041-8213/ae247a.
- Manoj Kaplinghat, Sean Tulin, and Hai-Bo Yu. Dark matter halos as particle colliders: Unified solution to small-scale structure puzzles from dwarfs to clusters. *Physical Review Letters*, 116: 041302, 2016. doi: 10.1103/PhysRevLett.116.041302.
- Jun Koda and Paul R. Shapiro. Gravothermal collapse of isolated self-interacting dark matter haloes: N-body simulation versus the fluid model. *Monthly Notices of the Royal Astronomical Society*, 415:1125–1137, 2011. doi: 10.1111/j.1365-2966.2011.18684.x.
- Donald Lynden-Bell and Roger Wood. The gravo-thermal catastrophe in isothermal spheres and the onset of red-giant structure for stellar systems. *Monthly Notices of the Royal Astronomical Society*, 138:495–525, 1968. doi: 10.1093/mnras/138.4.495.
- Piero Madau and Martin J. Rees. Massive black holes as population III remnants. *The Astrophysical Journal Letters*, 551:L27–L30, 2001. doi: 10.1086/319848.
- Zhe Meng, Tan Chen, Bocheng Zhu, Fan Zhou, Bin Hu, Liang Gao, and Rong-Gen Cai. Spherically symmetric fluid simulations of black hole accretion in self-interacting dark matter halos. *arXiv e-prints*, 2026. URL <https://arxiv.org/abs/2607.02151>.
- Julio F. Navarro, Carlos S. Frenk, and Simon D. M. White. A universal density profile from hierarchical clustering. *The Astrophysical Journal*, 490:493–508, 1997. doi: 10.1086/304888.
- Nadav Joseph Outmezguine, Kimberly K. Boddy, Sophia Gad-Nasr, Manoj Kaplinghat, and Laura Sagunski. Universal gravothermal evolution of isolated self-interacting dark matter halos for velocity-dependent cross sections. *Monthly Notices of the Royal Astronomical Society*, 523:4786–4800, 2023. doi: 10.1093/mnras/stad1705.
- Annika H. G. Peter, Miguel Rocha, James S. Bullock, and Manoj Kaplinghat. Cosmological simulations with self-interacting dark matter – ii. halo shapes versus observations. *Monthly Notices of the Royal Astronomical Society*, 430:105–120, 2013. doi: 10.1093/mnras/sts535.
- Andrew Robertson, Richard Massey, and Vincent Eke. Cosmic particle colliders: Simulations of self-interacting dark matter with anisotropic scattering. *Monthly Notices of the Royal Astronomical Society*, 467:4719–4730, 2017. doi: 10.1093/mnras/stx463.
- Miguel Rocha, Annika H. G. Peter, James S. Bullock, Manoj Kaplinghat, Shea Garrison-Kimmel, Jose Onorbe, and Leonidas A. Moustakas. Cosmological simulations with self-interacting dark matter – i. constant-density cores and substructure. *Monthly Notices of the Royal Astronomical Society*, 430:81–104, 2013. doi: 10.1093/mnras/sts514.
- Philip L. Roe. Approximate Riemann solvers, parameter vectors, and difference schemes. *Journal of Computational Physics*, 43:357–372, 1981. doi: 10.1016/0021-9991(81)90128-5.
- Edwin E. Salpeter. Accretion of interstellar matter by massive objects. *The Astrophysical Journal*, 140:796–800, 1964. doi: 10.1086/147973.
- N. I. Shakura and R. A. Sunyaev. Black holes in binary systems. observational appearance. *Astronomy and Astrophysics*, 24:337–355, 1973.
- Maya Silverman, Abdelaziz Hussein, Arpit Arora, Mariangela Lisanti, Manoj Kaplinghat, Lina Necib, Andreas Thoyas, Stephanie O’Neil, Robyn E. Sanderson, Xuejian Shen, and Jorge Moreno. Mergers matter: Gravothermal collapse in dwarf halos with self-interacting dark matter. *arXiv e-prints*, 2026. URL <https://arxiv.org/abs/2606.02566>.

- Britton D. Smith, Greg L. Bryan, Simon C. O. Glover, Nathan J. Goldbaum, Matthew J. Turk, John Regan, John H. Wise, Hsi-Yu Schive, Tom Abel, Andrew Emerick, Brian W. O'Shea, Peter Anninos, Cameron B. Hummels, and Sadegh Khochfar. GRACKLE: A chemistry and cooling library for astrophysics. *Monthly Notices of the Royal Astronomical Society*, 466: 2217–2234, 2017. doi: 10.1093/mnras/stw3291.
- David N. Spergel and Paul J. Steinhardt. Observational evidence for self-interacting cold dark matter. *Physical Review Letters*, 84:3760–3763, 2000. doi: 10.1103/PhysRevLett.84.3760.
- Eleuterio F. Toro. *Riemann Solvers and Numerical Methods for Fluid Dynamics: A Practical Introduction*. Springer, Berlin, 3 edition, 2009. doi: 10.1007/b79761.
- Sean Tulin and Hai-Bo Yu. Dark matter self-interactions and small scale structure. *Physics Reports*, 730:1–57, 2018. doi: 10.1016/j.physrep.2017.11.004.
- Bram van Leer. Towards the ultimate conservative difference scheme. IV. a new approach to numerical convection. *Journal of Computational Physics*, 23:276–299, 1977. doi: 10.1016/0021-9991(77)90095-X.
- Mark Vogelsberger, Jesus Zavala, and Abraham Loeb. Subhaloes in self-interacting galactic dark matter haloes. *Monthly Notices of the Royal Astronomical Society*, 423:3740–3752, 2012. doi: 10.1111/j.1365-2966.2012.21182.x.
- Marta Volonteri. Formation of supermassive black holes. *The Astronomy and Astrophysics Review*, 18:279–315, 2010. doi: 10.1007/s00159-010-0029-x.
- Yi-Ming Zhong, Daneng Yang, and Hai-Bo Yu. The impact of baryonic potentials on the gravothermal evolution of self-interacting dark matter haloes. *Monthly Notices of the Royal Astronomical Society*, 526:758–770, 2023. doi: 10.1093/mnras/stad2765.



(a) Compact-potential growth and transport enhancement.



(b) Refined assembly-time and initial-timescale comparison.

Figure 2: The assembled baryonic potential creates a broad dynamical-time window rather than a narrow resonance. Conductive transport sets the amplitude of the enhancement, and the heat-off curve is an adiabatic diagnostic rather than a collisionless model.






Figure 3: Physical cooling and feedback response. The top panels show that metallicity has a secondary effect on the final limiter state and sound speed. The lower panels show the large cooling suppression required by pure heating and the simultaneous dilution of the dark channel in the mixed expansion model.




(a) Resolved compactness peak.




(b) Grid and inner-boundary convergence.

Figure 4: The preferred heavy-seed model has an internal compactness maximum, but the inner boundary remains the leading terminal-mass systematic. The threshold crossing is more stable than the terminal mass.



(a) Conservative seed ladder and feeding-boundary closure.



(b) Boundary-bracketed intermediate-seed transition.

Figure 5: Light-seed closure. Stellar-remnant seeds remain capture limited even though the SIDM fluid delivers substantial mass to the feeding radius. The intermediate-seed transition is bracketed rather than uniquely located because it depends on the capture coefficient and feeding boundary.




Figure 6: Sparse λ_B – M_{seed} capture-closure map from existing calculations. Left: discrete threshold classes; triangles denote crossings tested at only one feeding boundary and must not be interpreted as boundary-robust successes. Right: final masses for the two capture-coefficient sensitivity sequences. No continuous phase boundary is inferred from these sparse points.




Figure 7: Velocity-dependent transport calibration. The top panels show the conservative final masses and boundary sensitivity as functions of w and σ_0/m . The lower panels demonstrate that virial-scale matching does not determine the final mass and that the local effective cross section evolves strongly during growth.




Figure 8: Main effects and joint response in the six-parameter applicability map. The map is a controlled numerical domain, not a cosmological abundance calculation.




Figure 9: Boundary-refined classifications near the $10^7 M_{\odot}$ threshold. Threshold crossing is the stable classifier; terminal masses remain provisional when the absorbing inner boundary is unresolved.