

Long-Term Resistance Trained Human Muscles Have More Fibers, More Myofibrils, and Tighter Myofilament Packing Than Untrained

SUMIAKI MAEO^{1,2}, THOMAS G. BALSHAW^{2,3}, BENJAMIN MÄRZ^{4,5}, ZHAOXIA ZHOU⁴, BILL HAUG², NEIL R. W. MARTIN², NICOLA MAFFULLI^{6,7,8}, and JONATHAN P. FOLLAND^{2,3,9}

¹Faculty of Sport and Health Science, Ritsumeikan University, Kusatsu, JAPAN; ²School of Sport, Exercise & Health Sciences, Loughborough University, Loughborough, UNITED KINGDOM; ³Versus Arthritis Centre for Sport, Exercise and Osteoarthritis Research, Loughborough University, Loughborough, UNITED KINGDOM; ⁴Loughborough Materials Characterization Centre, Department of Materials, Loughborough University, Loughborough, UNITED KINGDOM; ⁵Shared Instrumentation Facility, Louisiana State University, Baton Rouge, LA; ⁶Department of Trauma and Orthopaedic Surgery, School Medicine, Surgery and Dentistry, University of Salerno, Salerno, ITALY; ⁷School of Pharmacy and Bioengineering, Keele University School of Medicine, Stoke on Trent, UNITED KINGDOM; ⁸Barts and the London School of Medicine and Dentistry, Queen Mary University of London, London, UNITED KINGDOM; and ⁹National Institute for Health and Care Research (NIHR) Leicester Biomedical Research Centre, UNITED KINGDOM

ABSTRACT

MAEO, S., T. G. BALSHAW, B. MÄRZ, Z. ZHOU, B. HAUG, N. R. W. MARTIN, N. MAFFULLI, and J. P. FOLLAND. Long-Term Resistance Trained Human Muscles Have More Fibers, More Myofibrils, and Tighter Myofilament Packing Than Untrained. *Med. Sci. Sports Exerc.*, Vol. 56, No. 10, pp. 1906–1915, 2024. **Introduction:** Increases in skeletal muscle size occur in response to prolonged exposure to resistance training that is typically ascribed to increased muscle fiber size. Whether muscle fiber number also changes remains controversial, and a paucity of data exists about myofibrillar structure. This cross-sectional study compared muscle fiber and myofibril characteristics in long-term resistance-trained (LRT) versus untrained (UNT) individuals. **Methods:** The maximal anatomical cross-sectional area (ACSA_{max}) of the biceps brachii muscle was measured by magnetic resonance imaging in 16 LRT (5.9 ± 3.5 yr' experience) and 13 UNT males. A muscle biopsy was taken from the biceps brachii to measure muscle fiber area, myofibril area, and myosin spacing. Muscle fiber number, and myofibril number in total and per fiber were estimated by dividing ACSA_{max} by muscle fiber area or myofibril area, and muscle fiber area by myofibril area, respectively. **Results:** Compared with UNT, LRT individuals had greater ACSA_{max} (+70%, $P < 0.001$), fiber area (+29%, $P = 0.028$), fiber number (+34%, $P = 0.013$), and myofibril number per fiber (+49%, $P = 0.034$) and in total (+105%, $P < 0.001$). LRT individuals also had smaller myosin spacing (−7%, $P = 0.004$; i.e., greater packing density) and a tendency toward smaller myofibril area (−16%, $P = 0.074$). ACSA_{max} was positively correlated with fiber area ($r = 0.526$), fiber number ($r = 0.445$), and myofibril number (in total $r = 0.873$ and per fiber $r = 0.566$), and negatively correlated with myofibril area ($r = -0.456$) and myosin spacing ($r = -0.382$) (all $P < 0.05$). **Conclusions:** The larger muscles of LRT individuals exhibited more fibers in cross-section and larger muscle fibers, which contained substantially more total myofibrils and more packed myofilaments than UNT participants, suggesting plasticity of muscle ultrastructure. **Key Words:** MUSCLE HYPERTROPHY, HYPERPLASIA, PROLIFERATION

Address for correspondence: Sumiaki Maeo, Ph.D., Faculty of Sport and Health Science, Ritsumeikan University, 1-1-1 Nojihigashi, Kusatsu-shi, Shiga 525-8577, Japan; E-mail: s-maeo@fc.ritsumei.ac.jp.

Submitted for publication January 2024.

Accepted for publication May 2024.

0195-9131/24/5610-1906/0

MEDICINE & SCIENCE IN SPORTS & EXERCISE®

Copyright © 2024 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American College of Sports Medicine. This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

DOI: 10.1249/MSS.0000000000003495

Skeletal muscle size/mass is a key determinant of muscle function (i.e., strength and power) and sport performance (1–4), as well as health and quality of life (5,6). The size of a whole muscle is determined primarily by the number and size of individual muscle fibers (7), which are composed of thousands of myofibrils, with each myofibril in turn containing several hundred myofilaments (Fig. 1). Increases in skeletal muscle size, known as muscle hypertrophy, occur in response to functional overload, that is, resistance training (RT), especially after prolonged systematic exposure (8). Indeed, numerous studies spanning five decades indicate that long-term resistance-trained (LRT) individuals have substantially larger muscles (+40%–80%) than untrained (UNT) individuals (9–14), as well as larger muscle fibers (+20%–60%) (12–14). However, it




FIGURE 1—Illustration of skeletal muscle structure and obtained example images of ACSAmax of the biceps brachii (A), muscle fiber (B), myofibril (C), and myofilament (D). The arrows in D show how the myosin spacing (distance) was measured. The illustration of skeletal muscle structure was purchased from Dreamstime.com (file ID: 80735107 created by Legger) and adapted with permission.

remains controversial whether larger muscle size of LRT is accompanied by/associated with a greater number of muscle fibers (15,16). Furthermore, although structural adaptations to the size and number of myofibrils may contribute to increased muscle fiber size, and thus also whole muscle size (17,18), a paucity of data is available on the plasticity of myofibrils induced by RT (8,16). A better understanding of the adaptations to RT is a first step in elucidating how these adaptations are produced.

Generation of new muscle fibers, called hyperplasia, occurs during developmental growth in mammals (19). In human muscles, early work by MacCallum (20) in 1898 suggested that hyperplasia does not occur after postnatal growth. However, later studies have found evidence that the number of fibers may change over the long term with a relatively slow process, such that it increases in response to daily functional demands/overload (21) and decreases with aging/reduced physical activity (22). Nevertheless, whether hyperplasia occurs following RT remains an open question. For example, similar muscle fiber numbers have been found before and after 12 wk of RT in young adults (23) and 6 months of RT in older adults (24) when estimated by dividing muscle anatomical cross-sectional area (ACSA) by fiber CSA. Similarly, MacDougall et al. (12) found no significant difference in muscle fiber numbers between young LRT and UNT males. On the contrary, Sale et al. (14) found a significant correlation between muscle fiber number and muscle size. This inconsistency might arise from the fact that the training interventions of the longitudinal studies (23,24) (12 wk to 6 months) were not of sufficient duration to induce distinct changes in muscle structures and/or due to small sample sizes (groups with $n \leq 8$) in both longitudinal (23) and cross-sectional (12) studies that reported no change/difference in fiber numbers.

There is limited literature on whether RT changes the ultrastructure of skeletal muscle. In humans, MacDougall (25) observed a 16% increase in myofibril CSA after 6 months of RT. However, a subsequent study reported typically no changes in myofibril CSA after short-term RT (18 wk) (26). Furthermore, neither study reported any measure of myofibril number (per fiber or in total). Thus, even the most basic questions, such as whether larger muscles/fibers following RT are accounted for by larger size and/or increased number of myofibrils, have not been resolved (8,16). In addition, preliminary studies with small participant numbers and short-duration RT interventions have reported the distance between myofilaments within a myofibril, indicative of myofilament packing density, to decrease after 10 wk ($n = 3$) (27), but not after 6 wk ($n = 8$) (28), of RT. Reduced myosin spacing could indicate a packing strategy by which more contractile filaments are contained in a given area, which would be expected to increase specific tension and thus strength irrespective of muscle size (29). Based on the limited evidence to date, further research is clearly needed to examine this possibility. Furthermore, individuals who have completed several years of RT might be expected to exhibit pronounced adaptations, where these are physiologically possible, and thus may reveal the potential for human ultrastructural adaptations.

The present study compared muscle fiber and myofibril characteristics in LRT versus UNT individuals. We hypothesized that, compared with UNT, LRT individuals would have 1) larger and a greater number of muscle fibers, 2) larger and a greater number of myofibrils (per fiber and in total), and 3) smaller myofilament spacing (greater packing density). We also hypothesized that the size and number of muscle fibers and myofibrils, as well as myofilament spacing, would be correlated with muscle ACSA.

TABLE 1. Descriptive characteristics of LRT and UNT individuals.

Variable	LRT	UNT	<i>P</i>	Cohen's <i>d</i>
Age (yr)	23.2 ± 3.6	20.5 ± 2.3	0.012*	0.89 [large]
Height (m)	1.82 ± 0.08	1.77 ± 0.07	0.152	0.56 [medium]
Humerus length (cm)	33.9 ± 2.1	33.3 ± 1.4	0.394	0.33 [small]
Body mass (kg)	88.2 ± 12.9	74.6 ± 11.4	0.006*	1.12 [large]
IPAQ value (MET·min·wk ⁻¹)	5150 ± 2655	2689 ± 1452	0.006*	1.15 [large]

Data are group mean ± SD. *n* = 16 for LRT and *n* = 13 for UNT.

*Significant difference between groups.

was estimated by dividing muscle ACSAmax by average fiber area, where average fiber area was calculated as follows (12):

$$[(\% \text{ type I number})(\text{mean type I area}) + (\% \text{ type II number})(\text{mean type II area})]/100$$

Ultrastructure analysis. Fixed tissue was rinsed in phosphate buffer, secondary fixed in osmium tetroxide, dehydration and resin embedded before cross-sectioning using an ultramicrotome to prepare sections of 50 nm thickness for transmission electron microscopy examination. Full protocols by McMillan and Eisenback (34) were followed. Ultrastructural images of the sample were taken using a FEI Tecnai F20 transmission electron microscope at magnifications of ×4000 (for myofibril area, Fig. 1C) and ×19,500 (for myosin spacing, Fig. 1D). Image analysis was undertaken using Fiji software with the investigator blinded to the participant code of each sample, and only transversely sectioned fibers were included in the analysis. Myofibril cross-sectional area was assessed by manually drawing around the perimeter of 200 individual myofibrils per participant throughout the entire sections. The numbers of myofibrils per fiber and per muscle were calculated by dividing muscle fiber area or muscle ACSAmax, respectively, by the average myofibril area. Finally, myosin spacing was assessed for 300 interspace distances per participant, by measuring the average distance across six myosin filaments in a line (i.e., 5 interspace distances, the length of an arrow in Fig. 1D, divided by 5) in three separate directions starting from a randomly chosen point (i.e., 15 distances starting from one point), and two random points for one myofibril (i.e., 30 distances for one myofibril) and for 10 different myofibrils (i.e., 300 distances in total) (28).

Statistical Analysis

All data on the whole cohort of participants were first checked for outliers by the median absolute deviation method (35), using a threshold value of 3 (very conservative (36)). One participant (LRT) for myofibril number per fiber was found to be an outlier, but was nevertheless included in the analysis because inclusion/exclusion of these data resulted in the same interpretation, and the data seemed biologically plausible. All analyses were performed using SPSS (version 25: IBM, Armonk, NY). Significance was set at an alpha level of 0.05. Data were checked for normality using Shapiro–Wilk tests in each group. Age, type I fiber area, myofibril area, and myosin spacing were nonnormally distributed, and thus, we used Mann–Whitney *U* tests to compare groups for these variables. For the other variables, unpaired *t*-tests were used. The effect sizes of between-group differences were calculated as Cohen's *d* values and interpreted as large (≥ 0.80), medium (0.50–0.79), small (0.20–0.49), or trivial (< 0.20). Bivariate relationships were assessed with Pearson's product–moment correlations to assess the relationship of muscle ACSAmax with the structural variables of fibers and myofibrils on the whole cohort data. In addition, the coefficient of determination (R^2) was calculated for these relationships. All data are presented as mean ± SD in the text, and individual data together with box and whisker plots are shown within the figures (detailed in each figure).

RESULTS

Group characteristics. Descriptive characteristics of the groups are shown in Table 1. LRT individual were slightly (~2 yr) older than UNT ($P = 0.012$). There were no significant between-group differences in height or humerus length ($P = 0.152$ – 0.394), but body mass was 18% larger for LRT than UNT ($P = 0.006$). The IPAQ value was 92% greater for LRT than UNT ($P = 0.006$).

Muscle ACSAmax. Biceps brachii muscle ACSAmax was 70% greater for LRT than UNT (18.4 ± 2.7 vs 10.8 ± 2.2 cm², $P < 0.001$, $d = 3.07$ “large”; Fig. 2). The large difference in ACSAmax remained similar (63%) when normalized to




FIGURE 2—ACSAmax of LRT and UNT males. * denotes a significant difference between groups ($P < 0.05$) by a *t*-test. Individual plots together with box (lower–upper quartile) and whisker (SD) plots are shown. The cross mark and line in the box indicate the mean and median, respectively.

humerus length² (1.60 ± 0.22 vs 0.98 ± 0.20 , $P < 0.001$, $d = 2.90$ “large”) to account for any possible influence of body size on ACSAmax. Considering the similarity of between-group differences in absolute and normalized ACSAmax, data are only reported as absolute values (cm²) to avoid redundancy. In addition to ACSAmax, muscle size was also assessed as ACSA at the location the biopsy was taken, but these two measures of muscle size produced identical findings, and therefore, data are only reported as ACSAmax to avoid redundancy.

Muscle fiber type composition, size, and number.

Muscle fiber type composition did not significantly differ between LRT and UNT when expressed by the proportion of either number (type I: $48.3\% \pm 8.1\%$ vs $45.7\% \pm 5.8\%$, type II: $51.7\% \pm 8.1\%$ vs $54.3\% \pm 5.8\%$, $P = 0.348$, $d = 0.36$ “small”; Fig. 3A) or area (type I: $44.7\% \pm 9.0\%$ vs $43.9\% \pm 6.2\%$, type II: $55.3\% \pm 9.0\%$ vs $56.1\% \pm 6.2\%$, $P = 0.796$, $d = 0.10$, “trivial”; Fig. 3B). Fiber area was significantly larger in LRT than UNT for both type I ($+27\%$, 6707 ± 1613 vs $5281 \pm 1389 \mu\text{m}^2$, $P = 0.022$, $d = 0.95$ “large”; Fig. 3C) and type II ($+31\%$, 8422 ± 2760 vs $6419 \pm 2137 \mu\text{m}^2$, $P = 0.041$, $d = 0.81$, “large”; Fig. 3D) fibers. Consequently, the average fiber area (of both fiber types) was significantly larger for LRT than UNT ($+29\%$, 7588 ± 2134 vs $5881 \pm 1726 \mu\text{m}^2$, $P = 0.028$, $d = 0.88$, “large”; Fig. 4A). Fiber number (i.e., in total within the biceps brachii ACSAmx) was also greater for LRT than UNT ($+34\%$, $260 \pm 77 \times 10^3$ vs $194 \pm 50 \times 10^3$, $P = 0.013$, $d = 1.02$, “large”; Fig. 4B). Both the average fiber area ($r = 0.526$, $R^2 = 0.277$, $P = 0.003$) and fiber number ($r = 0.445$,

$R^2 = 0.198, P = 0.015$) were significantly correlated with muscle ACSAmax (Figs. 4C, D).

Myofibril size and number. There was a tendency toward smaller myofibril area (-16% , 0.628 ± 0.204 vs $0.748 \pm 0.192 \mu\text{m}^2$, $P = 0.074$, $d = 0.61$, “medium”; Fig. 5A) for LRT versus UNT. The number of myofibrils per fiber ($+49\%$, $13.1 \pm 5.6 \times 10^3$ vs $8.8 \pm 4.1 \times 10^3$, $P = 0.034$, $d = 0.87$, “large”; Fig. 5B) and the total number of myofibrils within the whole muscle (i.e., per ACSAmax) ($+105\%$, $32.0 \pm 10.7 \times 10^8$ vs $15.6 \pm 5.6 \times 10^8$, $P < 0.001$, $d = 1.91$, “large”; Fig. 5C) were greater for LRT than UNT. Myofibril area ($r = -0.456$, $R^2 = 0.208$, $P = 0.015$), number of myofibrils per fiber ($r = 0.566$, $R^2 = 0.320$, $P = 0.002$), and within the whole muscle ($r = 0.208$, $R^2 = 0.762$, $P < 0.001$) were all significantly correlated with ACSAmax (Figs. 5D–F).

Myosin spacing. LRT had a modest but significantly smaller myosin spacing than UNT (-7% , 39.2 ± 3.3 vs 42.4 ± 2.5 , $P = 0.010$, $d = 1.08$, “large”; Fig. 6A), and this was negatively correlated with ACSAmx ($r = -0.387$, $R^2 = 0.150$, $P = 0.042$; Fig. 6B).

DISCUSSION

The main findings of this study were that LRT individuals who had 70% larger muscles than UNT, also had 1) larger size (+29%) and a greater number (+34%) of muscle fibers within the ACSAmx, 2) a tendency for smaller myofibrils (-16%) but more myofibrils per fiber (+49%) and in total (+105%),




FIGURE 3—Muscle fiber type composition by number (A) and by area (B), and type I fiber area (C) and type II fiber area (D) of LRT and UNT males. * denotes a significant difference between groups ($P < 0.05$). To compare the groups, Mann–Whitney U test (C) and t -tests (A, B, D) were used. Individual plots together with box (lower–upper quartile) and whisker (SD) plots are shown. The cross mark and line in the box indicate the mean and median, respectively.




FIGURE 4—Average fiber area across type I and II fibers (A) and estimated fiber number (B), and relationships of ACSAmax with average fiber area (C) and fiber number (D) of LRT and UNT males. * denotes a significant difference between groups ($P < 0.05$) by a t -test. Individual plots together with box (lower–upper quartile) and whisker (SD) plots are shown. The cross mark and line in the box indicate the mean and median, respectively.

and 3) smaller myosin spacing (-7% , i.e., greater packing density), than UNT. In addition, considering the whole cohort (i.e., both groups), muscle size (ACSAmax) was positively correlated with muscle fiber size ($r = 0.526$) and number ($r = 0.445$), as well as myofibril number (per fiber $r = 0.566$ and in total $r = 0.873$), and negatively correlated with myofibril size ($r = -0.456$) and myosin spacing ($r = -0.382$). These findings

mostly supported our hypotheses and suggest that the larger muscles of LRT individuals were characterized as having muscle fibers of greater size and number in cross section, which likely contain smaller but more myofibrils as well as more tightly packed myofilaments.

Whole muscle and muscle fibers. Muscle ACSAmax was 70% greater for LRT than UNT (Fig. 2). This between-group difference was similar to previous studies (57%–76%




FIGURE 5—Myofibril area (A), myofibril number per fiber (B), and total myofibril number (C), and relationships of ACSAmax with myofibril area (D), myofibril number per fiber (E), and total myofibril number (F) of LRT and UNT males. * denotes a significant difference between groups ($P < 0.05$) by a t -test, and # denotes a tendency toward significance ($P = 0.072$) by Mann–Whitney U test. Individual plots together with box (lower–upper quartile) and whisker (SD) plots are shown. The cross mark and line in the box indicate the mean and median, respectively.




FIGURE 6—Myosin interspace (A) and its relationship with ACSAmax (B) of LRT and UNT males. * denotes a significant difference between groups ($P < 0.05$) by Mann–Whitney U test. Individual plots together with box (lower–upper quartile) and whisker (SD) plots are shown. The cross mark and line in the box indicate the mean and median, respectively.

difference) (12,14) that compared the size of the biceps brachii between LRT (elite bodybuilders) and UNT males. Thus, the LRT participants recruited in this study can be considered as having substantially larger biceps brachii muscles compared with UNT, a widely documented adaptation to prolonged RT.

Muscle fiber type composition was similar between LRT and UNT, with both groups having 44%–48% type I fibers and 52%–56% type II fibers when expressed either by number or area (Figs. 3A, B), which is in line with the previous reports indicating no clear effect of LRT on fiber type composition (12,14). The muscle fiber area of LRT was similarly greater than UNT for both type I and type II fibers (+27% and +31%, Figs. 3C, D). This supports the previous finding of consistent fiber type hypertrophy within the biceps brachii of LRT versus UNT males (14) and suggests that the common observation of preferential type II fiber hypertrophy after short-term RT (37,38) does not persist with prolonged RT. The average fiber area (of both fiber types) was +29% larger for LRT than UNT (Fig. 4A) and was positively correlated with muscle ACSAmax (Fig. 4C). This agrees with widely acknowledged findings that enlarged muscle fibers explain, at least partly, enlarged muscle size after RT (see reviews (16,39) for details).

Furthermore, the between-group difference of muscle fiber size (29%) was found to be much smaller than that of muscle ACSAmax (70%). As a result, the estimated muscle fiber number within the ACSAmax of LRT was +34% greater than UNT (Fig. 4B). Therefore, there was a bigger percentage difference and effect size for fiber number than fiber area between LRT and UNT (+34% vs +29%; $d = 1.02$ vs $d = 0.88$). Fiber number was also positively correlated with muscle ACSAmax in the whole cohort, but with a slightly weaker correlation coefficient than for fiber area ($r = 0.445$ vs $r = 0.526$; Figs. 4C, D). The current findings are in agreement with previous studies that found either a significant correlation ($r = 0.60$) (14) or, although not reported as such by the authors, a tendency (i.e., $r = 0.35$, $n = 25$, $P < 0.10$) (12) between muscle fiber number and muscle size. However, these same studies also reported no differences in muscle fiber number between LRT/elite bodybuilders and UNT individuals (12,14), perhaps because these studies

included three groups and smaller participant numbers per group than the current investigation, potentially reducing their statistical power for finding differences.

There are two possible adaptive explanations for the observation of greater fiber number in the muscle cross section of prolonged RT individuals. First, it could indicate human muscle fiber hyperplasia after prolonged RT (i.e., an increase in the total number of muscle fibers within the whole muscle), in accordance with the adaptations to years of daily functional demands (21). Alternatively, in pennate muscles such as the vastus lateralis, changes in fiber/fascicle length could also increase the apparent number of fibers with a given cross section of the muscle, as well as increasing muscle CSA and volume (40). Although the current study did not assess fascicle length, we deliberately chose to examine the biceps brachii owing to its fusiform architecture, to minimize the potential confounding influence of differences in muscle architecture between groups (12,14). Therefore, because of the limited pennation of the biceps brachii, a genuine increase in fiber number may be the most likely explanation for the greater number of fibers in the muscle cross section of LRT individuals. Finally, given the cross-sectional nature of this study, we cannot rule out the possibility that the LRT participants had an inherently greater number of muscle fibers than their UNT counterparts, perhaps a consequence of a selection bias, and longitudinal studies of sufficient RT duration are needed to resolve this uncertainty.

Myofibrils and myofilaments. Myofibril area showed a tendency to be smaller (–16%) in LRT than UNT individuals (Fig. 5A), and there was a significant negative correlation between myofibril area and muscle ACSAmax (i.e., larger muscle, smaller myofibrils; Fig. 5D). This contradicted our second hypothesis and a previous RT study (25) that found a 16% increase in myofibril size after 6 months of RT. This discrepancy cannot be explained with any certainty, although average myofibrillar size may be the result of the combined effects of both myofibril growth, which increases myofibril size, and proliferation/splitting, which instead reduces myofibril size. After LRT (mean, 6 yr in the current study), myofibrillar proliferation/splitting could be sufficient to counter any myofibrillar growth resulting

in no overall change in average myofibrillar area. As a result of the tendency toward smaller myofibrils in LRT, together with their larger muscle fiber area and muscle ACSAmax, myofibril number was greater for LRT than UNT by +49% per fiber and +108% in total (Figs. 5B, C), and both were correlated with ACSAmax (Figs. 5E, F). A recent study by Jorgenson et al. (41) used high-resolution immunofluorescence microscopy and revealed that a relatively brief 7-wk RT period increased the number of myofibrils in type II fibers without altering myofibril size, which agrees with the current study. Therefore, muscle hypertrophy after RT appears to be associated with a greater amount of contractile/myofibrillar material, mainly because of an increased number of myofibrils, and this may be a primary reason why LRT individuals are capable of producing significantly higher levels of force and power compared with UNT (11,14).

Myosin spacing was smaller in LRT than UNT participants and was negatively correlated with ACSAmax (Figs. 6A, B). Although the between-group difference in myosin spacing was modest (−7%), its effect size was large (Cohen's $d = 1.08$) because of the low interindividual variability of this variable. This suggests that LRT had greater myofilament packing density than UNT, and larger muscles were also associated with more packed myofilaments. As mentioned earlier, two preliminary studies have reported a decrease ($n = 3$) (27) and no change ($n = 8$) (28) in myosin spacing after short-term RT. Similar to the current findings, a recent pilot study (42) found a small group of LRT (10 ± 3 yr of experience, $n = 6$) had ~9% more myosin and actin protein content (on a per mg basis), suggestive of greater myofibrillar packing. Nonetheless, the current study provides the first visual evidence for the greater contractile filament density of prolonged RT and thus offers a mechanistic basis for increased specific tension that would be expected to contribute to strength gains irrespective of hypertrophy (29). It is also notable that small (+2%–6%) but consistent increases in muscle radiological density, suggestive

of increased contractile filament packing, have been found after RT (28,43,44). Furthermore, we have previously found greater *in vivo* whole muscle-specific tension of LRT versus UNT (11), with a similar magnitude (+9%) to the difference in myosin spacing found in the current study (−7%), although a relatively small single fiber study (groups of $n = 6$) found no significant difference in fiber-specific tension of LRT versus UNT (45). Nonetheless, the current study provides the first evidence of differences in the ultrastructure of skeletal muscle tissue in LRT versus UNT that would be expected to modestly enhance specific tension.

Limitations. There are some limitations in this study. First, and most importantly, this was a cross-sectional study and not a longitudinal one, which makes it impossible to attribute causality (i.e., which adaptations were due to RT and which were inherent to the participants). However, studying LRT versus UNT individuals has the advantage to be able to identify larger potential adaptations/differences (12,14) than in short/medium-term longitudinal studies (25,27,28), and this may better inform the capability for adaptation. Nevertheless, more direct longitudinal studies of sufficient RT duration (e.g., ≥ 1 yr), ideally with multiple measurements over the intervention period, are warranted to further develop our knowledge of RT adaptations.

In addition, it should be acknowledged that not all the muscle ACSA is muscle fiber material (12), nor is all the fiber CSA myofibrillar material (46), yet our calculations of muscle fiber and myofibril numbers assume that they are. Thus, the actual numbers of muscle fibers and myofibrils will be lower than the estimated numbers reported in this study. However, in young healthy participants, a majority (80%–85%) of both muscle ACSA and fiber CSA do consist of muscle fibers and myofibrillar material, respectively, and these proportions appear to be consistent with RT (12,25,41,44,46). Therefore, it seems likely that our findings of LRT individuals having more muscle fibers and myofibrils than UNT participants, as well as the correlations between muscle ACSA and muscle




FIGURE 7—Summary of the differences in whole muscle, muscle fiber and ultrastructural variables of LRT compared with UNT males. Data are percentage differences in group mean values.

fiber or myofibril numbers, remain robust even when the noncontractile components of the biceps brachii muscle are taken into account. Nevertheless, we recommend that future studies correct for noncontractile components to estimate muscle fiber and myofibril numbers more precisely.

Finally, elements of our correlation analysis may have been confounded by circularity. For example, the relationship between muscle ACSAmax and fiber number (derived from ACSAmax divided by average fiber area) involves circularity, which may confound this relationship (47). Therefore, the between-group difference in fiber number within the muscle ACSAmax between LRT versus UNT may be more robust evidence than the relationship of fiber number with ACSAmax. In addition, we are aware that our fiber number calculation was a crude estimation mainly because of innate difficulty in generalizing single biopsy data to the whole muscle level (48); however, currently, there appears no other way to estimate fiber number of human muscles *in vivo*. Therefore, we used the same approach as previous studies (12,14) but with a larger sample size per group, and found clearer between-group differences than the previous investigations and a large effect size ($d = 1.02$) for greater fiber number within the muscle cross section of LRT versus UNT. Furthermore, the disproportionately greater group difference in ACSAmax (70%) compared with fiber area (29%) alone suggests that fiber number was greater for LRT than UNT. Therefore, although our approach of indirect estimation of fiber number has an inherent methodological limitation, we consider that our overall findings including fiber number add important cross-sectional data to the existing literature. Although it is impossible to directly measure the size and number of all muscle fibers *in vivo*, advancing diffusion tensor magnetic resonance imaging techniques have been shown to be

promising tools to noninvasively estimate muscle fiber diameters and fiber types in areas of interest (49,50). Thus, future studies should be directed toward adopting/developing these emerging techniques, together with traditional biopsy methods, to better understand the plasticity of muscle fibers and myofibrils following RT.

CONCLUSIONS

The main findings of this study are summarized in Figure 7. Compared with UNT, LRT had a substantially larger biceps brachii muscle area, which was mediated by having not only larger muscle fiber area but also a greater number of fibers. LRT had more myofibrils per fiber and in total, with a tendency for smaller myofibrils. A novel finding of this study was the greater myofilament packing density of LRT, suggesting that skeletal muscle ultrastructure may be adaptable and could contribute to changes in specific tension and strength.

The authors would like to thank the School of Life Sciences Imaging Centre at the University of Nottingham for sectioning the muscle tissues for transmission electron microscopy analysis. The authors acknowledge the use of facilities within the Loughborough Materials Characterization Centre.

This work was partly supported by JSPS KAKENHI grant 21H03335 (S. M.). The authors declare no competing interests. S. M. and J. P. F. conceived, designed, and drafted the study. All authors contributed to data collection or analysis, as well as interpretation of the results. All authors have read and approved the final version of the manuscript, including the order of presentation of the authors. This study was approved by Loughborough University Ethical Advisory Committee (R17-P174), and written informed consent was obtained from each participant. All data are available in the main text. Additional data related to this study will be made available by the corresponding author upon reasonable request. The results of the study are presented clearly, honestly, and without fabrication, falsification, or inappropriate data manipulation. The results of the present study do not constitute endorsement by the American College of Sports Medicine.

REFERENCES

- Balshaw TG, Maden-Wilkinson TM, Massey GJ, Folland JP. The human muscle size and strength relationship: effects of architecture, muscle force, and measurement location. *Med Sci Sports Exerc.* 2021;53(10):2140–51.
- O'Brien TD, Reeves ND, Baltzopoulos V, Jones DA, Maganaris CN. Strong relationships exist between muscle volume, joint power and whole-body external mechanical power in adults and children. *Exp Physiol.* 2009;94(6):731–8.
- Miller R, Balshaw TG, Massey GJ, et al. The muscle morphology of elite sprint running. *Med Sci Sports Exerc.* 2021;53(4):804–15.
- Kordi M, Folland J, Goodall S, et al. Mechanical and morphological determinants of peak power output in elite cyclists. *Scand J Med Sci Sports.* 2020;30(2):227–37.
- Prado CM, Purcell SA, Alish C, et al. Implications of low muscle mass across the continuum of care: a narrative review. *Ann Med.* 2018;50(8):675–93.
- Wang H, Hai S, Liu Y, Liu Y, Dong B. Skeletal muscle mass as a mortality predictor among nonagenarians and centenarians: a prospective cohort study. *Sci Rep.* 2019;9(1):2420.
- Frontera WR, Ochala J. Skeletal muscle: a brief review of structure and function. *Calcif Tissue Int.* 2015;96(3):183–95.
- Roberts MD, McCarthy JJ, Hornberger TA, et al. Mechanisms of mechanical overload-induced skeletal muscle hypertrophy: current understanding and future directions. *Physiol Rev.* 2023;103(4):2679–757.
- Maeo S, Balshaw TG, Lanza MB, Hannah R, Folland JP. Corticospinal excitability and motor representation after long-term resistance training. *Eur J Neurosci.* 2021;53(10):3416–32.
- Casolo A, Del Vecchio A, Balshaw TG, et al. Behavior of motor units during submaximal isometric contractions in chronically strength-trained individuals. *J Appl Physiol (1985).* 2021;131(5):1584–98.
- Maden-Wilkinson TM, Balshaw TG, Massey GJ, Folland JP. What makes long-term resistance-trained individuals so strong? A comparison of skeletal muscle morphology, architecture, and joint mechanics. *J Appl Physiol (1985).* 2020;128(4):1000–11.
- MacDougall JD, Sale DG, Alway SE, Sutton JR. Muscle fiber number in biceps brachii in bodybuilders and control subjects. *J Appl Physiol Respir Environ Exerc Physiol.* 1984;57(5):1399–403.
- Monti E, Toniolo L, Marcucci L, et al. Are muscle fibres of body builders intrinsically weaker? A comparison with single fibres of aged-matched controls. *Acta Physiol (Oxf).* 2021;231(2):e13557.
- Sale DG, MacDougall JD, Alway SE, Sutton JR. Voluntary strength and muscle characteristics in untrained men and women and male bodybuilders. *J Appl Physiol (1985).* 1987;62(5):1786–93.
- Murach KA, Dungan CM, Peterson CA, McCarthy JJ. Muscle fiber splitting is a physiological response to extreme loading in animals. *Exerc Sport Sci Rev.* 2019;47(2):108–15.
- Jorgenson KW, Phillips SM, Hornberger TA. Identifying the structural adaptations that drive the mechanical load-induced growth of skeletal muscle: a scoping review. *Cells.* 2020;9(7):1658.
- Schoenfeld BJ. The mechanisms of muscle hypertrophy and their application to resistance training. *J Strength Cond Res.* 2010;24(10):2857–72.

18. Goldspink G, Harridge SD. Cellular and molecular aspects of adaptation in skeletal muscle. In: Komi PV, editor. *Strength and Power in Sport*. Hoboken: Wiley; 2003. pp. 231–51.
19. Rehfeldt C, Stickland NC, Fiedler I, Wegner J. Environmental and genetic factors as sources of variation in skeletal muscle fibre number. *Basic Appl Myol*. 1999;9(5):235–54.
20. MacCallum JB. On the histogenesis of the striated muscle fibers and the growth of the human sartorius muscle. *Johns Hopkins Hosp Bull*. 1898;9:208–15.
21. Sjoström M, Lexell J, Eriksson A, Taylor CC. Evidence of fibre hyperplasia in human skeletal muscles from healthy young men? A left-right comparison of the fibre number in whole anterior tibialis muscles. *Eur J Appl Physiol Occup Physiol*. 1991;62(5):301–4.
22. Lexell J, Taylor CC, Sjoström M. What is the cause of the ageing atrophy? Total number, size and proportion of different fiber types studied in whole vastus lateralis muscle from 15- to 83-year-old men. *J Neurol Sci*. 1988;84(2–3):275–94.
23. McCall GE, Byrnes WC, Dickinson A, Pattany PM, Fleck SJ. Muscle fiber hypertrophy, hyperplasia, and capillary density in college men after resistance training. *J Appl Physiol* (1985). 1996;81(5):2004–12.
24. Nilwik R, Snijders T, Leenders M, et al. The decline in skeletal muscle mass with aging is mainly attributed to a reduction in type II muscle fiber size. *Exp Gerontol*. 2013;48(5):492–8.
25. MacDougall JD. Morphological changes in human skeletal muscle following strength training and immobilization. In: Jones NL, McCartney N, McComas AL, editors. *Human Muscle Power*. Champaign (IL): Human Kinetics; 1986. pp. 269–88.
26. Wang N, Hikida RS, Staron RS, Simoneau JA. Muscle fiber types of women after resistance training—quantitative ultrastructure and enzyme activity. *Pflugers Arch*. 1993;424(5–6):494–502.
27. Penman KA. Human striated muscle ultrastructural changes accompanying increased strength without hypertrophy. *Res Q*. 1970;41(3):418–24.
28. Claassen H, Gerber C, Hoppeler H, Luthi JM, Vock P. Muscle filament spacing and short-term heavy-resistance exercise in humans. *J Physiol*. 1989;409:491–5.
29. Folland JP, Williams AG. The adaptations to strength training: morphological and neurological contributions to increased strength. *Sports Med*. 2007;37(2):145–68.
30. Skarabot J, Balshaw TG, Maeo S, et al. Neural adaptations to long-term resistance training: evidence for the confounding effect of muscle size on the interpretation of surface electromyography. *J Appl Physiol* (1985). 2021;131(2):702–15.
31. Craig CL, Marshall AL, Sjoström M, et al. International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc*. 2003;35(8):1381–95.
32. Pipe JG. Motion correction with PROPELLER MRI: application to head motion and free-breathing cardiac imaging. *Magn Reson Med*. 1999;42(5):963–9.
33. Schindelin J, Arganda-Carreras I, Frise E, et al. Fiji: an open-source platform for biological-image analysis. *Nat Methods*. 2012;9(7):676–82.
34. McMillan JD, Eisenback MA. Transmission electron microscopy for analysis of mitochondria in mouse skeletal muscle. *Bio Protoc*. 2018;8(10):e2455.
35. Leys C, Ley C, Klein O, Bernard P, Licata L. Detecting outliers: do not use standard deviation around the mean, use absolute deviation around the median. *J Exp Soc Psychol*. 2013;49(4):764–6.
36. Miller J. Reaction time analysis with outlier exclusion: bias varies with sample size. *Q J Exp Psychol A*. 1991;43(4):907–12.
37. Campos GE, Luecke TJ, Wendeln HK, et al. Muscular adaptations in response to three different resistance-training regimens: specificity of repetition maximum training zones. *Eur J Appl Physiol*. 2002;88(1–2):50–60.
38. Roman WJ, Fleckenstein J, Stray-Gundersen J, Alway SE, Peshock R, Gonyea WJ. Adaptations in the elbow flexors of elderly males after heavy-resistance training. *J Appl Physiol* (1985). 1993;74(2):750–4.
39. Conceicao MS, Vechin FC, Lixandrao M, et al. Muscle fiber hypertrophy and myonuclei addition: a systematic review and meta-analysis. *Med Sci Sports Exerc*. 2018;50(7):1385–93.
40. Jorgenson KW, Hornberger TA. The overlooked role of fiber length in mechanical load-induced growth of skeletal muscle. *Exerc Sport Sci Rev*. 2019;47(4):258–9.
41. Jorgenson KW, Hibbert JE, Sayed RKA, et al. A novel imaging method (FIM-ID) reveals that myofibrillogenesis plays a major role in the mechanically induced growth of skeletal muscle. *Elife*. 2024;12:RP92674.
42. Vann CG, Roberson PA, Osburn SC, et al. Skeletal muscle myofibrillar protein abundance is higher in resistance-trained men, and aging in the absence of training may have an opposite effect. *Sports (Basel)*. 2020;8(1):7.
43. Jones DA, Rutherford OM. Human muscle strength training: the effects of three different regimens and the nature of the resultant changes. *J Physiol*. 1987;391:1–11.
44. Ruple BA, Godwin JS, Mesquita PHC, et al. Myofibril and mitochondrial area changes in type I and II tibers following 10 weeks of resistance training in previously untrained men. *Front Physiol*. 2021;12:728683.
45. Shoepe TC, Stelzer JE, Garner DP, Widrick JJ. Functional adaptability of muscle fibers to long-term resistance exercise. *Med Sci Sports Exerc*. 2003;35(6):944–51.
46. Luthi JM, Howald H, Claassen H, Rosler K, Vock P, Hoppeler H. Structural changes in skeletal muscle tissue with heavy-resistance exercise. *Int J Sports Med*. 1986;7(3):123–7.
47. Curran-Everett D. Explorations in statistics: the analysis of ratios and normalized data. *Adv Physiol Educ*. 2013;37(3):213–9.
48. Ruple BA, Mesquita PHC, Godwin JS, et al. Changes in vastus lateralis fibre cross-sectional area, pennation angle and fascicle length do not predict changes in muscle cross-sectional area. *Exp Physiol*. 2022;107(11):1216–24.
49. Cameron D, Abbassi-Daloui T, Heezen LGM, et al. Diffusion-tensor magnetic resonance imaging captures increased skeletal muscle fibre diameters in Becker muscular dystrophy. *J Cachexia Sarcopenia Muscle*. 2023;14(3):1546–57.
50. Hata J, Nakashima D, Tsuji O, et al. Noninvasive technique to evaluate the muscle fiber characteristics using q-space imaging. *PLoS One*. 2019;14(4):e0214805.