

**AXON NEUROLOGY SPECIALITY CENTER PLC**

| TIN: 0075102586

Tel: +251940155606 / +251974788888 | Fax: 011 278 2075

Email: Info@ethiopiastroke.com | Web: https://www.ethiopiastroke.com

MEDICAL ORDER

ID: #63048 | 09/05/2026

PATIENT NAME MERU MITHA .	MRN / ID 0013196	AGE / GENDER 62 / Female	ORDERED DATE ☐ P0+r ☐ P0+r ☐ P0+r ☐ P0+r ☐ P0+r ☐ P0+r ☐ P0+r ☐ P0+r ☐ P0+r 09/05/2026
REQUESTED BY Dr. Michael Yirgalem Teka	REQUESTED FROM Emergency Phase 2	COLLECTION DATE 09/05/2026	PATIENT PHONE +251901924424
COLLECTED BY Dr. Alemayehu Bedane Worke	PROCESSED BY Dr. Alemayehu Bedane Worke	VERIFIED BY Dr. Alemayehu Bedane Worke	AUTHORIZED BY Dr. Alemayehu Bedane Worke
AUTHORIZED DATE 09/05/2026			

IMAGING**MRI****# PARAMETER**

1 MRI of Lumbar Spine

CLINICAL EVALUATION & FINDINGS

Indication: Lower back pain radiating to bilateral lower extremity and Cervical pain radiating to the left hand associated with left hand weakness. (3 years duration) Reading by Dr Alex

Mitha Meru

INDICATION:

TECHNIQUE: Multiplanar, multisequence MR images of the lumbar spine were acquired without intravenous contrast

FINDINGS:

-Alignement: Unremarkable

-Vertebral body height and marrow signal: unremarkable

-T2 disc signals and heights : decreased disc signal at all levels suggesting their dehydration.

-Conus: : unremarkable

-Paraspinal soft tissue: Unremarkable

-L1-2 level is normal.

-L2-3 level is normal.

-L3-4 level is normal.

-L4-5 level is normal

-L5-S1 level : right facet hypertrophy and edema. Increased epidural fat/epidural lipomatosis . Mild diffuse disc bulge with added small central disc protrusion. All causing mild to moderate central canal narrowing. Bilateral mild to moderate neural foraminal stenosis

IMPRESSION:

-L5-S1 level : right facet hypertrophy and edema. Increased epidural fat/epidural lipomatosis . Mild diffuse disc bulge with added small central disc protrusion. All causing mild to moderate central canal narrowing. Bilateral mild to moderate neural foraminal stenosis

-There is posterior subluxation of C2 over C1 coccygeal segment.

N.B., left gluteal subcutaneous T2 intermediate signal lesion- ? Injection granuloma.

This is a digitally verified medical report. Results should be correlated clinically by the attending physician.

**Dr. Alemayehu Bedane
Worke**