

Superintegrable Systems. II.


N. Reshetikhin

YMSC

Tsinghua University

Last time: superintegrable models

classical and quantum.



Algebraically:

$$\begin{array}{c}
 C(M_{2n}) \xrightarrow{\quad} J_{2n-k} = C(P_{2n-k}) \xrightarrow{\quad} I_k = C(B_k) \\
 \text{Poisson with} \\
 \text{the trivial} \\
 \text{Poisson center} \\
 \uparrow \\
 \text{Poisson} \\
 \uparrow \\
 I_k = Z_Q(J_{2n-k})
 \end{array}$$

$(M_{2n} = \text{symplectic})$


Quantum

$$\begin{array}{ccc}
 A & \xrightarrow{\quad} & J \\
 \downarrow & & \downarrow \\
 C_h(M_{2n}) & \xrightarrow{\quad} & C_h(P_{2n-k}) \\
 & & \downarrow \\
 & & I = Z(J) \\
 & & \downarrow \\
 & & C_h(B_k)
 \end{array}$$


in any representation V of A joint eigensubspaces
 of I are finite direct sums of irreducible
 representations of J .

Examples

Kepler system
 (classical)



Hydrogen atom
 (quantum)



Spin Calogero - Moser - Sutherland systems

and their generalizations

① Spin Calogero - Moser (Hamiltonian reduction)
 J. Gibbons, T. Hermseu, 1984
 N.R., 2003, 2015

$$\text{nonsp} H = \frac{1}{2} \sum_{i=1}^n p_i^2 + \sum_{i < j} \frac{c}{(x_i - x_j)^2}, \quad \overline{\sin^2(x_i - x_j)} \quad \underline{\text{B. Sutherland}}$$

Classical: General fact: assume that

- M is a symplectic manifold,
- $G \times M \rightarrow M$ Hamiltonian action of a lie group.
i.e. $X \in g$, $X \cdot f = \{H_X, f\}$,
 $H_X(x) = \langle X, \mu(x) \rangle$, $\mu: M \rightarrow \mathfrak{g}^*$ moment map

Then:

- I M/G - Poisson (if nonsingular), $C(M)^G \subset C(M)$
 - $S(O) \subset M/G$ symplectic leaf, $O \subset g^*$
 - II $S(O) = \bar{\mu}'(O)/G$ $\bar{\mu}'(O) \subset M$
coisotropic pt $\nearrow \bar{x}_i$
Hamilton reduction
with respect to O
- Consider $M = T^*G \simeq \mathfrak{g}^* \times G$ (right translations)

(i) e_i - basis in \mathfrak{g} , $\pi: G \rightarrow \text{Aut}(V)$, $\underline{\underline{\pi_{ab}(g)}}$
 e_i - dual in \mathfrak{g}^* ,
 x_i - coord. on \mathfrak{g}^*

$$\left\{ \begin{array}{l} \{x_i, x_j\} = -C_{ij}^k x_k, \quad \{x_i, \pi_{ab}(g)\} = \sum_c \pi_{ac}(e_i) \pi_{cb}(g) \\ \{\pi_{ab}(g), \pi_{cd}(g)\} = 0 \end{array} \right. \quad T^*G \sim \wedge^* \mathfrak{g}$$

$\cdot G - \underline{\underline{g^* \times G}}$

(ii) Left "action" $G : (g^* \times G) \ni$

$$\alpha_L(h)(x, g) = (\text{Ad}_h^*(x), hg)$$

is Hamiltonian

$$Xf = \frac{d}{dt} f(\alpha_L(e^{tX})(x, g)) = \langle H_x^L, f \rangle_{\mathfrak{g}}(x, g), x \in \mathfrak{g}$$

$$H_x^L(x, g) = \underbrace{\langle X, x \rangle}_{\substack{x \in \mathfrak{g} \\ x \in \mathfrak{g}^*}} = \langle X, \mu_L(x, g) \rangle$$

$$\mu_L(x, g) = x.$$

$$\mu_L : T^*G \simeq \mathfrak{g}^* \times G \rightarrow \mathfrak{g}^*$$

is the moment map for the left action

(iii) For the right action the left action by mult. from the right

$$\alpha_R(h)(x, g) = (x, gh^{-1})$$

$$H_x^R(x, g) = - \langle X, \text{Ad}_{g^{-1}}^*(x) \rangle$$

$$\mu_R : T^*G \rightarrow \mathfrak{g}^*,$$

$$\mu_R(x, g) = - \text{Ad}_{g^{-1}}^*(x)$$

(iv) The adjoint action $(T^*G/\text{Ad}G)$

$$\text{Ad } h : (x, g) \mapsto (\text{Ad}_h^*(x), hg h^{-1})$$

$$\mu : T^*G \rightarrow \mathfrak{g}^*,$$

$$\mu = \mu_L + \mu_R$$

(v) Symplectic leaves of $T^*G/\text{Ad}G \simeq (\mathfrak{g}^* \times G)/\text{Ad}G$

$$0 \in \mathfrak{g}^*, 0 \in \mathfrak{g}^*/G,$$

$$S(0) = \bar{\mu}^{-1}(0)/\text{Ad}G$$

$$S(0) = \{ (x, g) \in \mathfrak{g}^* \times G \mid x - \text{Ad}_{g^{-1}}^*(x) \in 0 \} / \text{Ad}G$$

$G = \text{SU}_n$, by conjugation $g = g' h g'^{-1}$ $x = (x_{ij})$

$$i \neq j \quad x_{ij} - h_i x_{ij} h_j^{-1} = \mu_{ij} \text{ "coordinates" on } 0, \quad \mu_{ii} = 0$$

x_{ii} - no condition, remaining conjugations

with respect to $H \subset \text{SU}_n$ (diagonal matrices, Cartan subgroup). $N(H)$

Functions on $S(0) = \{ \text{Functions in } \mu_{ij}, i \neq j, h_i \overset{11}{\sim}, \mu_{ii} = 0 \}$,
 $x_{ii} \in N(H)$,

$$S(0) \simeq \left(\underset{x_{ii}=0}{T^*_H} \times (0//H) \right) / W, \quad N(H) \supset H$$

Hamilton. red. with
respect to H -action $r+r$


Alg. functions on $S(O) =$
 $= \text{Pol} \left(\frac{m_{ij}}{1 - h_i h_j^{-1}}, h_i^{\pm 1}, x_{ii} \right)^{S_n}$

$$N(H)/H = W$$

$$W = S_n$$

$$\dim(S(O)) = 2r + \underbrace{\dim(O)}_{p, q} - 2r = \dim(O), \quad r = n-1$$

A superintegrable system on $S(O)$



$$\mathfrak{g}^* \times_{\mathfrak{g}/G} \mathfrak{g}^* = \{(x, y) \mid Gx = - Gy\}$$

$$\cdot \quad \mathcal{P}(O) = \{(x, y) \mid Gx = - Gy, x - y \in O\} / r$$

$$\dim(\Phi(O)) = \dim(O) - r \quad \underline{u \in O}$$

$\cdot B(O) = \{O' \mid \underline{m(O', -O', O) \neq \emptyset}\}$

$$m(O', -O', O) = \{x, y \in O' \mid x - y \in O\} / G$$

$$= \overbrace{(O' \times (-O') \times O) / G}^{\text{sympl.}} \quad , \quad \underline{\text{symplectic}}$$

$$\dim(B(O)) = r \quad \begin{matrix} \text{Hamiltonian with} \\ \text{resp. to the diag. action of } G \end{matrix}$$

$\cdot \tilde{p}_2^{-1}(O') = m(O', -O', O), \quad \boxed{\dim(\tilde{p}_2^{-1}(O')) = \dim(O) - 2r}$

$$\Phi(O) = \bigcup_{O' \in B(O)} m(O', -O', O)$$

$$\dim(O) \geq r$$

Poisson commuting Hamiltonians:

$$\underline{I = C(B(O)) = \left. C(O^*)^G \right|_{B(O)}}$$

Casimir functions on O^* .

For SU_n :

$$\boxed{H_k = \frac{1}{k} \text{tr}(x^k)} \quad k=2, \dots, n$$

$$x_{ii} = p_i,$$

$$x_{ij} = \frac{\mu_{ij}}{1 - e^{i(q_i - q_j)}} \quad h_f = e^{iq_j}$$

$$H_2 = \frac{1}{2} \sum_{i=1}^n p_i^2 + \sum_{i < j} \frac{\mu_{ij} \mu_{ji}}{2 \sin^2(\frac{q_i - q_j}{2})}$$

for SU_n , $\mu_{ij} \mu_{ji} = |\mu_{ij}|^2$ like n particles
 $1D$, with "spin variable"
 $\mu_{ij}, \mu_{jk}, \dots = \mu_{ik} \delta_{jk} \dots$

Rank 1 orbits

$$\mu_{ij} = \overline{\psi_i} \psi_j - \delta_{ij} \frac{(\overline{\psi}, \psi)}{n}$$


O is $(2n-2)$ -dimensional

$$O // H \simeq \frac{pnt}{2n-2-2(n-1)}$$

$|\mu_{ij}| = c$. This is the case of the CM
nonspin model (Kazhdan-Kostant-Sternberg)

Quantum spin Calogero-Moser system

Quantization :



The Hilbert space G - compact simple

$$\mathcal{H}_\mu = \langle L_2(G \rightarrow V_\mu) \mid f(gh\bar{g}^{-1}) = \pi_\mu(g)f(h), g \in G, \underline{f(h) \in V_\mu[0]} \subset V_\mu \rangle \simeq \underline{\underline{L^2(H \rightarrow V_\mu[0])}}$$

$$h \in H \subset G, \underline{f(h) \in V_\mu[0]} \subset V_\mu \rangle \simeq \underline{\underline{L^2(H \rightarrow V_\mu[0])}}$$

$\mathcal{D}\text{iff}(G)$ acts naturally on \mathcal{H}_μ :

$$x \in g \subset \mathcal{D}\text{iff}(G)$$

$$(x_L f)(g) = \frac{d}{dt} f(e^{-tX}g) \Big|_{t=0}$$

$$(x_R f)(g) = \frac{d}{dt} f(g e^{tX}) \Big|_{t=0}$$

if $f \in \mathcal{H}_\mu$

$$(x_L + x_R)f(g) = \pi_\mu(x)f(g)$$

For sl_n , $\{e_{ij}\}$ -basis ($\sum_{i=1}^n e_{ii} = 0$)

$$(e_{ij}^R f)(h) = \frac{d}{dt} f(h e^{t e_{ij}}) \Big|_{t=0} = -h_i h_j^{-1} e_{ij}^L f(h)$$

$$\Rightarrow (e_{ij}^L - h_i h_j^{-1} e_{ij}^L) f(h) = \pi_\mu(e_{ij}) f(h)$$

sun

$$e_{ij}^L f(h) = \frac{1}{1 - h_i h_j^{-1}} \pi_\mu(e_{ij}) f(h),$$

$$e_{ii}^L f(h) = \frac{d}{dt} f(e^{-t e_{ii}} h) = -h_i^{-1} \frac{\partial f}{\partial h_i}(h)$$

$h_j = e^{i q_j}$


Corollary We can compute Casimirs $\mu_{ij} \mu_{ji}$

$$H = C_2 = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial q_i^2} + \frac{1}{4} \sum_{i \neq j} \frac{\pi_\mu(e_{ij} e_{ji})}{\sin^2(\frac{q_i - q_j}{2})}$$

$$C_k = \frac{1}{k} \text{tr}((e^L)^k)$$

Commutative differential operators.

Superintegrability of quantum spin CM



Joint spectrum of $\langle H_2, H_3, \dots \rangle$

$$\mathcal{H}_M = \bigoplus_{\lambda \in D_\mu} \mathcal{H}_\mu[\lambda], \quad \mathcal{H}_\mu[\lambda] \simeq \text{Hom}_G(V_\lambda \otimes V_\lambda^*, V_\mu)$$

$$D_\mu = \{\lambda \mid \text{Hom}_G(V_\lambda \otimes V_\lambda^*, V_\mu) \neq \emptyset\}$$

Vectors in $\mathcal{H}_\mu[\lambda]$:

$$\Psi_a(g) = \text{Tr}_{V_\lambda}(\pi_\lambda(g)a), \quad \text{Trace functions}$$

$$a : V_\lambda \rightarrow V_\lambda \otimes V_\mu , \quad a \in \text{Hom}_G(V_\lambda \otimes V_\lambda^*, V_\mu)$$

$$\hat{H}_k \psi_a(g) = c_k(\lambda) \psi_a(g)$$

Proposition $H_\mu(\lambda)$ is irreducible for \mathfrak{sl}_n .

Special cases:

References

tqfts.com

1) $\underline{\mu = m\omega_1}$ ($\text{Sym}^m(\mathbb{C}^n)$) \mathfrak{sl}_n ,

$V_\mu[0] \neq \{0\}$ iff $\underline{m=nk}$,

$$H = -\Delta + \sum_{i < j} \frac{2k(k+1)}{\sin^2\left(\frac{q_i - q_j}{2}\right)},$$

2) $\underline{\mu \vdash n}$, $\mu = \begin{array}{c} n \\ \vdash \end{array}$ n boxes,

In this case

$\underline{V_\mu[0]}$ = an irreducible S_n -module
corresponding to the Young diagram μ .

For $\underline{m > n}$

$$(\mathbb{C}^m)^{\otimes n} \cong \bigoplus_{\mu \vdash n} V_\mu^{\text{SL}_m} \otimes V_\mu [\circ]$$

$$\mathcal{H}_n^{(m)} = \boxed{L^2(h \rightarrow (\mathbb{C}^m)^{\otimes n})^{S_n}} = \bigoplus_{\mu} V_\mu^{\text{SL}_m} \otimes \mathcal{H}_\mu$$

n Bose particles with m internal degrees of freedom.

$$H = -\Delta + \sum_{i < j} \frac{1 + P_{ij}}{\sin^2(\frac{q_i - q_j}{2})}, \quad \checkmark$$

P_{ij} - permutation in $(\mathbb{C}^m)^{\otimes n}$. $L = \infty$

Lax matrix $L = x = \begin{pmatrix} p_i & \pi_\mu(e_{ij}) \\ p_{i\mu} & 1 - h_{ij} \end{pmatrix}$, $\frac{dL}{dt} = [H, L] =$
Semiclassical limit $= [L, M]_{\text{matrix}}$

$$\hat{H} = -\hbar^2 \Delta + \sum_{i < j} \frac{\pi_\mu(e_{ij} e_{ji})}{\sin^2(\frac{q_i - q_j}{2})},$$

- acts on \mathcal{H}_μ ,

- $\mathcal{H}_\mu[\lambda]$ is an eigensubspace with

$$\hat{H} \mathcal{H}_\mu[\lambda] = \hbar^2 c_2(\lambda) \mathcal{H}_\mu[\lambda]$$

$$\text{As } \hbar \rightarrow 0, \quad \mu = \frac{\mu_c}{\hbar}, \quad \lambda = \frac{\lambda_c}{\hbar}$$

$$(i) A_\mu \supset J_\mu \supset I_\mu$$

$\downarrow \hbar \rightarrow 0 \qquad \downarrow \hbar \rightarrow 0 \qquad \downarrow \hbar \rightarrow 0$

$$C(S(\mathcal{O}_{\mu_c})) \supset C(F(\mathcal{O}_{\mu_c})) \supset C(B(\mathcal{O}_{\mu_c}))$$

$$\hbar^2 c_2(\lambda) \rightarrow (\lambda_c, \lambda_c)$$

$$(ii) \quad \dim(\mathcal{H}_\mu[\lambda]) = \dim(\text{Hom}_G(V_\lambda \otimes V_\lambda^*, V_\mu)) = \\ = \hbar^{-\frac{n}{2}} \text{Vol}(M(\mathcal{O}_{\lambda_c}, \mathcal{O}_{-\omega_0(\lambda_c)}, \mathcal{O}_{\mu_c})) (1 + O(\hbar))$$

$$n = \dim(M(\mathcal{O}_{\lambda_c}, \mathcal{O}_{-\omega_0(\lambda_c)}, \mathcal{O}_{\mu_c}))$$

$$m(\mathcal{O}_s, \mathcal{O}_t, \mathcal{O}_u) = (\mathcal{O}_s \times \mathcal{O}_t \times \mathcal{O}_u) // G = \\ = \{(x, y, z) \mid x \in \mathcal{O}_s, y \in \mathcal{O}_t, z \in \mathcal{O}_u, \\ x+y+z = 0\} / G$$

In a Liouville integrable system
the multiplicity of an eigensubspace
is stable as $t \rightarrow 0$ and

$$\text{mult}(I_i \nabla = E_i \nabla) \stackrel{\hbar \rightarrow 0}{=} \#\left\{ \begin{array}{l} \text{connected} \\ \text{component} \end{array} \right. \text{of the corresponding} \\ \text{level surface of classical} \\ \text{integrals} \left. \right\}.$$

Next time:

- N-spin CM type systems.
 - 2D QCD revisited again.