

RESEARCH STATEMENT

KYLE BINDER

1. INTRODUCTION

My research is in combinatorial algebraic geometry, where I study combinatorial and algebraic properties of various cohomology theories on toric varieties.

Toric varieties provide a fascinating source of interactions between algebraic geometry, combinatorics, and commutative algebra. For one, normal toric varieties are in correspondence with rational polyhedral fans. Combinatorial objects like polytopes and atomic lattices induce rational polyhedral fans and therefore toric varieties. These toric varieties often remember pieces of the original combinatorial structure, and this gives a way to import powerful tools from algebraic geometry into combinatorics. Conversely, many geometric invariants of a toric variety can be computed in terms of the combinatorics of the fan, often using tools from commutative algebra.

These connections between geometry, combinatorics, and commutative algebra have recently led to a number of exciting developments involving various cohomology theories on toric varieties. Starting with [AHK18], many long standing conjectures in matroid theory have been resolved using the Chow cohomology of toric varieties associated to matroids; see also [ADH23, BST23]. Adiprasito's [Adi19] proof of the g -conjecture for PL spheres essentially studies Lefschetz properties for the intersection cohomology of simplicial toric prevarieties. These are both variants of the singular cohomology rings of toric varieties. Similarly, recent attention has been paid to describing toric vector bundles combinatorially [DRJS18, KM26, Dev23], and sheaf cohomology and K -theory on toric varieties has begun to play a central role in matroid theory [BEST23, EHL23, EFL25, Eur24].

Both of these cohomology theories are in practice computed via commutative algebra. For singular cohomology, one resolves the Stanley–Reisner ring of the fan over the symmetric algebra of the character lattice of the dense torus. For sheaf cohomology, one resolves the Cox module of the vector bundle over the Cox ring of the toric variety. In both of these cases, the geometry, combinatorics, and commutative algebra involved influence and interact with each other. My research focuses on how the combinatorial structure of these toric varieties and vector bundles influence the algebraic properties of the respective singular or sheaf cohomology, as well as the combinatorial properties of these cohomologies themselves.

- (1) **Singular cohomology of toric varieties.** My work focuses on the case of smooth (and potentially non-complete) toric varieties, whereas previous work has typically focused on the complete (and potentially singular) case. The combinatorics here is more subtle than in the complete case (§2.1.1), but my work on the singular cohomology rings of matroids (§2.1.2) shows that the combinatorial structure of the associated toric varieties can lead to explicit combinatorial descriptions of the singular cohomology ring and new Lefschetz properties.
- (2) **Tropical toric vector bundles.** Tropical toric vector bundles are a recent combinatorial and tropical abstraction of equivariant vector bundles on a toric variety. Any toric vector bundle admits numerous choices of tropicalization, and the resulting combinatorial structure of the tropical toric vector bundles constrains cohomological properties of the original bundle, such as the dimension of global sections. My work centers on extracting these relations between the combinatorics and cohomology of classical toric vector bundles and using them to develop a combinatorial notion of sheaf cohomology for tropical toric vector bundles.

2. PREVIOUS WORK

2.1. Singular cohomology of smooth toric varieties. We give a brief overview of the singular cohomology rings of smooth toric varieties in order to describe my previous research. Let X_Σ be a smooth toric variety defined by the rational fan $\Sigma \subseteq N_{\mathbb{Q}} \cong \mathbb{Q}^n$ with dense open torus T . The singular cohomology ring of X_Σ is strongly related to combinatorial commutative algebra. Indeed, Franz–Fu exhibit a natural isomorphism of rings

$$(1) \quad H^\bullet(X_\Sigma) \cong \mathrm{Tor}_\bullet^{\mathrm{Sym} N_{\mathbb{Q}}^\vee}(\mathbb{Q}[\Sigma], \mathbb{Q}),$$

where $\mathbb{Q}[\Sigma]$ is the *Stanley–Reisner* ring of Σ [FF24]. Moreover, the natural bigrading of Tor has a geometric meaning. Under the *weight filtration* $0 = W_0 H^\bullet(X_\Sigma) \leq \dots \leq W_{2n} H^\bullet(X_\Sigma) = H^\bullet(X_\Sigma)$ of Deligne’s mixed Hodge structure, Weber showed that $\mathrm{Tor}_i^{\mathrm{Sym} N_{\mathbb{Q}}^\vee}(\mathbb{Q}[\Sigma], \mathbb{Q})_j \cong \mathrm{Gr}_{2j}^W H^{2j-i}(X_\Sigma)$ [Web04]. In other words, the bigraded *Betti numbers* of $\mathbb{Q}[\Sigma]$ are the *Hodge numbers* of X_Σ .

2.1.1. Top non-vanishing weight cohomology of smooth toric varieties. The relationship between the singular cohomology ring of X_Σ and the combinatorics of Σ is subtle. For example, when X_Σ is complete, the Hodge numbers are combinatorially determined by Σ and independent of the embedding in $N_{\mathbb{Q}}$. Here, they correspond to the *h-vector* of the associated simplicial complex $\Delta(\Sigma)$.¹ When X_Σ is not complete this does not hold, and the Hodge numbers depend on the particular embedding $\Sigma \subseteq N_{\mathbb{Q}}$.

However, even though the entire singular cohomology ring of X_Σ is not combinatorially determined by Σ , some pieces of cohomology still are. We show in [Bin] that the top non-vanishing weight of cohomology depends only on the simplicial cohomology of $\Delta(\Sigma)$ and its subcomplexes.

Theorem 1. *Let X_Σ be a smooth toric variety. Then the top non-vanishing weight of cohomology is*

$$(2) \quad \mathrm{Gr}_{2n-2i}^W H^\bullet(X_\Sigma) \cong \bigoplus_{\substack{I \subseteq \Delta(\Sigma)_0 \\ |I| = |\Delta(\Sigma)_0| - i}} \tilde{H}^\bullet(\Delta(\Sigma)|_I)$$

where i is the smallest integer for which (2) is non-trivial and $\Delta(\Sigma)_0$ is the set of vertices of $\Delta(\Sigma)$.

Previously, it was only known that the *top-weight* cohomology $\mathrm{Gr}_{2n}^W H^\bullet(X_\Sigma)$ is combinatorially determined, using work of Hacking [Hac08] and Payne [Pay13] on boundary complexes and Alexander duality. When the top-weight cohomology vanishes, boundary complexes are not enough to see any of the other associated gradeds. Our approach instead uses (1) and flat base change of Tor .

In [Bin] we also study the action of the toric automorphism group $\mathrm{Aut}_T(X_\Sigma)$ (which permutes rays in Σ , so may be viewed as a subgroup of $\mathfrak{S}_{\Delta(\Sigma)_0}$) on the top non-vanishing weight cohomology and show that the isomorphism (2) is equivariant after twisting the right-hand side by $\det \mathrm{Cl}(X_\Sigma)_{\mathbb{Q}}$ and a sign representation. Using this description, we show that $\mathrm{Gr}_{2n-2i}^W H^\bullet(X_\Sigma)$ is only combinatorially determined as a vector space, not as an $\mathrm{Aut}_T(X_\Sigma)$ -representation.

Corollary 2. *We construct two rational fans $\Sigma_1, \Sigma_2 \subseteq N_{\mathbb{Q}}$ such that*

- (1) $\Delta(\Sigma_1) = \Delta(\Sigma_2)$
- (2) As subgroups of $\mathfrak{S}_{\Delta(\Sigma_1)_0}$, $\mathrm{Aut}_T(X_{\Sigma_1}) = \mathrm{Aut}_T(X_{\Sigma_2})$.
- (3) $\mathrm{Gr}_{2n}^W H^\bullet(X_{\Sigma_1})$ and $\mathrm{Gr}_{2n}^W H^\bullet(X_{\Sigma_2})$ are not isomorphic as $\mathrm{Aut}_T(X_{\Sigma_1})$ -representations.

¹This $\Delta(\Sigma)$ can be viewed as the simplicial complex of the intersection $\Sigma \cap S^{n-1}$ or the link of the origin in Σ .

2.1.2. *Singular cohomology rings of matroids.* Much of my work on the singular cohomology of toric varieties studies particular toric varieties arising from matroids, which are combinatorial objects that abstract the notion of linear independence in vector spaces. In their field-defining paper, Adiprasito, Huh, and Katz [AHK18] introduce the *Chow ring* of a matroid M to be the Chow ring of the toric variety X_{Σ_M} associated to the order complex of the lattice of flats of M . These toric varieties and their Chow rings have since played a central role in the geometric theory of matroids.

However, the Chow ring misses a lot of the geometry of X_{Σ_M} . Indeed, X_{Σ_M} is generally not compact, so the Chow ring injects, but does not surject, into the singular cohomology ring. In [Bin26a], I began studying the *singular cohomology ring* of a matroid, $H^\bullet(X_{\Sigma_M})$, in order to understand these toric varieties better. My main result shows that these singular cohomology rings contain interesting combinatorial structure.

Theorem 3 ([Bin26a, Theorem 6.6] and [Bin]). *Let M be a loopless matroid of rank r on ground set $[n]$. Then*

- (1) $\mathrm{Gr}_j^W H^k(X_{\Sigma_M}) = 0$ for $j > n - r + k$ and $j < 2k - 2r + 2$.
- (2) $H^k(X_{\Sigma_M}) = 0$ for $k > n + r - 2$.

As an $\mathrm{Aut}(M)$ -representation, $\mathrm{Gr}_{2n-2}^W H^{n+r-2}(X_{\Sigma_M}) \cong \mathrm{OS}^r(M) \otimes \mathrm{sgn}$, where $\mathrm{OS}^\bullet$ is the Orlik–Solomon algebra of M .

For uniform matroids, I was able to give a complete description of the singular cohomology ring, and I showed that this ring has a number of interesting combinatorial properties ranging from Lefschetz properties to real-rooted generating functions [Bin26b].

One of the miraculous properties of the Chow ring of a matroid is that it satisfies the *Kähler package*, and in particular, Hard Lefschetz. This property says that multiplication by powers of a generic hyperplane class induces isomorphisms between cohomology in complementary degree. However, the Kähler package for the singular cohomology ring fails due to the dimension of the top-degree cohomology often not being 1 (this is a consequence of Theorem 3). Still, we show that the singular cohomology rings of uniform matroids satisfy a slightly weaker *strong Lefschetz property* with respect to the weight filtration.

Theorem 4 ([Bin26b, Theorem 5.5]). *Let $\ell \in \mathrm{Gr}_2^W H^2(X_{\Sigma_{U_{r,n}}})$ be a generic hyperplane class. Then the map*

$$\times \ell^d : \mathrm{Gr}_j^W H^k(X_{\Sigma_{U_{r,n}}}) \longrightarrow \mathrm{Gr}_{j+2d}^W H^{k+2d}(X_{\Sigma_{U_{r,n}}})$$

is injective for $0 \leq d \leq r + j - 2k - 1$ and surjective for $d > r + j - 2k + 1$.

To our knowledge, this is the only sort of Lefschetz result known for non-compact varieties.

Another surprising feature of the Chow ring of a uniform matroid is that its generating function—the *Chow polynomial*—is real-rooted, and it is an important conjecture that the Chow polynomial of any matroid is real-rooted [FS24, Ste21]. For the singular cohomology ring, the correct analogues of the Chow polynomial are the *refined Hodge–Poincaré polynomials*

$$\underline{H}_{U_{r,n}}^i(x) = \sum_{j=0}^{r-1} \dim \mathrm{Gr}_{2i+2j}^W H^{2j+i}(X_{\Sigma_{U_{r,n}}}) \cdot x^j$$

for $0 \leq i \leq n - r$. These polynomials are the “higher weight” analogues of the Chow polynomial, with $\underline{H}_{U_{r,n}}^0(x)$ being the Chow polynomial itself.

Theorem 5 ([BV]). *The refined Hodge–Poincaré polynomials of a uniform matroid are real-rooted.*

The proof uses the combinatorics of an explicit basis of the singular cohomology ring I constructed in [Bin26b] along with a criterion for polynomials to be real-rooted from [BV25, Theorem 4.16].

2.2. Tropical toric vector bundles and sheaf cohomology. A toric vector bundle on X_Σ is a vector bundle $\mathcal{E}$ such that the structure map $\mathcal{E} \rightarrow X_\Sigma$ is T -equivariant. Such bundles were characterized by Kaneyama [Kan75] in terms of a cocycle condition and Klyachko [Kly89] in terms of a compatible family of decreasing filtrations of a vector space.

An exciting recent development in this theory is a combinatorial and tropical abstraction of toric vector bundles called *tropical toric vector bundles*, defined independently and simultaneously by Khan–MacLagan [KM26] and Kaveh–Manon [KM24]. Tropical bundles may be thought of as replacing the decreasing filtrations of a vector space in Klyachko’s description with a decreasing chain of flats of a matroid. Currently, tropical toric vector bundles have well-defined notions of equivariant K - and Chern-classes, Euler characteristic, and rank of global sections.

My research on tropical toric vector bundles has focused on providing a well-defined, combinatorial theory of sheaf cohomology. One major obstacle in defining this theory is that there is not yet a good notion of morphisms for tropical bundles, so there is no way of using a Čech complex for cohomology as in the classical case of toric vector bundles. However, in joint work with Sergio Cristancho, Caitlin Davis, Christopher Manon, Emma Pickard, and Mahrud Sayrafi, we construct a well-defined notion of Betti numbers for the sheaf cohomology of tropical toric vector bundles over $\mathbb{P}^n$. The main idea is that the various formulas for the Betti numbers of toric vector bundles on $\mathbb{P}^n$ given by Klyachko [Kly89] can all be defined matroid-theoretically, and then one is left to check that these formulas are well-behaved.

Theorem 6 ([BCD⁺]). *Klyachko’s formulas are well-defined for tropical toric vector bundles on $\mathbb{P}^n$, assuming the underlying matroid is modular. Moreover,*

- (1) *The alternating sum of Betti numbers computes the Euler characteristic.*
- (2) *(Horrocks’ splitting criterion) A tropical toric vector bundle $\mathcal{E}$ over $\mathbb{P}^n$ splits if and only if, for every toric line bundle $\mathcal{L}$, the higher Betti numbers of $\mathcal{E} \otimes \mathcal{L}$ vanish.*

3. FUTURE PROJECTS

3.1. Lefschetz properties on non-compact toric varieties. Lefschetz properties on the Artinian and Gorenstein reductions of Stanley–Reisner rings has attracted much attention recently in connection to the g -theorem [Adi19]. Morally speaking, these reductions correspond to the cohomology of *toric prevarieties* whose minimal orbits are fixed points, which is a generalization of compact toric varieties. However, my work on Theorem 4 suggests there is also much to discover about Lefschetz properties in the case of non-compact toric varieties.

Problem 7. *Which classes of smooth (non-complete) toric varieties exhibit Lefschetz properties?*

Answers to this question would enhance our knowledge of the geometry of non-compact toric varieties as well give interesting examples of Lefschetz properties in combinatorial commutative algebra.

While every projective toric variety exhibits Hard Lefschetz, there are examples of quasi-projective toric varieties which exhibit no Lefschetz properties. For example, a fan whose maximal cones have wildly different dimensions often will not exhibit Lefschetz properties. An answer to

Problem 7 must then involve determining structural properties of fans and toric varieties which force Lefschetz properties.

One potential structural property that appears promising is the toric variety having a large collection of nice torus-invariant divisors. In fact, the argument at the heart of Theorem 4 relies on the fact that each $X_{\Sigma_{U_r,n}}$ possesses a set of torus-invariant divisors D_i which (1) look like projective space times a torus, and (2) generate the reduced singular cohomology under the Gysin morphisms $H^\bullet(D_i) \rightarrow \tilde{H}^\bullet(X_{\Sigma_{U_r,n}})$. This sort of argument is very similar to the “partition of unity” argument of Adiprasito [Adi19], and getting a combinatorial description of such toric varieties would be very interesting.

Another approachable problem along the same lines of Problem 7 is the following.

Problem 8. *Suppose that Σ' is a subdivision of the rational fan $\Sigma \subseteq N_{\mathbb{Q}}$. Do X_{Σ} and $X_{\Sigma'}$ possess the same Lefschetz properties?*

A similar question for toric varieties satisfying the Kähler package, along with a few other properties, was answered in the affirmative by [ADH23], but the question is open for weaker Lefschetz properties. We suspect that our work on toric blow-ups in [Bin26a] will lead to a clear answer.

3.2. Singular cohomology of braid matroids and projective geometries. Our work on uniform matroids shows that the singular cohomology rings of matroids have rich combinatorial structure that extends the well-studied combinatorics in the Chow ring. One goal of our research program is to understand these combinatorics for all matroids.

Long-Term Goal 9. *For an arbitrary loopless matroid, give combinatorial formulas for the refined Hodge–Poincaré polynomials of the singular cohomology ring and produce a combinatorial basis.*

Aside from understanding the singular cohomology ring better, achieving this goal would give a better understanding of the Chow ring and Chow polynomial of a matroid. Current formulas for the Chow polynomials of a general matroid are recursive and do not give much combinatorial insight into the matroid itself. In our work on the singular cohomology rings of uniform matroids, the extra information contained in the singular cohomology ring allowed us to construct a natural basis for the Chow ring in a way that gave combinatorial proofs for many of the existing formulas for the Chow polynomials of uniform matroids. We are hopeful that the additional information captured by the singular cohomology ring yields similar combinatorial structure for the Chow polynomial of a general matroid.

As a series of stepping-stones to achieving this long-term goal, we plan on considering the singular cohomology of other explicit families of matroids with nice symmetry.

Problem 10. *Describe the singular cohomology rings of braid matroids and projective geometries.*

Our techniques for the singular cohomology of uniform matroids relied on the fact that the class of uniform matroids is closed under truncation and restriction. As the classes of braid matroids and projective geometries also behave well under these operations, we expect to solve Problem 10 using generalizations of our techniques for uniform matroids.

3.3. Tropical toric vector bundles on permutohedral varieties. We would like to extend our work on the cohomology of tropical toric vector bundles over projective space to tropical toric vector bundles over a general base.

Long-Term Goal 11. *Develop a combinatorial notion of higher cohomology for tropical toric vector bundles.*

At the moment, developing such a cohomology theory seems far off, as the obvious morphisms for tropical toric vector bundles do not have abelian structure. For example, Jun, Sistko, and Wright [JSW25] construct a nice category for tropical toric reflexive sheaves using morphisms from the theory of matroids over idylls [BL21]. However, this category is only *proto*-abelian, and derived functor cohomology is subtle here [FLS17]. In future work, it may be profitable to apply ideas from cohomology in $\mathbb{F}_1$ -geometry to resolve these issues.

A more attainable interim goal is to define higher cohomology over families of toric varieties with well-understood structure.

Problem 12. *Define higher cohomology for tropical toric vector bundles over the permutohedral toric variety X_E .*

A promising approach to this problem, and in fact its main motivation, is to combinatorialize the methods used in [Eur24, Liu25] to compute the cohomology of tautological bundles of matroids. The idea there is to use the deletion map $f: X_E \rightarrow X_{E \setminus n}$, derived pushforwards, and the Leray spectral sequence to reduce the computation to the cohomology of bundles over $\mathbb{P}^1$. The key to combinatorializing this process is to compute the pushforward of a toric vector bundle in terms of the Klyachko data.

Problem 13. *Let $f: X_\Sigma \rightarrow X_{\Sigma'}$ be a toric morphism, and let $\mathcal{E}$ be a toric vector bundle on X_Σ such that $f_*\mathcal{E}$ is a reflexive sheaf. Describe the Klyachko data for $f_*\mathcal{E}$ in terms of that for $\mathcal{E}$.*

We suspect that one can apply the ideas from [Zot25], which describes an algorithm for computing higher direct images in terms of Cox modules, in order to solve this problem. Aside from helping resolve Problem 12, an interesting and important application of Problem 13 would be to understand the combinatorics of the Frobenius pushforward of vector bundles on toric varieties, as this functor on toric vector bundles plays a key role in the theory of toric vector bundles on $\mathbb{P}^n$ [Rac25].

REFERENCES

- [ADH23] Federico Ardila, Graham Denham, and June Huh, *Lagrangian geometry of matroids*, J. Amer. Math. Soc. **36** (2023), no. 3, 727–794. MR 4583774
- [Adi19] Karim Adiprasito, *Combinatorial Lefschetz theorems beyond positivity*, <https://arxiv.org/abs/1812.10454>, 2019.
- [AHK18] Karim Adiprasito, June Huh, and Eric Katz, *Hodge theory for combinatorial geometries*, Ann. of Math. (2) **188** (2018), no. 2, 381–452. MR 3862944
- [BCD⁺] Kyle Binder, Sergio Cristancho, Caitlin Davis, Chris Manon, Emma Pickard, and Mahrud Sayrafi, *Tropical toric vector bundles over projective space*, In preparation.
- [BEST23] Andrew Berget, Christopher Eur, Hunter Spink, and Dennis Tseng, *Tautological classes of matroids*, Invent. Math. **233** (2023), no. 2, 951–1039. MR 4607725
- [Bin] Kyle Binder, *Top non-vanishing weight cohomology of toric varieties*, In preparation.
- [Bin26a] ———, *The singular cohomology ring of a matroid*, <https://arxiv.org/abs/2412.05732>, 2026.
- [Bin26b] ———, *The singular cohomology ring of a uniform matroid: combinatorics and Lefschetz properties*, <https://arxiv.org/abs/2606.07490>, 2026.
- [BL21] Matthew Baker and Oliver Lorscheid, *The moduli space of matroids*, Advances in Mathematics **390** (2021), 107883.
- [BST23] Andrew Berget, Hunter Spink, and Dennis Tseng, *Log-concavity of matroid h -vectors and mixed Eulerian numbers*, Duke Math. J. **172** (2023), no. 18, 3475–3520. MR 4718433

- [BV] Kyle Binder and Lorenzo Vecchi, *Real-rootedness in the singular cohomology of uniform matroids*, In preparation.
- [BV25] Petter Brändén and Lorenzo Vecchi, *Chow polynomials of totally nonnegative matrices and posets*, <https://arxiv.org/abs/2509.17852>, 2025.
- [Dev23] Lucie Devey, *A combinatorial description of stability for toric vector bundles*, <https://arxiv.org/abs/2212.11020>, 2023.
- [DRJS18] Sandra Di Rocco, Kelly Jabbusch, and Gregory G. Smith, *Toric vector bundles and parliaments of polytopes*, Trans. Amer. Math. Soc. **370** (2018), no. 11, 7715–7741. MR 3852446
- [EFL25] Christopher Eur, Alex Fink, and Matt Larson, *Vanishing theorems for combinatorial geometries*, <https://arxiv.org/abs/2510.05207>, 2025.
- [EHL23] Christopher Eur, June Huh, and Matt Larson, *Stellahedral geometry of matroids*, Forum Math. Pi **11** (2023), Paper No. e24, 48. MR 4653766
- [Eur24] Christopher Eur, *Cohomologies of tautological bundles of matroids*, Selecta Math. (N.S.) **30** (2024), no. 5, Paper No. 85, 19. MR 4805085
- [FF24] Matthias Franz and Xin Fu, *Cohomology of smooth toric varieties: naturality*, J. Pure Appl. Algebra **228** (2024), no. 6, Paper No. 107590, 38. MR 4687383
- [FLS17] Jaret Flores, Oliver Lorscheid, and Matt Szczesny, *Čech cohomology over $\mathbb{F}_{12}$* , Journal of Algebra **485** (2017), 269–287.
- [FS24] Luis Ferroni and Benjamin Schröter, *Valuative invariants for large classes of matroids*, J. Lond. Math. Soc. (2) **110** (2024), no. 3, Paper No. e12984, 86. MR 4795885
- [Hac08] Paul Hacking, *The homology of tropical varieties*, Collect. Math. **59** (2008), no. 3, 263–273. MR 2452307
- [JSW25] Jaiung Jun, Alex Sistko, and Cameron Wright, *Proto-exact categories of matroids over idylls and tropical toric reflexive sheaves*, arXiv preprint arXiv:2509.08144 (2025).
- [Kan75] Tamafumi Kaneyama, *On equivariant vector bundles on an almost homogeneous variety*, Nagoya Math. J. **57** (1975), 65–86. MR 376680
- [Kly89] A. A. Klyachko, *Equivariant bundles over toric varieties*, Izv. Akad. Nauk SSSR Ser. Mat. **53** (1989), no. 5, 1001–1039, 1135. MR 1024452
- [KM24] Kiumars Kaveh and Christopher Manon, *Tropical vector bundles and matroids*, <https://arxiv.org/abs/2405.03576>, 2024.
- [KM26] Bivas Khan and Diane Maclagan, *Tropical toric vector bundles*, Adv. Math. **484** (2026), Paper No. 110646, 67. MR 4993396
- [Liu25] Ruizhen Liu, *Vanishing theorems on wonderful varieties*, <https://arxiv.org/abs/2511.00696>, 2025.
- [Pay13] Sam Payne, *Boundary complexes and weight filtrations*, Michigan Math. J. **62** (2013), no. 2, 293–322. MR 3079265
- [Rac25] Feliks Rączka, *Frobenius pushforwards of vector bundles on projective spaces*, manuscripta mathematica **176** (2025), no. 1, 6.
- [Ste21] Matthew Stevens, *Real-rootedness conjectures in matroid theory*, Bachelor’s thesis, 2021.
- [Web04] Andrzej Weber, *Weights in the cohomology of toric varieties*, Cent. Eur. J. Math. **2** (2004), no. 3, 478–492. MR 2113544
- [Zot25] Sasha Zotine, *Computing higher direct images in Macaulay2*, <https://arxiv.org/abs/2505.22835>, 2025.