

CHARACTERIZATION THEORY OF PARACOMPACT SPACES AND BASIC COVERING PROPERTIES

Ji-Guang Jiang (蒋继光)

Preface

Paracompact spaces were introduced by Dieudonné [1944] as a natural generalization of compact topological spaces. On the other hand, fully normal spaces were defined by Tukey [1940] as a generalization of metric spaces. It is perhaps surprising that these two generalizations coincide, since Stone [1948] proved that a Hausdorff space is paracompact iff it is fully normal. In fact, the paracompact spaces provide one of the most successful common generalizations of compact spaces and metric spaces, and thus constitute one of the most important classes of spaces in point-set topology.

Paracompact spaces and the five related classes of covering properties are fundamental. Their characterization theorems are rich and colorful. The study of characterization theory for these basic covering properties has continued for more than 80 years since the publication of Tukey's paper [1940].

This book presents a systematic account of this theory. In Chapters 2 – 4, we present results published between 1940 and 1985, while Chapters 5 – 8 present results obtained by the author. Some of the characterization theorems appeared in Jiang [1987] – [1992] or in the manuscript Jiang [2015]. But the newest results appear here for the first time.

Finally, my sincere thanks go to my former students and longtime friends, Bing LONG (龙冰) and Maokang LUO (罗懋康). Bing carefully read through the entire manuscript and undertook the painstaking task of typesetting it. Maokang took care of the logistical arrangements and communications with the publishing house.

Chengdu, September 2026

Contents

Chapter 1 Preliminaries	4
1.1 Basic Covering Properties	4
1.2 Induded Covers	4
1.3 Inrerior-Preserving Families and Semi-Open Covers	4
Chapter 2 jiang8	5
Index	7

Chapter 1

Preliminaries

1.1 Basic Covering Properties

We use the usual terms and notations^[1]. An ordinal is the set of all smaller ordinals, and a cardinal is an initial ordinal. We use κ, μ for cardinals; λ a for infinite cardinal; $\alpha, \beta, \gamma, \theta$ for ordinal numbers. $\omega = \{0, 1, 2, \dots\}$ is the smallest infinite cardinal. $|A|$ denotes the cardinality of a set A . The empty set ϕ is the unique set B such that $|B| = 0$. $\mathbb{R}$ denotes the real line $[0, 1]$ the closed unit interval in $\mathbb{R}$.

For any sets $Z, Y, A \subset Z$, we denote $\mathcal{P}(Z) = \{A : A \subset Z\}$; $Y^X = \{f : f \text{ is a function from } X \text{ into } Y\}$. If $f \in Y^X$, we write $f : X \rightarrow Y$.

If $f : \mathbb{Z} \rightarrow Y$, $A \subset \mathbb{Z}$ and $B \subset Y$, the $f(A) : A \rightarrow Y$ defined by $f(A(x)) = f(x)$ for each $x \in A$; $f(A) = \{f(x) : x \in A\}$. $f^{-1}B = \mathcal{F}(B) = \{x \in \mathbb{Z} : f(x) \in B\}$.

For a set A and $n \rightarrow \infty$, let $[A]^n = \{B \subset A : |B| = n\}$,

$[A]^{<w} = \cup\{[A]^n : n < w\}$.

Let $w^0 = \{\phi\}$, for $n = 1, \dots, w^n =$

$= \{f : f; n \text{ to } w\} = \{(k_0, t_1, \dots, t_{n-1}) : \forall i < n \forall i \in w\}$ where

$m_i = f(i)$. If $t < w$, let $\phi \oplus h = (\overline{t_1})$, $(k_0, \dots, k_{n-1}) \oplus h =$

$(\overline{k_0}, \dots, \overline{k_{n-1}}, \dots)$. Let $w^{<w} = \cup w^n$.

Let $U = \{U_\alpha : \alpha \in \Delta\}$ be any family of subsets of $\mathbb{Z}$. Without loss of generality we may assume the indexing set Δ be any set of ordinals. We call

1.2 Induded Covers

1.3 Inrerior-Preserving Families and Semi-Open Covers

Chapter 2

jiang8

Alexandroff, P. S. 119247 Sur les ensembles de la premiere classe at les ensembles abstraits, C.R, Acad, Paris, 178 (19247, 185-187

@articleAlexandroff1924, author = Alexandroff, P. S., title = Sur les ensembles de la premiere classe at les ensembles abstraits, journal = C.R, Acad, Paris, volume = 178, number = 1, pages = 185–187, year = 1924, type = Journal Article

articleArensDugandji, author = Arens, R., title = Remark on the concepse of compactness, journal = Portugalice Math., volume= 9, number = , pages = 141–143 year = 1950, type = Journal Article

Arhangel'skir, A.V. 119617 New criteria for paracompactress, and metrizable of an arbitrary Ti-space, Dukel. Abad. Naut 555R, 141019617, 1051-1055. " " C19667 Mappings and spaces, Russiar Math. s u r v e y s , 2 1 0 1 9 6 6 3 1 1 5 - 1 6 2 . Bing, R. M. 119517 Matrization of topological spaces, Canad. a , M a t h . 3 0 1 9 5 1 2 1 7 5 - 1 8 6 . Boona, J. R. 619133 A characterization of meta compaotness in the class of o refinable spaces, fen. Tupalogy Apal, 3 (19732 253-264. Burke, D. K. C19,69J On subparacompact spaces. Proc. Amor. Mat

华罗庚数论导引^{[2]p. 1}

FeiH2 FeiH3

,
Atiyah^[3]

KelleyJ, General Topology^[4]

蒋继光, 一般拓扑学专题选讲^[5]

Bibliography

- [1] ALEXANDROFF P S. Sur les ensembles de la premiere classe at les ensembles abstraits [J]. C.R, Acad, Paris, 1924, 178: 185-187.
- [2] 华罗庚. 数论导引[M]. 科学出版社, 1957.
- [3] ATIYAH M F, MACDONALD I G. Introduction to commutative algebra[M]. Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1969: ix+128.
- [4] KELLEY J. General Topology (科学出版社有两版中译本)[M]. Springer-Verlag, 1975.
- [5] 蒋继光. 一般拓扑学专题选讲[M]. 四川教育出版社, 1991.

Index

cardinal, §1.1, 4

closed unit interval, §1.1, 4

empty set, §1.1, 4

ordinal, §1.1, 4

real line, §1.1, 4