

ECE 490 (Introduction to Optimization) – Homework 2

Due: 11:59pm, Feb. 24

Problem 1. (15 points) Minimization of Quadratic Functions

- (a) Suppose $f(x_1, x_2, x_3) = 2x_1^2 + 2x_2^2 + x_3^2 + 2x_2x_3 - 2x_1 - 2x_2 - 2x_3 + 5$. Find the minimum and maximum of f over \mathbb{R}^3 if they exist.
- (b) Consider the positive definite quadratic minimization problem $\min_{x \in \mathbb{R}^n} \frac{1}{2}x^T Qx$ where Q is a positive definite matrix. Apply the gradient method with the stepsize α_k chosen by the direct line search. What is α_k ? Write out α_k as a function of Q and x_k .
- (c) Consider the ridge regression problem $\min_{x \in \mathbb{R}^n} \frac{1}{n} \sum_{i=1}^n \{(a_i^T x - b_i)^2 + \frac{\lambda}{2} \|x\|_2^2\}$, where $\lambda > 0$ and (a_i, b_i) (for $i = 1, 2, \dots, n$) are given. What is the optimal solution x^* ? is the optimal solution unique?

Problem 2. (20 points) Convexity:

- (a) Consider $f : \mathbb{R}^n \rightarrow \mathbb{R}$. For $y_1, y_2 \in \mathbb{R}^n$, define the function $g : \mathbb{R} \rightarrow \mathbb{R}$ by $g(x) = f(x(y_1 - y_2) + y_2)$, $x \in \mathbb{R}$. Prove that f is convex on \mathbb{R}^n if and only if g is convex on \mathbb{R} for all $y_1, y_2 \in \mathbb{R}^n$.
- (b) Is the following set convex?

$$\mathcal{S} = \{x = (x_1, x_2) \in \mathbb{R}^2 : x_1 > 0, x_2 > 0, \text{ and } x_1 \log(x_1) + x_2 \log(x_2) \leq 4\}.$$

- (c) Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be a concave function and $f : \mathbb{R} \rightarrow \mathbb{R}$ be a concave increasing function. Prove that $f(g(x))$ is a concave function.

Problem 3. (10 points) Consider the problem of minimizing the function $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ defined as:

$$f(x) = f(x_1, x_2) = 2x_1^2 + 2x_2^4$$

. Use steepest descent with Armijo's Rule, with parameters $\tilde{\alpha} = 1$, $\sigma = 0.05$, and $\beta = 0.5$. Find α_k if $x_1 = (1, 0)$.

Problem 4. (25 points) Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex function and assume that $f(x) \geq f_{\min}$, for all x . Consider the following version of gradient descent method with constant step size

$$x_{k+1} = x_k - \alpha D \nabla f(x_k),$$

where D is a positive definite matrix. Let $\lambda_{\min} > 0$ and $\lambda_{\max} > 0$ denote the minimum and maximum eigenvalues of D . Assume that f has Lipschitz gradients with Lipschitz constant L . Show that if

$$0 < \alpha < \frac{2\lambda_{\min}}{L\lambda_{\max}^2},$$

then

$$\lim_{k \rightarrow \infty} \nabla f(x_k) = 0.$$

Problem 5. Define $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ as

$$f(x_1, x_2) = x_1^2 + 2\frac{1-\epsilon}{1+\epsilon}x_1x_2 + x_2^2$$

with $0 < \epsilon < 1$. Now, we consider the minimization problem of f , i.e. $\min_{x_1, x_2} f(x_1, x_2)$.

- (a) (5 points) What are the minimizers of f ?
- (b) (10 points) Find the largest $m > 0$ and the smallest $M > 0$ in terms of ϵ such that

$$mI \preceq \nabla^2 f(x_1, x_2) \preceq MI$$

for all (x_1, x_2) , where I is the identity matrix. Find the condition number of $\nabla^2 f$ given by $\kappa := \frac{M}{m}$ in terms of ϵ .

- (c) (5 points) How does κ change as ϵ decreases to 0. Do you expect gradient descent to converge faster or slower as ϵ decreases to 0?
- (d) (20 points) Write Python code to implement gradient descent with constant step-size $\alpha = \frac{m}{2M^2}$ starting from $(x_1, x_2) = (1, 1)$. Plot the trajectories of gradient descent for $\epsilon = 1, 0.1, 0.01, 0.001$ in the parameter space, i.e., the space of (x_1, x_2) , along with the values of $f(x_1, x_2)$ during the optimization. Specifically, for each choice of ϵ , you need to plot two figures: (i) scatter plot showing (x_1, x_2) points for all iterations, where the x -axis is for x_1 and the y -axis is for x_2 ; you need to mark the initial and final iterations in the plot. (ii) plot the curve of $f(x_1, x_2)$ vs. the number of iterations. Does the variation of your observed rate of convergence with ϵ agree with your expectation in part (c)? What happens with the simulations if you try $\alpha = \frac{1}{M}$? Does the algorithm still converge?

Problem 6. (20 points) Consider a continuously differentiable function $f : \mathbb{R}^n \rightarrow \mathbb{R}$.

- (a) Assume that f has Lipschitz gradients with Lipschitz constant L . In addition, f is also m -strongly convex. Prove the following inequality holds for all $x, y \in \mathbb{R}^n$

$$(\nabla f(x) - \nabla f(y))^T(x - y) \geq \frac{mL}{m + L} \|x - y\|^2 + \frac{1}{m + L} \|\nabla f(x) - \nabla f(y)\|^2.$$

- (b) Use the above inequality to show that the gradient descent method $x_{k+1} = x_k - \alpha \nabla f(x_k)$ with $\alpha = \frac{1}{L}$ satisfies the following convergence rate bound:

$$\|x_k - x^*\| \leq \left(1 - \frac{m}{L}\right)^k \|x_0 - x^*\|,$$

where x^* is assumed to be the unique global min of f .