





		CONTROLLED FORM		
TITLE: GXP and Non-GXP Study Report Review and Approval Form				
DOCUMENT NO. GE003FM01	VERSION 02	ISSUE DATE 17 Oct 2014	EFFECTIVE DATE 20 Oct 2014	PAGE 2 of 2

☐ Vendor Study Protocol		☒ Vendor Study Report		☐ Lpath Study Report	
Lpath Study No.: LT3114-PHA-008-R		Version No.: 01		☒ Non GXP GCP	☐ GLP GMP
Product Name: LT3114		Test Article: B3; LT1015			
Study Title: Evaluation of selected test substances for analgesic activity using the spinal nerve ligation (Chung) model of neuropathic pain in the rat (i.v. administration)					
Study Vendor/Site: ☐ N/A Porsolt, France		Study Director: ☐ N/A Rosalia Matteo		Study Contributor(s): ☐ N/A Ann Gregus; Stephanie Knapik	
Vendor Study No./Version: 14.0061		☒ N/A		Study Dates: July - August 2014	Report Date: October 2014

Lab Notebook Reference: LNB#278 pp.36-37 (Plasma concentration analysis - Appendix 1)		☐ N/A
Executive summary of study results:		☐ See Attached
The efficacy of the murine antibody, B3, was examined in a model of spinal nerve ligation (Chung model) in rats. Rats were surgically prepared by ligation of the L5 nerve and were allowed to recover for 14 days. All animals were tested for responses to either mechanical (electronic von Frey measurement) or thermal (infrared radiant source) stimuli prior to surgery, prior to antibody administration (days 11-13) and then 3 days after each dose of antibody. Antibodies (B3 and control IgG; LT1015) were administered at 30mg/kg intravenously on day 14, 17, 21 and 24 following surgery. Morphine (12mg/kg ip; administered 30 minutes before testing on Day 14) and gabapentin (100mg/kg ip; administered 180 minutes before testing on Day 21) were used as comparator agents. Spinal ligation resulted in a marked decrease in the force needed to induce paw withdrawal for both mechanical and thermal stimulation. As expected, morphine and gabapentin increased the paw withdrawal thresholds for both tactile and thermal stimuli. Both LT 1015 and B3 transiently improved paw withdrawal following tactile stimulation following the first dose but overall there was no significant effects seen. However, B3 demonstrated significant improvement in the paw withdrawal latency following thermal stimulation at both doses in comparison with animals treated with control antibody. The benefit of B3 treatment at the lower dose (10mg/kg) was seen up to the last day of dosing (24 days) whereas at 30mg/kg the response decreased following the day 21 dose.		

LPATH APPROVAL			
Function	Name (Print)	Signature	Date
Study Director/Monitor	Rosalia G. Matteo		11-26-2014
Contributor	Stephanie Knapik Ann M. Gregus		11/26/14 11/26/14
Research	GARY WOODRUFF		26 th November 2014
Development	DARIO PAGGIARINO		12/1/2014
Quality Assurance	ELVIRA MARQUEZ		01 DEC 14

STUDY N° 14.0061
STUDY REPORT
EVALUATION OF SELECTED TEST SUBSTANCES FOR ANALGESIC ACTIVITY
USING THE SPINAL NERVE LIGATION (CHUNG) MODEL OF NEUROPATHIC PAIN IN THE RAT
(I.V. ADMINISTRATION)

Sponsor:

LPATH INC.
4025 Sorrento Valley Blvd.
San Diego, CA 92121
U.S.A.

Author:

Dr. Camille LE CUDENNEC

Sponsor Study Monitor:

Dr. Gary WOODNUTT

Test Facility:

PORSOLT
Research Laboratory
Z.A. de Glatigné
53940 Le Genest-Saint-Isle
FRANCE

<u>Study Plan Approval Date</u>	:	24 June 2014
<u>Experimental Start Date</u>	:	04 July 2014
<u>Experimental Termination Date</u>	:	04 August 2014
<u>Draft Report Date</u>	:	21 August 2014
<u>Final Report Date</u>	:	06 October 2014

TABULATED SUMMARY REPORT N° 14.0061
Page 1/2

<u>SPONSOR</u> : LPATH INC.	<u>SPONSOR STUDY REFERENCE</u> : Not communicated
<u>TEST SUBSTANCES</u> : 1015 and B3	<u>TITLE</u> : Evaluation of selected test substances for analgesic activity using the Spinal Nerve Ligation (Chung) model of Neuropathic Pain in the rat) (i.v. administration).
<u>GENERIC NAME</u> : Not communicated	
<u>INDICATION</u> : Not communicated	<u>STUDY PERIOD</u> : July - August 2014 <u>DRAFT REPORT DATE</u> : 21 August 2014 <u>FINAL REPORT DATE</u> : 06 October 2014
<u>AIM OF STUDY</u> : To evaluate the potential analgesic effects of 1015 and B3 using the Chung/SNL model of neuropathic pain in the rat.	
<u>ANIMALS</u>	: Male Sprague Dawley rats (Janvier Labs), 91-138 g (Day -3) body weight range.
<u>CONDITIONING OF ANIMALS</u>	: Stabilized for at least 5 days after delivery in macrolon cages (425 x 266 x 185 mm; 5 animals per cage) on wood litter with free access to food and water.
<u>TEST SUBSTANCES AND</u>	: 1015 (Lpath Inc.), batch n° HR121458, colorless solution, diluted in phosphate buffered saline (PBS). B3 (Lpath Inc.), batch n° HR121456, colorless solution, diluted in PBS.
<u>TREATMENT SCHEDULE</u>	30 mg/kg, administered i.v. on Day 14, Day 17, Day 21 and Day 24 (i.e. 72 hours or 96 hours before the test) (3 ml/kg). Doses expressed in mg/kg of supplied substance.
<u>CONTROL SUBSTANCE</u>	: Phosphate buffered saline (Sigma, reference n° P-5368, batch n° SLBH8322V), administered under the same experimental conditions.
<u>REFERENCE SUBSTANCE</u>	: Morphine hydrochloride (Coopération Pharmaceutique Française, reference n° 1515300), batch n° D3065/2, white powder, dissolved in physiological saline. 12 mg/kg i.p., administered 30 minutes before the test on Day 14 (3 ml/kg). Doses expressed in mg/kg of salt.
<u>COMPARISON SUBSTANCE</u>	: Gabapentin (Toronto Research Chemicals Inc., reference n° G117250), batch n° 2-YFD-160-1, white powder, dissolved in physiological saline. 100 mg/kg i.p., administered 180 minutes before the test on Day 21 (3 ml/kg). Doses expressed in mg/kg of supplied substance.
<u>METHOD OF EVALUATION</u>	: Effects in the test in comparison with vehicle controls.
<u>STATISTICAL ANALYSIS</u>	: One-way ANOVA followed by Dunnett's t test and Unpaired Student's t tests.
<u>STUDY CONDUCTED BY THE APPLICANT:</u> YES ☐ NO ☒	
<u>IF "NO", INDICATE THE NAME AND ADDRESS OF THE INSTITUTE THAT CONDUCTED THE STUDY:</u> Porsolt Research Laboratory, Z.A. de Glatigné, 53940 Le Genest-Saint-Isle, France	
<u>STUDY IN COMPLIANCE WITH GLP:</u> YES ☐ NO ☐ NOT REQUIRED ☒ N PAGES: 75	

TABULATED SUMMARY REPORT N° 14.0061
Page 2/2

<u>SPONSOR</u> : LPATH INC.	<u>SPONSOR STUDY REFERENCE</u> : Not communicated
<u>TEST SUBSTANCES</u> : 1015 and B3	<u>TITLE</u> : Evaluation of selected test substances for analgesic activity using the Spinal Nerve Ligation (Chung) model of Neuropathic Pain in the rat (i.v. administration).
<u>GENERIC NAME</u> : Not communicated	
<u>INDICATION</u> : Not communicated	<u>STUDY PERIOD</u> : July - August 2014 <u>DRAFT REPORT DATE</u> : 21 August 2014 <u>FINAL REPORT DATE</u> : 06 October 2014

RESULTS:

Pre-tests

- Before surgery, inter-groups comparison displayed similar results against tactile and thermal stimulations in both hindpaws. Values of the results in the von Frey Test increased in parallel of body weight between Day -3 and Day -1. A slight difference between hindpaws was noted in the Plantar Test on Day -1 in the group to be treated with 1015. This difference was not considered as biologically relevant.

Baseline on Day 14

Tactile stimulation (Electronic von Frey Test)

- Spinal nerve ligation induced a marked decrease in the force inducing paw-withdrawal after tactile stimulation in the lesioned paw of all groups, as compared with the non-lesioned paw (% change between -86% and -89%, $p < 0.001$), indicating tactile allodynia.

Thermal stimulation (Plantar Test)

- Spinal nerve ligation induced a marked decrease in the paw-withdrawal latency after tactile stimulation in the lesioned paw of all groups, as compared with the non-lesioned paw (% change between -43% and -57%, $p < 0.001$), indicating thermal hyperalgesia.

Test

Tactile stimulation (electronic von Frey Test)

- Analysis of the force inducing paw-withdrawal of lesioned and non-lesioned paws (delta from baseline – test) using ANOVA (with group as main factor) revealed the absence of significant effects of the test substances at each testing days.
- Morphine (12 mg/kg i.p.), administered 30 minutes before the first test (Day 14), markedly increased the force inducing paw-withdrawal of the lesioned paw, as compared with vehicle controls (+28.2, $p < 0.001$). It did not affect the non-lesioned paw.
- Gabapentin (100 mg/kg i.p.), administered 180 minutes before the second test (Day 21), markedly increased the force inducing paw-withdrawal of the lesioned paw, as compared with vehicle controls (+14.9, $p < 0.001$). It did not affect the non-lesioned paw.

Thermal stimulation (Plantar Test)

- Analysis of paw-withdrawal latency of lesioned and non-lesioned paws (delta from baseline – test) using ANOVA (with group as main factor) revealed the absence of significant effects of the test substances at each testing days.
- Morphine (12 mg/kg i.p.), administered 30 minutes before the first test (Day 14), markedly increased the paw-withdrawal latency of the lesioned paw, as compared with vehicle controls (+25.8, $p < 0.001$). It did not affect the non-lesioned paw.
- Gabapentin (100 mg/kg i.p.), administered 180 minutes before the second test (Day 21), did not significantly affect the paw-withdrawal latency of the lesioned or non-lesioned paws, as compared with vehicle controls.

NB: The apparent increases of some behavioral endpoints following treatment with the test substances was explored by an additional statistical analysis using unpaired Student's t test and the following effects were observed:

Tactile stimulation (Electronic von Frey Test)

- 1015 (30 mg/kg i.v.), administered on Day 14, 3 days before the test 1, significantly increased the force inducing paw-withdrawal of the lesioned paw, as compared with vehicle controls (+5.8, $p < 0.05$).
- B3 (30 mg/kg i.v.), administered on Day 14, 3 days before the test 1, significantly increased the force inducing paw-withdrawal of the lesioned paw, as compared with vehicle controls (+6.4, $p < 0.05$). A similar trend was observed on Test 2 (4 days after the second administration) (+4.6, $p = 0.0815$).

Thermal stimulation (Plantar Test)

- 1015 (30 mg/kg i.v.), administered on Day 14, 3 days before the test 1, tended to increase the paw-withdrawal latency of the lesioned paw, as compared with vehicle controls (+6.0, $p = 0.0896$).
- B3 (30 mg/kg i.v.), administered on Day 14, 3 days before the test 1, significantly increased the paw-withdrawal latency of the lesioned paw, as compared with vehicle controls (+11.2, $p < 0.05$). This effect persisted on Test 2 (4 days after the second administration) (+11.5, $p < 0.05$). At 10 mg/kg, B3 clearly increased the paw-withdrawal latency of the lesioned paw, on test 3 (3 days after the third dosing) (+16.9, $p < 0.01$).

CONCLUSION:

These results suggest the absence of significant analgesic effects of 1015 (30 mg/kg) and B3 (10 and 30 mg/kg) administered i.v. at 4 opportunities against tactile allodynia and thermal hyperalgesia in the Neuropathic Pain Test (Chung model) in the rat.

However, some trends towards analgesic effects against tactile allodynia were observed 3 days after the first administration of 1015 and the higher dose of B3. Against thermal hyperalgesia, the higher dose of B3 showed a potential analgesic effect after the first (test on Day 17) and the second administration (test 2 on Day 21); the lower dose seemed more active after the third administration (test 3 on Day 24). This latter effect was confirmed by a complementary statistical analysis using two-way ANOVA.

Gabapentin (100 mg/kg i.p.) displayed analgesics effects against tactile allodynia and morphine (12 mg/kg i.p.) was active against tactile allodynia and thermal hyperalgesia in the same model.

TABLE OF CONTENTS

1.	REPORT APPROVAL	5
2.	EXPERIMENTAL PROCEDURE.....	6
	Neuropathic Pain, Chung/SNL model in the rat (<i>reference PI 13R</i>)	6
3.	RESULTS	8
	Neuropathic Pain, Chung/SNL model in the rat.....	8
4.	CONCLUSION	10
5.	TABLES.....	11
6.	FIGURES (produced by the Sponsor).....	18
7.	INDIVIDUAL DATA	23
8.	COMPLEMENTARY STATISTICAL ANALYSIS (produced by the Sponsor)	73

1. REPORT APPROVAL

Camille Le CUDENNEC, Ph.D.
Study Director



Date: 06 October 2014

2. EXPERIMENTAL PROCEDURE

Neuropathic Pain, Chung/SNL model in the rat (reference PI 13R)

The method, which detects antihyperalgesic activity in rats with neuropathic pain, follows that described by Kim and Chung (Pain, 50, 355-363, 1992).

Tight ligation of spinal nerves in rats is associated with hyperalgesia, allodynia and spontaneous pain, and therefore constitutes a model for peripheral neuropathic pain in humans. Antihyperalgesics reduce these chronic signs of pain hypersensitivity.

Pre-test (Days -3 to -1): Pre-surgery baseline von Frey and thermal thresholds for both hindpaws was measured.

Surgery (Day 0): Rats received an s.c. injection of Duphamox® LA 15-30 minutes before the surgery. Rats were anesthetized (combination of Ketamine and medetomidine) and an incision at the L4-S2 level was performed to expose the left L5 nerve. A ligature was tied tightly around the L5 nerve. The wound was then sutured and the rats were allowed to recover.

Pre-test (Days 11 or 12 or 13): when the chronic pain state was fully installed, rats were submitted to a pre-test and pseudo-randomly assigned to treatment groups matched on the basis of their pain response to tactile stimulation (electronic von Frey). Only rats who responded on the lesioned paw to forces between 0 and 25% of that of the non-lesioned paw were included in the experiment.

Testing phase (Days 14 - 27)

Baseline before treatment / Within-subject comparisons: On the first testing day (day 14), all animals were evaluated before treatment for their pain response consecutively to tactile (electronic von Frey test) and thermal (plantar test) stimulations of their lesioned and non-lesioned hindpaws. Dosing of groups 1 to 4 was then performed and rats were evaluated 3 days post dosing. Each testing phase was followed by a new dosing and rats were again evaluated 3 or 4 days thereafter. In the group 5, rats were again evaluated 30 minutes after administration of morphine. They were evaluated again on Day 21 180 minutes after administration of gabapentin and on days 24 and 27 similarly to the other groups.

Tactile allodynia evaluation: Electronic von Frey Test

The animal was placed under an inverted acrylic plastic box (18 x 11.5 x 14 cm) on a grid floor and left free to habituate for 5-8 minutes. The tip of an electronic Von Frey probe was then applied with increasing force first to the non-lesioned and to the lesioned hindpaw. The force required to induce paw-withdrawal was automatically recorded. This procedure was carried out 3 times and the mean force per paw was calculated.

Thermal hyperalgesia evaluation: Plantar Test

The apparatus consisted of individual acrylic plastic boxes (18 x 11.5 x 14 cm) placed upon an elevated glass floor. A rat was placed in the box and left free to habituate for 5-8 minutes. A mobile infrared radiant source (132 ± 10 mW/cm²) was then focused first under the non-lesioned and then the lesioned hindpaw. The paw-withdrawal latency was automatically recorded. In order to prevent tissue damage the heat source was automatically turned off after 45 seconds.

10 rats were studied per group. The test was performed blind.

The test substance 1015 was evaluated at 1 dose (30 mg/kg), B3 was evaluated at 2 doses (10 and 30 mg/kg). These substances were administered i.v. 3 or 4 days before each testing phase, and compared with a vehicle control group.

Morphine (12 mg/kg i.p.), administered 30 minutes before the test or Gabapentin (100 mg/kg i.p. or s.c.) administered 180 minutes before the test, were used as reference substance and comparison substance respectively. Dosing and testing of these substances were performed only on days 14 for morphine and for gabapentin on Day 21 after a washout period.

The experiment therefore included 5 groups.


Group	Number of animals	Treatment	Route	Dose-level (mg/kg)	Concentration (mg/ml)	Administration Volume (ml/kg)
1	10 males	PBS	i.v.	-	-	3
2	10 males	1015	i.v.	30	10	3
3	10 males	B3	i.v.	10	3.3	3
4	10 males	B3	i.v.	30	10	3
5	10 males	Morphine	i.p.	12	4	3
		Gabapentin (after washout)	i.p.	100	33.3	

Inter-group comparison will be performed for the test substances using ANOVA followed by post hoc analysis for each measure on delta (test-baseline) of lesioned and non-lesioned paw. Data with the reference/comparison substances will be compared with the pre-dose baseline using ANOVA followed by post hoc analysis.

At the end of the testing phase, the animals were anesthetized and then whole blood samples were collected by cardiocentesis into blood collection tubes containing K2 EDTA. Blood samples were kept cold and immediately spun in a centrifuge (5000 rpm for 10 minutes at 4°C). After centrifugation, the entire resultant plasma obtained was immediately transferred to 2 suitably labeled polypropylene tubes (2 aliquots of approximately 200-250 µl if possible). The tubes were stored upright at -80°C ± 10°C until shipped for analysis.

Immediately following blood collection, the animals were euthanized by CO₂ asphyxiation.

Planning (except for group 5):



3. RESULTS

Neuropathic Pain, Chung/SNL model in the rat

Pre-tests

- Before surgery, inter-groups comparison displayed similar results against tactile and thermal stimulations in both hindpaws. Values of the results in the von Frey test increased in parallel of body weight between Day -3 and Day -1. A slight difference between hindpaws was noted in the plantar test on Day -1 in the group to be treated with 1015. This difference was not considered as biologically relevant.

Baseline on Day 14

Tactile stimulation (Electronic von Frey)

- Spinal nerve ligation induced a marked decrease in the force inducing paw-withdrawal after tactile stimulation in the lesioned paw of all groups, as compared with the non-lesioned paw (% change between -86% and -89%, $p < 0.001$), indicating tactile allodynia.

Thermal stimulation (Plantar Test)

- Spinal nerve ligation induced a marked decrease in the paw-withdrawal latency after tactile stimulation in the lesioned paw of all groups, as compared with the non-lesioned paw (% change between -43% and -57%, $p < 0.001$), indicating thermal hyperalgesia.

Test

Tactile stimulation (Electronic von Frey Test)

- Analysis of the force inducing paw-withdrawal of lesioned and non-lesioned paws (delta from baseline – test) using ANOVA (with group as main factor) revealed the absence of significant effects of the test substances at each testing days.
- Morphine (12 mg/kg i.p.), administered 30 minutes before the first test (Day 14), markedly increased the force inducing paw-withdrawal of the lesioned paw, as compared with vehicle controls (+28.2, $p < 0.001$). It did not affect the non-lesioned paw.
- Gabapentin (100 mg/kg i.p.), administered 180 minutes before the second test (Day 21), markedly increased the force inducing paw-withdrawal of the lesioned paw, as compared with vehicle controls (+14.9, $p < 0.001$). It did not affect the non-lesioned paw.

Thermal stimulation (Plantar Test)

- Analysis of paw-withdrawal latency of lesioned and non-lesioned paws (delta from baseline – test) using ANOVA (with group as main factor) revealed the absence of significant effects of the test substances at each testing days.
- Morphine (12 mg/kg i.p.), administered 30 minutes before the first test (Day 14), markedly increased the paw-withdrawal latency of the lesioned paw, as compared with vehicle controls (+25.8, $p < 0.001$). It did not affect the non-lesioned paw.
- Gabapentin (100 mg/kg i.p.), administered 180 minutes before the second test (Day 21), did not significantly affect the paw-withdrawal latency of the lesioned or non-lesioned paws, as compared with vehicle controls.

- The apparent increases of some behavioral endpoints following treatment with the test substances was explored by an additional statistical analysis using unpaired Student's t test and the following effects were observed.

Tactile stimulation (Electronic von Frey Test)

- 1015 (30 mg/kg i.v.), administered on Day 14, 3 days before the test 1, significantly increased the force inducing paw-withdrawal of the lesioned paw, as compared with vehicle controls (+5.8, $p < 0.05$).
- B3 (30 mg/kg i.v.), administered on Day 14, 3 days before the test 1, significantly increased the force inducing paw-withdrawal of the lesioned paw, as compared with vehicle controls (+6.4, $p < 0.05$). A similar trend was observed on Test 2 (4 days after the second administration) (+4.6, $p = 0.0815$).

Thermal stimulation (Plantar Test)

- 1015 (30 mg/kg i.v.), administered on Day 14, 3 days before the test 1, tended to increase the paw-withdrawal latency of the lesioned paw, as compared with vehicle controls (+6.0, $p = 0.0896$).
- B3 (30 mg/kg i.v.), administered on Day 14, 3 days before the test 1, significantly increased the paw-withdrawal latency of the lesioned paw, as compared with vehicle controls (+11.2, $p < 0.05$). This effect persisted on Test 2 (4 days after the second administration) (+11.5, $p < 0.05$). At 10 mg/kg, B3 clearly increased the paw-withdrawal latency of the lesioned paw, on test 3 (3 days after the third dosing) (+16.9, $p < 0.01$).

(See Sections [Tables](#), [Figures](#) and [Individual Data](#)).

4. CONCLUSION

These results suggest the absence of significant analgesic effects of 1015 (30 mg/kg) and B3 (10 and 30 mg/kg) administered i.v. at 4 opportunities against tactile allodynia and thermal hyperalgesia in the Neuropathic Pain Test (Chung model) in the rat.

However, some trends towards analgesic effects against tactile allodynia were observed 3 days after the first administration of 1015 and the higher dose of B3. Against thermal hyperalgesia, the higher dose of B3 showed a potential analgesic effect after the first (test on Day 17) and the second administration (test 2 on Day 21); the lower dose seemed more active after the third administration (test 3 on Day 24). This latter effect was confirmed by a complementary statistical analysis using two-way ANOVA.

Gabapentin (100 mg/kg i.p.) displayed analgesic effects against tactile allodynia and morphine (12 mg/kg i.p.) was active against tactile allodynia and thermal hyperalgesia in the same model.

5. TABLES

TABLES**Neuropathic Pain Test (Chung) in the rat****Table 1: Tactile stimulation (electronic Von Frey test) - Baseline on Day 14**

Treatment	Force inducing paw-withdrawal (g) - Baseline on Day 14		
	Lesioned paw		Non-lesioned paw
	(mean $\pm$ s.e.m.)	Compared with non-lesioned paw	(mean $\pm$ s.e.m.)
Vehicle (i.v.)	9.1 $\pm$ 1.0	*** -89%	80.5 $\pm$ 3.2
1015 (30 mg/kg i.v.)	10.0 $\pm$ 0.7	*** -87%	78.7 $\pm$ 2.6
B3 (10 mg/kg i.v.)	10.9 $\pm$ 0.6	*** -86%	76.1 $\pm$ 2.2
B3 (30 mg/kg i.v.)	8.6 $\pm$ 1.3	*** -89%	80.0 $\pm$ 2.9
Morphine (12 mg/kg i.p.) and Gabapentin (100 mg/kg i.p.)	9.2 $\pm$ 1.2	*** -88%	78.1 $\pm$ 2.3

Paired Student's t test: NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

Neuropathic Pain Test (Chung) in the rat**Table 2: Effects of test substances on tactile stimulation (electronic Von Frey test) - lesioned paw**

Treatment	Force inducing paw-withdrawal (g) - delta from baseline - lesioned paw							
	Test 1 on Day 17 (#1) (3 days post dosing)		Test 2 on Day 21 (#2) (4 days post dosing)		Test 3 on Day 24 (#3) (3 days post dosing)		Test 4 on Day 27 (#3) (3 days post dosing)	
	(mean ± s.e.m.)	Compared with Vehicle (i.v.)	(mean ± s.e.m.)	Compared with Vehicle (i.v.)	(mean ± s.e.m.)	Compared with Vehicle (i.v.)	(mean ± s.e.m.)	Compared with Vehicle (i.v.)
		(1) Δ		(1) Δ		(1) Δ		(1) Δ
Vehicle (i.v.)	-1.6 ± 1.2		+0.5 ± 1.4		+3.5 ± 1.9		+1.1 ± 2.3	
1015 (30 mg/kg i.v.)	+4.2 ± 2.1	+5.8	+6.0 ± 3.3	+5.5	+3.6 ± 2.2	+0.1	+7.4 ± 3.2	+6.3
B3 (10 mg/kg i.v.)	+0.8 ± 1.7	NS +2.4	+4.3 ± 2.1	NS +3.8	+6.4 ± 2.7	NS +2.9	+3.8 ± 3.1	NS +2.7
B3 (30 mg/kg i.v.)	+4.8 ± 2.2	+6.4	+5.1 ± 2.0	+4.6	+5.9 ± 1.6	+2.4	+4.6 ± 2.7	+3.5
Morphine (12 mg/kg i.p.) and Gabapentin (100 mg/kg i.p.)	+26.6 ± 3.3	*** +28.2	+15.4 ± 2.8	*** +14.9	+5.5 ± 2.9	NS +2.0	+4.1 ± 1.8	NS +3.0

(#1): for morphine/gabapentin treated group: Test 1 on Day 14 (30 minutes post dosing of morphine (12 mg/kg i.p.)).

(#2): for morphine/gabapentin treated group: Test 2 on Day 21 (180 minutes post dosing of gabapentin (100 mg/kg i.p.)).

(#3): for morphine/gabapentin treated group: No treatment on these testing days.

Inter-group comparison: NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(1): For test substances treated groups: One-way ANOVA with group as factor.

For morphine/gabapentin treated group: Unpaired Student's t test.

(2): Dunnett's t test when One-way ANOVA is significant.

NB: for vehicle (i.v.) control group: comparison of lesioned paw versus non-lesioned paw using paired Student's t test:

Baseline: $p < 0.0001$ *** - Test 1: $p < 0.0001$ *** - Test 2: $p < 0.0001$ *** - Test 3: $p < 0.0001$ *** - Test 4: $p < 0.0001$ ***.

Neuropathic Pain Test (Chung) in the rat**Table 3: Effects of test substances on tactile stimulation (electronic Von Frey test) - non-lesioned paw**

Treatment	Force inducing paw-withdrawal (g) - delta from baseline - non-lesioned paw							
	Test 1 on Day 17 (#1) (3 days post dosing)		Test 2 on Day 21 (#2) (4 days post dosing)		Test 3 on Day 24 (#3) (3 days post dosing)		Test 4 on Day 27 (#3) (3 days post dosing)	
	(mean ± s.e.m.)	Compared with Vehicle (i.v.)	(mean ± s.e.m.)	Compared with Vehicle (i.v.)	(mean ± s.e.m.)	Compared with Vehicle (i.v.)	(mean ± s.e.m.)	Compared with Vehicle (i.v.)
		(1) Δ		(1) Δ		(1) Δ		(1) Δ
Vehicle (i.v.)	-5.0 ± 3.9		-0.3 ± 3.6		-6.5 ± 2.7		-1.7 ± 4.2	
1015 (30 mg/kg i.v.)	-1.1 ± 3.7	+3.9	+2.6 ± 4.0	+2.9	-5.0 ± 5.1	+1.5	-0.2 ± 4.7	+1.5
B3 (10 mg/kg i.v.)	+3.3 ± 4.2	NS +8.3	-5.1 ± 3.6	NS -4.8	+0.5 ± 3.4	NS +7.0	-1.1 ± 3.8	NS +0.6
B3 (30 mg/kg i.v.)	-5.6 ± 3.4	-0.6	-4.3 ± 3.3	-4.0	-3.1 ± 5.3	+3.4	-4.9 ± 3.3	-3.2
Morphine (12 mg/kg i.p.) and Gabapentin (100 mg/kg i.p.)	+0.6 ± 3.7	NS +5.6	-2.4 ± 3.7	NS -2.1	+2.1 ± 4.3	NS +8.6	-1.0 ± 4.8	NS +0.7

(#1): for morphine/gabapentin treated group: Test 1 on Day 14 (30 minutes post dosing of morphine (12 mg/kg i.p.)).

(#2): for morphine/gabapentin treated group: Test 2 on Day 21 (180 minutes post dosing of gabapentin (100 mg/kg i.p.)).

(#3): for morphine/gabapentin treated group: No treatment on these testing days.

Inter-group comparison: NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(1): For test substances treated groups: One-way ANOVA with group as factor.

For morphine/gabapentin treated group: Unpaired Student's t test.

(2): Dunnett's t test when One-way ANOVA is significant.

Neuropathic Pain Test (Chung) in the rat**Table 4: Thermal stimulation (plantar test) - Baseline on Day 14**

Treatment	Paw-withdrawal latency (s) - Baseline on Day 14		
	Lesioned paw		Non-lesioned paw
	(mean $\pm$ s.e.m.)	Compared with non-lesioned paw	(mean $\pm$ s.e.m.)
Vehicle (i.v.)	15.6 $\pm$ 2.3	*** -49%	30.5 $\pm$ 3.4
1015 (30 mg/kg i.v.)	15.0 $\pm$ 2.8	*** -57%	35.0 $\pm$ 2.8
B3 (10 mg/kg i.v.)	20.9 $\pm$ 2.1	*** -43%	36.4 $\pm$ 2.7
B3 (30 mg/kg i.v.)	19.2 $\pm$ 2.9	*** -50%	38.6 $\pm$ 2.3
Morphine (12 mg/kg i.p.) and Gabapentin (100 mg/kg i.p.)	20.0 $\pm$ 3.0	*** -48%	38.7 $\pm$ 2.8

Paired Student's t test: NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

Neuropathic Pain Test (Chung) in the rat**Table 5: Effects of test substances on thermal stimulation (plantar test) - lesioned paw**

Treatment	Paw-withdrawal latency (s) - delta from baseline - lesioned paw							
	Test 1 on Day 17 (#1) (3 days post dosing)		Test 2 on Day 21 (#2) (4 days post dosing)		Test 3 on Day 24 (#3) (3 days post dosing)		Test 4 on Day 27 (#3) (3 days post dosing)	
	(mean ± s.e.m.)	Compared with Vehicle (i.v.)	(mean ± s.e.m.)	Compared with Vehicle (i.v.)	(mean ± s.e.m.)	Compared with Vehicle (i.v.)	(mean ± s.e.m.)	Compared with Vehicle (i.v.)
		(1) Δ		(1) Δ		(1) Δ		(1) Δ
Vehicle (i.v.)	-2.3 ± 2.5		+2.1 ± 3.6		-0.2 ± 4.2		-1.3 ± 2.5	
1015 (30 mg/kg i.v.)	+3.7 ± 2.3	+6.0	+5.4 ± 3.6	+3.3	+9.7 ± 4.3	+9.9	+7.0 ± 6.0	+8.3
B3 (10 mg/kg i.v.)	+6.6 ± 5.1	NS +8.9	+11.7 ± 4.5	NS +9.6	+16.7 ± 2.8	NS +16.9	+6.7 ± 4.8	NS +8.0
B3 (30 mg/kg i.v.)	+8.9 ± 3.6	+11.2	+13.6 ± 4.1	+11.5	+3.3 ± 6.4	+3.5	+0.6 ± 5.3	+1.9
Morphine (12 mg/kg i.p.) and Gabapentin (100 mg/kg i.p.)	+23.5 ± 3.6	*** +25.8	+9.3 ± 4.6	NS +7.2	+3.3 ± 5.7	NS +3.5	+3.5 ± 5.1	NS +4.8

(#1): for morphine/gabapentin treated group: Test 1 on Day 14 (30 minutes post dosing of morphine (12 mg/kg i.p.)).

(#2): for morphine/gabapentin treated group: Test 2 on Day 21 (180 minutes post dosing of gabapentin (100 mg/kg i.p.)).

(#3): for morphine/gabapentin treated group: No treatment on these testing days.

Inter-group comparison: NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(1): For test substances treated groups: One-way ANOVA with group as factor.

For morphine/gabapentin treated group: Unpaired Student's t test.

(2): Dunnett's t test when One-way ANOVA is significant.

NB: for vehicle (i.v.) control group: comparison of lesioned paw versus non-lesioned paw using paired Student's t test:

Baseline: $p = 0.0003$ *** - Test 1: $p = 0.0001$ *** - Test 2: $p = 0.0004$ *** - Test 3: $p = 0.0002$ *** - Test 4: $p = 0.0018$ **.

Neuropathic Pain Test (Chung) in the rat**Table 6: Effects of test substances on thermal stimulation (plantar test) - non-lesioned paw**

Treatment	Paw-withdrawal latency (s) - delta from baseline - non-lesioned paw							
	Test 1 on Day 17 (#1) (3 days post dosing)		Test 2 on Day 21 (#2) (4 days post dosing)		Test 3 on Day 24 (#3) (3 days post dosing)		Test 4 on Day 27 (#3) (3 days post dosing)	
	(mean ± s.e.m.)	Compared with Vehicle (i.v.)	(mean ± s.e.m.)	Compared with Vehicle (i.v.)	(mean ± s.e.m.)	Compared with Vehicle (i.v.)	(mean ± s.e.m.)	Compared with Vehicle (i.v.)
		(1) Δ		(1) Δ		(1) Δ		(1) Δ
Vehicle (i.v.)	+6.2 ± 4.1		+7.6 ± 3.4		+11.8 ± 3.2		-1.9 ± 4.3	
1015 (30 mg/kg i.v.)	+0.3 ± 3.6	-5.9	-2.0 ± 4.5	-9.6	+5.3 ± 4.4	-6.5	-2.2 ± 5.4	-0.3
B3 (10 mg/kg i.v.)	+6.7 ± 3.1	NS +0.5	+4.9 ± 3.5	NS -2.7	+4.4 ± 2.7	NS -7.4	+2.8 ± 3.5	NS +4.7
B3 (30 mg/kg i.v.)	+4.0 ± 3.6	-2.2	+5.0 ± 2.3	-2.6	+6.1 ± 2.3	-5.7	-0.7 ± 4.5	+1.2
Morphine (12 mg/kg i.p.) and Gabapentin (100 mg/kg i.p.)	+6.3 ± 2.8	NS +0.1	+3.6 ± 2.5	NS -4.0	+3.2 ± 3.8	NS -8.6	-1.9 ± 1.9	NS 0.0

(#1): for morphine/gabapentin treated group: Test 1 on Day 14 (30 minutes post dosing of morphine (12 mg/kg i.p.)).

(#2): for morphine/gabapentin treated group: Test 2 on Day 21 (180 minutes post dosing of gabapentin (100 mg/kg i.p.)).

(#3): for morphine/gabapentin treated group: No treatment on these testing days.


Inter-group comparison: NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.


(1): For test substances treated groups: One-way ANOVA with group as factor.


For morphine/gabapentin treated group: Unpaired Student's t test.


(2): Dunnett's t test when One-way ANOVA is significant.

6. FIGURES (produced by the Sponsor)









7. INDIVIDUAL DATA

INDIVIDUAL DATA**Neuropathic Pain Test (Chung) in the rat****Test substance..... : 1015**

Supplier..... : Lpath Inc.

Batch number..... : HR121458

Appearance..... : Colorless solution

Preparation..... : diluted in phosphate buffered saline (PBS)

Route of administration..... : i.v.

Pretreatment time..... : on Day 14, Day 17, Day 21 and Day 24 (i.e. 72 h or 96 h before the test)

Dose expression..... : mg/kg of supplied substance

Volume of administration..... : 3 ml/kg body weight

Test substance..... : B3

Supplier..... : Lpath Inc.

Batch number..... : HR121456

Appearance..... : Colorless solution

Preparation..... : diluted in PBS

Route of administration..... : i.v.

Pretreatment time..... : on Day 14, Day 17, Day 21 and Day 24 (i.e. 72 h or 96 h before the test)

Dose expression..... : mg/kg of supplied substance

Volume of administration..... : 3 ml/kg body weight

Control substance..... : phosphate buffered saline

Supplier..... : Sigma

Reference number..... : P-5368

Batch number..... : SLBH8322V

Route of administration..... : i.v.

Pretreatment time..... : on Day 14, Day 17, Day 21 and Day 24 (i.e. 72 h or 96 h before the test)

Volume of administration..... : 3 ml/kg body weight

Reference substance..... : MORPHINE hydrochloride

Supplier..... : Coopération Pharmaceutique Française

Reference number..... : 1515300

Batch number..... : D3065/2

Appearance..... : white powder

Preparation..... : dissolved in physiological saline

Route of administration..... : i.p.

Pretreatment time..... : 30 min before the test (Day 14)

Dose expression..... : mg/kg of salt

Volume of administration..... : 3 ml/kg body weight

Comparison substance..... : GABAPENTIN

Supplier..... : Toronto Research Chemicals Inc.

Reference number..... : G117250

Batch number..... : 2-YFD-160-1

Appearance..... : white powder

Preparation..... : dissolved in physiological saline

Route of administration..... : i.p.

Pretreatment time..... : 180 min before the test (Day 21)

Dose expression..... : mg/kg of supplied substance

Volume of administration..... : 3 ml/kg body weight

Neuropathic Pain Test (Chung) in the rat

Animals..... : male Sprague Dawley rats (Janvier Labs)

Body weight range..... : 91-138 g (Day -3)

Delivery date..... : 27 June 2014

Number per group..... : 10

Experimental start date..... : 4 July 2014

Experimental termination date..... : 4 August 2014

Technician reference..... : BH9-298

Neuropathic Pain Test (Chung) in the rat**Pre-test on Day -3: Tactile stimulation (electronic Von Frey test)**

To be treated with	Rat number	Weight (g)	Force inducing paw-withdrawal (g)	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
Vehicle	1	125	28.3	26.4
	2	123	40.0	34.7
	3	125	21.2	28.9
	4	123	32.9	20.9
	5	126	21.5	27.9
	29	126	28.3	39.3
	31	113	31.0	24.6
	32	108	26.9	21.2
	33	121	26.1	18.6
	34	106	30.3	29.0
Mean ± s.e.m.		120 2	28.7 1.7	27.2 2.0
t value (a) p value (a)			0.633 0.5427 NS	
Percent change (a)			+6%	

To be treated with	Rat number	Weight (g)	Force inducing paw-withdrawal (g)	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
1015 (30 mg/kg i.v.)	6	125	39.0	31.1
	7	123	26.4	33.0
	8	125	25.9	27.2
	9	124	23.6	38.2
	10	126	27.5	27.4
	35	115	29.5	21.7
	36	104	28.8	31.3
	37	117	26.4	21.4
	38	111	32.3	31.8
	44	120	23.3	32.9
Mean ± s.e.m.		119 2	28.3 1.5	29.6 1.7
t value (a) p value (a)			0.571 0.5817 NS	
Percent change (a)			-4%	

Paired Student's t test (a): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(a): compared with non-lesioned paw.

Neuropathic Pain Test (Chung) in the rat**Pre-test on Day -3: Tactile stimulation (electronic Von Frey test)**

To be treated with	Rat number	Weight (g)	Force inducing paw-withdrawal (g)	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
B3 (10 mg/kg i.v.)	13	111	43.0	31.0
	14	113	38.2	28.3
	15	118	36.2	35.1
	16	113	22.7	26.9
	17	133	30.2	25.9
	40	118	31.9	32.2
	41	134	38.1	24.7
	42	133	25.3	39.4
	55	123	23.0	26.6
	39	102	26.1	32.9
Mean ± s.e.m.		120 3	31.5 2.3	30.3 1.5
t value (a) p value (a)			0.419 0.6854 NS	
Percent change (a)			+4%	

To be treated with	Rat number	Weight (g)	Force inducing paw-withdrawal (g)	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
B3 (30 mg/kg i.v.)	18	123	28.0	32.6
	19	131	24.8	35.3
	20	126	33.4	31.3
	21	122	30.9	43.4
	22	102	23.1	17.8
	45	126	30.4	27.5
	46	125	29.0	26.8
	47	130	23.1	25.1
	48	123	29.4	19.5
	49	116	32.2	24.4
Mean ± s.e.m.		122 3	28.4 1.2	28.4 2.4
t value (a) p value (a)			0.026 0.9802 NS	
Percent change (a)			0%	

Paired Student's t test (a): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(a): compared with non-lesioned paw.

Neuropathic Pain Test (Chung) in the rat**Pre-test on Day -3: Tactile stimulation (electronic Von Frey test)**

To be treated with	Rat number	Weight (g)	Force inducing paw-withdrawal (g)	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
Morphine (12 mg/kg i.p.) and Gabapentin (100 mg/kg i.p.)	23	107	38.2	27.0
	24	91	19.9	25.3
	25	117	25.3	22.2
	27	112	28.5	29.4
	28	111	28.2	18.9
	50	114	28.0	29.8
	51	124	22.4	22.5
	52	138	34.5	38.6
	53	131	39.1	35.8
	54	124	35.0	31.8
Mean ± s.e.m.		117 4	29.9 2.1	28.1 2.0
t value (a)			1.045	
p value (a)			0.3232	
			NS	
Percent change (a)			+6%	

Paired Student's t test (a): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(a): compared with non-lesioned paw.

Neuropathic Pain Test (Chung) in the rat**Pre-test on Day -3: Thermal stimulation (plantar test)**

To be treated with	Rat number	Weight (g)	Paw-withdrawal latency (s)	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
Vehicle	1	125	13.3	10.9
	2	123	7.3	4.2
	3	125	13.5	13.3
	4	123	9.3	12.3
	5	126	26.3	23.5
	29	126	14.4	13.9
	31	113	30.2	45.0
	32	108	45.0	39.0
	33	121	6.8	18.9
	34	106	12.8	7.8
Mean ± s.e.m.		120 2	17.9 3.9	18.9 4.2
t value (a) p value (a)			0.443 0.6681 NS	
Percent change (a)			-5%	

To be treated with	Rat number	Weight (g)	Paw-withdrawal latency (s)	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
1015 (30 mg/kg i.v.)	6	125	13.4	10.4
	7	123	24.4	16.0
	8	125	26.9	42.9
	9	124	25.2	19.8
	10	126	35.7	32.8
	35	115	31.9	28.5
	36	104	14.9	15.3
	37	117	15.8	27.3
	38	111	39.1	15.5
	44	120	9.6	9.2
Mean ± s.e.m.		119 2	23.7 3.2	21.8 3.4
t value (a) p value (a)			0.566 0.5855 NS	
Percent change (a)			+9%	

Paired Student's t test (a): NS = Not Significant; * = p < 0.05; ** = p < 0.01; *** = p < 0.001.

(a): compared with non-lesioned paw.

Neuropathic Pain Test (Chung) in the rat**Pre-test on Day -3: Thermal stimulation (plantar test)**

To be treated with	Rat number	Weight (g)	Paw-withdrawal latency (s)	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
B3 (10 mg/kg i.v.)	13	111	20.2	18.1
	14	113	7.9	17.1
	15	118	45.0	45.0
	16	113	45.0	32.3
	17	133	15.6	13.8
	40	118	28.1	20.6
	41	134	11.4	21.6
	42	133	24.4	23.2
	55	123	21.8	13.6
	39	102	36.1	45.0
Mean ± s.e.m.		120 3	25.6 4.1	25.0 3.7
t value (a) p value (a)			0.209 0.8394 NS	
Percent change (a)			+2%	

To be treated with	Rat number	Weight (g)	Paw-withdrawal latency (s)	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
B3 (30 mg/kg i.v.)	18	123	19.8	9.2
	19	131	38.7	45.0
	20	126	10.6	13.8
	21	122	12.8	25.8
	22	102	45.0	38.3
	45	126	21.0	15.8
	46	125	21.4	35.0
	47	130	12.9	16.4
	48	123	18.1	12.5
	49	116	19.0	25.1
Mean ± s.e.m.		122 3	21.9 3.6	23.7 3.9
t value (a) p value (a)			0.661 0.5252 NS	
Percent change (a)			-8%	

Paired Student's t test (a): NS = Not Significant; * = p < 0.05; ** = p < 0.01; *** = p < 0.001.

(a): compared with non-lesioned paw.

Neuropathic Pain Test (Chung) in the rat**Pre-test on Day -3: Thermal stimulation (plantar test)**

To be treated with	Rat number	Weight (g)	Paw-withdrawal latency (s)	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
Morphine (12 mg/kg i.p.) and Gabapentin (100 mg/kg i.p.)	23	107	15.3	23.7
	24	91	45.0	41.8
	25	117	31.7	14.3
	27	112	29.9	37.1
	28	111	29.4	45.0
	50	114	26.6	45.0
	51	124	15.1	12.1
	52	138	40.6	45.0
	53	131	45.0	45.0
	54	124	14.9	16.9
Mean ± s.e.m.		117 4	29.4 3.7	32.6 4.5
t value (a)			1.000	
p value (a)			0.3434	
			NS	
Percent change (a)			-10%	

Paired Student's t test (a): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(a): compared with non-lesioned paw.

Neuropathic Pain Test (Chung) in the rat**Pre-test on Day -1: Tactile stimulation (electronic Von Frey test)**

To be treated with	Rat number	Weight (g)	Force inducing paw-withdrawal (g)	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
Vehicle	1	157	36.8	35.6
	2	148	35.6	25.9
	3	153	34.2	31.6
	4	156	31.3	34.5
	5	157	48.7	49.7
	29	151	40.6	37.4
	31	137	37.6	46.3
	32	131	30.1	25.1
	33	149	26.1	22.4
	34	125	30.0	25.4
Mean ± s.e.m.		146 4	35.1 2.0	33.4 2.9
t value (a) p value (a)			1.070 0.3127 NS	
Percent change (a)			+5%	

To be treated with	Rat number	Weight (g)	Force inducing paw-withdrawal (g)	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
1015 (30 mg/kg i.v.)	6	160	42.2	40.9
	7	156	36.1	44.4
	8	149	36.3	50.6
	9	152	51.9	43.9
	10	158	39.1	30.2
	35	145	32.8	32.5
	36	124	17.8	33.2
	37	143	42.8	50.1
	38	133	60.1	29.8
	44	146	27.9	36.1
Mean ± s.e.m.		147 4	38.7 3.7	39.2 2.5
t value (a) p value (a)			0.108 0.9161 NS	
Percent change (a)			-1%	

Paired Student's t test (a): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(a): compared with non-lesioned paw.

Neuropathic Pain Test (Chung) in the rat**Pre-test on Day -1: Tactile stimulation (electronic Von Frey test)**

To be treated with	Rat number	Weight (g)	Force inducing paw-withdrawal (g)	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
B3 (10 mg/kg i.v.)	13	132	45.7	33.9
	14	136	36.3	35.5
	15	139	44.4	43.9
	16	135	46.7	37.2
	17	165	31.2	41.0
	40	147	49.4	36.2
	41	166	27.3	33.1
	42	159	31.0	34.7
	55	152	38.6	40.3
	39	127	39.6	26.3
Mean ± s.e.m.		146 4	39.0 2.4	36.2 1.5
t value (a) p value (a)			1.047 0.3226 NS	
Percent change (a)			+8%	

To be treated with	Rat number	Weight (g)	Force inducing paw-withdrawal (g)	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
B3 (30 mg/kg i.v.)	18	151	30.3	30.3
	19	163	44.3	53.3
	20	155	54.9	35.7
	21	145	35.7	52.1
	22	123	25.2	19.2
	45	154	27.9	30.1
	46	146	41.5	38.8
	47	151	40.0	38.6
	48	153	47.4	42.5
	49	143	33.9	51.8
Mean ± s.e.m.		148 3	38.1 2.9	39.2 3.5
t value (a) p value (a)			0.323 0.7539 NS	
Percent change (a)			-3%	

Paired Student's t test (a): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(a): compared with non-lesioned paw.

Neuropathic Pain Test (Chung) in the rat**Pre-test on Day -1: Tactile stimulation (electronic Von Frey test)**

To be treated with	Rat number	Weight (g)	Force inducing paw-withdrawal (g)	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
Morphine (12 mg/kg i.p.) and Gabapentin (100 mg/kg i.p.)	23	133	35.6	45.2
	24	114	34.2	20.2
	25	143	62.7	65.4
	27	134	37.6	33.7
	28	132	18.8	24.7
	50	141	20.1	30.2
	51	147	47.7	45.9
	52	163	46.3	31.9
	53	160	39.6	37.0
	54	152	56.4	39.8
Mean ± s.e.m.		142 5	39.9 4.5	37.4 4.0
t value (a)			0.800	
p value (a)			0.4441	
			NS	
Percent change (a)			+7%	

Paired Student's t test (a): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(a): compared with non-lesioned paw.

Neuropathic Pain Test (Chung) in the rat**Pre-test on Day -1: Thermal stimulation (plantar test)**

To be treated with	Rat number	Weight (g)	Paw-withdrawal latency (s)	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
Vehicle	1	157	45.0	41.8
	2	148	9.8	11.6
	3	153	26.9	29.2
	4	156	23.1	45.0
	5	157	27.5	31.5
	29	151	18.6	22.3
	31	137	19.7	17.6
	32	131	45.0	25.5
	33	149	45.0	17.4
	34	125	28.5	8.4
Mean ± s.e.m.		146 4	28.9 3.9	25.0 3.8
t value (a) p value (a)			0.838 0.4236 NS	
Percent change (a)			+16%	

To be treated with	Rat number	Weight (g)	Paw-withdrawal latency (s)	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
1015 (30 mg/kg i.v.)	6	160	32.3	25.7
	7	156	18.3	23.8
	8	149	18.0	44.5
	9	152	10.8	19.4
	10	158	14.1	25.6
	35	145	27.7	27.8
	36	124	12.2	15.5
	37	143	15.9	16.2
	38	133	31.9	45.0
	44	146	17.2	23.2
Mean ± s.e.m.		147 4	19.8 2.5	26.7 3.3
t value (a) p value (a)			2.391 0.0405 *	
Percent change (a)			-26%	

Paired Student's t test (a): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(a): compared with non-lesioned paw.

Neuropathic Pain Test (Chung) in the rat**Pre-test on Day -1: Thermal stimulation (plantar test)**

To be treated with	Rat number	Weight (g)	Paw-withdrawal latency (s)	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
B3 (10 mg/kg i.v.)	13	132	24.4	19.9
	14	136	12.2	14.8
	15	139	14.5	10.7
	16	135	28.8	23.3
	17	165	45.0	28.0
	40	147	13.8	16.0
	41	166	23.4	18.1
	42	159	18.9	20.5
	55	152	12.8	10.2
	39	127	20.8	18.1
Mean ± s.e.m.		146 4	21.5 3.2	18.0 1.7
t value (a) p value (a)			1.957 0.0820 NS	
Percent change (a)			+19%	

To be treated with	Rat number	Weight (g)	Paw-withdrawal latency (s)	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
B3 (30 mg/kg i.v.)	18	151	38.4	45.0
	19	163	14.0	21.3
	20	155	10.4	14.5
	21	145	18.4	18.8
	22	123	45.0	45.0
	45	154	23.5	32.2
	46	146	26.2	23.1
	47	151	12.0	13.7
	48	153	23.3	36.9
	49	143	20.6	13.7
Mean ± s.e.m.		148 3	23.2 3.5	26.4 3.9
t value (a) p value (a)			1.693 0.1247 NS	
Percent change (a)			-12%	

Paired Student's t test (a): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(a): compared with non-lesioned paw.

Neuropathic Pain Test (Chung) in the rat**Pre-test on Day -1: Thermal stimulation (plantar test)**

To be treated with	Rat number	Weight (g)	Paw-withdrawal latency (s)	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
Morphine (12 mg/kg i.p.) and Gabapentin (100 mg/kg i.p.)	23	133	39.0	28.0
	24	114	38.4	33.9
	25	143	31.0	18.4
	27	134	45.0	40.1
	28	132	45.0	45.0
	50	141	20.4	19.6
	51	147	16.5	21.3
	52	163	19.0	28.9
	53	160	45.0	45.0
	54	152	12.9	20.0
Mean ± s.e.m.		142 5	31.2 4.1	30.0 3.3
t value (a)			0.519	
p value (a)			0.6165	
			NS	
Percent change (a)			+4%	

Paired Student's t test (a): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(a): compared with non-lesioned paw.

Neuropathic Pain Test (Chung) in the rat**Pre-test on Day 11, 12 or 13 (post surgery): Tactile stimulation (electronic Von Frey test)**

To be treated with	Rat number	Weight (g)	Mean of pre-tests 1 et 2	Force inducing paw-withdrawal (g)	
			<i>Lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
Vehicle	1	261	32.6	9.7	73.7
	2	252	37.8	10.9	73.3
	3	284	27.7	9.8	70.9
	4	270	32.1	14.4	73.4
	5	252	35.1	10.7	84.8
	29	269	34.5	11.7	82.9
	31	259	34.3	5.6	74.9
	32	244	28.5	11.4	94.7
	33	263	26.1	10.8	70.6
	34	225	30.2	8.7	73.7
Mean ± s.e.m.		258 5	31.9 1.2	10.4 0.7	77.3 2.4
t value (a) p value (a)				27.740 <0.0001 ***	
Percent change (a)				-87%	

To be treated with	Rat number	Weight (g)	Mean of pre-tests 1 et 2	Force inducing paw-withdrawal (g)	
			<i>Lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
1015 (30 mg/kg i.v.)	6	254	40.6	9.6	94.8
	7	280	31.3	15.4	73.8
	8	265	31.1	7.7	72.5
	9	260	37.8	12.0	94.0
	10	283	33.3	10.1	70.4
	35	276	31.2	10.7	92.1
	36	243	23.3	9.6	79.6
	37	273	34.6	15.0	90.5
	38	251	46.2	2.0	75.3
	44	265	25.6	15.0	93.3
Mean ± s.e.m.		265 4	33.5 2.1	10.7 1.3	83.6 3.2
t value (a) p value (a)				24.684 <0.0001 ***	
Percent change (a)				-87%	

Paired Student's t test (a): NS = Not Significant; * = p < 0.05; ** = p < 0.01; *** = p < 0.001.

(a): compared with non-lesioned paw.

Neuropathic Pain Test (Chung) in the rat**Pre-test on Day 11, 12 or 13 (post surgery): Tactile stimulation (electronic Von Frey test)**

To be treated with	Rat number	Weight (g)	Mean of pre-tests 1 et 2	Force inducing paw-withdrawal (g)	
			<i>Lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
B3 (10 mg/kg i.v.)	13	261	44.4	15.0	82.5
	14	238	37.3	5.1	74.7
	15	238	40.3	12.4	84.7
	16	245	34.7	13.0	74.7
	17	287	30.7	12.0	99.5
	40	267	40.7	11.8	76.8
	41	287	32.7	14.9	92.3
	42	273	28.2	7.5	78.4
	55	270	30.8	6.4	89.3
	39	239	32.9	6.8	67.8
Mean ± s.e.m.		261 6	35.3 1.7	10.5 1.2	82.1 3.0
t value (a) p value (a)				25.854 <0.0001 ***	
Percent change (a)				-87%	

To be treated with	Rat number	Weight (g)	Mean of pre-tests 1 et 2	Force inducing paw-withdrawal (g)	
			<i>Lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
B3 (30 mg/kg i.v.)	18	254	29.2	11.2	70.1
	19	293	34.6	12.1	72.3
	20	283	44.2	14.1	80.6
	21	250	33.3	9.7	86.0
	22	225	24.2	7.9	67.0
	45	273	29.2	14.1	83.4
	46	267	35.3	7.9	73.4
	47	285	31.6	5.2	87.1
	48	274	38.4	12.7	75.1
	49	270	33.1	7.1	79.2
Mean ± s.e.m.		267 6	33.3 1.7	10.2 1.0	77.4 2.2
t value (a) p value (a)				27.513 <0.0001 ***	
Percent change (a)				-87%	

Paired Student's t test (a): NS = Not Significant; * = p < 0.05; ** = p < 0.01; *** = p < 0.001.

(a): compared with non-lesioned paw.

Neuropathic Pain Test (Chung) in the rat**Pre-test on Day 11, 12 or 13 (post surgery): Tactile stimulation (electronic Von Frey test)**

To be treated with	Rat number	Weight (g)	Mean of pre-tests 1 et 2	Force inducing paw-withdrawal (g)	
			<i>Lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
Morphine (12 mg/kg i.p.) and Gabapentin (100 mg/kg i.p.)	23	237	36.9	8.7	66.2
	24	216	27.1	14.1	74.4
	25	265	44.0	9.0	76.3
	27	246	33.1	13.6	79.0
	28	248	23.5	11.9	72.8
	50	261	24.1	8.0	79.4
	51	278	35.1	7.6	71.3
	52	266	40.4	11.5	88.1
	53	274	39.4	6.7	81.1
	54	260	45.7	9.4	82.1
Mean ± s.e.m.		255 6	34.9 2.5	10.1 0.8	77.1 2.0
t value (a) p value (a)				32.418 <0.0001 ***	
Percent change (a)				-87%	

Paired Student's t test (a): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(a): compared with non-lesioned paw.

Neuropathic Pain Test (Chung) in the rat**Baseline on Day 14: Tactile stimulation (electronic Von Frey test)**

To be treated with	Rat number	Weight (g)	Force inducing paw-withdrawal (g)	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
Vehicle	1	266	11.1	69.2
	2	256	14.1	65.5
	3	293	4.4	85.2
	4	273	11.5	86.2
	5	259	12.7	77.6
	29	268	6.0	93.5
	31	261	9.5	77.9
	32	245	7.4	79.6
	33	264	6.7	96.9
	34	224	7.9	73.4
Mean ± s.e.m.		261 6	9.1 1.0	80.5 3.2
t value (a) p value (a)			18.256 <0.0001 ***	
Percent change (a)			-89%	

To be treated with	Rat number	Weight (g)	Force inducing paw-withdrawal (g)	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
1015 (30 mg/kg i.v.)	6	258	11.7	88.2
	7	283	14.9	69.6
	8	261	7.2	85.0
	9	253	8.8	68.7
	10	283	9.1	91.1
	35	280	10.4	77.5
	36	241	8.4	76.6
	37	273	8.5	75.5
	38	249	9.0	85.2
	44	272	11.5	69.3
Mean ± s.e.m.		265 5	10.0 0.7	78.7 2.6
t value (a) p value (a)			23.422 <0.0001 ***	
Percent change (a)			-87%	

Paired Student's t test (a): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(a): compared with non-lesioned paw.

Neuropathic Pain Test (Chung) in the rat**Baseline on Day 14: Tactile stimulation (electronic Von Frey test)**

To be treated with	Rat number	Weight (g)	Force inducing paw-withdrawal (g)	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
B3 (10 mg/kg i.v.)	13	269	11.6	72.6
	14	242	10.0	66.9
	15	239	9.1	83.3
	16	243	13.9	77.6
	17	291	13.4	74.5
	40	266	11.0	91.5
	41	294	10.7	78.7
	42	271	8.7	72.3
	55	270	7.8	73.1
	39	244	12.5	70.3
Mean ± s.e.m.		263 6	10.9 0.6	76.1 2.2
t value (a) p value (a)			28.066 <0.0001 ***	
Percent change (a)			-86%	

To be treated with	Rat number	Weight (g)	Force inducing paw-withdrawal (g)	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
B3 (30 mg/kg i.v.)	18	262	8.7	95.6
	19	297	14.1	68.4
	20	289	4.8	86.7
	21	251	5.2	83.7
	22	227	2.2	64.9
	45	266	10.1	76.0
	46	266	16.1	76.3
	47	282	8.5	87.6
	48	271	6.5	84.2
	49	272	9.7	76.5
Mean ± s.e.m.		268 6	8.6 1.3	80.0 2.9
t value (a) p value (a)			20.955 <0.0001 ***	
Percent change (a)			-89%	

Paired Student's t test (a): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(a): compared with non-lesioned paw.

Neuropathic Pain Test (Chung) in the rat**Baseline on Day 14: Tactile stimulation (electronic Von Frey test)**

To be treated with	Rat number	Weight (g)	Force inducing paw-withdrawal (g)	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
Morphine (12 mg/kg i.p.) and Gabapentin (100 mg/kg i.p.)	23	246	13.3	84.5
	24	216	11.5	78.5
	25	263	9.0	75.7
	27	251	13.3	81.7
	28	242	6.8	71.3
	50	262	5.4	75.3
	51	280	13.5	82.9
	52	266	4.7	76.3
	53	276	4.5	64.2
	54	257	10.0	90.7
Mean ± s.e.m.		256 6	9.2 1.2	78.1 2.3
t value (a) p value (a)			40.101 <0.0001 ***	
Percent change (a)			-88%	

Paired Student's t test (a): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(a): compared with non-lesioned paw.

Neuropathic Pain Test (Chung) in the rat**Baseline on Day 14: Thermal stimulation (plantar test)**

To be treated with	Rat number	Weight (g)	Paw-withdrawal latency (s)	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
Vehicle	1	266	4.8	25.0
	2	256	7.8	30.1
	3	293	10.2	22.4
	4	273	20.6	44.2
	5	259	28.1	40.2
	29	268	19.8	38.4
	31	261	20.5	26.8
	32	245	10.8	15.6
	33	264	19.2	45.0
	34	224	13.8	17.2
Mean ± s.e.m.		261 6	15.6 2.3	30.5 3.4
t value (a) p value (a)			5.714 0.0003 ***	
Percent change (a)			-49%	

To be treated with	Rat number	Weight (g)	Paw-withdrawal latency (s)	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
1015 (30 mg/kg i.v.)	6	258	7.7	33.0
	7	283	16.7	39.2
	8	261	22.2	26.9
	9	253	4.8	19.3
	10	283	22.8	45.0
	35	280	3.5	27.2
	36	241	16.5	32.9
	37	273	4.8	36.8
	38	249	26.2	45.0
	44	272	24.4	45.0
Mean ± s.e.m.		265 5	15.0 2.8	35.0 2.8
t value (a) p value (a)			8.728 <0.0001 ***	
Percent change (a)			-57%	

Paired Student's t test (a): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(a): compared with non-lesioned paw.

Neuropathic Pain Test (Chung) in the rat**Baseline on Day 14: Thermal stimulation (plantar test)**

To be treated with	Rat number	Weight (g)	Paw-withdrawal latency (s)	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
B3 (10 mg/kg i.v.)	13	269	17.4	33.2
	14	242	15.6	29.0
	15	239	12.3	45.0
	16	243	31.2	45.0
	17	291	28.6	38.3
	40	266	22.1	45.0
	41	294	27.8	45.0
	42	271	22.8	34.3
	55	270	14.3	28.4
	39	244	17.0	20.8
Mean ± s.e.m.		263 6	20.9 2.1	36.4 2.7
t value (a) p value (a)			6.270 0.0001 ***	
Percent change (a)			-43%	

To be treated with	Rat number	Weight (g)	Paw-withdrawal latency (s)	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
B3 (30 mg/kg i.v.)	18	262	9.8	29.1
	19	297	7.2	45.0
	20	289	21.9	39.1
	21	251	13.1	45.0
	22	227	15.9	42.1
	45	266	17.7	30.9
	46	266	29.4	45.0
	47	282	36.4	45.0
	48	271	24.9	26.8
	49	272	15.5	37.8
Mean ± s.e.m.		268 6	19.2 2.9	38.6 2.3
t value (a) p value (a)			5.734 0.0003 ***	
Percent change (a)			-50%	

Paired Student's t test (a): NS = Not Significant; * = p < 0.05; ** = p < 0.01; *** = p < 0.001.

(a): compared with non-lesioned paw.

Neuropathic Pain Test (Chung) in the rat**Baseline on Day 14: Thermal stimulation (plantar test)**

To be treated with	Rat number	Weight (g)	Paw-withdrawal latency (s)	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
Morphine (12 mg/kg i.p.) and Gabapentin (100 mg/kg i.p.)	23	246	10.1	38.2
	24	216	4.9	45.0
	25	263	33.1	34.9
	27	251	27.9	45.0
	28	242	15.7	45.0
	50	262	17.3	45.0
	51	280	12.1	17.7
	52	266	22.3	38.1
	53	276	30.6	45.0
	54	257	26.0	32.9
Mean ± s.e.m.		256 6	20.0 3.0	38.7 2.8
t value (a)			4.796	
p value (a)			0.0010 ***	
Percent change (a)			-48%	

Paired Student's t test (a): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(a): compared with non-lesioned paw.

Neuropathic Pain Test (Chung) in the rat**Test 1 on Day 17 (3 days post dosing): Tactile stimulation (electronic Von Frey test)**

Treatment	Rat number	Weight (g)	Force inducing paw-withdrawal (g)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
Vehicle	1	284	8.6	75.3	-2.5	+6.1
	2	275	5.8	73.4	-8.3	+7.9
	3	315	4.7	62.3	+0.3	-22.9
	4	299	6.1	68.6	-5.4	-17.6
	5	282	10.5	90.4	-2.2	+12.8
	29	291	10.3	82.4	+4.3	-11.1
	31	283	10.7	76.2	+1.2	-1.7
	32	263	8.6	82.6	+1.2	+3.0
	33	280	6.1	81.4	-0.6	-15.5
	34	243	4.4	62.4	-3.5	-11.0
Mean ± s.e.m.		282 6	7.6 0.8	75.5 2.9	-1.6 1.2	-5.0 3.9
t value (a) p value (a)			28.986 <0.0001 ***			
Percent change (a)			-90%			

Treatment	Rat number	Weight (g)	Force inducing paw-withdrawal (g)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
1015 (30 mg/kg i.v.)	6	276	7.5	95.9	-4.2	+7.7
	7	308	13.0	70.7	-1.9	+1.1
	8	293	10.1	89.9	+2.9	+4.9
	9	283	11.5	77.7	+2.7	+9.0
	10	298	19.9	100.0	+10.8	+8.9
	35	312	12.2	62.0	+1.8	-15.5
	36	270	11.8	71.0	+3.4	-5.6
	37	302	23.7	84.5	+15.2	+9.0
	38	274	6.7	60.5	-2.3	-24.7
	44	294	24.6	63.9	+13.1	-5.4
Mean ± s.e.m.		291 5	14.1 2.0	77.6 4.5	+4.2 2.1	-1.1 3.7
t value (b) p value (b)					2.364 0.0295 *	0.737 0.4704 NS
Mean change (b)					+5.8	+3.9

Paired Student's t test (a): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

Unpaired Student's t test (b): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(a): compared with non-lesioned paw.

(b): compared with vehicle.

Neuropathic Pain Test (Chung) in the rat**Test 1 on Day 17 (3 days post dosing): Tactile stimulation (electronic Von Frey test)**

Treatment	Rat number	Weight (g)	Force inducing paw-withdrawal (g)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
B3 (10 mg/kg i.v.)	13	292	5.1	66.8	-6.5	-5.8
	14	262	16.8	90.8	+6.8	+23.9
	15	259	18.1	92.9	+9.0	+9.6
	16	268	14.8	84.0	+0.9	+6.4
	17	320	7.7	63.1	-5.7	-11.4
	40	294	16.0	72.4	+5.0	-19.1
	41	317	8.2	93.5	-2.5	+14.8
	42	294	7.7	72.1	-1.0	-0.2
	55	297	11.9	87.7	+4.1	+14.6
	39	269	10.1	70.6	-2.4	+0.3
Mean ± s.e.m.		287 7	11.6 1.4	79.4 3.7	+0.8 1.7	+3.3 4.2
t value (b) p value (b)					1.143 0.2681 NS	1.464 0.1604 NS
Mean change (b)					+2.4	+8.3

Treatment	Rat number	Weight (g)	Force inducing paw-withdrawal (g)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
B3 (30 mg/kg i.v.)	18	280	11.7	73.1	+3.0	-22.5
	19	326	11.1	54.4	-3.0	-14.0
	20	317	9.1	85.4	+4.3	-1.3
	21	275	17.2	64.6	+12.0	-19.1
	22	257	19.7	68.1	+17.5	+3.2
	45	285	11.1	87.2	+1.0	+11.2
	46	286	19.9	77.7	+3.8	+1.4
	47	308	16.1	77.5	+7.6	-10.1
	48	297	14.6	83.0	+8.1	-1.2
	49	287	3.7	73.2	-6.0	-3.3
Mean ± s.e.m.		292 7	13.4 1.6	74.4 3.2	+4.8 2.2	-5.6 3.4
t value (b) p value (b)					2.587 0.0186 *	0.111 0.9125 NS
Mean change (b)					+6.4	-0.6

Unpaired Student's t test (b): NS = Not Significant; * = p < 0.05; ** = p < 0.01; *** = p < 0.001.

(b): compared with vehicle.

Neuropathic Pain Test (Chung) in the rat**Test 1 on Day 14 (30 minutes post dosing): Tactile stimulation (electronic Von Frey test)**

Treatment	Rat number	Weight (g)	Force inducing paw-withdrawal (g)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
Morphine (12 mg/kg i.p.)	23	246	25.5	71.3	+12.2	-13.2
	24	216	31.4	72.6	+19.9	-5.9
	25	263	47.5	71.4	+38.5	-4.3
	27	251	30.6	89.6	+17.3	+7.9
	28	242	19.2	73.9	+12.4	+2.6
	50	262	32.9	84.5	+27.5	+9.2
	51	280	52.9	79.8	+39.4	-3.1
	52	266	37.0	69.7	+32.3	-6.6
	53	276	37.5	91.1	+33.0	+26.9
	54	257	43.2	82.8	+33.2	-7.9
Mean ± s.e.m.		256 6	35.8 3.2	78.7 2.5	+26.6 3.3	+0.6 3.7
t value (b) p value (b)					8.107 <0.0001 ***	1.042 0.3111 NS
Mean change (b)					+28.2	+5.6

Unpaired Student's t test (b): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(b): compared with vehicle.

Neuropathic Pain Test (Chung) in the rat**Test 1 on Day 17 (3 days post dosing): Thermal stimulation (plantar test)**

Treatment	Rat number	Weight (g)	Paw-withdrawal latency (s)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
Vehicle	1	284	4.3	45.0	-0.5	+20.0
	2	275	16.8	40.5	+9.0	+10.4
	3	315	14.5	17.6	+4.3	-4.8
	4	299	21.8	45.0	+1.2	+0.8
	5	282	10.2	27.6	-17.9	-12.6
	29	291	14.1	45.0	-5.7	+6.6
	31	283	19.4	27.0	-1.1	+0.2
	32	263	15.2	45.0	+4.4	+29.4
	33	280	9.8	40.8	-9.4	-4.2
	34	243	6.1	33.3	-7.7	+16.1
Mean ± s.e.m.		282 6	13.2 1.8	36.7 3.1	-2.3 2.5	+6.2 4.1
t value (a) p value (a)			6.516 0.0001 ***			
Percent change (a)			-64%			

Treatment	Rat number	Weight (g)	Paw-withdrawal latency (s)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
1015 (30 mg/kg i.v.)	6	276	14.1	45.0	+6.4	+12.0
	7	308	8.2	17.0	-8.5	-22.2
	8	293	24.3	33.5	+2.1	+6.6
	9	283	4.8	25.8	0.0	+6.5
	10	298	15.7	45.0	-7.1	0.0
	35	312	13.3	39.4	+9.8	+12.2
	36	270	26.3	28.0	+9.8	-4.9
	37	302	9.5	45.0	+4.7	+8.2
	38	274	39.0	45.0	+12.8	0.0
	44	294	31.6	29.6	+7.2	-15.4
Mean ± s.e.m.		291 5	18.7 3.5	35.3 3.2	+3.7 2.3	+0.3 3.6
t value (b) p value (b)					1.794 0.0896 NS	1.077 0.2957 NS
Mean change (b)					+6.0	-5.9

Paired Student's t test (a): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.**Unpaired Student's t test (b):** NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(a): compared with non-lesioned paw.

(b): compared with vehicle.

Neuropathic Pain Test (Chung) in the rat**Test 1 on Day 17 (3 days post dosing): Thermal stimulation (plantar test)**

Treatment	Rat number	Weight (g)	Paw-withdrawal latency (s)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
B3 (10 mg/kg i.v.)	13	292	37.7	38.5	+20.3	+5.3
	14	262	22.1	45.0	+6.5	+16.0
	15	259	45.0	45.0	+32.7	0.0
	16	268	45.0	45.0	+13.8	0.0
	17	320	37.8	32.4	+9.2	-5.9
	40	294	5.3	45.0	-16.8	0.0
	41	317	6.4	45.0	-21.4	0.0
	42	294	31.6	45.0	+8.8	+10.7
	55	297	16.3	45.0	+2.0	+16.6
	39	269	27.9	45.0	+10.9	+24.2
Mean ± s.e.m.		287 7	27.5 4.6	43.1 1.4	+6.6 5.1	+6.7 3.1
t value (b) p value (b)					1.583 0.1309 NS	0.098 0.9230 NS
Mean change (b)					+8.9	+0.5

Treatment	Rat number	Weight (g)	Paw-withdrawal latency (s)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
B3 (30 mg/kg i.v.)	18	280	26.3	45.0	+16.5	+15.9
	19	326	17.1	23.4	+9.9	-21.6
	20	317	30.8	45.0	+8.9	+5.9
	21	275	26.9	42.8	+13.8	-2.2
	22	257	40.6	45.0	+24.7	+2.9
	45	285	22.3	45.0	+4.6	+14.1
	46	286	44.8	45.0	+15.4	0.0
	47	308	17.9	45.0	-18.5	0.0
	48	297	31.5	45.0	+6.6	+18.2
	49	287	22.4	45.0	+6.9	+7.2
Mean ± s.e.m.		292 7	28.1 2.9	42.6 2.1	+8.9 3.6	+4.0 3.6
t value (b) p value (b)					2.566 0.0194 *	0.394 0.6981 NS
Mean change (b)					+11.2	-2.2

Unpaired Student's t test (b): NS = Not Significant; * = p < 0.05; ** = p < 0.01; *** = p < 0.001.

(b): compared with vehicle.

Neuropathic Pain Test (Chung) in the rat**Test 1 on Day 14 (30 minutes post dosing): Thermal stimulation (plantar test)**

Treatment	Rat number	Weight (g)	Paw-withdrawal latency (s)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
Morphine (12 mg/kg i.p.)	23	246	45.0	45.0	+34.9	+6.8
	24	216	45.0	45.0	+40.1	0.0
	25	263	45.0	45.0	+11.9	+10.1
	27	251	45.0	45.0	+17.1	0.0
	28	242	45.0	45.0	+29.3	0.0
	50	262	45.0	45.0	+27.7	0.0
	51	280	45.0	45.0	+32.9	+27.3
	52	266	45.0	45.0	+22.7	+6.9
	53	276	45.0	45.0	+14.4	0.0
	54	257	29.7	45.0	+3.7	+12.1
Mean ± s.e.m.		256 6	43.5 1.5	45.0 0.0	+23.5 3.6	+6.3 2.8
t value (b) p value (b)					5.830 <0.0001 ***	0.026 0.9792 NS
Mean change (b)					+25.8	+0.1

Unpaired Student's t test (b): NS = Not Significant; * = p < 0.05; ** = p < 0.01; *** = p < 0.001.

(b): compared with vehicle.

Neuropathic Pain Test (Chung) in the rat**Test 2 on Day 21 (4 days post dosing): Tactile stimulation (electronic Von Frey test)**

Treatment	Rat number	Weight (g)	Force inducing paw-withdrawal (g)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
Vehicle	1	311	13.9	79.0	+2.8	+9.8
	2	305	12.2	74.2	-1.9	+8.7
	3	354	7.2	76.5	+2.8	-8.7
	4	317	4.9	66.8	-6.6	-19.4
	5	317	11.1	96.0	-1.6	+18.4
	29	330	14.5	81.1	+8.5	-12.4
	31	316	9.5	81.0	0.0	+3.1
	32	292	5.7	77.0	-1.7	-2.6
	33	310	12.4	95.4	+5.7	-1.5
	34	255	5.2	75.0	-2.7	+1.6
Mean ± s.e.m.		311 8	9.7 1.2	80.2 2.9	+0.5 1.4	-0.3 3.6
t value (a) p value (a)			28.336 <0.0001 ***			
Percent change (a)			-88%			

Treatment	Rat number	Weight (g)	Force inducing paw-withdrawal (g)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
1015 (30 mg/kg i.v.)	6	296	7.0	98.4	-4.7	+10.2
	7	333	9.9	72.9	-5.0	+3.3
	8	321	10.9	73.6	+3.7	-11.4
	9	307	12.4	66.8	+3.6	-1.9
	10	325	23.9	82.5	+14.8	-8.6
	35	345	22.7	67.9	+12.3	-9.6
	36	300	7.4	87.6	-1.0	+11.0
	37	320	34.6	100.0	+26.1	+24.5
	38	304	4.6	77.1	-4.4	-8.1
	44	328	26.5	86.3	+15.0	+17.0
Mean ± s.e.m.		318 5	16.0 3.2	81.3 3.7	+6.0 3.3	+2.6 4.0
t value (b) p value (b)					1.520 0.1460 NS	0.552 0.5880 NS
Mean change (b)					+5.5	+2.9

Paired Student's t test (a): NS = Not Significant; * = p < 0.05; ** = p < 0.01; *** = p < 0.001.

Unpaired Student's t test (b): NS = Not Significant; * = p < 0.05; ** = p < 0.01; *** = p < 0.001.

(a): compared with non-lesioned paw.

(b): compared with vehicle.

Neuropathic Pain Test (Chung) in the rat**Test 2 on Day 21 (4 days post dosing): Tactile stimulation (electronic Von Frey test)**

Treatment	Rat number	Weight (g)	Force inducing paw-withdrawal (g)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
B3 (10 mg/kg i.v.)	13	330	7.5	82.6	-4.1	+10.0
	14	282	12.2	68.6	+2.2	+1.7
	15	287	13.3	70.6	+4.2	-12.7
	16	296	9.2	67.9	-4.7	-9.7
	17	356	9.5	56.9	-3.9	-17.6
	40	327	20.8	72.9	+9.8	-18.6
	41	352	17.9	61.3	+7.2	-17.4
	42	321	22.1	75.0	+13.4	+2.7
	55	327	17.3	73.8	+9.5	+0.7
	39	293	21.7	80.0	+9.2	+9.7
Mean ± s.e.m.		317 8	15.2 1.7	71.0 2.5	+4.3 2.1	-5.1 3.6
t value (b) p value (b)					1.486 0.1546 NS	0.955 0.3523 NS
Mean change (b)					+3.8	-4.8

Treatment	Rat number	Weight (g)	Force inducing paw-withdrawal (g)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
B3 (30 mg/kg i.v.)	18	310	12.2	89.7	+3.5	-5.9
	19	349	12.8	81.3	-1.3	+12.9
	20	354	10.9	85.8	+6.1	-0.9
	21	292	20.7	67.0	+15.5	-16.7
	22	283	7.8	61.7	+5.6	-3.2
	45	312	25.8	60.5	+15.7	-15.5
	46	310	23.7	85.7	+7.6	+9.4
	47	340	9.7	77.1	+1.2	-10.5
	48	330	6.8	68.2	+0.3	-16.0
	49	320	6.6	79.6	-3.1	+3.1
Mean ± s.e.m.		320 7	13.7 2.2	75.7 3.3	+5.1 2.0	-4.3 3.3
t value (b) p value (b)					1.845 0.0815 NS	0.826 0.4198 NS
Mean change (b)					+4.6	-4.0

Unpaired Student's t test (b): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(b): compared with vehicle.

Neuropathic Pain Test (Chung) in the rat**Test 2 on Day 21 (180 minutes post dosing): Tactile stimulation (electronic Von Frey test)**

Treatment	Rat number	Weight (g)	Force inducing paw-withdrawal (g)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
Gabapentin (100 mg/kg i.p.)	23	294	24.2	71.7	+10.9	-12.8
	24	271	20.3	77.1	+8.8	-1.4
	25	307	34.8	74.7	+25.8	-1.0
	27	305	31.9	86.9	+18.6	+5.2
	28	294	17.7	72.3	+10.9	+1.0
	50	321	38.1	74.7	+32.7	-0.6
	51	347	28.8	69.1	+15.3	-13.8
	52	305	23.2	81.5	+18.5	+5.2
	53	332	12.4	81.7	+7.9	+17.5
	54	303	14.4	67.3	+4.4	-23.4
Mean ± s.e.m.		308 7	24.6 2.7	75.7 1.9	+15.4 2.8	-2.4 3.7
t value (b) p value (b)					4.787 0.0001 ***	0.413 0.6842 NS
Mean change (b)					+14.9	-2.1

Unpaired Student's t test (b): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(b): compared with vehicle.

Neuropathic Pain Test (Chung) in the rat**Test 2 on Day 21 (4 days post dosing): Thermal stimulation (plantar test)**

Treatment	Rat number	Weight (g)	Paw-withdrawal latency (s)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
Vehicle	1	311	5.1	24.7	+0.3	-0.3
	2	305	13.1	33.5	+5.3	+3.4
	3	354	13.2	35.7	+3.0	+13.3
	4	317	6.1	45.0	-14.5	+0.8
	5	317	45.0	45.0	+16.9	+4.8
	29	330	18.9	33.0	-0.9	-5.4
	31	316	19.8	34.8	-0.7	+8.0
	32	292	32.5	45.0	+21.7	+29.4
	33	310	5.8	45.0	-13.4	0.0
	34	255	17.5	38.9	+3.7	+21.7
Mean ± s.e.m.		311 8	17.7 4.0	38.1 2.2	+2.1 3.6	+7.6 3.4
t value (a) p value (a)			5.465 0.0004 ***			
Percent change (a)			-54%			

Treatment	Rat number	Weight (g)	Paw-withdrawal latency (s)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
1015 (30 mg/kg i.v.)	6	296	10.6	13.7	+2.9	-19.3
	7	333	3.2	39.4	-13.5	+0.2
	8	321	10.0	14.6	-12.2	-12.3
	9	307	7.2	45.0	+2.4	+25.7
	10	325	39.8	45.0	+17.0	0.0
	35	345	11.9	45.0	+8.4	+17.8
	36	300	24.3	19.0	+7.8	-13.9
	37	320	20.2	27.1	+15.4	-9.7
	38	304	31.6	36.9	+5.4	-8.1
	44	328	45.0	45.0	+20.6	0.0
Mean ± s.e.m.		318 5	20.4 4.6	33.1 4.2	+5.4 3.6	-2.0 4.5
t value (b) p value (b)					0.647 0.5260 NS	1.684 0.1095 NS
Mean change (b)					+3.3	-9.6

Paired Student's t test (a): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

Unpaired Student's t test (b): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(a): compared with non-lesioned paw.

(b): compared with vehicle.

Neuropathic Pain Test (Chung) in the rat**Test 2 on Day 21 (4 days post dosing): Thermal stimulation (plantar test)**

Treatment	Rat number	Weight (g)	Paw-withdrawal latency (s)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
B3 (10 mg/kg i.v.)	13	330	45.0	45.0	+27.6	+11.8
	14	282	21.4	45.0	+5.8	+16.0
	15	287	37.5	45.0	+25.2	0.0
	16	296	45.0	45.0	+13.8	0.0
	17	356	30.9	44.4	+2.3	+6.1
	40	327	40.8	45.0	+18.7	0.0
	41	352	8.3	23.8	-19.5	-21.2
	42	321	45.0	45.0	+22.2	+10.7
	55	327	31.6	45.0	+17.3	+16.6
	39	293	20.6	29.5	+3.6	+8.7
Mean ± s.e.m.		317 8	32.6 4.0	41.3 2.5	+11.7 4.5	+4.9 3.5
t value (b) p value (b)					1.671 0.1120 NS	0.549 0.5899 NS
Mean change (b)					+9.6	-2.7

Treatment	Rat number	Weight (g)	Paw-withdrawal latency (s)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
B3 (30 mg/kg i.v.)	18	310	22.1	45.0	+12.3	+15.9
	19	349	40.3	45.0	+33.1	0.0
	20	354	37.5	35.0	+15.6	-4.1
	21	292	37.2	45.0	+24.1	0.0
	22	283	43.1	45.0	+27.2	+2.9
	45	312	22.4	45.0	+4.7	+14.1
	46	310	45.0	45.0	+15.6	0.0
	47	340	40.5	45.0	+4.1	0.0
	48	330	35.2	40.7	+10.3	+13.9
	49	320	4.2	45.0	-11.3	+7.2
Mean ± s.e.m.		320 7	32.8 4.0	43.6 1.0	+13.6 4.1	+5.0 2.3
t value (b) p value (b)					2.111 0.0490 *	0.624 0.5407 NS
Mean change (b)					+11.5	-2.6

Unpaired Student's t test (b): NS = Not Significant; * = p < 0.05; ** = p < 0.01; *** = p < 0.001.

(b): compared with vehicle.

Neuropathic Pain Test (Chung) in the rat**Test 2 on Day 21 (180 minutes post dosing): Thermal stimulation (plantar test)**

Treatment	Rat number	Weight (g)	Paw-withdrawal latency (s)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
Gabapentin (100 mg/kg i.p.)	23	294	45.0	45.0	+34.9	+6.8
	24	271	33.1	45.0	+28.2	0.0
	25	307	22.8	45.0	-10.3	+10.1
	27	305	45.0	45.0	+17.1	0.0
	28	294	29.7	45.0	+14.0	0.0
	50	321	24.0	45.0	+6.7	0.0
	51	347	17.8	38.6	+5.7	+20.9
	52	305	21.9	45.0	-0.4	+6.9
	53	332	37.0	45.0	+6.4	0.0
	54	303	16.8	24.5	-9.2	-8.4
Mean ± s.e.m.		308 7	29.3 3.3	42.3 2.1	+9.3 4.6	+3.6 2.5
t value (b) p value (b)					1.222 0.2374 NS	0.923 0.3681 NS
Mean change (b)					+7.2	-4.0

Unpaired Student's t test (b): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(b): compared with vehicle.

Neuropathic Pain Test (Chung) in the rat**Test 3 on Day 24 (3 days post dosing): Tactile stimulation (electronic Von Frey test)**

Treatment	Rat number	Weight (g)	Force inducing paw-withdrawal (g)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
Vehicle	1	328	16.9	73.2	+5.8	+4.0
	2	325	6.1	52.8	-8.0	-12.7
	3	388	6.9	81.2	+2.5	-4.0
	4	344	22.1	76.7	+10.6	-9.5
	5	341	9.8	73.9	-2.9	-3.7
	29	351	14.4	76.0	+8.4	-17.5
	31	334	21.9	85.1	+12.4	+7.2
	32	313	11.0	80.7	+3.6	+1.1
	33	332	8.3	82.6	+1.6	-14.3
	34	286	9.1	58.1	+1.2	-15.3
Mean ± s.e.m.		334 8	12.7 1.9	74.0 3.3	+3.5 1.9	-6.5 2.7
t value (a) p value (a)			19.922 <0.0001 ***			
Percent change (a)			-83%			

Treatment	Rat number	Weight (g)	Force inducing paw-withdrawal (g)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
1015 (30 mg/kg i.v.)	6	313	3.8	59.2	-7.9	-29.0
	7	364	13.5	86.6	-1.4	+17.0
	8	346	4.1	78.6	-3.1	-6.4
	9	343	10.8	58.5	+2.0	-10.2
	10	342	24.7	63.0	+15.6	-28.1
	35	378	18.7	67.8	+8.3	-9.7
	36	329	15.8	70.7	+7.4	-5.9
	37	357	15.3	94.5	+6.8	+19.0
	38	327	17.6	82.4	+8.6	-2.8
	44	357	11.4	75.4	-0.1	+6.1
Mean ± s.e.m.		346 6	13.6 2.0	73.7 3.8	+3.6 2.2	-5.0 5.1
t value (b) p value (b)					0.034 0.9732 NS	0.254 0.8027 NS
Mean change (b)					+0.1	+1.5

Paired Student's t test (a): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

Unpaired Student's t test (b): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(a): compared with non-lesioned paw.

(b): compared with vehicle.

Neuropathic Pain Test (Chung) in the rat**Test 3 on Day 24 (3 days post dosing): Tactile stimulation (electronic Von Frey test)**

Treatment	Rat number	Weight (g)	Force inducing paw-withdrawal (g)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
B3 (10 mg/kg i.v.)	13	353	19.0	62.1	+7.4	-10.5
	14	304	8.7	77.5	-1.3	+10.6
	15	306	20.2	78.7	+11.1	-4.6
	16	324	7.7	76.6	-6.2	-1.0
	17	379	21.8	63.9	+8.4	-10.6
	40	347	30.1	75.5	+19.1	-16.0
	41	377	30.7	82.0	+20.0	+3.3
	42	343	10.3	86.3	+1.6	+14.0
	55	357	8.8	81.5	+1.0	+8.4
	39	318	15.0	81.8	+2.5	+11.5
Mean ± s.e.m.		341 9	17.2 2.7	76.6 2.5	+6.4 2.7	+0.5 3.4
t value (b) p value (b)					0.852 0.4056 NS	1.608 0.1253 NS
Mean change (b)					+2.9	+7.0

Treatment	Rat number	Weight (g)	Force inducing paw-withdrawal (g)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
B3 (30 mg/kg i.v.)	18	335	13.9	74.5	+5.2	-21.1
	19	375	14.8	82.2	+0.7	+13.8
	20	380	10.4	64.9	+5.6	-21.8
	21	308	22.4	80.7	+17.2	-3.0
	22	301	12.9	91.1	+10.7	+26.2
	45	334	15.8	76.3	+5.7	+0.3
	46	332	21.1	88.6	+5.0	+12.3
	47	367	14.5	66.8	+6.0	-20.8
	48	354	9.8	69.4	+3.3	-14.8
	49	350	9.7	74.6	0.0	-1.9
Mean ± s.e.m.		344 8	14.5 1.4	76.9 2.8	+5.9 1.6	-3.1 5.3
t value (b) p value (b)					0.968 0.3460 NS	0.569 0.5764 NS
Mean change (b)					+2.4	+3.4

Unpaired Student's t test (b): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(b): compared with vehicle.

Neuropathic Pain Test (Chung) in the rat**Test 3 on Day 24 (3 days post dosing): Tactile stimulation (electronic Von Frey test)**

Treatment	Rat number	Weight (g)	Force inducing paw-withdrawal (g)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
Morphine (12 mg/kg i.p.) and Gabapentin (100 mg/kg i.p.)	23	310	15.6	74.0	+2.3	-10.5
	24	298	11.3	87.1	-0.2	+8.6
	25	328	5.4	71.1	-3.6	-4.6
	27	327	8.2	73.9	-5.1	-7.8
	28	303	19.0	75.6	+12.2	+4.3
	50	347	32.0	85.6	+26.6	+10.3
	51	370	21.8	64.9	+8.3	-18.0
	52	324	13.2	76.5	+8.5	+0.2
	53	364	7.0	94.8	+2.5	+30.6
	54	324	13.3	99.0	+3.3	+8.3
Mean ± s.e.m.		330 8	14.7 2.5	80.3 3.5	+5.5 2.9	+2.1 4.3
t value (b) p value (b)					0.560 0.5821 NS	1.685 0.1093 NS
Mean change (b)					+2.0	+8.6

Unpaired Student's t test (b): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(b): compared with vehicle.

Neuropathic Pain Test (Chung) in the rat**Test 3 on Day 24 (3 days post dosing): Thermal stimulation (plantar test)**

Treatment	Rat number	Weight (g)	Paw-withdrawal latency (s)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
Vehicle	1	328	7.1	44.2	+2.3	+19.2
	2	325	38.7	45.0	+30.9	+14.9
	3	388	8.0	45.0	-2.2	+22.6
	4	344	6.6	45.0	-14.0	+0.8
	5	341	38.2	41.1	+10.1	+0.9
	29	351	6.0	45.0	-13.8	+6.6
	31	334	17.5	36.5	-3.0	+9.7
	32	313	5.3	45.0	-5.5	+29.4
	33	332	22.4	45.0	+3.2	0.0
	34	286	3.5	31.2	-10.3	+14.0
Mean ± s.e.m.		334 8	15.3 4.3	42.3 1.5	-0.2 4.2	+11.8 3.2
t value (a) p value (a)			6.151 0.0002 ***			
Percent change (a)			-64%			

Treatment	Rat number	Weight (g)	Paw-withdrawal latency (s)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
1015 (30 mg/kg i.v.)	6	313	24.8	39.2	+17.1	+6.2
	7	364	35.6	45.0	+18.9	+5.8
	8	346	7.4	45.0	-14.8	+18.1
	9	343	34.9	45.0	+30.1	+25.7
	10	342	41.2	38.5	+18.4	-6.5
	35	378	11.6	45.0	+8.1	+17.8
	36	329	28.2	45.0	+11.7	+12.1
	37	357	3.9	37.3	-0.9	+0.5
	38	327	42.0	37.9	+15.8	-7.1
	44	357	16.6	25.3	-7.8	-19.7
Mean ± s.e.m.		346 6	24.6 4.4	40.3 2.0	+9.7 4.3	+5.3 4.4
t value (b) p value (b)					1.633 0.1199 NS	1.206 0.2434 NS
Mean change (b)					+9.9	-6.5

Paired Student's t test (a): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

Unpaired Student's t test (b): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(a): compared with non-lesioned paw.

(b): compared with vehicle.

Neuropathic Pain Test (Chung) in the rat**Test 3 on Day 24 (3 days post dosing): Thermal stimulation (plantar test)**

Treatment	Rat number	Weight (g)	Paw-withdrawal latency (s)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
B3 (10 mg/kg i.v.)	13	353	33.2	45.0	+15.8	+11.8
	14	304	23.3	29.1	+7.7	+0.1
	15	306	32.0	31.8	+19.7	-13.2
	16	324	45.0	45.0	+13.8	0.0
	17	379	35.6	45.0	+7.0	+6.7
	40	347	45.0	45.0	+22.9	0.0
	41	377	45.0	45.0	+17.2	0.0
	42	343	26.8	45.0	+4.0	+10.7
	55	357	45.0	44.6	+30.7	+16.2
	39	318	45.0	32.5	+28.0	+11.7
Mean ± s.e.m.		341 9	37.6 2.7	40.8 2.1	+16.7 2.8	+4.4 2.7
t value (b) p value (b)					3.327 0.0038 **	1.766 0.0944 NS
Mean change (b)					+16.9	-7.4

Treatment	Rat number	Weight (g)	Paw-withdrawal latency (s)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
B3 (30 mg/kg i.v.)	18	335	27.8	44.4	+18.0	+15.3
	19	375	45.0	45.0	+37.8	0.0
	20	380	21.3	45.0	-0.6	+5.9
	21	308	15.5	45.0	+2.4	0.0
	22	301	14.5	45.0	-1.4	+2.9
	45	334	45.0	45.0	+27.3	+14.1
	46	332	37.0	45.0	+7.6	0.0
	47	367	8.8	45.0	-27.6	0.0
	48	354	5.0	45.0	-19.9	+18.2
	49	350	5.2	42.4	-10.3	+4.6
Mean ± s.e.m.		344 8	22.5 4.9	44.7 0.3	+3.3 6.4	+6.1 2.3
t value (b) p value (b)					0.463 0.6491 NS	1.468 0.1595 NS
Mean change (b)					+3.5	-5.7

Unpaired Student's t test (b): NS = Not Significant; * = p < 0.05; ** = p < 0.01; *** = p < 0.001.

(b): compared with vehicle.

Neuropathic Pain Test (Chung) in the rat**Test 3 on Day 24 (3 days post dosing): Thermal stimulation (plantar test)**

Treatment	Rat number	Weight (g)	Paw-withdrawal latency (s)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
Morphine (12 mg/kg i.p.) and Gabapentin (100 mg/kg i.p.)	23	310	38.2	45.0	+28.1	+6.8
	24	298	18.0	45.0	+13.1	0.0
	25	328	25.9	43.2	-7.2	+8.3
	27	327	5.7	42.0	-22.2	-3.0
	28	303	45.0	30.8	+29.3	-14.2
	50	347	30.1	45.0	+12.8	0.0
	51	370	12.6	45.0	+0.5	+27.3
	52	324	5.6	45.0	-16.7	+6.9
	53	364	37.8	32.8	+7.2	-12.2
	54	324	13.8	45.0	-12.2	+12.1
Mean ± s.e.m.		330 8	23.3 4.5	41.9 1.7	+3.3 5.7	+3.2 3.8
t value (b)					0.494	1.735
p value (b)					0.6270	0.0999
					NS	NS
Mean change (b)					+3.5	-8.6

Unpaired Student's t test (b): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(b): compared with vehicle.

Neuropathic Pain Test (Chung) in the rat**Test 4 on Day 27 (3 days post dosing): Tactile stimulation (electronic Von Frey test)**

Treatment	Rat number	Weight (g)	Force inducing paw-withdrawal (g)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
Vehicle	1	352	3.9	84.7	-7.2	+15.5
	2	346	7.6	47.4	-6.5	-18.1
	3	404	12.4	82.1	+8.0	-3.1
	4	363	12.0	63.0	+0.5	-23.2
	5	361	7.7	94.3	-5.0	+16.7
	29	371	20.8	88.2	+14.8	-5.3
	31	351	17.6	75.9	+8.1	-2.0
	32	332	5.0	91.4	-2.4	+11.8
	33	348	6.6	87.5	-0.1	-9.4
	34	299	8.4	73.2	+0.5	-0.2
Mean ± s.e.m.		353 9	10.2 1.7	78.8 4.6	+1.1 2.3	-1.7 4.2
t value (a) p value (a)			13.858 <0.0001 ***			
Percent change (a)			-87%			

Treatment	Rat number	Weight (g)	Force inducing paw-withdrawal (g)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
1015 (30 mg/kg i.v.)	6	328	10.3	90.2	-1.4	+2.0
	7	380	29.9	71.7	+15.0	+2.1
	8	361	5.2	81.7	-2.0	-3.3
	9	358	33.0	79.3	+24.2	+10.6
	10	356	8.8	66.0	-0.3	-25.1
	35	405	10.2	97.5	-0.2	+20.0
	36	355	32.2	76.2	+23.8	-0.4
	37	379	12.3	93.6	+3.8	+18.1
	38	349	10.3	63.3	+1.3	-21.9
	44	373	21.1	65.6	+9.6	-3.7
Mean ± s.e.m.		364 7	17.3 3.4	78.5 3.9	+7.4 3.2	-0.2 4.7
t value (b) p value (b)					1.595 0.1282 NS	0.248 0.8069 NS
Mean change (b)					+6.3	+1.5

Paired Student's t test (a): NS = Not Significant; * = p < 0.05; ** = p < 0.01; *** = p < 0.001.**Unpaired Student's t test (b):** NS = Not Significant; * = p < 0.05; ** = p < 0.01; *** = p < 0.001.

(a): compared with non-lesioned paw.

(b): compared with vehicle.

Neuropathic Pain Test (Chung) in the rat**Test 4 on Day 27 (3 days post dosing): Tactile stimulation (electronic Von Frey test)**

Treatment	Rat number	Weight (g)	Force inducing paw-withdrawal (g)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
B3 (10 mg/kg i.v.)	13	369	8.2	64.8	-3.4	-7.8
	14	324	21.9	79.5	+11.9	+12.6
	15	327	33.8	69.6	+24.7	-13.7
	16	347	7.6	78.6	-6.3	+1.0
	17	395	11.1	79.4	-2.3	+4.9
	40	363	15.6	72.6	+4.6	-18.9
	41	400	16.1	70.5	+5.4	-8.2
	42	365	12.1	90.3	+3.4	+18.0
	55	374	15.6	82.7	+7.8	+9.6
	39	338	4.5	62.0	-8.0	-8.3
Mean ± s.e.m.		360 8	14.7 2.7	75.0 2.7	+3.8 3.1	-1.1 3.8
t value (b) p value (b)					0.709 0.4874 NS	0.113 0.9109 NS
Mean change (b)					+2.7	+0.6

Treatment	Rat number	Weight (g)	Force inducing paw-withdrawal (g)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
B3 (30 mg/kg i.v.)	18	349	12.2	85.6	+3.5	-10.0
	19	395	6.0	69.8	-8.1	+1.4
	20	404	19.1	71.9	+14.3	-14.8
	21	320	24.7	73.0	+19.5	-10.7
	22	315	7.3	72.3	+5.1	+7.4
	45	350	17.0	58.4	+6.9	-17.6
	46	346	13.5	76.6	-2.6	+0.3
	47	388	10.2	78.6	+1.7	-9.0
	48	370	16.2	73.8	+9.7	-10.4
	49	367	5.2	91.2	-4.5	+14.7
Mean ± s.e.m.		360 10	13.1 2.0	75.1 2.8	+4.6 2.7	-4.9 3.3
t value (b) p value (b)					0.988 0.3361 NS	0.585 0.5658 NS
Mean change (b)					+3.5	-3.2

Unpaired Student's t test (b): NS = Not Significant; * = p < 0.05; ** = p < 0.01; *** = p < 0.001.

(b): compared with vehicle.

Neuropathic Pain Test (Chung) in the rat**Test 4 on Day 27 (6 days post dosing): Tactile stimulation (electronic Von Frey test)**

Treatment	Rat number	Weight (g)	Force inducing paw-withdrawal (g)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
Morphine (12 mg/kg i.p.) and Gabapentin (100 mg/kg i.p.)	23	325	11.3	70.1	-2.0	-14.4
	24	308	19.1	83.4	+7.6	+4.9
	25	343	5.8	64.5	-3.2	-11.2
	27	348	13.6	94.4	+0.3	+12.7
	28	331	12.1	67.6	+5.3	-3.7
	50	370	7.5	56.1	+2.1	-19.2
	51	392	25.6	69.2	+12.1	-13.7
	52	342	9.0	81.2	+4.3	+4.9
	53	378	5.9	94.9	+1.4	+30.7
	54	339	23.4	89.5	+13.4	-1.2
Mean ± s.e.m.		348 8	13.3 2.2	77.1 4.3	+4.1 1.8	-1.0 4.8
t value (b) p value (b)					1.065 0.3010 NS	0.111 0.9127 NS
Mean change (b)					+3.0	+0.7

Unpaired Student's t test (b): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(b): compared with vehicle.

Neuropathic Pain Test (Chung) in the rat**Test 4 on Day 27 (3 days post dosing): Thermal stimulation (plantar test)**

Treatment	Rat number	Weight (g)	Paw-withdrawal latency (s)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
Vehicle	1	352	14.8	45.0	+10.0	+20.0
	2	346	13.0	13.4	+5.2	-16.7
	3	404	17.0	23.7	+6.8	+1.3
	4	363	9.0	36.8	-11.6	-7.4
	5	361	16.4	36.3	-11.7	-3.9
	29	371	16.4	38.5	-3.4	+0.1
	31	351	23.2	38.1	+2.7	+11.3
	32	332	8.5	20.2	-2.3	+4.6
	33	348	9.3	16.9	-9.9	-28.1
	34	299	14.8	17.1	+1.0	-0.1
Mean ± s.e.m.		353 9	14.2 1.4	28.6 3.6	-1.3 2.5	-1.9 4.3
t value (a) p value (a)			4.365 0.0018 **			
Percent change (a)			-50%			

Treatment	Rat number	Weight (g)	Paw-withdrawal latency (s)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
1015 (30 mg/kg i.v.)	6	328	9.8	40.1	+2.1	+7.1
	7	380	44.2	45.0	+27.5	+5.8
	8	361	6.0	20.5	-16.2	-6.4
	9	358	14.3	24.0	+9.5	+4.7
	10	356	18.7	45.0	-4.1	0.0
	35	405	45.0	45.0	+41.5	+17.8
	36	355	26.6	45.0	+10.1	+12.1
	37	379	27.6	37.7	+22.8	+0.9
	38	349	13.0	14.5	-13.2	-30.5
	44	373	14.4	11.3	-10.0	-33.7
Mean ± s.e.m.		364 7	22.0 4.3	32.8 4.3	+7.0 6.0	-2.2 5.4
t value (b) p value (b)					1.279 0.2172 NS	0.048 0.9624 NS
Mean change (b)					+8.3	-0.3

Paired Student's t test (a): NS = Not Significant; * = p < 0.05; ** = p < 0.01; *** = p < 0.001.**Unpaired Student's t test (b):** NS = Not Significant; * = p < 0.05; ** = p < 0.01; *** = p < 0.001.

(a): compared with non-lesioned paw.

(b): compared with vehicle.

Neuropathic Pain Test (Chung) in the rat**Test 4 on Day 27 (3 days post dosing): Thermal stimulation (plantar test)**

Treatment	Rat number	Weight (g)	Paw-withdrawal latency (s)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
B3 (10 mg/kg i.v.)	13	369	45.0	45.0	+27.6	+11.8
	14	324	24.7	34.0	+9.1	+5.0
	15	327	33.6	45.0	+21.3	0.0
	16	347	9.8	45.0	-21.4	0.0
	17	395	14.4	15.6	-14.2	-22.7
	40	363	26.5	43.7	+4.4	-1.3
	41	400	45.0	43.2	+17.2	-1.8
	42	365	27.3	45.0	+4.5	+10.7
	55	374	20.1	45.0	+5.8	+16.6
	39	338	29.5	30.5	+12.5	+9.7
Mean ± s.e.m.		360 8	27.6 3.7	39.2 3.1	+6.7 4.8	+2.8 3.5
t value (b) p value (b)					1.492 0.1530 NS	0.849 0.4068 NS
Mean change (b)					+8.0	+4.7

Treatment	Rat number	Weight (g)	Paw-withdrawal latency (s)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
B3 (30 mg/kg i.v.)	18	349	19.7	28.7	+9.9	-0.4
	19	395	16.1	45.0	+8.9	0.0
	20	404	8.9	45.0	-13.0	+5.9
	21	320	14.2	45.0	+1.1	0.0
	22	315	32.0	45.0	+16.1	+2.9
	45	350	40.9	45.0	+23.2	+14.1
	46	346	45.0	45.0	+15.6	0.0
	47	388	11.0	14.0	-25.4	-31.0
	48	370	4.5	45.0	-20.4	+18.2
	49	367	5.8	21.5	-9.7	-16.3
Mean ± s.e.m.		360 10	19.8 4.6	37.9 3.8	+0.6 5.3	-0.7 4.5
t value (b) p value (b)					0.333 0.7431 NS	0.199 0.8447 NS
Mean change (b)					+1.9	+1.2

Unpaired Student's t test (b): NS = Not Significant; * = p < 0.05; ** = p < 0.01; *** = p < 0.001.

(b): compared with vehicle.

Neuropathic Pain Test (Chung) in the rat**Test 4 on Day 27 (6 days post dosing): Thermal stimulation (plantar test)**

Treatment	Rat number	Weight (g)	Paw-withdrawal latency (s)		Delta from baseline	
			<i>Lesioned paw</i>	<i>Non-lesioned paw</i>	<i>Lesioned paw</i>	<i>Non-lesioned paw</i>
Morphine (12 mg/kg i.p.) and Gabapentin (100 mg/kg i.p.)	23	325	35.3	45.0	+25.2	+6.8
	24	308	13.1	45.0	+8.2	0.0
	25	343	8.9	26.0	-24.2	-8.9
	27	348	32.1	45.0	+4.2	0.0
	28	331	24.5	45.0	+8.8	0.0
	50	370	45.0	45.0	+27.7	0.0
	51	392	21.8	24.4	+9.7	+6.7
	52	342	19.8	29.5	-2.5	-8.6
	53	378	18.8	40.2	-11.8	-4.8
	54	339	16.1	22.8	-9.9	-10.1
Mean ± s.e.m.		348 8	23.5 3.5	36.8 3.1	+3.5 5.1	-1.9 1.9
t value (b)					0.854	0.000
p value (b)					0.4042	1.0000
					NS	NS
Mean change (b)					+4.8	0.0

Unpaired Student's t test (b): NS = Not Significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$.

(b): compared with vehicle.

Blood sampling in the rat

Rats used in the Neuropathic Pain Test (Chung), sampled at the end of the experiment

Vehicle	Rat number									
	A1	A2	A3	A4	A5	A6	A7	A8	A9	A10
Date (dd/mm/yy)	04/08/14	04/08/14	04/08/14	04/08/14	04/08/14	04/08/14	04/08/14	04/08/14	04/08/14	04/08/14
Body weight (g)	352	346	404	363	361	371	351	332	348	-
Time of sampling	00h00	00h10	00h20	00h32	00h42	00h52	01h57	02h09	02h18	-

1015 (30 mg/kg i.v.)	Rat number									
	B1	B2	B3	B4	B5	B6	B7	B8	B9	B10
Date (dd/mm/yy)	04/08/14	04/08/14	04/08/14	04/08/14	04/08/14	04/08/14	04/08/14	04/08/14	04/08/14	04/08/14
Body weight (g)	328	380	361	358	356	405	355	379	349	373
Time of sampling	00h02	00h12	00h23	00h34	00h44	00h54	01h59	02h11	02h20	02h31

B3 (10 mg/kg i.v.)	Rat number									
	C1	C2	C3	C4	C5	C6	C7	C8	C9	C10
Date (dd/mm/yy)	04/08/14	04/08/14	04/08/14	04/08/14	04/08/14	04/08/14	04/08/14	04/08/14	04/08/14	04/08/14
Body weight (g)	369	324	327	347	395	363	400	365	374	338
Time of sampling	00h04	00h14	00h25	00h36	00h46	00h56	02h01	02h13	02h25	02h33

B3 (30 mg/kg i.v.)	Rat number									
	D1	D2	D3	D4	D5	D6	D7	D8	D9	D10
Date (dd/mm/yy)	04/08/14	04/08/14	04/08/14	04/08/14	04/08/14	04/08/14	04/08/14	04/08/14	04/08/14	04/08/14
Body weight (g)	349	395	404	320	315	350	346	388	370	367
Time of sampling	00h06	00h16	00h27	00h38	00h48	01h53	02h03	02h15	02h27	02h35

Morphine (12 mg/kg i.p.) and Gabapentin (100 mg/kg i.p.)	Rat number									
	E1	E2	E3	E4	E5	E6	E7	E8	E9	E10
Date (dd/mm/yy)	04/08/14	04/08/14	04/08/14	04/08/14	04/08/14	04/08/14	04/08/14	04/08/14	04/08/14	04/08/14
Body weight (g)	325	308	343	348	331	370	392	342	378	339
Time of sampling	00h08	00h18	00h29	00h40	00h50	01h55	02h06	02h17	02h29	02h37

Blood sampling in the rat

Remarks:

Hemolysis for rats B7, D7 and D8.

-: missing value due to technical problem.

8. COMPLEMENTARY STATISTICAL ANALYSIS (produced by the Sponsor)

Within each row, compare columns (simple effects within rows)

Number of families 7

Number of comparisons per family 4

Alpha 0.05

Bonferroni's multiple comparisons test Mean Diff. 95% CI of diff. Significant? Summary

Row 1

1015 (30 mg/kg) Lesioned vs. VEHICLE	5.800	-6.847 to 18.45	No	ns		
1015 (30 mg/kg) Lesioned vs. 1015 (30 mg/kg) Nonlesioned	1.920	-10.73 to 14.57	No	ns		
1015 (30 mg/kg) Lesioned vs. B3 (10 mg/kg)	-1.860	-14.51 to 10.79	No	ns		
1015 (30 mg/kg) Lesioned vs. B3 (30 mg/kg)	1.760	-10.89 to 14.41	No	ns		

Row 2

1015 (30 mg/kg) Lesioned vs. VEHICLE	-9.070	-21.72 to 3.577	No	ns		
1015 (30 mg/kg) Lesioned vs. 1015 (30 mg/kg) Nonlesioned	-6.830	-19.48 to 5.817	No	ns		
1015 (30 mg/kg) Lesioned vs. B3 (10 mg/kg)	-1.620	-14.27 to 11.03	No	ns		
1015 (30 mg/kg) Lesioned vs. B3 (30 mg/kg)	-3.340	-15.99 to 9.307	No	ns		

Row 3

1015 (30 mg/kg) Lesioned vs. VEHICLE	-0.6000	-13.25 to 12.05	No	ns		
1015 (30 mg/kg) Lesioned vs. 1015 (30 mg/kg) Nonlesioned	-20.07	-32.72 to -7.423	Yes	***		
1015 (30 mg/kg) Lesioned vs. B3 (10 mg/kg)	-5.950	-18.60 to 6.697	No	ns		
1015 (30 mg/kg) Lesioned vs. B3 (30 mg/kg)	-4.220	-16.87 to 8.427	No	ns		

Row 4

1015 (30 mg/kg) Lesioned vs. VEHICLE	5.460	-7.187 to 18.11	No	ns		
1015 (30 mg/kg) Lesioned vs. 1015 (30 mg/kg) Nonlesioned	-16.65	-29.30 to -4.003	Yes	**		
1015 (30 mg/kg) Lesioned vs. B3 (10 mg/kg)	-8.830	-21.48 to 3.817	No	ns		
1015 (30 mg/kg) Lesioned vs. B3 (30 mg/kg)	-9.380	-22.03 to 3.267	No	ns		

Row 5

1015 (30 mg/kg) Lesioned vs. VEHICLE	2.680	-9.967 to 15.33	No	ns		
1015 (30 mg/kg) Lesioned vs. 1015 (30 mg/kg) Nonlesioned	-12.69	-25.34 to -0.04344	Yes	*		
1015 (30 mg/kg) Lesioned vs. B3 (10 mg/kg)	-12.23	-24.88 to 0.4166	No	ns		
1015 (30 mg/kg) Lesioned vs. B3 (30 mg/kg)	-12.37	-25.02 to 0.2766	No	ns		

Row 6

1015 (30 mg/kg) Lesioned vs. VEHICLE	9.290	-3.357 to 21.94	No	ns		
1015 (30 mg/kg) Lesioned vs. 1015 (30 mg/kg) Nonlesioned	-15.70	-28.35 to -3.053	Yes	**		
1015 (30 mg/kg) Lesioned vs. B3 (10 mg/kg)	-12.97	-25.62 to -0.3234	Yes	*		
1015 (30 mg/kg) Lesioned vs. B3 (30 mg/kg)	2.110	-10.54 to 14.76	No	ns		

Row 7

1015 (30 mg/kg) Lesioned vs. VEHICLE	7.720	-4.927 to 20.37	No	ns		
1015 (30 mg/kg) Lesioned vs. 1015 (30 mg/kg) Nonlesioned	-10.85	-23.50 to 1.797	No	ns		
1015 (30 mg/kg) Lesioned vs. B3 (10 mg/kg)	-5.630	-18.28 to 7.017	No	ns		
1015 (30 mg/kg) Lesioned vs. B3 (30 mg/kg)	2.150	-10.50 to 14.80	No	ns		

Test details Mean 1 Mean 2 Mean Diff. SE of diff. N1 N2 t DF

Row 1												
1015 (30 mg/kg) Lesioned vs. VEHICLE	23.69	17.89	5.800	5.034	10	10	1.152	315				
1015 (30 mg/kg) Lesioned vs. 1015 (30 mg/kg) Nonlesioned	23.69	21.77	1.920	5.034	10	10			0.3814	315		
1015 (30 mg/kg) Lesioned vs. B3 (10 mg/kg)	23.69	25.55	-1.860	5.034	10	10	0.3695	315				
1015 (30 mg/kg) Lesioned vs. B3 (30 mg/kg)	23.69	21.93	1.760	5.034	10	10	0.3496	315				
Row 2												
1015 (30 mg/kg) Lesioned vs. VEHICLE	19.84	28.91	-9.070	5.034	10	10	1.802	315				
1015 (30 mg/kg) Lesioned vs. 1015 (30 mg/kg) Nonlesioned	19.84	26.67	-6.830	5.034	10	10			1.357	315		
1015 (30 mg/kg) Lesioned vs. B3 (10 mg/kg)	19.84	21.46	-1.620	5.034	10	10	0.3218	315				
1015 (30 mg/kg) Lesioned vs. B3 (30 mg/kg)	19.84	23.18	-3.340	5.034	10	10	0.6635	315				
Row 3												
1015 (30 mg/kg) Lesioned vs. VEHICLE	14.96	15.56	-0.6000	5.034	10	10	0.1192	315				
1015 (30 mg/kg) Lesioned vs. 1015 (30 mg/kg) Nonlesioned	14.96	35.03	-20.07	5.034	10	10			3.987	315		
1015 (30 mg/kg) Lesioned vs. B3 (10 mg/kg)	14.96	20.91	-5.950	5.034	10	10	1.182	315				
1015 (30 mg/kg) Lesioned vs. B3 (30 mg/kg)	14.96	19.18	-4.220	5.034	10	10	0.8383	315				
Row 4												
1015 (30 mg/kg) Lesioned vs. VEHICLE	18.68	13.22	5.460	5.034	10	10	1.085	315				
1015 (30 mg/kg) Lesioned vs. 1015 (30 mg/kg) Nonlesioned	18.68	35.33	-16.65	5.034	10	10			3.307	315		
1015 (30 mg/kg) Lesioned vs. B3 (10 mg/kg)	18.68	27.51	-8.830	5.034	10	10	1.754	315				
1015 (30 mg/kg) Lesioned vs. B3 (30 mg/kg)	18.68	28.06	-9.380	5.034	10	10	1.863	315				
Row 5												
1015 (30 mg/kg) Lesioned vs. VEHICLE	20.38	17.70	2.680	5.034	10	10	0.5324	315				
1015 (30 mg/kg) Lesioned vs. 1015 (30 mg/kg) Nonlesioned	20.38	33.07	-12.69	5.034	10	10			2.521	315		
1015 (30 mg/kg) Lesioned vs. B3 (10 mg/kg)	20.38	32.61	-12.23	5.034	10	10	2.429	315				
1015 (30 mg/kg) Lesioned vs. B3 (30 mg/kg)	20.38	32.75	-12.37	5.034	10	10	2.457	315				
Row 6												
1015 (30 mg/kg) Lesioned vs. VEHICLE	24.62	15.33	9.290	5.034	10	10	1.845	315				
1015 (30 mg/kg) Lesioned vs. 1015 (30 mg/kg) Nonlesioned	24.62	40.32	-15.70	5.034	10	10			3.119	315		
1015 (30 mg/kg) Lesioned vs. B3 (10 mg/kg)	24.62	37.59	-12.97	5.034	10	10	2.576	315				
1015 (30 mg/kg) Lesioned vs. B3 (30 mg/kg)	24.62	22.51	2.110	5.034	10	10	0.4191	315				
Row 7												
1015 (30 mg/kg) Lesioned vs. VEHICLE	21.96	14.24	7.720	5.034	10	10	1.534	315				
1015 (30 mg/kg) Lesioned vs. 1015 (30 mg/kg) Nonlesioned	21.96	32.81	-10.85	5.034	10	10			2.155	315		
1015 (30 mg/kg) Lesioned vs. B3 (10 mg/kg)	21.96	27.59	-5.630	5.034	10	10	1.118	315				
1015 (30 mg/kg) Lesioned vs. B3 (30 mg/kg)	21.96	19.81	2.150	5.034	10	10	0.4271	315				

APPENDIX 1**Antibody (B3) concentrations in plasma at the termination of Study No: 14.0061****Methods****ELISA**

Ninety six well plates were coated with 100 µl of 1 µg/ml of (C12)LPA-BSA-SMCC for 1 hour at 37 degrees. The plates were washed 4 times with PBS and blocked with 150µl of PBS-BSA-tween for 1 hour at room temperature. 100µl of B3 antibody was added starting at 4 µg/ml and diluted 2-fold across the plate for the standard curve. Rat plasma samples were diluted to 1:100, 1:1000, 1:2000, 1:4000, 1:8000, and 1:16000 and 100µl of each dilution was run in duplicate. Plates were incubated at room temperature for 1 hour. Next, plates were washed 4 times with PBS and 100µl of a peroxidase conjugated anti-mouse IgG (H+L) (Jackson immunoresearch 115-035-062) was added at a 1:40K dilution and incubated for 1 hour at room temperature. Finally, the plates were washed four times with PBS and developed with 100µl cold TMB substrate for 12 minutes at room temperature. The reaction was stopped with 100µl of 1M H₂SO₄. Plates were read at 450 nm.

Data Archive

LPath Notebook # 278 pages 36-37

L:\WORKING FOLDERS-LPATH/Stephanie/2014/PK/9-11-14 PK for Ann.xls

RESULTS**B3 Concentration in plasma samples**

Tratment	Rat number	[Ab] (ug/mL)	tube label
Vehicle	1	0.00	A1
	2	0.00	A2
	3	0.00	A3
	4	0.00	A4
	5	0.00	A5
	29	0.00	A6
	31	0.00	A7
	32	0.00	A8
	33	0.00	A9
	34		no A10?
1015 (30 mg/kg i.v.)	6	0.00	B1
	7	0.00	B2
	8	0.00	B3
	9	0.00	B4
	10	0.00	B5
	35	0.00	B6
	36	0.00	B7
	37	0.00	B8
	38	0.00	B9
	44	0.00	B10
B3 (10 mg/kg i.v.)	13	432.82	C1
	14	151.58	C2
	15	67.00	C3
	16	190.15	C4
	17	156.24	C5
	40	151.29	C6
	41	212.48	C7
	42	0.00	C8
	55	182.34	C9
	39	60.64	C10

Tratment	Rat number	[Ab] (ug/mL)	tube label
B3 (30 mg/kg i.v.)	18	318.47	D1
	19	166.42	D2
	20	221.25	D3
	21	50.32	D4
	22	67.52	D5
	45	751.15	D6
	46	468.13	D7
	47	426.15	D8
	48	1106.06	D9
	49	643.11	D10
Morphine (12 mg/kg i.p.) and Gabapentin (100 mg/kg i.p.)	23	0.00	E1
	24	0.00	E2
	25	0.00	E3
	27	0.00	E4
	28	0.00	E5
	50	0.00	E6
	51	0.00	E7
	52	0.00	E8
	53	0.00	E9
	54	0.00	E10