Brain Health Maintenance:

Discovery of New Drugs affordably – Re-purposing Principles

This is used here to Discover Novel Drugs that can treat a neuro-inflammatory disease

(Multiple sclerosis or thinning of white matter)

Inclusion criteria

- Reporting of primary data from clinical trial(s).
- Qualitative or quantitative data on either safety or efficacy.

Initial screening of candidate drugs

Eligibility criteria for publications

Exclusion criteria

- Studies reporting secondary analyses of previously published data.
- Reports of protocols for potential clinical trials.
- Studies in patients with relapsing-remitting MS.
- Use of interventions already licensed for clinical application in MS.
- Studies reporting interventions not administered orally.
- Studies reporting combination interventions.
- Studies reporting only non-pharamacological interventions (e.g. acupuncture, aromatherapy, physiotherapy, exercise).
- Studies reporting use of levodopa treatment for Parkinson's disease.

Systematic evaluation: Candidate drugs

Meta-Analysis

	CAMARADES	Delphi	GRADE
Binary response items:			
Yes (1 point); No (0 points)			
Peer reviewed publication	X		
Statement of potential conflicts of interest	Χ		
Sample size calculation	X	X	
Random allocation to group	Χ	X	Χ
Ternary response items:			
Yes (1 point); No (0 points); Not Clear (0.5 points)			
Were the groups similar at baseline regarding the most important prognostic indicators?		X	
Were the eligibility criteria specified?		X	
Were point estimates and measures of variability presented for the primary outcome measures?		X	
Was there an intention to treat analysis?		X	
Quinary response items:			
N/A; Definitely yes (1 point); Probably yes (0.75 points); Probably no (0.25 points); Definitely no (0 points)			
Was selection of treatment and control groups drawn from the same population?			X
Can we be confident that patients received the allocation treatment?			X


Selection process:

Publications & Interventions

Phase I Systematic Iiterature review

Systematic evaluation of interventions

Phase III
Committee
evaluation of
interventions



Phase-1:

Systematic Survey of Literature

Candidate drugs excluded during short-listing, with reason of Rejection

[MOA = mechanism of action]

Drug	Reason for exclusion
15+- Dexoxyspergualin	Immunosuppressive MOA
4 Ammonium Phosphate	Limited biological plausibility
Adrenocorticotropic Hormone	Immunosuppressive MOA
Adrenocorticotropic Homone 1–17	Immunosuppressive MOA
Amantidine/Isoprinosine	Symptomatic benefit
Anastal	Combination therapy
Antithymocyte Globulin	Immunosuppressive MOA
Arachidonic Acid	Limited biological plausibility
Azathioprine	Immunosuppressive MOA
Azathioprine & Prednisolone	Immunosuppressive MOA
Azathioprine/6-mercaptopurine	Immunosuppressive MOA
Baclofen	Symptomatic benefit

Short-listed drugs ranked by overall drug scores

	Number of publications	Efficacy Score	Safety Score	Quality Score	Patient Sample Size Score	Overall Drug Score
Dextromethorphan + Quinidine	3	3.3	2.3	4	3	56.2
Amantadine	57	3⋅1	2.4	2.2	1.9	55-2
Memantine	34	2.7	2	2.6	2.4	51.6
Gabapentin	8	2.7	2.5	2.9	2.5	46.7
4-Aminopyridine	10	2.7	2.5	3	2.1	44-4
Modafinil	8	2.9	2.4	3.1	2.1	44.4
Creatine	12	2.1	2.3	2.7	2.4	35.9
Selegiline	11	2.7	1.9	2.7	2.3	34.2
L-amphetamine sulfate	1	3	3	4	3	32.5
Minocycline	11	2.3	2.6	2.3	2.2	31.9
Vitamin E	9	2.1	2	3	2.4	31.1
Coenzyme Q ₁₀	9	2.2	1.9	3.2	2.3	30.6

Phase-2:

Evaluation of Interventions

Neurobehavioral Pathological efficacy outcomes


Neurobehavioural score

Expanded in Next Slide


Symbol estimates of efficiency of drugs.


Symbol's diameter denote: log_{10} (No. of animals in that study

Vertical line represents: Effect = nil or zero



Improvement in Outcome (SD)





Phase-3:

Assessment by Domain Knowledge Committee

Drugs Rejected

Committee review phase

[EAE = experimental allergic encephalomyelitis, i.e. Experimental Multiple sclerosis disease in animals]

	·			
	Drug	Reason for exclusion		
Excluded after round 1	4-Aminopyridine	Risk of seizures, limited efficacy data		
	Atomoxetine	Safety profile, no efficacy data in MS		
	Clofibrate	Safety profile, neutral efficacy data		
	Bromocriptine	Safety profile		
	Dextromethorphan	Requires co-administration with quinidine		
	Diazepam	Safety profile, limited evidence for efficacy		
	Ginseng	Limited clarity around MOA, no efficacy data in MS		
	Imipramine	Limited evidence for efficacy		
	Isoniazid	Likely symptomatic benefit only, no efficacy data in MS		
	Levamisole	Safety profile		
	Linomide	Safety profile		
Excluded after round 2	Amantadine	Limited evidence for neuroprotective MOA		
	Aspirin	No evidence in EAE, insufficient basis to trial as neuroprotective agent in SPMS		
	Carbamazepine	Better safety and efficacy data for same-class alternative (oxcarbazepine)		
	Coenzyme Q ₁₀	Limited efficacy data in MS		
	Creatine	Safety profile, limited efficacy data		
	Creatine	Safety profile, limited efficacy data		

Final recommendations as candidate drugs for Multiple Sclerosis (MS)

Intervention	Current main clinical application and mechanism of action
Ibudilast	Anti-inflammatory use in asthma: non-selective phosphodiesterase (PDE 3,4,10,11) inhibitor and macrophage Migration Inhibitor Factor (MIF) inhibitor.
Riluzole	MND/ALS: glutamate release inhibitor/inactivation of voltage-dependent sodium channels.
Amiloride	Diuretic: acid sensing ion channel blocker.
Pirfenidone	Pulmonary fibrosis: antagonises synthesis of TGF-beta & TNF-alpha; antifibrotic/anti-inflammatory activity.
Fluoxetine	Antidepressant: selective serotonin reuptake inhibitor.
Oxcarbazepine	Anticonvulsant: voltage sensitive sodium channel blocker.
PUFA class (Linoleic Acid, Lipoic acid; Omega-3 fatty acid, Max EPA oil)	None / dietary supplements: mechanism of action unclear.

Limitations of this Approach

- Possibility of incomplete ascertainment of relevant publications, and publication bias favouring studies with positive findings.
- Does not overcome problems that were inherent in the design and execution of the primary studies.
- Exclusion of drugs based on their mode of action will have overlooked drugs with more than one relevant mode of action.