Hand Assessment

Walaa Abu-Taleb







Observation

- Bouchard's nodes
- Heberden's nodes
- Claw hand
- Club nails
- Dupuytren's contracture

FIGURE 1

The nodule, cord, and contracture seen in Dupuytren's disease




PHOTOS COURTESY OF: JOANNE R. WERNTZ, MD

Skin dimpling and a palpable nodule are evident near the base of a patient's ring finger (**A**). A well-defined cord associated with the little finger is causing minimal misalignment (**B**). Severe contracture at the proximal interphalangeal joint (**C**) can interfere with routine daily activities.


Observation


- Mallet finger
- Boutonniere's deformity
- Swan’s neck deformity












Observation













Palpation

- Temperature
- Swelling
- Tenderness
- Nodules or Masses
- Pulse

Hand Movements


A to D, Finger motion. E to I, Thumb motion.
(A, Finger flexion;
B, finger extension;
C, finger abduction;
D, finger adduction;
E, thumb flexion;
F, thumb extension;
G, thumb abduction;
H, thumb adduction;
and I, thumb opposition.)








(A)



(B)




ROM Assessment

Muscle Testing


1. Grip Strength: dynamometer
2. Pinch Strength: tip-to-tip, lateral, three-point pinch
3. Individual muscles and muscle groups of the hand: Flexors, extensors, intrinsics
4. Functional Tasks: grasping objects, picking up small items, or writing.

Hand Grasps



Power
Precision
Cylindrical
Spherical
Hook
Pinch
Tripod
Lumbrical


Finkelstein Test- De Quervain's Tenosynovitis

- Step 1: Active radial deviation against gravity
- Step 2: passive radial deviation against gravity
- Step 3: passive thumb pressure towards the palm




Modefied Allen's Test- Vascular Assessment





Neurological testing

- Sensory
- Motor
- Reflex





Thank you