Known phyla, unknown proteome: a controlled census of the Rub' al-Khali soil metagenome

A. Lopez Velazquez *et al.*
King Abdullah University of Science and Technology

Abstract

Reference coverage limits interpretation of microbial gene content. We analysed 551 soil metagenomes from the Rub' al-Khali and recovered 5,438 species-level genome representatives. Among 43.4 million protein clusters linked to genome bins, 9.89% lacked a detected UniRef50 match. A recurrent subset contained 106,042 families and 12.5% of the dark representatives. Annotated frames had lower dN/dS than shifted frames, but most artifact-flagged families also had dN/dS below one. Between-group contrasts depended on divergence and population weighting. Culture-derived genomes revealed dark calls with matching DNA and no corresponding annotated gene. Structural reference coverage depended on score normalisation and retrieval. Per-genome darkness varied by lineage: 237 Thermoplasmatota representatives had a median dark fraction of 31.0%. The combined desert catalogue and SPIRE recovered more dark families than an E-value-corrected MGnify90 rarefaction expectation. The catalogue connects reference-absent gene content to genomes and repeated sampling, supporting investigation of the unknown proteome of hyperarid soils.

Introduction

Drylands cover roughly two fifths of the land surface and are projected to expand [1, 2], and their hot hyperarid interiors are the low-biomass case in which contamination approaches the detection limit [3]. The Rub' al-Khali, the largest contiguous sand sea, had no genome-resolved microbial survey. A companion 16S rRNA survey of the same transect mapped its bacterial ecology by landscape, soil position and rainfall [4], and listed what amplicons cannot reach: gene content without primer bias, function tied to a genome, and the archaea.

What genomes reveal is mostly unknown. Across global catalogues a large share of microbial protein families match no characterised family, and the size of that share depends on the reference and the rule used to call it [5–7]. The unknown families are numerous, span species, evolve under purifying selection and concentrate in uncultivated lineages [8]; most are restricted to one ecosystem [9]; and large structural surveys extend domain assignments beyond sequence-based annotations [10, 11]. Interpreting the remaining reference-absent proteins requires separating evidence for coding status from evidence for structural or functional novelty. Repeated environmental sampling provides an opportunity to connect these questions to the persistence and genomic context of individual families.

A reference no-hit depends on gene prediction, database composition and search sensitivity. Recurrent families can support evolutionary comparisons, while culture-derived assemblies offer an independent reconstruction of DNA from related organisms. Known-fold proteins processed through a structural pipeline provide a retrieval control. Combining these comparisons in one sampling programme can delimit which parts of an unknown repertoire are supported by each form of evidence.

Here we combine these comparisons in one biome. We shotgun sequenced 551 soil samples from 61 sampled locations across the Rub' al-Khali on five expeditions over 31 months, at three soil positions and with elemental chemistry, recovered 5,438 species-level genomes, and measured the unknown fraction of their proteome against a control at every layer: annotated-family controls for selection and reference absence; long-read genomes of strains cultured from soils of the same basin; blinded known-fold controls for structure; and matched comparisons for a lineage with a high dark fraction. The comparisons distinguish frame-dependent evolutionary estimates in recurrent families from variation in reference coverage across lineages, genomes and structural search criteria.

Results

Genome recovery and taxonomic coverage

We sampled 59 core transect sites and two auxiliary locations along a 1,043 km transect of the Rub' al-Khali on five expeditions between March 2023 and October 2025, at the soil surface, in the subsurface and beside plant roots, and shotgun sequenced 551 soil samples to a median depth of 21.58 Gbp per sample (Fig. 1a,b). Of the 549 samples that assembled, two SemiBin2 binnings (pretrained model and self-supervised) refined by DAS_Tool yielded 14,426 bins, 4,225 near-complete and 8,209 additional medium-completeness bins under the operational CheckM2 thresholds; dereplication at 95% average nucleotide identity gave 5,438 species representatives, including 2,259 near-complete genomes (Fig. 1c). Full MIMAG assessment classified 126 as high quality and 5,312 as medium quality. Taxonomy throughout is GTDB release 232.

Every representative was placed to class rank, within 51 GTDB phylum labels. The first empty taxonomic rank was order for 11 representatives, family for 142, genus for 1,343 and species for 3,755; 187 carry a species assignment. These are mutually exclusive depths of the first empty rank, not counts of validated new taxa. Of the 11 order-level gaps, the 10 with adequate marker coverage were chimera-clean by GUNC (clade separation 0.01 to 0.32 against a 0.45 threshold), and de novo trees with release-calibrated relative evolutionary divergence supported no new order: six are single-genome lineages and the two multi-genome clades sit at genus and family depth. Empty-rank depth is distinct from a naming heuristic that flags an empty or digit-bearing GTDB label (Table 1). The latter is a proxy for reference naming coverage.

Table 1: Mutually exclusive taxonomy depths among 5,438 representatives. The domain count is zero for both rules.

First flagged rank	Phylum	Class	Order	Family	Genus	Species	None
Empty label	0	0	11	142	1,343	3,755	187
Empty or digit-bearing label	10	560	272	1,006	1,613	1,855	122

Recurrent-family comparisons depend on reading frame and gene calling

Gene prediction on the 552 assembly files gave 753,305,533 proteins, which collapsed to 449,265,306 cluster representatives at 90% identity over 80% of the length. A cluster was called sequence-dark when none of its searched members returned a UniRef50 hit under a sensitive search with an E-value threshold and no identity or coverage filter (Methods); 69,938,628 representatives (15.6%) were dark. Restricting to clusters with at least one member on a contig of the 37,949 bins meeting the operational CheckM2 medium-or-better thresholds in the two binnings before refinement gives the genome-resolved census: 4,292,446 of 43,423,322 representatives (9.89%) have no reference homolog. Both percentages are shares of cluster representatives, not of proteins or of species. Within the genome-resolved dark set, 65.8% of representatives are shorter than 100 residues and 56.4% are partial gene calls. Across the assembly-wide catalogue,

including unbinned sequence, the corresponding dark-set values are 75.7% and 86.7%; among all 449 million representatives, including dark clusters, 29.7% are shorter than 100 residues and 87.5% are partial. These distributions use different catalogue scopes.

We selected recurrent families to examine sequence constraint and distribution. Clustering the 4,292,446 genome-resolved dark representatives at 50% identity produced 3,658,797 clusters: 3,353,521 singletons, 196,450 two-member clusters and 108,826 clusters with at least three members. Requiring at least three samples and two sites retained 106,042 families with 536,437 members, or 12.5% of the dark representatives. The other 2,784 clusters with at least three members failed the spread criterion. An additional 782,616 singleton or two-member clusters recurred across the required samples and sites but remained outside the admitted family set. The median admitted family was assembled in 12 samples at 9 sites. Subsequent family analyses are conditional on this recurrence-based selection.

Persistence: with reads from all 551 samples mapped to the member genes, the median family is detected in 98 samples, four families are confined to one sample, and 79.4% of families are detected in all five expeditions (94.8% in at least four). Under the stricter criterion of independent assembly, 28.9% were assembled in all five. Detection by covered fraction favours short genes, and these frequencies describe prevalence under the mapping rule. A prevalence-conditioned null and competitive mapping would be needed to interpret temporal persistence beyond repeated DNA detection.

Selection: the median pairwise dN/dS across 97,200 evaluable dark families is 0.425, with 96.7% below one, against 0.388 (99.6% below one, $n = 95{,}647$) for 106,042 UniRef50-matched families selected using representative length and family size and passed through the same clustering, alignment and estimation steps (Fig. 2a). The pooled distributions differ, and the groups also differ in sequence divergence: median family mean dS is 0.607 in the dark arm and 0.863 in the control. Comparisons therefore depend on divergence strata. For families with mean dS below 0.10, median dN/dS is 0.582 in the dark arm and 0.321 in the control; for mean dS between 0.5 and 1.0, the medians are 0.435 and 0.452. The control provides a reference distribution for the estimation pipeline, with divergence and family-admission differences that remain relevant to interpretation.

The difference between stratum medians was +0.261 (95% family-bootstrap interval 0.197–0.288) at mean dS below 0.1 and −0.0163 (−0.0187 to −0.0142) at dS 0.5–1.0 (Fig. 2c). These conditional intervals treat families as independent. Time dependence, population sampling and correlated estimation error limit direct interpretation of pairwise dN/dS as selection strength [12, 13].

Coarsened matching on mean synonymous divergence, median coding-sequence length, GC content and sequence count retained 61,427 families in each arm, with medians 0.386 for dark and 0.423 for annotated families. This subset retained 7.5% of dark families in the lowest divergence stratum and 98.8% in the highest; 88.4% had mean dS at least 0.5. Using all families in the 553 common strata before equal-count sampling (93,220 dark and 95,541 annotated families), weighted mean differences of family medians were +0.012 using the dark-arm distribution, −0.022 using the annotated-arm distribution and −0.018 using overlap weights. These means and the pooled medians are different estimands. The direction of the contrast depends on divergence and the target population. Residual dS imbalance remained after matching (standardised mean difference −0.124), and dS shares estimation noise with dN/dS.

For a seeded input subset of 10,000 families, shifting member genes by one nucleotide and masking stop codons produced 9,148 evaluable families, with median dN/dS 1.086 and 37.6% below one. The corresponding masked annotated-frame analysis yielded 9,182 families, with median 0.427 and 97.0% below one (Fig. 2b). Both arms were evaluable for 9,029 families, and the shifted frame gave the higher estimate in 92.4%. This contrast shows that the estimated dN/dS depends on reading frame. Interpreting coding status also requires tests of overlapping gene calls and of the effects of sequence ascertainment. An AntiFam screen flagged 141 admitted member proteins in 56 families. A coordinate-verified screen found 70 members in 46 families overlapping

an opposite-strand non-dark Pyrodigal call by at least half their length. Excluding the union of 102 flagged families left 97,109 evaluable dark families, with median dN/dS 0.4251 and 96.7% below one. The 91 evaluable flagged families themselves had median dN/dS 0.762 (interquartile range 0.53–0.97), and 76.9% were below one; among the 47 AntiFam-flagged families, the median was 0.831 and 76.6% were below one. Thus a below-one estimate does not distinguish a coding gene from an artifact-flagged family. A flag on any member need not invalidate every member, and overlapping frames can inherit sequence constraints [14, 15]. These checks identify known spurious-family matches and overlaps among existing calls; they do not detect every uncalled alternative ORF or establish translation [16]. In particular, the overlap audit depends on both calls surviving Prodigal's gene-selection step, which suppresses heavily overlapping alternatives [17].

Genomes recovered from cultures provide a complementary comparison. Long-read assemblies of 722 cultured genomes from saline pans, oases and campground soils of the basin were assessed with the same UniRef50 search rule. They shared a species-level ANI match with 25 catalogue representatives, yielding 23 representative–culture groups after assignment to the best-ANI representative. Excluding the *Escherichia coli* representative detected in the no-template control left 22 groups. Cultured genomes had a lower dark fraction in 18 of these groups, with a median difference of -0.36 percentage points (Wilcoxon $p = 0.0066$). The cultured assemblies had a median of two contigs, and 49% were single-contig assemblies. Across all cultured genomes, dark proteins had median length 59 residues, compared with 70 for genome-resolved dark representatives.

In the 22 retained representative–culture pairs, reciprocal protein matches covered 65,478 of 81,723 UniRef50-matched MAG proteins (80.1%) and 238 of 1,507 dark MAG proteins (15.8%). The annotated fraction was more frequently shared in every pair; shared matches had median identity 99.7%. Among the 166 complete dark calls of at least 100 residues, 26 had a reciprocal partner. These comparisons measure sharing between two predicted proteomes. Gene-caller sensitivity, assembly fragmentation, bin membership and strain variation can each contribute to unmatched calls. Base-level nucleotide alignments covering at least 90% of a coding interval at 95% identity supported 516 of 1,105 eligible dark calls without a protein partner (46.7%) and 4,877 of 16,056 annotated calls without a partner (30.4%). Among protein-shared calls, the corresponding counts were 219 of 236 dark calls (92.8%) and 62,925 of 65,477 annotated calls (96.1%). The union of reciprocal-protein and DNA support covered 752/1,341 eligible dark calls (56.1%) and 70,354/81,533 annotated calls (86.3%). This union includes protein-supported calls that failed the nucleotide threshold; it is not a DNA-only detection rate. Protein sharing therefore understates the combined evidence for shared sequence in both groups (Fig. 2d–f). Across pairs, recovery of protein-shared calls ranged from 67.9% to 99.6%. In the post hoc subset of 20 pairs with at least 90% recovery, DNA support for protein-unshared calls was 496/940 (52.8%) for dark calls and 4,254/11,240 (37.8%) for annotated calls. A nucleotide-search negative still depends on divergence and alignment sensitivity. Among the 516 DNA-supported dark calls without protein partners, 343 had no Bakta CDS overlapping at least half the aligned target span, 81 overlapped a different frame or orientation, and 92 had a same-orientation CDS compatible with the phase at the alignment start. The corresponding annotated counts were 606, 183 and 4,088. This local diagnostic separates possible gene-call and protein-match effects; it does not establish full codon correspondence or validate a gene. Across all DNA-supported calls, including those with protein partners, 344 of 735 dark calls (46.8%) and 607 of 67,802 annotated calls (0.90%) lacked a Bakta CDS covering half the aligned span. This diagnostic provides direct evidence of gene-call disagreement at shared DNA. Only 65 of the 1,341 eligible dark calls belonged, through their 90%-identity clusters, to 47 admitted recurrent families. The culture result therefore describes a broader bacterial call set and cannot be transferred wholesale to the recurrent-family analysis. Harmonised gene prediction would further distinguish these cases.

The cultured comparison covers well-referenced bacteria with low dark fractions; archaeal lineages require separate validation.

Per-genome darkness varies strongly among lineages

Per genome, the dark fraction is the share of protein instances on a representative's contigs whose cluster has no UniRef50 homolog; across the 5,438 representatives its median is 6.3% (Fig. 3a) (interquartile range 3.7–10.4%), a per-genome rate. Assembly quality (completeness, contamination, N50 and length together) explains 11.5% of the variance in that fraction. Adding phylum raises the explained share to 76.8% and genus to 83.5%; 72.0% of the variance is unique to lineage and 2.0% unique to quality. Larger genomes are slightly less dark (-0.096 per tenfold increase in corrected size, R^2 0.067), so genome size does not explain darkness. Taxonomic novelty predicts part of it: the median dark fraction falls from 0.217 for genomes first flagged at phylum to 0.017 for genomes with no flagged rank (Fig. 3b), and within Actinomycetota, Pseudomonadota and Bacteroidota genomes with a placeholder genus are darker than those with a named genus (medians 0.075 against 0.022, 0.051 against 0.030, 0.068 against 0.053). That much is expected, because a lineage absent from UniRef50 is dark by construction. Novelty does not absorb the lineage term: with first empty or digit-bearing rank in the model, phylum still adds 0.340 to R^2, 52% of its unconditioned increment. Phylum and genus enter with far more degrees of freedom than the four quality covariates and the model treats genomes as independent draws, so these increments are descriptive upper bounds; the like-for-like test is the matched contrast in the next section.

The residual is concentrated. After the quality model, Thermoplasmatota genomes are 0.21 darker than predicted, Cyanobacteriota 0.14 and Bdellovibrionota 0.10, while Halobacteriota, Bacillota, Pseudomonadota and Actinomycetota sit below the line. Thermoplasmatota have the largest positive residual among these phyla; Actinomycetota contribute more dark protein instances in absolute terms.

Thermoplasmatota carry a large share of the attributable dark families

The catalogue holds 237 Thermoplasmatota species: 185 in class SW-10-69-26, 51 in class E2 and one in Thermoplasmata, none with a named genus. GTDB release 232 holds 21 and 197 species representatives in the two classes, fewer together than this one sand sea. Their median dark fraction is 0.310, against 0.046 for the other 354 archaea and 0.062 for the 4,847 bacteria, and 236 of 237 lie above the bacterial 95th percentile.

The obvious alternative is that rare, poorly assembled genomes look unknown. We therefore matched each Thermoplasmatota representative 1:1 without replacement to a comparator on completeness, contamination, corrected genome size, N50, mean relative abundance and prevalence. Against other archaea, 218 of 237 found a match within one standardised unit; the matched pairs have median dark fractions of 0.308 and 0.048, a 6.5-fold difference, with the Thermoplasmatota genome the darker in 216 of 218 pairs. Against bacteria all 237 matched within half a unit, with residual imbalance below 0.05 standardised units on every covariate; medians 0.310 against 0.069, 4.5-fold, darker in 237 of 237 pairs (Fig. 3c). The contrast persists after this quality and abundance matching. In a separate comparison restricted to genomes whose first empty or digit-bearing label occurs at class rank, the 236 Thermoplasmatota representatives have median darkness 0.310, compared with 0.125 in 324 other representatives, a 2.5-fold contrast. These descriptive comparisons share phylogenetic dependence within lineages.

Thermoplasmatota contribute a small share of community abundance and a larger share of the attributable unknown repertoire. They are 4.4% of species and 0.8% of mapped reads (median across samples), yet carry 19.8% of the 77,069 single-phylum dark families and 15.2% of dark protein instances, an enrichment of 4.6-fold by families and 4.0-fold by proteins; Actinomycetota, half of the mapped community, carry dark families in proportion to their species share (1.1-fold).

These are not rare organisms. Thermoplasmatota representatives are detected in 481 of 549 samples, a median of 21 species per sample, at a median 0.39% of reads, and they are the cosmopolitan archaea of the sand sea where Halobacteriota are patchy (81 of 549 samples, in bulk where present). Depth is their axis: their share is 0.55% of reads in subsurface samples against

0.17% at the surface, higher below than above at 40 of 45 sites with both positions (Wilcoxon $p = 4 \times 10^{-7}$, unchanged after dividing by the mapped fraction), and indifferent to plant roots (26 of 53 sites, $p = 0.66$). Annotations in the remaining approximately 70% provide a provisional metabolic profile. Against Halobacteriota, other archaea and bacteria matched one to one on completeness, the 237 encode glycolysis (82%), a TCA cycle run through 2-oxoglutarate:ferredoxin oxidoreductase (96%), a glyoxylate cycle (79% against 35% in matched Halobacteriota), complex I and II, cytochrome c oxidase (59%) and an archaeal ATPase (91%), with no detected marker annotations for methanogenesis, ammonia oxidation or rhodopsin (0 of 237, against 51 of 269 Halobacteriota for rhodopsin). The annotations include more peptide-associated than sugar-associated uptake functions: branched-chain amino-acid transporters in 79% and sugar transporters in 2% (73% in matched Halobacteriota), an extracellular serine protease in 96%, endoglucanase in 79% and chitinase in 44%. They lack the carbon skeleton of branched-chain amino-acid synthesis (2%, against 91%) while keeping the terminal transaminase (93%), a pattern consistent with uptake dependence, and they have few annotations for cobalamin, thiamine or F420 synthesis and little catalase (29% against 85%). All carry an archaellum and 79% a CRISPR-Cas system. The smaller class E2 (2.9 against 3.8 Mb at the same completeness) has shed importers and several amino-acid pathways. These annotations suggest peptide and polysaccharide use in subsurface archaea. Pathway completeness, auxotrophy and rhodopsin absence require further validation with lineage-sensitive models; incomplete pathways and missing orthologues in these genomes cannot establish physiological absence. Approximately a third of their proteins remain without a UniRef50 match.

Whole-protein and local structural matches give different reference-coverage estimates

Sequence search misses remote homology, so the 1,466,943 genome-resolved dark representatives of at least 100 residues were screened by predicted structure (Methods). A 3Di search against the AlphaFold Database (v6, 241,070,489 entries) and the PDB (release 250101) left 997,646 (68.0%) with no hit in either, a triage rather than a result. Structural clustering of those candidates and admission at three or more members gave 17,221 fold families with one model each, 13,770 folded here with ESMFold and 3,451 reused from an earlier run. Only 2,838 models (16.5%) passed the confidence gate (mean pLDDT ≥ 0.70, ≥ 100 residues); the gate is on confidence.

Co-searched with them under the same command were 3,684 blinded controls: complete chains and deterministic fragments of experimentally solved PDB chains, matched to the candidates' length, composition and partiality and renamed before folding, of which 3,013 passed the same gate. Against both references, 1,031 of 2,838 dark models (36.3%) fell below a query-normalised TM-score of 0.5, compared with 101 of 3,013 controls (3.35%; Fig. 4a). Raw-hit inspection separates retrieval from scoring: 22 dark models and 66 controls returned no rows from either reference. Among models with a retrieved alignment, 1,009 of 2,816 dark models and 35 of 2,947 controls fell below the query threshold. This conditional comparison selects on retrieval and does not calibrate all search failures.

Alignment-normalised scores give a different result. Across all qualified models, 95 dark models and 68 controls fell below 0.5; among retrieved models, the counts were 73 of 2,816 (2.59%) and two of 2,947 (0.068%). Thus missing retrievals account for most control failures under the alignment-normalised score. Treating all failures as a single control rate obscures this difference. The query and alignment scores are independent maxima across reported hits and may come from different reference targets.

Local matches remain frequent under alignment-length gates. Requiring at least 50 aligned residues leaves 97 dark models below the alignment-normalised threshold. Of the 1,031 models below the query-normalised threshold, 797 have an alignment-normalised score of at least 0.5 over at least 100 residues and half the query length. These are models whose full length is poorly matched under the query-normalised metric; they can contain substantial familiar regions. At

query thresholds of 0.45 and 0.55, dark-model counts are 780 and 1,300, compared with 85 and 118 controls (Fig. 4b,c). Domain arrangement, extensions and uncertain inter-domain placement can affect this comparison. Confidence-qualified predictions alone do not validate a novel fold [18].

Pfam recognises a domain in 8.75% of the 4,880 member proteins of the 1,031 families and in 14.0% of those families, compared with 90.7% of the controls. Sequence and structural searches therefore provide different descriptions of reference coverage. The controls derive from PDB chains present in the references and serve as positive retrieval controls. Homology-excluded controls and domain-resolved analysis would be needed to calibrate remote homology and delimit unmatched structural regions.

Reference composition affects recovery of dark families

Two ceilings measure how far existing references reach into this proteome. The first is annotation: of the 19,186,853 protein-coding genes in the 5,438 representatives, 55.6% carry no KEGG orthologue, and the value is 53.9% in the near-complete representatives alone, showing that incomplete annotation persists in this subset. The second is prediction. In the 448 samples with both a genome profile and a 16S profile from the companion survey, marker-gene prediction agreed with genome-encoded content (median weighted nearest-taxon index 0.159, recall of genome-encoded orthologues 86.0%, Spearman ρ 0.839 on shared orthologues; Fig. 5a). Its reach is fixed by its reference: 2,517 of the 11,374 orthologues the genomes encode (22%) lie outside the 10,543 families it can emit, and in 420 of the 448 samples no orthologue inside that set went unpredicted, placing this component of the shortfall in reference coverage.

We next tested whether sequence-dark families have detectable homologues in additional catalogues. We searched the 106,042 dark family representatives together with the 106,042 matched UniRef50-matched families against MGnify90, GEM, SPIRE and five desert catalogues, 1.258 billion sequences, in one DIAMOND pass: 28.74% of dark families (30,479) had no hit in any of them, compared with 0.41% of the control (435) (Fig. 5b). Absence here is defined by the catalogue and the search rule, not by geography; MGnify90 biome labels carry no coordinates. What the control isolates is which reference is missing. The five desert catalogues together hold 147.5 million sequences, a fifth of MGnify90, yet leave 42.60% of dark families unexplained against MGnify90's 52.68%. Compared with an analytical expectation for random retention of MGnify90 sequences at the same size (a rarefaction model whose full-catalogue endpoint agrees with the observed MGnify90 absence to 0.15 percentage points), the desert catalogues leave 24.33 fewer points of dark families unexplained and SPIRE 21.30 fewer, while the same comparison for known proteins moves by about one point (Fig. 5b). The larger-release sensitivity search recovered a small additional fraction. Against the complete MGnify90 2026_07 release (1.66 billion sequences, searched with a different aligner, so a robustness check), 5.8% of the absent dark families gain a hit while 34.3% of the control's residual absences are resolved, leaving 27.08% of dark families unmatched. The absent families are not rare or local within the basin: their median occupancy is 37 of 61 sampled locations against 38 for families with a hit, and the medians range from 44 versus 50 locations in the shortest stratum to the values shown in Fig. 5c. The combined desert catalogue and SPIRE recover more dark families than the analytical MGnify90 expectation. Their redundancy, assembly basis and biome composition differ, so the comparison combines these properties. Searches of actual random subsets and matched non-desert soil catalogues would separate their contributions.

Exploratory functional and environmental associations are reported in Supplementary Note S1.

Discussion

This catalogue links reference-absent protein clusters to genomes and repeated sampling of a hyperarid environment. Evidence is strongest for the recurrent subset: 106,042 families hold 12.5% of genome-resolved dark representatives. Most evaluable families have median pairwise dN/dS below one, and the annotated frame usually has a lower estimate than the shifted frame. The annotated-frame contrast is evidence about reading-frame-dependent sequence patterns. Because most artifact-flagged families also have dN/dS below one, the estimate alone cannot establish coding status. Recurrence and recruitment describe the persistence of homologous DNA; expression measurements and overlap-aware gene validation would further resolve coding status and biological activity.

Darkness also varies strongly among lineages. Quality- and abundance-matched comparisons retain a large Thermoplasmatota contrast, while a comparison at the same placeholder rank gives a smaller 2.5-fold difference. Reference proximity and shared ancestry therefore remain relevant to the lineage pattern. Maeke et al. [19] examined the relationship between rarity and unknown-gene content in other Thermoplasmatota; direct searches against close relatives would test how much of the desert catalogue's darkness reflects representation in UniRef50.

The culture comparison directly exposes gene-call dependence: 343 protein-unmatched dark calls have nucleotide support in the culture assembly but no Bakta CDS covering half the aligned region. Including nucleotide support narrows the dark-versus-annotated sharing contrast. Gene prediction and annotation filtering therefore make a substantial contribution to the protein-level gap in these bacterial pairs; the residual gap remains sensitive to strain divergence, assembly and alignment thresholds. Re-calling both partners with one pipeline would distinguish these contributions further. The current control evidence also has limited campaign coverage: one no-template library was sequenced with expedition four, and the catalogue-level lineage and environmental summaries still include the flagged representatives. Sensitivity analyses across these summaries are required to assess their contribution.

The global family catalogues established that unknown families are numerous, span species, evolve under purifying selection and concentrate in uncultivated taxa [5, 6, 8], that restriction to one ecosystem is the norm (63.4% of 608,258 families; 9), and that structural surveys recover many familiar folds and domain combinations [10, 11]. Our 28.74% reference absence is unremarkable as a fraction, and 36.3% of confident models fall below the whole-query similarity threshold, while local similarity is detected much more frequently. The known-fold comparison provides a positive-control retrieval baseline; domain boundaries and remote-homology controls would refine the biological interpretation of the unmatched regions. This study links families to repeated sampling within one biome and measures a reference-composition effect: the desert-derived comparison explains 24 percentage points more of the dark families than the analytical E-value-corrected expectation at the same size, against about one point for the annotated control. SPIRE gives a 21.30-point gain and GEM a 4.27-point gain, while Atacama has 4.67 points more absence than expected. These differences motivate tests of reference composition, assembly basis and redundancy using matched sequence sets.

Supplementary functional profiles provide exploratory context for the genome resource. The companion amplicon comparison gives a median rank correlation of 0.839 on shared orthologues, alongside a reference-coverage limit: 55.6% of protein-coding genes have no KEGG orthologue assignment.

These measurements describe predicted gene content and DNA-based abundance. Incomplete genomes reduce sensitivity to gene presence, with consequences that depend on how genome quality varies among the compared communities; half the representatives are below 90% complete. The frame-shifted null calibrates whether the annotated frame is the selected one, not neutral evolution in general. What any dark family does is not measured here.

The two archaeal classes are culture and fold targets, and none of the 24 shared species remaining after contaminant exclusion already in hand as strains is archaeal. Strain-level persistence

across five expeditions is the next study on these reads.

Methods

Sampling, sequencing and environmental data

Soil was sampled on five expeditions along a west to east transect through the Saudi Rub' al-Khali from the western escarpment (19.15°N, 45.17°E) to the Omani border (20.71°N, 54.73°E), a cumulative path of 1,043 km with 60 numbered sites at predetermined coordinates and ten named locations. The expeditions ran on 17–20 March 2023, 14 July 2023 (restricted by heat to eight sites), 17–21 February 2024, 19–21 August 2024 and 12–15 October 2025. At each site and visit three soil positions were sampled in three co-located replicates: surface soil at 0 cm, subsurface soil at 5–10 cm, and soil from the root zone at 5–10 cm, termed root-adjacent because soil was not consistently attached to roots. Soil was held at −20°C in the field and at −80°C in the laboratory. Sampling design and field handling are shared with the companion 16S rRNA survey and field-data descriptor [4, 20]. The 551 soil samples with complete metadata (117 surface, 166 subsurface, 268 root-adjacent) are the cohort used throughout; they cover 59 of the 60 transect sites and two named locations, the 61 sampled locations used in occupancy statistics. Narrower cohorts are defined where used.

DNA was extracted from 0.9–1.2 g of soil with the QIAGEN PowerLyzer PowerSoil kit (12855-50) or the DNeasy PowerSoil Pro kit (47014), homogenised on a PowerLyzer 24, eluted in 50 μL and quantified with the Qubit high-sensitivity double-stranded DNA assay. DNA was fragmented on a Covaris instrument to 350 bp, purified with AMPure XP beads at 1.6:1, prepared with the QIAseq FX and NEBNext Ultra II kits and sequenced at KAUST in 150-nt paired-end mode on Illumina NovaSeq 6000 and NovaSeq X instruments. The submission comprises 554 runs for 551 samples: 221 NovaSeq 6000 runs and 333 NovaSeq X runs across four sequencing batches. The sample-to-run manifest records instrument, library kit and batch; three samples have two submitted runs. The exact commands combining those runs remain unverified, and sequencing-batch sensitivity remains unresolved for the exploratory environmental analyses. Reads were processed with fastp v1.0.1 [21] with base correction in overlapping pairs, a qualified-base threshold of Q20, at most 40% unqualified bases per read, a minimum length of 50 nt and a low-complexity filter at 20%; surviving pairs and orphaned reads were both retained. The median sample retained 21.58 Gbp (156.6 million reads; IQR 16.53–25.08 Gbp).

Soil chemistry and climate covariates are those of the companion survey restricted to the samples profiled here: X-ray fluorescence total elemental contents of archived soil (248 profiled samples with chemistry; 251 in the family-level cohort) and reanalysis climate series (453 and 456 samples). The retained laboratory XRF table is distinct from the fifth-expedition handheld field measurements. Units, instrument metadata and detection-limit coding require confirmation. Climate inputs were retrieved from the Open-Meteo archive as daily temperature and precipitation series over twelve-month windows centred on each visit; retained tables record the start and end dates. The provider's model selection and original API responses were not archived. The de Martonne index divides annualised precipitation by mean temperature plus ten. The environmental analysis did not include pH. The complete analysis table rebuilds byte-for-byte from the preserved metadata, laboratory and climate summary tables; regenerating the original climate retrieval is a separate provenance gap.

Assembly, binning, dereplication and taxonomy

Each sample was assembled with MEGAHIT v1.2.9 [22] (`meta-large` preset, contigs $\geq$ 500 nt, k-mers seen at least twice, pruning level 1, low-local-coverage ratio 0.3), supplying pairs and orphaned reads; 549 of the 551 samples produced an assembly (`51Sr1` produced no contigs and `F5Dr1` an empty file). Assembly statistics used QUAST v5.2.0 [23]. Every assembly was binned twice

with SemiBin2 v2.2.1 [24]: with the pretrained global model (`single_easy_bin --environment global --sequencing-type short_read`) on minimap2 [25] coverage of the paired reads, and with a self-supervised per-sample model (`single_easy_bin --self-supervised -m 1000`) on BWA-MEM [26] coverage of pairs and orphans. They produced 16,333 and 21,616 bins of at least medium quality, 37,949 in the union of the two unrefined binnings. The two bin sets were reconciled per sample with DAS_Tool v1.1.7 [27] at its defaults (score threshold 0.5, DIAMOND search); of 542 samples binned by both runs, 539 yielded refined bins, 14,426 in total. Quality was estimated with CheckM2 v1.1.0 [28] and tiered by completeness and contamination alone: near-complete at completeness $\geq 90\%$ and contamination $\leq 5\%$ (4,225), medium at $\geq 50\%$ and $\leq 10\%$ inclusive (8,209; two bins sit at exactly 10.0%), the rest below these thresholds (1,992). These operational tiers use CheckM2 alone. Subsequent barrnap v0.9 and tRNAscan-SE v2.0.12 [29] assessment of representatives found all three rRNAs in 157 genomes and at least 18 tRNA amino-acid types in 3,629. Full MIMAG classification gave 126 high-quality and 5,312 medium-quality representatives [30]. No chimerism screen was run on the 14,426 bins; GUNC v1.1.1 [31] against ProGenomes 3 was run on the ten order-gap candidates only.

The 12,434 bins of at least medium quality were dereplicated with dRep v3.6.2 [32] (fastANI [33], primary and secondary ANI 0.90 and 0.95, completeness $\geq 50\%$, contamination $\leq 10\%$, other parameters default; one bin was dropped internally), giving 5,438 species representatives (2,259 near-complete, 3,179 additional genomes meeting the medium-completeness thresholds). Representatives were classified with GTDB-Tk v2.7.2 [34] (`classify_wf`, ANI pre-screen on) against GTDB release 232 [35]; 1,898 classifications changed relative to an earlier run against release 226, and every reported taxon is from release 232. The first-empty-rank census assigns each representative to the first of the seven ranks from domain to species carrying an empty label; classes are mutually exclusive. The ten order-gap candidates with adequate marker coverage were placed on de novo trees (GTDB-Tk `de_novo_wf`) and assessed by relative evolutionary divergence against release-232 rank intervals; the eleventh candidate (46% of marker positions aligned) was excluded from adjudication.

Protein catalogue and the sequence-dark call

Genes were called on assemblies rather than on bins, so that the catalogue covers unbinned sequence; the 552 assembly files present at that step are a superset of the 549 canonical assemblies, a catalogue-level basis that enters no per-sample cohort. Prediction used Pyrodigal v3.7.1 [17, 36] (`-p meta --no-stop-codon`, partial genes retained): 753,305,533 proteins, clustered with MMseqs2 v18.8cc5c [37, 38] (`linclust --min-seq-id 0.9 -c 0.8 --cov-mode 0`) into 449,265,306 representatives, the unit counted in every dark rate. Every representative was searched against UniRef50 release 2026_01 [39] with `mmseqs search -s 7.5 -e 1e-3 --max-seqs 10`, a pure E-value criterion with no identity or coverage filter; a cluster is sequence-dark when no searched member returned a hit. A cluster is genome-resolved when any member lies on a contig assigned to one of the 37,949 unrefined bins of at least medium quality, tested over every member; this membership rule, not read mapping, defines the 43,423,322 genome-resolved representatives and the 4,292,446 dark ones among them. Functional annotation of the 5,438 representatives used Bakta v1.12.0 [40] for gene calling and eggNOG-mapper v2.1.13 [41] with DIAMOND v2.0.15 [42] against the v2.1.12 reference data for KEGG orthologues. Representatives that refinement emitted as strict contig subsets of a parent bin were annotated by restricting the parent's gene calls to their own contigs; the resulting table (`rep_ko_v2`) has no empty rows.

Read recruitment and detection

Two recruitment runs underlie the paper. The representative-level run mapped each sample's paired reads to the 5,438 representatives with CoverM v0.7.0 [43] in genome mode (`-p minimap2-sr -m relative_abundance`, all other parameters default). The default minimum

covered fraction of 10% defines detection: a representative is detected when its relative abundance exceeds zero, that is, when at least a tenth of its genome is covered. This is an inherited default, not a chosen threshold. The census and stress analyses use the frozen 548-sample profile cohort available when those analyses were run; subsequent profiles are outside that cohort. The later Thermoplasmatota ecology analysis used a separate 550-column snapshot: F13PRr1 was absent and F5Dr1 contained no usable abundance values, leaving 549 samples. Consequently its surface/subsurface comparison has 45 paired sites, whereas the earlier stress analysis has 44. These are analysis-specific frozen cohorts. Community values use two denominators: the share of all reads (organisms outside the catalogue counted as non-carriers, a lower bound) and the share of reads mapping to any representative (median unmapped fraction near 51%). The member-aware run mapped the reads of all 551 samples to the 536,437 dark-family member genes in contig mode (`-m tpm covered_fraction`). A family is present when any member gene reaches a covered fraction of 0.5 and its abundance is the summed member TPM. That rule is easier to satisfy for a short gene and reverses the length to occupancy relation when replaced by any-read presence, so occupancy is compared within length strata. The two shotgun control libraries sequenced with the fourth expedition, a no-template extraction control (335,190 read pairs after filtering) and a positive control of *Hafnia paralvei* (119,683,914 pairs; GTDB-Tk R232 on its assembly, ANI 98.99%), were profiled with the identical CoverM invocation and index. Representatives detected in the no-template control are annotated as reagent contaminants in the catalogue table and excluded from the pathogen census, which is therefore not reported; the no-template assembly (11 contigs, 11 kb) was identified by UniRef50 search of its predicted proteins.

Dark families, matched control and selection

The 4,292,446 genome-resolved dark representatives were clustered with MMseqs2 `easy-cluster` (`--min-seq-id 0.5 -c 0.8 --cov-mode 0`, default sensitivity 4) and each cluster mapped through the 90%-identity membership table to the samples and sites in which its instances were assembled. A cluster was admitted as a dark family when it held at least three distinct 90%-identity representatives assembled in at least three samples and two sites (106,042 families, 536,437 member genes; 2,784 three-member clusters failing the spread gate were dropped because they failed the spread criterion). Member nucleotide sequences were excised from the assemblies using the prediction coordinates. The control pool comprised every non-dark representative on a contig of the same 37,949 bins, with darkness excluded against the assembly-wide dark set. Because dark representatives are short, the pool was sampled per 10-residue length bin to reproduce the dark length distribution (seed 20260813; the sample ran 12.5% short of target, in the 20–39 residue bins), clustered identically, and control families were drawn to match dark families by size bin (3–9 exactly, 10–19, 20 or more), one per dark family. Two asymmetries are disclosed: control families were admitted on member count alone, without the sample and site gate, and after admission the dark query set is longer than the control (median 105 against 75 residues). These differences, together with the divergent dS distributions, affect comparability; their direction and size depend on the endpoint. Family admission uses the unrefined bins; phylum attribution traces each member through its contig to the refined bin, the dereplicated representative and its release-232 classification, which attributes 78,727 families.

For each family with at least three usable coding sequences ($\geq$ 90 nt, translatable under table 11 without internal stops, protein $\geq$ 25 residues), proteins were aligned with MAFFT v7.526 (`--auto`) [44] and back-translated to codons. Pairwise dN and dS were computed by the Nei–Gojobori method [45] (Biopython v1.87 [46]) over at most 80 deterministically thinned pairs per family, retaining pairs with at least 30 codons after removal of gapped or ambiguous positions, excluding identical pairs and requiring $0 < \mathrm{dS} \leq 2$. The family statistic is the median pairwise dN/dS (97,200 dark and 95,647 control families). For the joint sensitivity comparison, we extracted median coding-sequence length and GC fraction from all indexed family FASTA sequences of at least 90 nt, before the usable-sequence filters for dN/dS. We formed strata us-

ing mean dS boundaries 0, 0.1, 0.25, 0.5, 1 and 2; coding-sequence length boundaries 0, 60, 100, 200 and 400 amino-acid equivalents; GC bins of width 0.05; and aligned sequence-count boundaries 0, 5, 10 and 20. The final length and count bins were open-ended. Within each stratum represented in both arms, we sampled the smaller arm's count from each arm without replacement (seed 20261013). We repeated selection for seeds 20261013–20261032 and reported covariate balance and median differences. Seed ranges measure selection sensitivity, not sampling uncertainty. Divergence-stratum median differences used 1,000 independent family bootstrap replicates per arm (seed 20261014), with 2.5th and 97.5th percentiles. These intervals condition on the observed strata and assume family independence. Strata were left-closed; the final dS bin included 2. Weighted mean contrasts averaged within-stratum mean differences using dark-arm counts, control-arm counts or the smaller arm count. This comparison conditions on the retained common-support families; shared ancestry and the original family-construction process remain sources of dependence.

The null arm reads the same member genes in the +1 frame: the first nucleotide is dropped, the sequence trimmed to a codon multiple, every codon translating to a stop under table 11 masked to NNN (removed by the existing pairwise gap and ambiguity stripping), and all other filters left identical. It was run on a seeded subset of 10,000 families with at least three indexed coding sequences (seed 20260920), together with the same input subset in the annotated frame. Stop codons were masked in both calibration arms. An input audit found terminal stops in 43,953 of 50,898 sequences and internal stops in 21 sequences from 16 families. The null, annotated-frame and paired output sets contain 9,148, 9,182 and 9,029 families, respectively. A shifted frame inherits the composition and substitution pattern of the real gene, so this null tests whether the annotated frame is the selected one, not neutral evolution in general. Expedition persistence counts expeditions with presence under the member-aware rule; re-assembly persistence counts expeditions in which an instance was independently assembled. Threshold robustness reclustered the same representatives at 30% identity.

AntiFam v8.0 (278 profiles, release 12 October 2023) was searched with HMMER `hmmsearch --cut_ga` against all genome-resolved dark representatives and the control protein pool. Matches flagged 3,385 dark proteins, including 141 members of admitted families. A family was flagged when any admitted member matched. For the overlap audit, original Pyrodigal FASTA-header coordinates were verified for all 536,437 admitted members, then intersected with opposite-strand calls from the same sample and contig. A candidate required overlap covering at least 50% of the dark call. The opposing call's darkness was resolved through the complete 90%-identity cluster membership and assembly-wide dark-ID tables. This audit is conditional on the gene caller retaining both ORFs. The original Prodigal algorithm limits opposite-strand overlaps to 200 bp between 3' ends and excludes 5'-end overlap [17]; the largest observed co-retained overlap here was 195 nt. The audit therefore has little power for alternatives suppressed by the caller. Sensitivity exclusions used the union of family flags.

Cultured-genome comparison

Long-read genomes recovered from 722 cultured assemblies from saline pans, oases and campground soils of the same basin (companion pangenome catalogue) were compared with the 5,438 representatives by skani v0.3.1 [47] (`sketch`, `search -s 75`); a shared species required ANI $\geq$ 95% and aligned fraction $\geq$ 30%. The Bakta proteomes of all 722 cultured genomes (3,024,784 proteins) were searched against UniRef50 release 2026_01 with the identical MMseqs2 rule, so cultured darkness is per protein under the same call as catalogue darkness, which is per cluster representative. The 4,292,446 dark representatives, together with the 106,042 dark and 106,042 matched control family representatives of the reference-absence search, were searched against the cultured proteomes with DIAMOND v2.0.15 (`blastp --very-sensitive -e 1e-5 -k 500`), a homolog requiring reciprocal query and subject coverage $\geq$ 50%. Because several cultured strains map to one representative, the rate comparison was paired by species: one cultured aggregate

against one representative, each cultured genome assigned to its best-ANI representative, giving 23 pairs from the 25 shared species, before contaminant exclusion. We excluded representative `V42PRr2__single_SemiBin_33`, identified as *E. coli* in the no-template control, leaving 22 groups for the paired rate comparison by Wilcoxon signed-rank. Protein sharing used the best-ANI cultured genome of each representative and reciprocal best DIAMOND hits (`--very-sensitive -e 1e-5 -k 5`), requiring at least 50% identity and 50% coverage of both proteins. Sharing percentages pool protein counts across the 22 pairs. The complete-call sensitivity subset contains 166 dark MAG proteins of at least 100 residues. Gene callers differ (Bakta on cultured genomes, Pyrodigal on assemblies).

For the nucleotide sensitivity search, we extracted each MAG gene's recorded interval from its dereplicated representative assembly and retained intervals of at least 90 nucleotides. Each was aligned to the paired culture-derived assembly using minimap2 v2.17-r943-dirty (`-c -x sr -k 11 -w 5 -n 2 -m 20 --secondary=yes -N 20`). Support required a single alignment spanning at least 90% of the query with at least 95% nucleotide identity, calculated as matching bases divided by alignment block length. Counts were stratified by darkness and reciprocal protein-hit status. Input assemblies, gene-coordinate tables and protein-search outputs are identified in the analysis checksum manifest. Recovery of the protein-shared subset provides a positive comparison for this search's sensitivity. Per-pair recovery and a post hoc restriction to pairs with at least 90% recovery were reported. All qualifying alignments were checked at identity thresholds 0.85, 0.90 and 0.95. For all DNA-supported calls, culture Bakta CDS coordinates were intersected with each alignment's target span. A CDS had to cover at least half that span; orientation and phase at the first aligned position defined a diagnostic category, with same-orientation phase compatibility taking precedence if several alignments qualified. Eligible dark calls were traced through the complete 90%-identity cluster table into the admitted-family membership table. Indels can change phase downstream, so this is not a full coding-sequence correspondence test.

Lineage darkness and the matched null

For each representative, every protein on its contigs was traced to its 90%-identity cluster and scored dark or not by the cluster call above; the dark fraction is the share of dark protein instances, computed on the refined winner contigs of all 5,438 representatives (native key, no renaming). Ordinary least squares (statsmodels v0.14.2) modelled the dark fraction on assembly quality (completeness, contamination, $\log_{10}$ N50, $\log_{10}$ length), then adding phylum and genus (genera with fewer than five representatives pooled; unnamed genera one level); unique and shared variance are differences of R^2 between nested models. Placeholder-name depth was coded as the first rank with an empty or digit-bearing label, separately from the taxonomy census of empty ranks and added to the quality model before phylum. No phylogenetic mixed model was fitted, so the representatives enter as independent draws. Genome size was completeness-corrected length. The archaeal contrast is a 1:1 nearest-neighbour match without replacement on standardised completeness, contamination, $\log_{10}$ corrected size, $\log_{10}$ N50, $\log_{10}$ mean relative abundance and prevalence (samples detected), against two pools reported separately, other archaea and bacteria, at calipers of 0.5 to 2.0 standardised units (0.25 also against bacteria); residual imbalance is reported as standardised mean differences. Dark-family attribution by phylum uses the 78,727 attributable families; enrichment is the phylum's share of single-phylum dark families divided by its share of species, and its share of dark protein instances divided by its share of all protein instances. The class placement of the Thermoplasmatota representatives was displayed on an approximate-maximum-likelihood tree (FastTree, LG with gamma; 48) of the GTDB-Tk release-232 archaeal marker alignment, with the 1,647 release-232 Thermoplasmatota species representatives and 30 Halobacteriota and Thermoproteota outgroups; the tree supports no branching claim.

Structural analysis

Queries were the 1,466,943 genome-resolved dark representatives of at least 100 residues. 3Di strings were predicted with ProstT5 [49] through Foldseek [50] (`createdb --prostt5-model`) and searched (`--alignment-type 2 -s 9.5 -e 1e-3 --max-seqs 50`) against the AlphaFold Database v6 on UniProt 2025_03 (241,070,489 entries) [51] and the PDB release of 1 January 2025 (324,204 chains in the Foldseek database) [52]. Queries with no hit in either were clustered in structure space (`cluster --single-step-clustering --cluster-mode 2 -s 4`) and clusters of at least three members admitted as fold families (17,221). One model per family was folded with ESMFold [53] (`esmfold_v1`, transformers v4.40.2, PyTorch v2.2.2, half-precision trunk, sequences truncated at 1,022 residues, four recycles), except that 3,451 families reused a member's existing ESMFold model from an earlier candidate set; those reused representatives come from a novelty-enriched prior cohort and their dual no-hit rate is higher (42.8% against 35.3% for models folded here), and the reported rate holds when they are excluded. Models were screened as coordinates (`--alignment-type 1 -e 10 -s 9.5 --max-seqs 50`) against the same references. 3Di prediction, triage and clustering ran on a GPU build of Foldseek (commit `8dc75c74`); the validation screen ran on the packaged build `10-941cd33`, with candidates and controls in one query set under one command. A model enters at length ≥ 100 and mean pLDDT ≥ 0.70; a dual no-hit is a query-normalised TM-score below 0.5 against both references [54], the alignment-normalised score is reported alongside, and a model returning no row scores zero. The 3,684 blinded controls are 1,777 complete PDB chains and 1,907 deterministic crops matched to an earlier candidate set on length, composition and partiality, renamed before folding by the same procedure; 3,013 passed the gate. Retrieval failures were counted separately from retrieved alignments with scores below threshold. The primary comparison retains all qualified models; a secondary comparison conditions on at least one returned alignment. For each query, maxima under the two normalisations were computed independently across both references. Sensitivity searches of the retained raw-hit tables imposed at least 50 aligned residues, or at least 100 residues together with 50% query coverage.

The 4,880 member proteins of the candidate families, the 1,807 other confident dark representatives and the 3,013 gate-passing controls were searched in one run against Pfam-A 38.2 with HMMER v3.3.2 (`hmmsearch --cut_ga`); per protein we recorded the number of domains, the fraction of the sequence covered by Pfam envelopes and the longest uncovered stretch.

Reference ceilings and absence from catalogues

The orthologue ceiling is one minus the pooled ratio of Bakta protein-coding genes carrying a KEGG orthologue to all Bakta protein-coding genes over the 5,438 representatives (19,186,853 genes); the CheckM2 gene-call denominator gives 58.3%. Predicted profiles are PICRUSt2 v2.4.1 [55] KEGG-orthologue predictions from the companion V3–V4 amplicon survey (10,543 families, 1,270 profiles, byte-identical to that study's input) with weighted nearest-sequenced-taxon indices. Genome-encoded profiles join the orthologue table to the CoverM profiles, presence being relative abundance above zero; 448 samples carry both profiles (four amplicon duplicates summed and renormalised; no amplicon-side depth or NSTI filter was applied, and 29 profiles return fewer than the full family set). Per sample we report presence-recall of genome-encoded orthologues, Spearman correlation over orthologues both methods report, and the count of genome-encoded orthologues outside the predictor's emitted set; the last is a reference-vintage gap, not prediction error.

Reference absence searched the 106,042 dark and 106,042 control family representatives in one file with DIAMOND v2.0.15 (`blastp --sensitive --evalue 1e-3 -k 1`) against MGnify90 (717,738,164 cluster representatives [56]), GEM (122,978,191 proteins, 52,515 genomes [57]), the SPIRE species representatives (269,496,580 proteins from 107,008 genomes, Prodigal v2.6.3 `-p meta` [58]) and the gene calls of five desert metagenome studies assembled in SPIRE (147,531,178 proteins: Atacama PRJNA484015, 6,193,359; desert trace-gas autotrophy PR-

JNA657906, 7,200,701; JGI arid soils, 30,569,562; JGI Antarctic cold desert, 49,206,550; JGI Mojave, 54,361,006). A representative with no reported hit is absent under this rule. Searching the desert catalogues separately rather than concatenated returns 2,544 further families (2.40%), so cross-catalogue ordering is descriptive and the content-versus-depth comparison rests on a rarefaction. Both query sets were searched once against MGnify90 at `--evalue 1 -k 500 --max-hsps 1`. Under random retention of subjects with probability p a hit remains reportable at the 10^{-3} rule only if $E \leq 10^{-3}/p$, so the probability of no reportable hit is $(1-p)^{h(p)}$ over the $h(p)$ subjects meeting that bound, and expected absence is its mean over families at p matched to each catalogue's size. The `-k` cap inflates the expectation by at most $(1-p)^{500}$, about 1.3% at the smallest catalogue. The depth check searched the absent families against the complete MGnify90 2026_07 set (1,661,142,385 sequences) with MMseqs2 `search -s 7.5 -e 1e-3 --max-seqs 10`; it uses a different aligner and release and is never unioned with the DIAMOND sets. Occupancy of the absent families uses the member-aware presence matrix over the 61 sampled locations, within length strata.

Environmental structure and stress censuses

Family abundance (TPM) over the 547 environmentally annotated samples was Hellinger-transformed and compared by Bray–Curtis dissimilarity, excluding families present in fewer than five samples (105,977 retained). Each block was tested by distance-based PERMANOVA [59] with 999 permutations of samples: soil position (dummy-coded), space (standardised latitude and longitude), climate (temperature, precipitation, aridity index; $n = 456$) and geochemistry (eleven standardised axes: S, Cl, Na, Ca, K, Fe, Mn, V, Si, Ti and a summed salinity proxy; $n = 251$), the latter two also conditioned on space and on space and position. Every block was tested twice, with permutations free over samples and restricted within site (59 sites); each environmental block kept $p \leq 0.002$ under the restricted test, geography is untestable within site by construction, and the vanadium control stayed null (0.88), so nesting does not carry the result. Sequencing depth ($\log_{10}$ filtered bases per sample, from the read-filtering reports) explains 1.1% of composition and leaves the geochemistry share beyond space unchanged (0.089). Variance partitioning [60] over position, space and geochemistry was run on the 251-sample chemistry cohort (499 permutations); a Mantel test [61] compared Bray–Curtis with great-circle distance. A single vanadium axis run through the identical PERMANOVA is the negative control. At family level, abundance was correlated by Spearman rank with each of thirteen elements, the aridity index and a plant-root contrast over families present in at least 25 samples of the axis subset, Benjamini–Hochberg corrected within axis; the five most associated axes were re-tested as partial Spearman controlling ranked latitude and longitude. The salinity proxy (unweighted Na + Cl + S) retained 41% of its associations under conditioning and is not used as a family-level axis.

Stress censuses screened a 28-orthologue DNA-damage panel in eight groups and a 13-orthologue compatible-solute synthesis panel (Supplementary Table S3) against the orthologue table, prevalences being fractions of the 5,438 representatives. Community values per sample used occurrence, whole-community and mapped-community metrics over the 548-sample cohort; the unmapped fraction does not differ between surface and subsurface (54.3% against 53.7%, Mann–Whitney $p = 0.50$), and root-adjacent comparisons use the mapped denominator. Surface against subsurface was tested on site medians by Wilcoxon signed-rank over the 44 sites with both positions profiled, Holm-corrected within metric; effects were re-tested within each of the eight most abundant phyla (a phylum tested at ≥ 15 paired sites with ≥ 15 detected species per sample) and interpreted only if the direction replicated, which the mismatch-repair contrast failed. Within-species differentiation compared surface and subsurface member genomes of the 327 species recovered from both positions (324 with annotations) by Wilcoxon signed-rank. Gradient tests were partial Spearman on ranked latitude and longitude against five salt axes ($n = 248$) and three climate axes ($n = 453$), Benjamini–Hochberg corrected within block; the

Halobacteriota association was re-tested on centred log-ratio fractions and on detection.

Differential encoded function by soil position used the carriage of each feature, the share of the mapped community whose representatives encode it, over the 549 profiled samples: 282 KEGG modules in DRAM's step form (present when at least two thirds of steps are encoded; 117 modules absent everywhere or universal were excluded) and the 4,127 orthologues at 5 to 95% prevalence. Replicates were collapsed to the median per site, position and visit; root-adjacent minus bulk (the median of surface and subsurface at the same visit) was taken at 52 sites and surface minus subsurface at 37, each difference then medianed over visits per site. A feature passed at Benjamini–Hochberg $q < 0.05$ within basis and contrast, an absolute effect of at least 5 percentage points, and the same sign when carriage was recomputed within Actinomycetota alone and within Pseudomonadota alone (samples where the phylum held under 1% of mapped abundance blanked); within-phylum magnitudes are reported beside each effect. One-step modules are single orthologues or any-of sets and are named by enzyme. Size normalisation (per Mb of completeness-corrected genome) and restriction to representatives at least 90% complete were run as sensitivity analyses. Chemistry and expedition contrasts were computed but are not reported, the former because X-ray fluorescence coverage differs by expedition and the latter because only eight site pairs qualify.

Biosynthetic and resistance censuses

Biosynthetic gene clusters were detected with antiSMASH v8.0.4 [62] in minimal mode on the 5,438 genomes, counting per-species presence of a product type; 85.3% of 44,002 regions abut a contig edge, so completeness and novelty are not estimated. Resistance and stress determinants used AMRFinderPlus v4.2.7 (database 2026-05-15.1, plus set, no organism option, so no point mutations) [63]. Because the no-template control assembled into a laboratory plasmid carrying aph(3')-Ia, a TEM β-lactamase and a chloramphenicol acetyltransferase, resistance hits were checked against contigs matching that plasmid: two aph(3')-Ia hits sit on such contigs, and aph(3')-Ia occurs in seven species in all, five of them on an identical 7,838-bp contig at 100% identity. Removing every aph(3')-Ia hit lowers aminoglycoside prevalence from 11.9% to 11.8%; no other reported class moves. Ectoine synthesis calls from the orthologue panel and from antiSMASH were compared as two independent methods. None of these censuses has a matched genome set or size normalisation and they are reported as description; the pathogen census of the representatives is not reported because two of its hits are reagent contaminants.

Statistics

Tests are two-sided and rank-based: Wilcoxon signed-rank for paired designs, Mann–Whitney U for two groups, Kruskal–Wallis for more. Associations are Spearman or partial Spearman on ranked coordinates. Multiple-testing control is applied within named blocks (Holm across traits within a metric [64]; Benjamini–Hochberg within an axis or block [65]) and never across the paper. Confidence intervals for proportions are Wilson intervals. Stochastic steps are seeded. Estimators used NumPy v1.26 [66] and SciPy v1.11 [67]. The independent unit is the species representative for genome-level statistics, the family for family-level statistics, the site for paired compartment contrasts, and the sample for PERMANOVA, where non-independence within site is acknowledged above.

Supplementary Note S1: exploratory functional and environmental profiles

These analyses provide descriptive context for the catalogue. Laboratory XRF units, detection limits and the meaning of recorded zeros remain undocumented. Family-level sample correlations reuse sites and visits; relative-abundance effects, sequencing batch and extraction kit remain

potential confounders. An exploratory sensitivity analysis aggregated the 251 chemistry samples to 59 sites and retained the 94,568 families present in at least 25 measured samples and five sites. Within-site mean TPM and median chemistry gave 59,120 sulfur, 49,387 calcium, 48,567 chloride and 12,054 iron associations at 5% FDR. This changes the observational unit but leaves spatial, compositional, batch and measurement concerns unresolved. The larger sulfur count after aggregation can reflect between-site structure and is not evidence that spatial confounding has been removed. The tests below do not establish chemical exclusion or physiological responses.

Encoded function shifts with soil position, mostly by lineage turnover

The annotated fraction of the catalogue carries the toolkit expected of a hyperarid soil (Extended Data Table 1). DNA-damage repair is near-universal (base and nucleotide excision repair in 99.6% and 99.5% of species) and compatible-solute synthesis is encoded by 79.5%. Biosynthetic gene clusters occur in 97.5% of species but 85.3% of the 44,002 regions abut a contig edge, and acquired resistance determinants occur in 37.4%, led by β-lactam (24.6%) and arsenic (17.3%) determinants, the latter rising with calcium and falling with precipitation after conditioning on geography.

Encoded-function contrasts were evaluated by soil position. For 282 KEGG modules and 4,127 orthologues we compared the share of the mapped community encoding each feature between paired samples of one site visit, root-adjacent soil against bulk soil at 52 sites and surface against subsurface at 37, and kept a feature only if it cleared a 5-percentage-point floor at 5% false-discovery rate and kept its sign when recomputed within Actinomycetota alone and within Pseudomonadota alone. Fourteen modules pass beside roots. The root-adjacent community encodes more cytochrome bd oxidase (+13.0 points; +17.0 within Actinomycetota, +1.9 within Pseudomonadota), serine-to-cysteine synthesis (+11.0), the Entner-Doudoroff pathway (+9.4), heme synthesis (+6.9 and +7.2 by the two routes) and glutathione synthesis (+6.6), and less F420 synthesis (−10.1), complex I (−7.8) and its NDH-1-type relative (−10.3). Nine modules pass at the surface, where putrescine synthesis (+8.7), formaldehyde assimilation (+8.7) and acetate kinase-phosphotransacetylase (+7.0) rise and F420 synthesis again falls (−10.1); photolyase (+8.8) and glycine betaine transport (+10.4) rise at the surface. The within-phylum effect is a median 0.16 of the whole-community effect, so the shifts ride mostly on lineage turnover; the mapped fraction is 5.6 points higher beside roots, so root-adjacent values describe the mapped half of the community; and no sign reverses when the analysis is restricted to representatives at least 90% complete. These patterns describe differences in encoded capacity and community composition.

Exploratory environmental associations

We explored associations between environmental covariates and the distribution of dark families. Environmental covariates were associated with repertoire composition under the fitted models. Across the 547 samples with environmental annotation, composition of the 106,042-family repertoire (105,977 families present in at least five samples) is structured by soil position (PERMANOVA $R^2 = 0.044$) and by geographic position (0.079), and communities decay with distance (Mantel $r = 0.341$). In the 251 samples with X-ray fluorescence chemistry, eleven geochemical axes explain 13.2% of compositional variance, of which 8.9% survives conditioning on latitude and longitude and 9.2% conditioning on position as well; climate (temperature, precipitation, aridity index, $n = 456$) explains 3.0% beyond space. Partitioning the three blocks on the chemistry cohort gives geochemistry 0.106 of unique variance against 0.055 for space and 0.057 for position. Every environmental block holds when permutations are restricted within site ($p \leq 0.002$; geography itself cannot be tested that way) and when sequencing depth is conditioned out, and the same design run on a single vanadium axis returns $R^2 = 0.003$, $p = 0.78$, under unrestricted permutations (within-site restricted $p = 0.884$). Each R^2 is a share of compositional variance across families, not of any one family's abundance.

At family resolution, 94,568 families were testable against the chemistry axes and 100,293 against aridity. Sulfur associates with 54,227 families at 5% false-discovery rate, calcium with 47,174, chloride with 46,972, iron with 24,110 and the aridity index with 23,031; in each case 90 to 96% of associations are negative, a pattern in relative abundances whose interpretation depends on compositionality and non-detect handling. Vanadium, molybdenum, nickel, cobalt, manganese and zinc associate with no family. Conditioning on geography retains 82 to 86% of the sulfur, chloride and aridity associations and 71% of the calcium ones; a summed salinity proxy retains 41% and is not used as an axis. Aridity is associated with relative family abundance under this model, although the index remains within the hyperarid class along the transect.

Soil-position associations differ among the phyla to which families are attributed. Of the 102,005 dark families testable against the root-adjacent contrast, 19,818 are more abundant beside roots and 40,495 less, and this is the least spatially confounded axis in the panel: 94.1% of the associations survive conditioning on geography, against 71 to 86% for the chemical axes. The root-depleted families belong to the desert core, 52.7% Actinomycetota; the root-enriched families do not. Actinomycetota fall to 16.8% of them and Thermoplasmatota to 11.2%, while Pseudomonadota (19.2% against 9.6% of all attributable families), Myxococcota (11.3% against 4.9%), Planctomycetota (8.3% against 2.4%), Bacillota, Bdellovibrionota, Cyanobacteriota and Verrucomicrobiota are two to four times over-represented. Both sets have higher unadjusted five-expedition detection frequencies than the average dark family (86.8% and 89.8% detected in all five expeditions against 79.4%). Beside desert plants the unknown proteome shifts from the phyla that own it in bulk sand to lineages known from rhizospheres elsewhere, and the detection-frequency comparison is conditional on prevalence and the mapping rule. Effects are modest (median ρ of $+0.15$ and -0.19), no host plant is identified in the metadata, and root-adjacent soil was not consistently attached to roots.

The annotated fraction sorts along the same axes, in two modes. Soil-position contrasts include the following: across 44 sites with paired surface and subsurface samples, the share of the mapped community carrying photolyase is 24.3% at the surface against 12.1% below, carotenoid synthesis 70.3% against 61.0%, and non-homologous end joining runs the other way (80.0% against 85.7%) . The photolyase shift replicates in the same direction inside each of the three phyla with enough paired sites to test, and a within-species test across 324 species present in both positions is null (all $p > 0.17$), consistent with species turnover in this comparison. Elemental associations include the following: after conditioning on geography, the share of Halobacteriota rises with sulfur ($\rho = 0.451$, $p = 7 \times 10^{-14}$, $n = 248$) and chloride ($\rho = 0.385$), robust to centred log-ratio and to detection-only coding, while the near-universal DNA-repair floor moves with no gradient. These exploratory associations motivate analyses that account jointly for batch, spatial dependence, compositionality and measurement limits.

Data and code availability

Successful private ENA submission receipts map 554 runs from 551 soil samples to study PRJEB125153. The accession map is retained in the review evidence package; the public filereport returned zero runs on 13 October 2026. The recorded release date is 1 November 2026; public release or reviewer access remains to be confirmed before submission. The 14,426 refined bins, 5,438 representatives, analysis inputs and raw outputs remain on IBEX at the paths in the provenance inventories. A MAG deposition with MIMAG metadata and an archived release DOI are pending. The code repository is https://github.com/bio-ontology-research-group/empty-quarter-mags; this review revision, its compact evidence tables and figures are held locally and on IBEX and have not yet been published there. The compact driver reproduces the stated table calculations and figures; complete raw-to-final reproduction is limited by 78 missing raw-file paths, 302 permission failures, unverified duplicate-run merge commands and incomplete historical claim-registry migration. These are documented submission prerequisites. Reference database releases are named where they are used: UniRef50 2026_01, GTDB release

232, AlphaFold Database v6 on UniProt 2025_03, PDB release 250101, MGnify90 and MGnify90 2026_07, GEM, SPIRE, and the five desert studies listed in the Methods.

References

[1] Jianping Huang, Haipeng Yu, Xiaodan Guan, Guoyin Wang, and Ruixia Guo. Accelerated dryland expansion under climate change. *Nature Climate Change*, 6(2):166–171, 2016. doi: 10.1038/nclimate2837.

[2] Fernando T. Maestre, Manuel Delgado-Baquerizo, Thomas C. Jeffries, David J. Eldridge, Victoria Ochoa, Beatriz Gozalo, José L. Quero, Miguel García-Gómez, Antonio Gallardo, Werner Ulrich, Matthew A. Bowker, et al. Increasing aridity reduces soil microbial diversity and abundance in global drylands. *Proceedings of the National Academy of Sciences USA*, 112(51):15684–15689, 2015. doi: 10.1073/pnas.1516684112.

[3] Noah Fierer, Pok Man Leung, Rachael Lappan, Raphael Eisenhofer, Francesco Ricci, Sophie I. Holland, Nicholas Dragone, Linda L. Blackall, Xiyang Dong, Cristina Dorador, Belinda C. Ferrari, Jacqueline Goordial, Susan P. Holmes, Fumio Inagaki, Tal Korem, Simone S. Li, Thulani P. Makhalanyane, Jessica L. Metcalf, Niranjan Nagarajan, William D. Orsi, Erin R. Shanahan, Alan W. Walker, Laura S. Weyrich, Jack A. Gilbert, Amy D. Willis, Benjamin J. Callahan, Ashley Shade, Julian Parkhill, Jillian F. Banfield, and Chris Greening. Guidelines for preventing and reporting contamination in low-biomass microbiome studies. *Nature Microbiology*, 10(7):1570–1580, 2025. doi: 10.1038/s41564-025-02035-2.

[4] Rund Tawfiq et al. Landscape-scale bacterial biogeography across the Rub' al-Khali reveals recurring spatial and soil-position structure. Companion manuscript in preparation, 2026.

[5] Georgios A. Pavlopoulos, Fotis A. Baltoumas, Sirui Liu, Oguz Selvitopi, Antonio Pedro Camargo, Stephen Nayfach, Ariful Azad, Simon Roux, Lee Call, Natalia N. Ivanova, I-Min A. Chen, David Paez-Espino, Evangelos Karatzas, Ioannis Iliopoulos, Konstantinos Konstantinidis, James M. Tiedje, Jennifer Pett-Ridge, David Baker, Axel Visel, Christos A. Ouzounis, Sergey Ovchinnikov, Aydin Buluc, and Nikos C. Kyrpides. Unraveling the functional dark matter through global metagenomics. *Nature*, 622(7983):594–602, 2023. doi: 10.1038/s41586-023-06583-7.

[6] Chiara Vanni, Matthew S. Schechter, Silvia G. Acinas, Albert Barberán, Pier Luigi Buttigieg, Emilio O. Casamayor, Tom O. Delmont, Carlos M. Duarte, A. Murat Eren, Robert D. Finn, Renzo Kottmann, Alex Mitchell, Pablo Sánchez, Kimmo Siren, Martin Steinegger, Frank Oliver Glöckner, and Antonio Fernàndez-Guerra. Unifying the known and unknown microbial coding sequence space. *eLife*, 11:e67667, 2022. doi: 10.7554/eLife.67667.

[7] Luis Pedro Coelho, Renato Alves, Álvaro Rodríguez Del Río, Pernille Neve Myers, Carlos P. Cantalapiedra, Joaquin Giner-Lamia, Thomas S. B. Schmidt, Daniel R. Mende, Askarbek Orakov, Ivica Letunic, Falk Hildebrand, Thea Van Rossum, Sofia K. Forslund, Supriya Khedkar, Oleksandr M. Maistrenko, Shaojun Pan, Longhao Jia, Pamela Ferretti, Shinichi Sunagawa, Xing-Ming Zhao, Henrik Bjørn Nielsen, Jaime Huerta-Cepas, and Peer Bork. Towards the biogeography of prokaryotic genes. *Nature*, 601(7892):252–256, 2022. doi: 10.1038/s41586-021-04233-4.

[8] Álvaro Rodríguez del Río, Joaquín Giner-Lamia, Carlos P. Cantalapiedra, Jorge Botas, Ziqi Deng, Ana Hernández-Plaza, Martí Munar-Palmer, Saray Santamaría-Hernando, José J. Rodríguez-Herva, Hans-Joachim Ruscheweyh, Lucas Paoli, Thomas S. B. Schmidt, Shinichi Sunagawa, Peer Bork, Emilia López-Solanilla, Luis Pedro Coelho, and Jaime Huerta-Cepas. Functional and evolutionary significance of unknown genes from uncultivated taxa. *Nature*, 626(7998):377–384, 2024. doi: 10.1038/s41586-023-06955-z.

[9] Eleni Aplakidou, Fotis A. Baltoumas, Maria N. Chasapi, Efthimia Lamari, Ilias Georgakopoulos-Soares, Evangelos Karatzas, Ioannis Iliopoulos, Aydin Buluç, Robert D. Finn, Antonio Pedro Camargo, Nikos C. Kyrpides, and Georgios A. Pavlopoulos. Quadrupling the protein family space with global metagenomics. *Nucleic Acids Research*, 2026. doi: 10.1093/nar/gkag938.

[10] Andy M. Lau, Nicola Bordin, Shaun M. Kandathil, Ian Sillitoe, Vaishali P. Waman, Jude Wells, Christine A. Orengo, and David T. Jones. Exploring structural diversity across the protein universe with The Encyclopedia of Domains. *Science*, 386(6721):eadq4946, 2024. doi: 10.1126/science.adq4946.

[11] Jingi Yeo, Yewon Han, Nicola Bordin, Andy M. Lau, Shaun M. Kandathil, Hyunbin Kim, Eli Levy Karin, Milot Mirdita, David T. Jones, Christine Orengo, and Martin Steinegger. Metagenomic-scale analysis of the predicted protein structure universe. *bioRxiv*, 2025. doi: 10.1101/2025.04.23.650224. AFESM: 821M AFDB+ESMatlas structures; near-saturation of domain-fold space (12+33 novel folds).

[12] Sergey Kryazhimskiy and Joshua B. Plotkin. The population genetics of dn/ds. *PLoS Genetics*, 4(12):e1000304, 2008. doi: 10.1371/journal.pgen.1000304.

[13] Carina F. Mugal, Jochen B.W. Wolf, and Ingemar Kaj. Why time matters: Codon evolution and the temporal dynamics of dn/ds. *Molecular Biology and Evolution*, 2014. doi: 10.1093/molbev/mst192.

[14] Chase W Nelson, Zachary Ardern, Xinzhu Wei, and Koichiro Tamura. Olgenie: Estimating natural selection to predict functional overlapping genes. *Molecular Biology and Evolution*, 2020. doi: 10.1093/molbev/msaa087.

[15] Shibu Yooseph, Granger Sutton, Douglas B Rusch, Aaron L Halpern, Shannon J Williamson, Karin Remington, Jonathan A Eisen, Karla B Heidelberg, Gerard Manning, Weizhong Li, Lukasz Jaroszewski, Piotr Cieplak, Christopher S Miller, Huiying Li, Susan T Mashiyama, Marcin P Joachimiak, Christopher van Belle, John-Marc Chandonia, David A Soergel, Yufeng Zhai, Kannan Natarajan, Shaun Lee, Benjamin J Raphael, Vineet Bafna, Robert Friedman, Steven E Brenner, Adam Godzik, David Eisenberg, Jack E Dixon, Susan S Taylor, Robert L Strausberg, Marvin Frazier, J. Craig Venter, and Sean Eddy. The sorcerer ii global ocean sampling expedition: Expanding the universe of protein families. *PLoS Biology*, 2007. doi: 10.1371/journal.pbio.0050016.

[16] Ruth Y. Eberhardt, Daniel H. Haft, Marco Punta, Maria Martin, Claire O'Donovan, and Alex Bateman. Antifam: a tool to help identify spurious orfs in protein annotation. *Database: The Journal of Biological Databases and Curation*, 2012. doi: 10.1093/database/bas003.

[17] Doug Hyatt, Gwo-Liang Chen, Philip F. LoCascio, Miriam L. Land, Frank W. Larimer, and Loren J. Hauser. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics*, 11:119, 2010. doi: 10.1186/1471-2105-11-119.

[18] Ailsa K Orr, Alex Bateman, and Dmitrij Frishman. Folding the unfoldable 2: using alphafold and esmfold to explore spurious proteins. *Bioinformatics Advances*, 2026. doi: 10.1093/bioadv/vbag160.

[19] Mara D. Maeke, Xiuran Yin, Lea C. Wunder, Chiara Vanni, Tim Richter-Heitmann, Samuel Miravet-Verde, Hans-Joachim Ruscheweyh, Shinichi Sunagawa, Jenny Fabian, Judith Piontek, Michael W. Friedrich, and Christiane Hassenrück. Extensive data mining uncovers novel diversity among members of the rare biosphere within the thermoplasmatota. *Microbiome*, 13(1):155, 2025. doi: 10.1186/s40168-025-02140-8.

[20] Rund Tawfiq, Marwa Abdelhakim, Mohammed Alarawi, Susana Martinez Arbas, Sulaiman M. Alajel, Abderahmane Derouiche, Raik Grünberg, Maxat Kulmanov, Kexin Niu, Alejandra Lopez Velazquez, Tiannyu Wang, Magnus Rueping, and Robert Hoehndorf. A multimodal field dataset and semantic knowledge graph for the Rub' al-Khali desert microbiome. Companion data descriptor in preparation, 2026.

[21] Shifu Chen, Yanqing Zhou, Yaru Chen, and Jia Gu. fastp: an ultra-fast all-in-one fastq preprocessor. *Bioinformatics*, 34(17):i884–i890, 2018. doi: 10.1093/bioinformatics/bty560.

[22] Dinghua Li, Chi-Man Liu, Ruibang Luo, Kunihiko Sadakane, and Tak-Wah Lam. Megahit: an ultra-fast single-node solution for large and complex metagenomics assembly via succinct de bruijn graph. *Bioinformatics*, 31(10):1674–1676, 2015. doi: 10.1093/bioinformatics/btv033.

[23] Alexey Gurevich, Vladislav Saveliev, Nikolay Vyahhi, and Glenn Tesler. Quast: quality assessment tool for genome assemblies. *Bioinformatics*, 29(8):1072–1075, 2013. doi: 10.1093/bioinformatics/btt086.

[24] Shaojun Pan, Xing-Ming Zhao, and Luis Pedro Coelho. SemiBin2: self-supervised contrastive learning leads to better MAGs for short- and long-read sequencing. *Bioinformatics*, 39 (Supplement 1):i21–i29, 2023. doi: 10.1093/bioinformatics/btad209.

[25] Heng Li. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics*, 34(18): 3094–3100, 2018. doi: 10.1093/bioinformatics/bty191.

[26] Heng Li. Aligning sequence reads, clone sequences and assembly contigs with bwa-mem. *arXiv*, 1303.3997, 2013. doi: 10.48550/arXiv.1303.3997.

[27] Christian M. K. Sieber, Alexander J. Probst, Allison Sharrar, Brian C. Thomas, Matthias Hess, Susannah G. Tringe, and Jillian F. Banfield. Recovery of genomes from metagenomes via a dereplication, aggregation and scoring strategy. *Nature Microbiology*, 3:836–843, 2018. doi: 10.1038/s41564-018-0171-1.

[28] Alex Chklovski, Donovan H. Parks, Ben J. Woodcroft, and Gene W. Tyson. CheckM2: a rapid, scalable and accurate tool for assessing microbial genome quality using machine learning. *Nature Methods*, 20(8):1203–1212, 2023. doi: 10.1038/s41592-023-01940-w.

[29] Patricia P Chan, Brian Y Lin, Allysia J Mak, and Todd M Lowe. trnascan-se 2.0: improved detection and functional classification of transfer rna genes. *Nucleic Acids Research*, 2021. doi: 10.1093/nar/gkab688.

[30] Robert M. Bowers, Nikos C. Kyrpides, Ramunas Stepanauskas, Miranda Harmon-Smith, Devin Doud, T. B. K. Reddy, Frederik Schulz, Jessica Jarett, Adam R. Rivers, Emiley A. Eloe-Fadrosh, Susannah G. Tringe, Natalia N. Ivanova, Alex Copeland, Alicia Clum, Eric D. Becraft, Rex R. Malmstrom, Bruce Birren, others, and Tanja Woyke. Minimum information about a single amplified genome (MISAG) and a metagenome-assembled genome (MIMAG) of bacteria and archaea. *Nature Biotechnology*, 35(8):725–731, 2017. doi: 10.1038/nbt.3893.

[31] Askarbek Orakov, Anthony Fullam, Luis Pedro Coelho, Supriya Khedkar, Damian Szklarczyk, Daniel R. Mende, Thomas S. B. Schmidt, and Peer Bork. GUNC: detection of chimerism and contamination in prokaryotic genomes. *Genome Biology*, 22:178, 2021. doi: 10.1186/s13059-021-02393-0.

[32] Matthew R. Olm, Christopher T. Brown, Brandon Brooks, and Jillian F. Banfield. drep: a tool for fast and accurate genomic comparisons that enables improved genome recovery from metagenomes through de-replication. *The ISME Journal*, 11(12):2864–2868, 2017. doi: 10.1038/ismej.2017.126.

[33] Chirag Jain, Luis M. Rodriguez-R, Adam M. Phillippy, Konstantinos T. Konstantinidis, and Srinivas Aluru. High throughput ANI analysis of 90k prokaryotic genomes reveals clear species boundaries. *Nature Communications*, 9(1):5114, 2018. doi: 10.1038/s41467-018-07641-9.

[34] Pierre-Alain Chaumeil, Aaron J. Mussig, Philip Hugenholtz, and Donovan H. Parks. GTDB-Tk v2: memory friendly classification with the genome taxonomy database. *Bioinformatics*, 38(23):5315–5316, 2022. doi: 10.1093/bioinformatics/btac672.

[35] Donovan H. Parks, Maria Chuvochina, Christian Rinke, Aaron J. Mussig, Pierre-Alain Chaumeil, and Philip Hugenholtz. GTDB: an ongoing census of bacterial and archaeal diversity through a phylogenetically consistent, rank normalized and complete genome-based taxonomy. *Nucleic Acids Research*, 50(D1):D785–D794, 2022. doi: 10.1093/nar/gkab776.

[36] Martin Larralde. Pyrodigal: Python bindings and interface to prodigal, an efficient method for gene prediction in prokaryotes. *Journal of Open Source Software*, 7(72):4296, 2022. doi: 10.21105/joss.04296.

[37] Martin Steinegger and Johannes Söding. MMseqs2 enables sensitive protein sequence searching for the analysis of massive data sets. *Nature Biotechnology*, 35(11):1026–1028, 2017. doi: 10.1038/nbt.3988.

[38] Martin Steinegger and Johannes S"oding. Clustering huge protein sequence sets in linear time. *Nature Communications*, 9:2542, 2018. doi: 10.1038/s41467-018-04964-5.

[39] Baris E. Suzek, Yuqi Wang, Hongzhan Huang, Peter B. McGarvey, and Cathy H. Wu. UniRef clusters: a comprehensive and scalable alternative for improving sequence similarity searches. *Bioinformatics*, 31(6):926–932, 2015. doi: 10.1093/bioinformatics/btu739.

[40] Oliver Schwengers, Lukas Jelonek, Marius Alfred Dieckmann, Sebastian Beyvers, Jochen Blom, and Alexander Goesmann. Bakta: rapid and standardized annotation of bacterial genomes via alignment-free sequence identification. *Microbial Genomics*, 7(11):000685, 2021. doi: 10.1099/mgen.0.000685.

[41] Carlos P. Cantalapiedra, Ana Hernández-Plaza, Ivica Letunic, Peer Bork, and Jaime Huerta-Cepas. eggNOG-mapper v2: functional annotation, orthology assignments, and domain prediction at the metagenomic scale. *Molecular Biology and Evolution*, 38(12):5825–5829, 2021. doi: 10.1093/molbev/msab293.

[42] Benjamin Buchfink, Klaus Reuter, and Hajk-Georg Drost. Sensitive protein alignments at tree-of-life scale using DIAMOND. *Nature Methods*, 18(4):366–368, 2021. doi: 10.1038/s41592-021-01101-x.

[43] Samuel T. N. Aroney, Rhys J. P. Newell, Jakob N. Nissen, Antonio Pedro Camargo, Gene W. Tyson, and Ben J. Woodcroft. CoverM: read alignment statistics for metagenomics. *Bioinformatics*, 41(4):btaf147, 2025. doi: 10.1093/bioinformatics/btaf147.

[44] Kazutaka Katoh and Daron M. Standley. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution*, 30 (4):772–780, 2013. doi: 10.1093/molbev/mst010.

[45] Masatoshi Nei and Takashi Gojobori. Simple methods for estimating the numbers of synonymous and nonsynonymous nucleotide substitutions. *Molecular Biology and Evolution*, 3 (5):418–426, 1986. doi: 10.1093/oxfordjournals.molbev.a040410.

[46] Peter J. A. Cock, Tiago Antao, Jeffrey T. Chang, Brad A. Chapman, Cymon J. Cox, Andrew Dalke, Iddo Friedberg, Thomas Hamelryck, Frank Kauff, Bartek Wilczynski, and Michiel J. L. de Hoon. Biopython: freely available python tools for computational molecular biology and bioinformatics. *Bioinformatics*, 25(11):1422–1423, 2009. doi: 10.1093/bioinformatics/btp163.

[47] Jim Shaw and Yun William Yu. Fast and robust metagenomic sequence comparison through sparse chaining with skani. *Nature Methods*, 2023. doi: 10.1038/s41592-023-02018-3.

[48] Morgan N. Price, Paramvir S. Dehal, and Adam P. Arkin. FastTree 2 — approximately maximum-likelihood trees for large alignments. *PLoS ONE*, 5(3):e9490, 2010. doi: 10.1371/journal.pone.0009490.

[49] Michael Heinzinger, Konstantin Weissenow, Joaquin Gomez Sanchez, Adrian Henkel, Milot Mirdita, Martin Steinegger, and Burkhard Rost. Bilingual language model for protein sequence and structure. *NAR Genomics and Bioinformatics*, 6(4):lqae150, 2024. doi: 10.1093/nargab/lqae150.

[50] Michel van Kempen, Stephanie S. Kim, Charlotte Tumescheit, Milot Mirdita, Jeongjae Lee, Cameron L. M. Gilchrist, Johannes Söding, and Martin Steinegger. Fast and accurate protein structure search with Foldseek. *Nature Biotechnology*, 42(2):243–246, 2024. doi: 10.1038/s41587-023-01773-0.

[51] Mihaly Varadi, Damian Bertoni, Paulyna Magana, Urmila Paramval, Ivanna Pidruchna, Malarvizhi Radhakrishnan, Maxim Tsenkov, Sreenath Nair, Milot Mirdita, Jingi Yeo, Oleg Kovalevskiy, Kathryn Tunyasuvunakool, Agata Laydon, Augustin Žídek, Hamish Tomlinson, Dhavanthi Hariharan, Josh Abrahamson, Tim Green, John Jumper, Ewan Birney, Martin Steinegger, Demis Hassabis, and Sameer Velankar. AlphaFold protein structure database in 2024: providing structure coverage for over 214 million protein sequences. *Nucleic Acids Research*, 52(D1):D368–D375, 2024. doi: 10.1093/nar/gkad1011.

[52] Helen M. Berman, John Westbrook, Zukang Feng, Gary Gilliland, T. N. Bhat, Helge Weissig, Ilya N. Shindyalov, and Philip E. Bourne. The protein data bank. *Nucleic Acids Research*, 28(1):235–242, 2000. doi: 10.1093/nar/28.1.235.

[53] Zeming Lin, Halil Akin, Roshan Rao, Brian Hie, Zhongkai Zhu, Wenting Lu, Nikita Smetanin, Robert Verkuil, Ori Kabeli, Yaniv Shmueli, Allan dos Santos Costa, Maryam Fazel-Zarandi, Tom Sercu, Salvatore Candido, and Alexander Rives. Evolutionary-scale prediction of atomic-level protein structure with a language model. *Science*, 379(6637):1123–1130, 2023. doi: 10.1126/science.ade2574.

[54] Yang Zhang and Jeffrey Skolnick. Scoring function for automated assessment of protein structure template quality. *Proteins: Structure, Function, and Bioinformatics*, 57(4):702–710, 2004. doi: 10.1002/prot.20264.

[55] Gavin M. Douglas, Vincent J. Maffei, Jesse R. Zaneveld, Svetlana N. Yurgel, James R. Brown, Christopher M. Taylor, Curtis Huttenhower, and Morgan G. I. Langille. PICRUSt2 for prediction of metagenome functions. *Nature Biotechnology*, 38(6):685–688, 2020. doi: 10.1038/s41587-020-0548-6.

[56] Lorna Richardson, Ben Allen, Germana Baldi, Martin Beracochea, Maxwell L. Bileschi, others, and Robert D. Finn. MGnify: the microbiome sequence data analysis resource in 2023. *Nucleic Acids Research*, 51(D1):D753–D759, 2023. doi: 10.1093/nar/gkac1080.

[57] Stephen Nayfach, Simon Roux, Rekha Seshadri, Daniel Udwary, others, Emiley A. Eloe-Fadrosh, and Nikos C. Kyrpides. A genomic catalog of Earth's microbiomes. *Nature Biotechnology*, 39(4):499–509, 2021. doi: 10.1038/s41587-020-0718-6.

[58] Thomas S. B. Schmidt, Anthony Fullam, Pamela Ferretti, Askarbek Orakov, Oleksandr M. Maistrenko, Hans-Joachim Ruscheweyh, Ivica Letunic, Yiqian Duan, Thea Van Rossum, Shinichi Sunagawa, Daniel R. Mende, Robert D. Finn, Michael Kuhn, Luis Pedro Coelho, and Peer Bork. SPIRE: a searchable, planetary-scale microbiome resource. *Nucleic Acids Research*, 52(D1):D777–D783, 2024. doi: 10.1093/nar/gkad943.

[59] Brian H. McArdle and Marti J. Anderson. Fitting multivariate models to community data: A comment on distance-based redundancy analysis. *Ecology*, 82(1):290–297, 2001. doi: 10.1890/0012-9658(2001)082[0290:FMMTCD]2.0.CO;2.

[60] Daniel Borcard, Pierre Legendre, and Pierre Drapeau. Partialling out the spatial component of ecological variation. *Ecology*, 73(3):1045–1055, 1992. doi: 10.2307/1940179.

[61] Nathan Mantel. The detection of disease clustering and a generalized regression approach. *Cancer Research*, 27(2):209–220, 1967. PMID: 6018555.

[62] Kai Blin, Simon Shaw, Lisa Vader, Judit Szenei, Zachary L Reitz, Hannah E Augustijn, José D D Cediel-Becerra, Valérie de Crécy-Lagard, Robert A Koetsier, Sam E Williams, et al. antismash 8.0: extended gene cluster detection capabilities and analyses of chemistry, enzymology, and regulation. *Nucleic Acids Research*, 53(W1):W32–W38, 2025. doi: 10.1093/nar/gkaf334.

[63] Michael Feldgarden, Vyacheslav Brover, Narjol Gonzalez-Escalona, Jonathan G. Frye, Julie Haendiges, Daniel H. Haft, Maria Hoffmann, James B. Pettengill, Arjun B. Prasad, Glenn E. Tillman, Gregory H. Tyson, and William Klimke. AMRFinderPlus and the reference gene catalog facilitate examination of the genomic links among antimicrobial resistance, stress response, and virulence. *Scientific Reports*, 11(1):12728, 2021. doi: 10.1038/s41598-021-91456-0.

[64] Sture Holm. A simple sequentially rejective multiple test procedure. *Scandinavian Journal of Statistics*, 6(2):65–70, 1979. JSTOR 4615733; no live DOI — 10.2307/4615733 returns 404 at both Crossref and doi.org (checked 2026-08-18).

[65] Yoav Benjamini and Yosef Hochberg. Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 57(1):289–300, 1995. doi: 10.1111/j.2517-6161.1995.tb02031.x.

[66] Charles R. Harris, K. Jarrod Millman, Stéfan J. van der Walt, Ralf Gommers, Pauli Virtanen, David Cournapeau, Eric Wieser, Julian Taylor, Sebastian Berg, Nathaniel J. Smith, et al. Array programming with numpy. *Nature*, 585(7825):357–362, 2020. doi: 10.1038/s41586-020-2649-2.

[67] Pauli Virtanen, Ralf Gommers, Travis E. Oliphant, Matt Haberland, Tyler Reddy, David Cournapeau, Evgeni Burovski, Pearu Peterson, Warren Weckesser, Jonathan Bright, et al. Scipy 1.0: fundamental algorithms for scientific computing in python. *Nature Methods*, 17 (3):261–272, 2020. doi: 10.1038/s41592-019-0686-2.




Figure 1: **Sampling and genome quality.** **a**, Counts of the 551 soil samples across five expeditions. **b**, Sample counts by soil position and expedition. **c**, CheckM2 completeness and contamination for the 5,438 species representatives. Sampling covers 59 core sites and two auxiliary locations; expedition and position counts describe samples, while the quality panel describes dereplicated genomes.




Figure 2: **Evolutionary comparisons of recurrent dark families and sharing in culture comparisons.** **a**, Empirical cumulative distributions of family median dN/dS for 97,200 dark and 95,647 annotated families. The display extends to dN/dS = 2; cumulative fractions use all families. **b**, Annotated- and shifted-frame calibration distributions from a 10,000-family input, with 9,182 and 9,148 evaluable outputs; the display extends to dN/dS = 3. **c**, Difference between dark and annotated family medians in left-closed divergence strata (final bin includes 2), with 95% intervals from 1,000 family bootstrap replicates. Labels give dark/control counts. Shared ancestry is not modelled. **d**, Percentage of annotated and dark MAG protein calls with reciprocal protein hits in the paired culture-derived assembly, for 22 species groups after exclusion of a confirmed *E. coli* contaminant. Each line joins the two fractions for one MAG. **e**, DNA support by protein-hit status, with counts above bars. Eligible coding intervals are at least 90 nt; support requires a base-level alignment at 95% identity and 90% query coverage. **f**, Recovery of protein-shared calls by genome pair; the dotted line marks a post hoc 90% diagnostic threshold.




Figure 3: **Lineage-associated variation in per-genome darkness.** **a**, The twelve highest median dark fractions among phyla with at least 20 representatives. **b**, Dark fraction by the first empty or digit-bearing label: 1, phylum; 2, class; 3, order; 4, family; 5, genus; 6, species; 7, no flagged rank. **c**, Thermoplasmatota (T) and matched other archaea (A) or bacteria (B), matched on completeness, contamination, corrected genome size, N50, abundance and prevalence. The archaeal and bacterial matching sets have separate target subsets (218 and 237 pairs). Boxes show the median and interquartile range; whiskers extend to 1.5 interquartile ranges, and points beyond the whiskers are omitted.




Figure 4: **Structural reference coverage depends on score normalisation and retrieval.** **a**, Best query- and alignment-normalised TM-scores across the PDB and AlphaFold Database for 2,838 confidence-qualified dark models and 3,013 controls. Dashed lines mark 0.5. **b**, Fractions below thresholds from 0.4 to 0.6 under both normalisations. **c**, Query-score no-hits at 0.5 separated into no returned alignment rows and retrieved scores below threshold. Zero-score counts are 22 dark models and 66 controls; nonzero below-threshold counts are 1,009 and 35. The controls derive from PDB chains represented in the search references.




Figure 5: **Reference coverage and catalogue composition.** **a**, Recall of genome-encoded orthologues and rank correlation on shared orthologues across 448 paired profiles. Boxes show medians and interquartile ranges, with 1.5-interquartile-range whiskers and outliers omitted. **b**, Observed absence versus the analytical E-value-corrected MGnify90 rarefaction expectation at each catalogue's size. The diagonal indicates agreement with the size-only expectation. Each point is a catalogue, coloured by query group; points below the diagonal have lower observed absence. Atacama has 4.67 percentage points more dark-family absence and 29.97 points more annotated-family absence than the size-only expectation, demonstrating a large reference-composition residual. The combined desert and SPIRE annotated residuals are +0.93 and +1.23 points in expected-minus-observed absence. **c**, Median occupied locations out of 61, stratified by representative length and absence from every searched catalogue. Absence is relative to the database and search rule.