
A. Order & quality-method composition
(* = 3 shared orders: 234 EQ, 59 GTDB, 8 MGnify)
o__JACQPN01 *
o__JACPAO01 *
o__SW-10-69-26 *
o__DHVEG-1
o__UBA202
o__
o__UBA9212
o__GCA-002838935
o__RBG-16-68-12
EQ_MAG (CheckM2, n=237)
GTDB_r232_rep (CheckM2, n=218)
MGnify (unknown method, n=29)
0
20
40
60
80
100
120
140
160
Number of genomes (total = 484)
B. Shared-order CheckM2 quality overlap (n=293)
(MGnify n=29 unknown method kept separate)
EQ_MAG shared orders (n=234)
GTDB_r232_rep shared orders (n=59)
CheckM2 contamination (%) [bins: [0,5), [5,10)]
0
2
4
6
8
10
50
60
70
80
90
100
CheckM2 completeness (%) [bins: [50,70), [70,90), [90,101]]
C. Within-stratum cohort medians (CheckM2 overlap cells)
(Descriptive differences vary across order/quality strata)
JACPAO01 [50,70) [0,5)
(EQ=8, GTDB=2) [n<5 sparse]
JACPAO01 [70,90) [0,5)
(EQ=19, GTDB=25)
JACPAO01 [70,90) [5,10)
(EQ=4, GTDB=1) [n<5 sparse]
JACPAO01 [90,101] [0,5)
(EQ=15, GTDB=10)
JACQPN01 [70,90) [0,5)
(EQ=25, GTDB=9)
JACQPN01 [90,101] [0,5)
(EQ=35, GTDB=5)
JACQPN01 [90,101] [5,10)
(EQ=23, GTDB=1) [n<5 sparse]
SW-10-69-26 [70,90) [0,5)
(EQ=14, GTDB=5)
SW-10-69-26 [90,101] [0,5)
(EQ=6, GTDB=1) [n<5 sparse]
EQ median (S3, B0)
GTDB median (S3, B0)
160
170
180
190
200
Median min added reactions in S3 (B0 baseline)