
A. Masked-reaction recovery across 7 reference genomes
(7 genomes; 9 masking conditions/replicates per genome: 3 fractions × 3 seeds)
MGYG000495096
(o__DHVEG-1, comp=91.12%, cont=2.58%)
MGYG000510876
(o__DHVEG-1, comp=97.69%, cont=1.44%)
MGYG000519748
(o__UBA202, comp=95.78%, cont=0.33%)
MGYG000519815
(o__UBA202, comp=98.56%, cont=0.27%)
MGYG000520046
(o__UBA202, comp=99.6%, cont=0.78%)
MGYG000523080
(o__DHVEG-1, comp=94.33%, cont=4.18%)
MGYG000526805
(o__JACQPN01, comp=98.06%, cont=4.39%)
0.0
0.1
0.2
0.3
0.4
0.5
Masked GPR reaction recovery rate under matched S4 + B2 medium
Mask 10% (3 random seeds/genome)
Mask 20% (3 random seeds/genome)
Mask 30% (3 random seeds/genome)
Per-genome mean (9 conditions)
Per-genome median
7-genome mean = 0.168
B. Alternative non-masked pathway fraction (7 reference genomes)
(Alternative reactions account for 76.7% of new additions)
0.4
0.5
0.6
0.7
0.8
0.9
1.0
Alternative candidate fraction among newly induced gap-fill reactions
Mask 10% (3 random seeds/genome)
Mask 20% (3 random seeds/genome)
Mask 30% (3 random seeds/genome)
Per-genome mean (9 conditions)
Per-genome median
7-genome mean = 0.767