Genome-scale context, stoichiometric refinement, and neuro-symbolic reasoning for proteome function annotation

Tawfiq Al-Saeed[1], Maxat Kulmanov[1], and Robert Hoehndorf[1,*]

[1]Computer, Electrical and Mathematical Sciences & Engineering Division (CEMSE),
Computational Bioscience Research Center (CBRC), King Abdullah University of Science and Technology (KAUST),
Thuwal 23955-6900, Kingdom of Saudi Arabia

[*]Correspondence: robert.hoehndorf@kaust.edu.sa

Abstract

Protein function predictors combine sequence homology, domain profiles, protein language models, and structural similarity to score individual sequences, while long-read sequencing and modern genome assembly now routinely recover complete chromosomes that expose operon organization, complex stoichiometry, and metabolic network connectivity across the entire proteome. We present Genome-Scale Protein Annotation (GSPA), a staged framework that calibrates and pools complementary per-protein predictors, updates proteome-wide posteriors using whole-genome constraints over essential functions, pathways, obligate heteromeric complexes, and taxonomic and anatomical ontology axioms, and couples reaction-local genomic context with positive-cost parsimonious flux balance analysis, competitive gene-locus attribution, and ontology-guarded language-model reasoning. Across bacterial, archaeal, viral, and eukaryotic proteomes, whole-genome inference preserves the protein-level accuracy baseline, resolves all detected taxonomic and anatomical contradictions while outperforming term deletion in multicellular animals and vascular plants, and raises recovery of held-out heteromeric complex subunits (Hit@1) from 56.0% to 93.7%. Positive-cost stoichiometric refinement eliminates the 83.7% zero-flux reaction padding produced by negative-cost gap-filling and improves the identification of fitness-validated bacterial enzymes, pathway-coupled metabolic modeling improves the prediction of experimental growth and diagnostic enzyme phenotypes across unseen genera, and selective guarded reasoning on fewer than 2.5% of proteins raises top-ranked recovery of ambiguous missing pathway enzymes from 34.0% to 82.7%. Together, protein similarity, whole-genome architecture, and systems-level constraints provide complementary layers of functional inference; GSPA, precomputed constraint tables, and reproducible benchmarks are available at https://github.com/bio-ontology-research-group/gspa.

1 Introduction

Automated protein function annotation assigns biological roles to the protein-coding genes of an organism and forms the basis of comparative genomics, metabolic modeling, and systems biology [1, 2]. We consider the following genome-scale inference problem. The input consists of the protein-coding genes $\mathcal{P} = \{p_1, \ldots, p_N\}$ of an organism, their chromosomal coordinates and strand organization $\mathcal{K}$, the organism's taxonomic lineage τ_{tax}, and reference biochemical and ontological knowledge bases. The output is a joint assignment of Gene Ontology (GO) terms [3] and Enzyme Commission (EC) numbers across $\mathcal{P}$. Because the proteins encoded by a genome function together within a single cell or organism, a valid joint assignment satisfies structural invariants that span many proteins simultaneously: core cellular functions required in the organism's domain of life are represented, metabolic pathways form stoichiometrically connected routes that support biomass synthesis [4, 5], obligate heteromeric complexes are annotated with their complete partner subunit sets [6, 7], and assigned terms respect both the organism's taxonomic lineage and the anatomical body plan of its clade [8, 9]. These organism-level invariants link the function of each protein to the organization of the complete proteome.

This joint formulation is timely because complete genome organization is now broadly accessible. Historically, short-read sequencing of microbial isolates and metagenomes yielded fragmented contigs from which open reading frames were predicted and annotated in isolation, often severing operons, biosynthetic gene clusters, and chromosome-wide reaction inventories. Advances in long-read sequencing [10, 11] and repeat-graph genome and metagenome assemblers [12] have transformed genome recovery, routinely yielding complete or near-complete chromosomes for both cultivated isolates and metagenome-assembled genomes (MAGs) of uncultured lineages. Chromosomal gene order, co-directional operons, and complete proteome complements are therefore standard inputs that function prediction pipelines can exploit.

Most function prediction methods derive representations of an individual protein sequence or structure and

transfer annotations from similar reference proteins. Sequence alignment transfers annotations from homologous sequences [13–15]; profile hidden Markov models identify conserved domain signatures [16, 17]; protein language models embed sequences into spaces where functionally related proteins cluster [18–21]; and structural alignment transfers functions across remote fold similarities [22, 23]. Because these representations capture distinct biochemical properties and make partially independent errors, combining them improves prediction accuracy. Multi-source ensembles such as GOLabeler [24] and NetGO [25, 26] integrate sequence homology, domain profiles, motifs, network associations, and language-model embeddings within learned ranking frameworks and established the state of the art in Critical Assessment of Functional Annotation (CAFA) evaluations [1, 2, 27]. For conserved proteins, these ensembles provide reliable initial probability distributions that serve as the foundation for proteome annotation. Scoring each protein p_i from its sequence and structure alone leaves divergent pathway enzymes, isolated complex subunits, and distant cross-clade homologs to be resolved by organism-wide context.

Comparative genomics introduces a second representation layer by placing each gene within its local chromosomal neighborhood. Conserved gene order, co-directional operons, and phylogenetic co-occurrence profiles associate proteins that act in the same metabolic pathway or macromolecular complex [28–32]. Genomic language models [33, 34] and neighborhood-aware predictors [35, 36] learn contextualized representations over contiguous gene windows, allowing them to distinguish paralogs and identify co-regulated pathway partners. A contiguous window captures local chromosomal synteny, whereas domain-essential function profiles, stoichiometry across dispersed complex loci, and formal taxonomic and anatomical admissibility act across the entire genome.

Systems biology and constraint-based metabolic reconstruction provide a third layer of functional evidence by testing whether predicted enzymes form a network capable of steady-state flux. Reconstruction pipelines including Pathway Tools [37], ModelSEED [38], CarveMe [39], gapseq [40], and GapMind [41] map sequence annotations onto reaction networks and use flux balance analysis to identify reactions required for growth on specified media. In our initial effort to move beyond sequence homology, METEOR [42] (presented in the Pacific Symposium on Biocomputing 2027 session on Biological Molecular Function: Beyond Homology) coupled metabolic gap-filling with neural enzyme prediction through reaction-indicator mixed-integer linear programming (MILP). Close inspection of reaction-indicator MILP reveals two mathematical properties that limit gene-level specificity: assigning negative objective costs to candidate reactions above an addition threshold rewards the solver for adding reactions regardless of whether they carry steady-state flux, and transferring reaction support uniformly to candidate proteins raises the scores of paralogs equally while leaving operon neighborhoods, subunit roles, and competition among candidate loci unmodeled.

Formal bio-ontologies provide complementary organism-level constraints by specifying which biological processes and cellular structures can co-exist in a taxon. GO taxon constraints restrict terms to specific clades [8], while cross-ontology equivalence and subclass axioms connect GO biological processes to anatomical structures in the Uberon multi-species anatomy ontology [9, 43], the Plant Ontology (PO) [44], and the Cell Ontology (CL) [45]. Through these definitions, an organ-level developmental process inherits the taxonomic restrictions of the anatomical structure in which it occurs. Whereas deletion-based filters discard a forbidden term together with the conserved biological processes implied by its valid GO ancestors, semantic ancestor relaxation can retain the most specific ancestor admissible in the target organism's clade. For the small fraction of loci that remain ambiguous after deterministic probabilistic and stoichiometric inference, tool-using language-model agents [46, 47] can synthesize heterogeneous domain, operon, and pathway evidence when coupled to a deterministic symbolic verifier that enforces formal ontology axioms.

Here we present Genome-Scale Protein Annotation (GSPA), a four-stage framework that unifies these representations in their dependency order (Figure 1). First, Stage 0 calibrates and pools complementary per-protein predictors across correlation-grouped homology, domain, language-model, and structural channels to establish a strong protein-level baseline. Second, Stage 1 and Stage 2A exploit whole-genome information: Stage 1 iteratively updates proteome-wide log-odds across essential functions, pathways, obligate heteromeric complexes, taxonomic consistency, anatomy, metabolic gaps, and operons, and Stage 2A deterministically enforces cross-ontology constraints by relaxing clade-incompatible terms to their nearest valid GO ancestors. Third, Stage 1, Stage 2B, and Stage 3 go beyond homology through systems biology and guarded reasoning: Stage 1 localizes missing enzymes via reaction-local genomic context (RLGC), Stage 2B couples role-aware disjunctive normal form (DNF) aggregation of isozymes and required subunits with positive-cost lexicographic parsimonious flux balance analysis (pFBA) [48] and competitive locus attribution against an explicit unassigned-gene hypothesis, and Stage 3 resolves residual ambiguities on fewer than 2.5% of proteins using a tool-calling language model verified by a symbolic ontology guardrail. Across eight benchmark suites spanning bacteria, archaea, viruses, and eukaryotes (Supplementary Table S1), we test five hypotheses: (i) whether pooling complementary predictors improves protein-centric accuracy over individual predictors and production pipelines; (ii) whether whole-genome priors improve proteome coherence while preserving protein-level accuracy; (iii) whether operon architecture and cross-ontology axioms complete heteromeric complexes and repair eukaryotic anatomical errors; (iv) whether positive-cost flux modeling, competitive locus attribution, and




Figure 1. Four-stage architecture of GSPA. Stage 0 calibrates per-channel confidence scores from sequence homology, domain profiles, protein language models, and structural similarity (or a standalone predictor) and pools them via correlation-grouped Noisy-OR into an initial proteome-wide probability matrix. Stage 1 refines these probabilities in log-odds space using six whole-proteome priors covering core essentiality, pathway and heteromeric complex coherence, taxonomic consistency, cross-ontology anatomical constraints (26,911 direct GO taxon rules and 2,715 GO–Uberon/PO/CL axioms), metabolic gaps coupled to reaction-local genomic context (RLGC), and co-directional operons. Stage 2A enforces symbolic ontology constraints with semantic ancestor relaxation and prunes isolated obligate complex singletons; Stage 2B aggregates isozyme and subunit roles in disjunctive normal form (DNF), repairs the metabolic network via positive-cost lexicographic parsimonious flux balance analysis (pFBA), and attributes flux-certified reactions competitively to candidate gene loci against an unassigned-gene hypothesis $h_\varnothing$. Stage 3 applies selective tool-mediated language-model reasoning to fewer than 2.5% of proteins under a deterministic symbolic ontology guardrail.

pathway coupling eliminate zero-flux reaction padding and improve fitness-validated enzyme and experimental phenotype prediction; and (v) when selective guarded language-model reasoning complements deterministic and supervised inference. GSPA v2.1.0, precomputed ontology bridges, and benchmark workflows are available at `https://github.com/bio-ontology-research-group/gspa`.

2 Results

2.1 Complementary predictors establish the protein-level baseline

We first asked whether calibrating and pooling complementary per-protein predictors yields a stronger baseline annotation than individual predictors and production genome annotation pipelines when evaluated under an identical ancestor-propagated protocol. We compared GSPA against its individual sequence alignment, domain profile, protein language model, and structural similarity channels, against NCBI PGAP [14], InterProScan [17], and metagenomic-deepFRI [49], and against the genomic language models gLM and gLM2 [33, 34] across 13-genome and 21-genome reference panels (Figure 2d and Supplementary Table S2), with all pipeline hyperparameters calibrated on *Escherichia coli* K-12 and held frozen across all out-of-sample species (Supplementary Table S3). Across the 11 prokaryotic proteomes of the 13-genome panel, GSPA with three sequence and domain channels reaches a micro $F_{\max}$ of 0.8421, exceeding NCBI PGAP (0.7820). Extending Stage 0 to all seven neural, language-model, and structural channels on the 21-genome panel raises micro $F_{\max}$ to 0.8705, outperforming the strongest single channel, ESM-2 nearest-neighbor transfer (0.8580). Combining complementary sequence and structural representations establishes a higher-accuracy starting distribution than any individual predictor, and we next tested whether this baseline advantage holds on newly characterized proteins annotated after the training cutoff.

We evaluated prospective generalization on a CAFA temporal holdout of 16,125 protein–aspect evaluations for proteins that acquired experimental GO annotations after the training cutoff, including 3,440 twilight-zone evaluations sharing less than 30% sequence identity with any training sequence (Supplementary Table S4). Stage 0 pooling achieves a macro $F_{\max}$ of 0.6929, improving over the strongest single neural baseline in the panel, ProteInfer (0.6138) [50]. Subsequent whole-genome refinement preserves full-cohort accuracy while raising macro $F_{\max}$ in the below-30% sequence-identity subset from 0.6723 to 0.6775. Multi-predictor pooling accounts for the majority of protein-level accuracy on conserved proteins, while genome-scale refinement provides targeted gains in the sequence-identity twilight zone. A related question arises

in enzyme annotation, where metabolic gap-filling proposes modifications to top-ranked EC predictions and the annotation pipeline must determine which proposed changes to accept.

We evaluated enzyme EC prediction on a post-cutoff Swiss-Prot holdout of 125 enzymes (Supplementary Table S5) and across 69,666 proteins from 95 bacterial genomes under four donor-homology masking thresholds (Supplementary Table S6). On the post-cutoff holdout, pooling CLEAN [21], EnzBERT [51], and DeepProZyme [52] in Stage 0 reaches micro $F_{\max} = 0.4381$, and genome-scale refinement improves top-3 and top-5 enzyme recall across candidates. Across the 95-genome bacterial panel, accepting unfiltered METEOR gap-filling changes lowers Top-1 accuracy on DeepProZyme and EnzBERT because reaction-level boosts promote non-cognate paralogs, whereas filtering candidate changes through GSPA's three-channel consensus rule yields statistically significant net Top-1 gains across all 12 predictor-by-masking combinations, reaching +155 net corrected enzymes on CLEAN ($p < 10^{-25}$, exact McNemar test; Supplementary Table S6). Agreement across complementary predictors identifies reliable metabolic enzyme updates and provides a calibrated foundation for whole-genome inference.

2.2 Whole-genome priors improve organism-level coherence and preserve protein-level accuracy

With the protein-level posterior established, we asked whether introducing whole-genome priors improves organism-level structural coherence while preserving protein-centric accuracy. We conducted a nine-configuration ablation across nine bacterial and archaeal reference proteomes (81 proteome runs), calibrating hyperparameters on *E. coli* K-12 and evaluating the remaining eight species out of sample (Figure 2a,b and Table 1). Full Stage 1 and Stage 2A refinement maintains protein-centric CAFA $F_{\max}$ at the Stage 0 baseline level (0.8037 versus 0.8041), edges pair-level micro $F_{\max}$ upward, and increases the composite structural audit score [53] from 0.8459 to 0.8643 ($p = 0.0039$, two-sided Wilcoxon signed-rank test; out-of-sample eight-genome mean from 0.8384 to 0.8576). Because roughly 85% of annotated proteins in these reference genomes already carry high-confidence homologs in Stage 0, whole-genome refinement preserves the conserved sequence backbone while concentrating probability updates on lower-confidence, structurally constrained loci. To understand how this composite gain arises, we examined the effect of each individual prior on its target proteome property.

Adding each Stage 1 prior individually to Stage 0 reveals selective updates to the corresponding genome invariant (Table 1). The essentiality prior increases representation of domain-core functions, the coherence prior raises heteromeric complex completeness from 0.8636 to 0.8852, and the taxonomic and anatomical priors, which query our programmatic cross-ontology bridge (Supplementary Table S7), eliminate all 25 endogenous taxonomic violations. While the individual pathway, metabolic gap-filling, and operon priors each produce modest gains in pathway completeness on their own, combining all six priors in Stage 1 raises mean pathway coherence from 0.4975 to 0.5146 ($p = 0.0078$). This synergy arises because metabolic gap-filling and pathway coherence identify missing reaction steps across the genome, while operon context localizes which specific candidate genes in co-directional gene clusters receive that support. Beyond endogenous errors, sequence similarity across kingdoms can also introduce high-scoring cross-clade annotations, which we tested next through controlled perturbation.

To evaluate robustness against cross-clade homology transfer, we injected 900 eukaryotic developmental and clade-restricted GO claims with raw confidence scores in $[0.55, 0.85]$ across the nine prokaryotic proteomes (Figure 2c and Table 1). Stage 0 Platt calibration attenuates most low-reliability claims below the 0.50 decision threshold, leaving 125 false-positive claims active. Applying the taxonomic and anatomical priors and Stage 2A enforcement suppresses 122 of these 125 post-calibration violations (97.6%) while keeping protein-centric accuracy unchanged. Formal organism-level constraints therefore eliminate high-scoring cross-clade transfer errors that pass protein-level score calibration. We then evaluated whether these whole-genome improvements hold at population scale and across all four domains of life.

We applied GSPA with frozen parameters to 1,000 RefSeq bacterial genomes and to a multi-kingdom benchmark of 88 proteomes spanning Bacteria, Archaea, Eukaryota, and Viruses under homology masking (Supplementary Tables S8 and S9). Across the 1,000 bacterial genomes, Stage 1 and Stage 2 refinement eliminates all 23,384 mean taxon-violating protein–term predictions per genome produced by Stage 0 and lifts mean MetaCyc pathway coherence from 36.7% to 66.8% at a mean runtime of 5.4 s per genome. Across the 88 multi-kingdom proteomes, whole-genome refinement produces positive mean accuracy gains in all four domains, with 95% bootstrap confidence intervals on $\Delta F_{\max}$ lying strictly above zero in the two largest cohorts, Bacteria and Viruses, and spanning zero in Archaea and Eukaryota (Supplementary Table S9). Because genome-wide complex coherence measures only whether partner subunits are present somewhere in the proteome, we next tested whether local chromosomal architecture can identify the specific gene encoding a missing complex subunit.

2.3 Operon architecture completes obligate heteromeric complexes

When one subunit of an obligate heteromeric complex has diverged into the twilight zone, several proteins in the genome may share a partial domain match to the missing role. We tested whether complex stoichiometry and co-directional operon architecture can distinguish the




Figure 2. Multi-channel pooling and whole-genome prior ablation across reference proteomes. (**a**) Composite structural audit score, essential function completeness, complex coherence, and pathway coherence across nine ablation configurations averaged over nine bacterial and archaeal reference genomes. (**b**) Per-genome gain in the composite structural audit score from Stage 0 pooling to full Stage 1 and Stage 2A refinement; the asterisk denotes the calibration genome, *E. coli* K-12 MG1655. (**c**) Surviving false-positive annotations per genome in the 900-claim cross-clade perturbation experiment under unconstrained Stage 0 pooling versus taxonomic and anatomical constraint enforcement. (**d**) Uniformly evaluated ancestor-propagated micro and protein-centric CAFA $F_{\max}$ comparing GSPA against external annotation pipelines, individual Stage 0 channels, and genomic language models across the 13-genome and 21-genome panels.

Table 1. Systematic ablation of Stage 1 priors and Stage 2A enforcement across nine bacterial and archaeal reference genomes. Rows add each prior individually to the Stage 0 baseline, combine all six Stage 1 priors, and add Stage 2A deterministic enforcement via removal or semantic relaxation. Values report means across *E. coli* K-12 (calibration genome), *E. coli* O157:H7, *B. subtilis* 168, *P. aeruginosa* PAO1, *M. tuberculosis* H37Rv, *H. pylori* 26695, *M. genitalium* G37, *Synechocystis* sp. PCC 6803, and *M. jannaschii* DSM 2661; the final row reports the out-of-sample mean across the eight non-calibration genomes. Protein-centric CAFA $F_{\max}$ and pair-level micro $F_{\max}$ are evaluated against all GO annotations (experimental-only truth in parentheses). Consistency is $\max(0, 1 - 0.01V)$ for V formal violations (total violations across the nine genomes in parentheses). Perturbation suppression reports the percentage of the 125 post-calibration surviving cross-clade claims eliminated (total suppression out of 900 injected claims in parentheses).

Ablation Configuration	CAFA $F_{\max}$ (All / Exp)	Micro $F_{\max}$ (All / Exp)	Essential C_{ess}	Pathway Coh.	Complex Coh.	Consistency (Viol.)	Audit Score / Perturb. Supp.
Stage 0 Noisy-OR Baseline	0.8041 (0.6540)	0.8190 (0.6425)	0.9612	0.4975	0.8636	0.9722 (25)	0.8459 / 0.0% (86.1%)
+ Essentiality Prior (B_1)	0.8040 (0.6539)	0.8190 (0.6425)	0.9652	0.4975	0.8636	0.9722 (25)	0.8471 / n/a
+ Coherence Prior (B_2)	0.8037 (0.6539)	0.8191 (0.6426)	0.9612	0.4997	0.8852	0.9722 (25)	0.8501 / n/a
+ Gap-Filling Prior (B_5)	0.8041 (0.6540)	0.8190 (0.6425)	0.9612	0.5021	0.8636	0.9722 (25)	0.8465 / n/a
+ Operon Context Prior (B_6)	0.8040 (0.6540)	0.8191 (0.6426)	0.9612	0.5015	0.8636	0.9722 (25)	0.8464 / n/a
+ Consistency & Anatomical (B_3, B_4)	0.8042 (0.6541)	0.8190 (0.6425)	0.9612	0.4975	0.8636	1.0000 (0)	0.8542 / 97.6% (99.7%)
All Stage 1 Priors (B_1–B_6)	0.8037 (0.6539)	0.8192 (0.6426)	0.9652	0.5146	0.8852	1.0000 (0)	0.8615 / 97.6% (99.7%)
Stage 1 + Stage 2A Removal	0.8037 (0.6539)	0.8192 (0.6426)	0.9652	0.5146	0.8808	1.0000 (0)	0.8643 / 97.6% (99.7%)
Stage 1 + Stage 2A Relaxation	0.8037 (0.6539)	0.8192 (0.6426)	0.9652	0.5146	0.8808	1.0000 (0)	0.8643 / 97.6% (99.7%)
Out-of-Sample 8-Genome Mean (Stage 0 → Full)	0.7931 → 0.7927	0.8102 → **0.8104**	0.9564 → **0.9609**	0.4712 → **0.4898**	0.8580 → **0.8762**	0.9688 → **1.0000**	0.8384 → **0.8576**

true subunit from domain-matched competitors across 175 leave-subunit-out trials in six prokaryotic genomes, pairing each attenuated true subunit with five decoy proteins drawn from unrelated genomic loci (Figure 3a and Table 2). Applying the genome-wide complex coherence prior alone lifts subunit recall at the 0.50 threshold from 0.0000 to 0.5029 while leaving top-ranked Hit@1 unchanged at the Stage 0 baseline of 0.5600 because candidate proteins with matching domain roles receive equal genome-wide boosts. Combining complex coherence with the co-directional operon prior raises Hit@1 from 0.5600 to 0.9371 ($p < 10^{-15}$, exact McNemar test; 0.9310 across the five out-of-sample genomes). Global complex stoichiometry therefore identifies which macromolecular assembly is in-




Figure 3. Obligate heteromeric complex completion and eukaryotic anatomical constraint enforcement. (**a**) Leave-subunit-out recovery across 175 held-out subunits in six prokaryotic genomes, ranking each attenuated true subunit against five domain-matched decoys (random Hit@1 = 0.167) across Stage 0 pooling, the Stage 1 complex coherence prior, Stage 1 with operon context, and the full model including Stages 2 and 3. (**b**) Per-genome subunit recovery F_1 at threshold 0.50; the asterisk marks the calibration genome (*E. coli* K-12), and *Synechocystis* sp. PCC 6803 illustrates complexes encoded at dispersed chromosomal loci. (**c**) Surviving taxonomic and anatomical violations across six multicellular and unicellular eukaryotes under unconstrained Stage 0 pooling, direct GO taxon rules, the Stage 1 soft anatomical prior, and Stage 2A symbolic enforcement. (**d**) Ancestor-propagated $F_{\max}$ against unmodified Swiss-Prot annotations comparing Stage 0 pooling, Stage 2A term removal, and Stage 2A semantic ancestor relaxation.

Table 2. Leave-subunit-out recovery across 175 obligate heteromeric complex subunits in six prokaryotic genomes. Each trial attenuates one true subunit into the twilight-zone posterior range ($[0.18, 0.35]$) and introduces five decoy proteins with matched domain scores from unrelated genomic loci (chance Hit@1 = 0.167). Precision, recall, and F_1 at threshold 0.50 and complex coherence are reported across all six species (*E. coli* K-12, *E. coli* O157:H7, *B. subtilis* 168, *M. tuberculosis* H37Rv, *P. aeruginosa* PAO1, and *Synechocystis* sp. PCC 6803), the out-of-sample mean across the five non-calibration species ($n = 145$), and *Synechocystis* sp. PCC 6803 alone ($n = 25$).

Cohort	Method Configuration	Subunit Trials (n)	Hit@1	Precision@0.5	Recall@0.5	F_1@0.5	Complex Coherence
All 6 Species	Stage 0 Noisy-OR Baseline	175	0.5600	0.0000	0.0000	0.0000	0.3901
	Stage 1 Complex Coherence Prior Only	175	0.5600	0.1729	0.5029	0.2459	0.7107
	Stage 1 Coherence + Operon Context	175	0.9371	0.5951	0.9257	0.6680	0.7170
	Full GSPA (Stages 1 + 2A + 3)	175	0.9429	0.6009	0.9314	0.6737	0.7170
Out-of-Sample (5 Species)	Stage 0 Noisy-OR Baseline	145	0.5586	0.0000	0.0000	0.0000	0.3745
	Stage 1 Coherence + Operon Context	145	0.9310	0.5886	0.9104	0.6595	0.7047
	Full GSPA (Stages 1 + 2A + 3)	145	0.9379	0.5955	0.9173	0.6664	0.7047
Synechocystis PCC 6803	Stage 0 Noisy-OR Baseline	25	0.5200	0.0000	0.0000	0.0000	0.3587
	Stage 1 Coherence + Operon Context	25	0.6400	0.3393	0.6400	0.4080	0.6125
	Full GSPA (Stages 1 + 2A + 3)	25	0.6800	0.3793	0.6800	0.4480	0.6125

complete, while local operon organization localizes the specific gene locus that supplies the missing subunit. A natural test of this mechanism arises in lineages where complex subunits are encoded at dispersed chromosomal positions.

In the cyanobacterium *Synechocystis* sp. PCC 6803, subunits of photosystem I and II, phycobilisomes, and NAD(P)H dehydrogenase are frequently distributed across separate genomic regions (Figure 3b and Table 2). Across the 25 trials in *Synechocystis*, operon context lifts Hit@1

from 0.5200 to 0.6400, a smaller increase than in the five enterobacterial, firmicute, and actinobacterial species where operons are tightly clustered. Adding Stage 2A singleton pruning and Stage 3 neuro-symbolic disambiguation, which inspects Pfam domain architectures and pathway neighborhood evidence across dispersed candidates, raises *Synechocystis* Hit@1 to 0.6800 and subunit F_1 from 0.4080 to 0.4480 (overall six-species Hit@1 = 0.9429 and $F_1 = 0.6737$). Local operon synteny resolves contiguous prokaryotic assemblies, and selective domain and pathway reasoning complements synteny when complex subunits are dispersed across the chromosome. Moving from prokaryotes to multicellular eukaryotes introduces a broader structural invariant: functional annotations must also conform to the anatomical body plan of the organism's clade.

2.4 Cross-ontology anatomical constraints correct eukaryotic annotations and preserve valid ancestral terms

We first tested whether ordinary multi-channel predictions in unperturbed eukaryotic proteomes contain endogenous taxonomic and anatomical contradictions. Evaluating the seven-channel Stage 0 ensemble across 6,155 proteins from six eukaryotes (*Saccharomyces cerevisiae*, *Caenorhabditis elegans*, *Drosophila melanogaster*, *Physcomitrium patens*, *Arabidopsis thaliana*, and *Mus musculus*) revealed 1,009 endogenous violations, concentrated in *C. elegans* and *D. melanogaster* where strong sequence and structural similarity transfers vertebrate organ-development and angiosperm terms onto invertebrate orthologs (Figure 3c and Table 3). Full GSPA eliminates all 1,009 endogenous violations while raising six-species mean CAFA $F_{\max}$ from 0.7096 to 0.7117. Because standard homology and language-model transfer across eukaryotic clades introduces frequent anatomical contradictions, we next asked which ontological representations are required to detect and eliminate them.

We compared direct GO taxon constraints, the Stage 1 soft anatomical prior, and Stage 2A deterministic cross-ontology enforcement on a stress-test cohort injecting 1,546 cross-clade developmental claims (1,452 post-calibration violations across 7,655 proteins; Figure 3c and Table 3). Direct GO taxon constraints leave 44.2% of the violations unresolved (642 of 1,452) because organ-level processes such as *heart development* (`GO:0007507`) and *neural crest cell migration* (`GO:0001755`) carry their taxonomic restrictions on the linked Uberon, PO, or CL anatomical classes instead of on the GO term itself. The Stage 1 soft anatomical prior eliminates 80.7% of violations on average across the six species, and Stage 2A deterministic enforcement over the GO–Uberon/PO/CL bridge eliminates 100% of violations while rescuing 81.2% of attenuated species-appropriate developmental annotations (502 of 618, reaching 90.1% in *M. musculus*). Enforcing anatomical validity across eukaryotes therefore requires traversing the logical definitions that link biological processes to clade-restricted anatomical structures. Once a term is flagged as anatomically inadmissible, the pipeline can either delete the term or relax it to its nearest valid GO ancestor, and we compared these two operations next.

When an invertebrate or plant protein receives a forbidden organ-specific annotation from a conserved signaling or transcriptional homolog, its participation in a broader developmental or morphogenetic process often remains valid. Evaluated against unmodified Swiss-Prot reference annotations, Stage 2A semantic ancestor relaxation retains 96.7% of the baseline annotation information content and raises six-species mean $F_{\max}$ to 0.9605 compared with 0.9568 under term deletion, outperforming deletion across all four multicellular animals and vascular plants and reaching $+0.0157$ $F_{\max}$ in *D. melanogaster* (Figure 3d and Table 3). In the unicellular fungus *S. cerevisiae* and the non-vascular moss *P. patens*, where injected organ-development claims land on housekeeping enzymes, deletion yields slightly higher precision and $F_{\max}$, whereas relaxation yields higher recall and information content. Semantic ancestor relaxation preserves valid ancestral developmental biology in tissue-forming eukaryotes while maintaining zero formal ontology violations. Having evaluated whole-genome constraints on complexes and ontological admissibility, we next examined how genomic architecture combines with systems biology to localize missing metabolic enzymes.

2.5 Genomic context and protein language models locate missing pathway enzymes

Reaction-level metabolic gap-filling identifies which reactions a network lacks for growth, leaving open which gene in the chromosome catalyzes each missing step. We first tested whether reaction-local genomic context (RLGC), which scores candidate genes by their base-pair proximity and co-directionality relative to anchor genes of pathway-adjacent reactions, can localize held-out enzymes prior to introducing protein language-model centroids. Across 256 strict leave-reaction-out trials in *E. coli* K-12 with sequence homologs of the held-out enzyme masked (Figure 4a and Table 4), sequence-only ranking places the true enzyme at Rank 1 for 1.95% of reactions, and METEOR's uniform reaction-level boost leaves this ordering unchanged because every candidate of a reaction receives the same score increment. Combining RLGC kernel density with the Stage 1 pathway coherence prior raises unconditional Hit@1 twelve-fold to 23.44% (34.29% within the candidate pool, $p < 10^{-12}$, exact McNemar test). Physical proximity to pathway-neighboring genes on the chromosome therefore provides locus-specific discrimination beyond reaction-level gap-filling. We next examined how local genomic context interacts with alignment-free protein language-model representations across metabolic roles and species.

When Swiss-Prot ESM-2 reaction embedding centroids are included alongside masked sequence alignments (Track B; Figure 4a–c and Table 4), centroid similarity alone

Table 3. Taxonomic and anatomical constraint enforcement and semantic ancestor relaxation across six eukaryotes. Violation columns report surviving formal contradictions under unconstrained Stage 0 pooling (stress-test total across 7,655 proteins with 1,546 injected cross-clade developmental claims, and endogenous unperturbed violations across 6,155 proteins in parentheses), direct GO taxon constraints (percentage of stress-test violations eliminated in parentheses), the Stage 1 soft anatomical prior, and Stage 2A symbolic enforcement. $F_{\max}$, propagated recall, and mean information content (IC, bits) are evaluated against unmodified Swiss-Prot annotations comparing Stage 0 pooling, Stage 2A term removal, and Stage 2A semantic ancestor relaxation. Developmental rescue reports the fraction of attenuated species-appropriate developmental claims (618 total) recovered above 0.50.

Species (Clade)	Proteins (Inj.)	Stage 0 Viol. (Endog.)	Direct GO Taxon Viol.	Stage 1 Soft Viol.	Stage 2A Viol.	Stage 0 $F_{\max}$	Removal $F_{\max}$ (Rec. / IC)	Relaxation $F_{\max}$ (Rec. / IC)	Dev. Rescue
S. cerevisiae (Fungi)	1,500 (300)	66 (3)	0 (100.0%)	0 (100.0%)	0 (100%)	0.9033	0.9604 (0.9249/12.02)	0.9592 (0.9349/12.07)	66.7% (2/3)
C. elegans (Nematoda)	1,500 (300)	389 (289)	190 (51.2%)	201 (48.3%)	0 (100%)	0.9015	0.9567 (0.9268/12.15)	0.9628 (0.9476/12.09)	66.7% (44/66)
D. melanogaster (Arthropoda)	1,500 (300)	894 (717)	426 (52.3%)	559 (37.5%)	0 (100%)	0.9076	0.9238 (0.8608/12.55)	0.9395 (0.8913/12.35)	70.7% (140/198)
P. patens (Bryophyta)	155 (46)	3 (0)	0 (100.0%)	0 (100.0%)	0 (100%)	0.8949	0.9828 (0.9697/9.64)	0.9792 (0.9796/9.79)	n/a (0/0)
A. thaliana (Magnoliopsida)	1,500 (300)	59 (0)	26 (55.9%)	1 (98.3%)	0 (100%)	0.8884	0.9497 (0.9054/11.13)	0.9547 (0.9232/11.18)	89.9% (62/69)
M. musculus (Mammalia)	1,500 (300)	41 (0)	0 (100.0%)	0 (100.0%)	0 (100%)	0.9271	0.9672 (0.9478/12.84)	0.9678 (0.9545/12.85)	90.1% (254/282)
6-Eukaryote Total / Mean	7,655 (1,546)	1,452 (1,009)	642 (55.8%)	761 (80.7% **mean**)	0 (100%)	0.9038	0.9568 (0.9226/11.72)	0.9605 (0.9385/11.72)	81.2% (502/618)
Endogenous Unperturbed Cohort	6,155 (0)	1,009	n/a	n/a	0 (100%)	0.7096	n/a	0.7117 **(Iter 1)** / 0.7109 **(Iter 3)**	n/a




Figure 4. Leave-reaction-out metabolic enzyme attribution and uncultured microbial dark-matter discovery. (**a**) Recovery of 256 held-out metabolic reactions in *E. coli* K-12 (calibration genome), reporting unconditional Hit@1 across all 256 reactions and conditional Hit@1 and Hit@3 within the Stage 0 candidate pool ($n = 175$–181) across sequence-only/METEOR uniform boosting, strict homology- and embedding-masked GSPA (Track A), RLGC operon kernel density alone, Stage 1 coherence with RLGC, ESM-2 reaction centroids alone, and full GSPA (Track B). (**b**) Role-stratified Top-1 recovery across 414 *E. coli* reactions spanning core biomass (Stratum A, $n = 307$), alternative catabolic (Stratum B, $n = 49$), peripheral/secondary metabolism (Stratum C, $n = 58$), and twilight-zone core reactions ($\leq 50\%$ identity, $n = 23$). (**c**) Out-of-sample transfer to *B. subtilis* 168 ($n = 144$ reactions) with parameters frozen on *E. coli*, alongside Stage 3 disambiguation of 50 ambiguous multi-candidate LRO gaps. (**d**) Mean candidate pathway gaps resolved per genome at posterior thresholds $q \geq 0.30$ and $q \geq 0.50$ across 97 uncultured MAGs from 16 bacterial and archaeal phyla.

recovers 62.11% of the 256 held-out *E. coli* enzymes unconditionally, and integrating RLGC and pathway coherence lifts recovery to 63.67% (90.06% within the candidate pool) by resolving ties among sequence-similar paralogs. Stratifying the broader 414-reaction *E. coli* benchmark across core biomass, alternative catabolic pathways, and peripheral or secondary metabolism shows that RLGC alone achieves its highest standalone pool recovery on alternative catabolic pathways (55.6%), whose specialized enzymes and transporters are tightly organized into co-regulated operons, while full GSPA maintains 82.8% to 93.5% pool Hit@1 across all three strata (Figure 4b and Table 4). Transferred with frozen hyperparameters to 144 held-out reactions in the Gram-positive bacterium *Bacil-*

Table 4. Leave-reaction-out (LRO) enzyme gene attribution in *E. coli* K-12, out-of-sample transfer to *B. subtilis* 168, and Stage 3 re-ranking of ambiguous metabolic gaps. *E. coli* K-12 served as the calibration genome. "All" evaluates every held-out reaction unconditionally (scoring reactions whose true enzyme is absent from the Stage 0 candidate pool as misses); "Pool" evaluates conditional ranking across reactions where the true gene enters the candidate pool (n_{cov}). Track A masks both sequence-alignment donors and embedding donors with cosine similarity > 0.50 and excludes reaction centroids; Track B masks sequence-alignment donors and retains Swiss-Prot ESM-2 reaction centroids.

Benchmark Cohort	Method / Ablation Variant	Reactions (n/n_{cov})	Hit@1 (All)	Hit@3 (All)	Hit@1 (Pool)	Hit@3 (Pool)	MRR (Pool)
E. coli K-12 Strict LRO	Sequence-Only / METEOR Uniform Boost	256/175	1.95% (5)	7.03%	2.86%	10.29%	0.1354
	GSPA Strict Homology-Masked (Track A)	256/181	11.33% (29)	34.77%	16.02%	49.17%	0.3685
	RLGC Operon Kernel Density Only	256/181	23.05% (59)	37.89%	32.60%	53.59%	0.4745
	GSPA Stage 1 Coherence + RLGC (No PLM)	256/175	23.44% (60)	36.33%	34.29%	53.14%	0.4864
	ESM-2 PLM Reaction Centroids Only	256/181	62.11% (159)	67.58%	87.85%	95.58%	0.9157
	Full GSPA (Track B: RLGC + Coherence + PLM)	256/181	63.67% (163)	68.36%	90.06%	96.69%	0.9330
E. coli K-12 Role-Stratified ($n = 414$)	Stratum A: Core Biomass (Full GSPA)	307/170	51.79% (159)	53.42%	93.53%	96.47%	0.9557
	Stratum B: Alternative Catabolic (Full GSPA)	49/27	48.98% (24)	53.06%	88.89%	96.30%	0.9216
	Stratum C: Peripheral / Secondary (Full GSPA)	58/29	41.38% (24)	48.28%	82.76%	96.55%	0.8793
	Stratum A Twilight Zone ($\leq 50\%$ Homology)	23/7	26.09% (6)	30.43%	85.71%	100.00%	0.9286
B. subtilis 168 Out-of-Sample LRO	GSPA Strict Homology-Masked (Track A)	144/76	11.11% (16)	31.94%	21.05%	60.53%	0.4221
	Full GSPA (Track B, Frozen Hyperparameters)	144/76	43.06% (62)	50.00%	81.58%	94.74%	0.8933
Ambiguous Multi-Candidate LRO	Stage 1 RLGC Operon Kernel Alone	50/50	34.00% (17)	86.00%	34.00%	86.00%	0.6020
	Stage 3 Guarded Re-Ranking (3-Seed Mean)	50/50	82.67% (41.3)	93.33%	82.67%	93.33%	0.8792

lus subtilis 168, full GSPA reaches 43.06% unconditional Hit@1 and 81.58% within the candidate pool (Figure 4c). Protein language models supply the primary candidate representation when reference enzymes exist for a reaction, while chromosomal context disambiguates competing paralogs and catabolic operons. Because flux balance on minimal medium leaves secondary metabolism unexercised, we next tested whether local genomic context extends to biosynthetic gene clusters and uncultured metagenomic dark matter.

Secondary metabolite pathways operate conditionally and carry zero steady-state flux under minimal glucose growth. Evaluated across 106 curated biosynthetic gene clusters from 12 bacterial species, GSPA's operon and pathway synteny scoring raises cluster gene recovery F_1 from 0.7395 to 0.8959 while maintaining 99.6% specificity across non-cluster chromosomal regions (Supplementary Table S10). Applying the same frozen pipeline to 97 high-quality metagenome-assembled genomes (MAGs) of uncultured bacteria and archaea spanning 16 phyla resolves 568 high-confidence enzyme attributions at posterior $q \geq 0.50$, lifting mean draft pathway completeness from 0.511 to 0.656 (Figure 4d and Supplementary Table S11). Local chromosomal synteny extends functional attribution to secondary metabolism and provides ranked, testable hypotheses for missing pathway steps in uncultivated microbial lineages. While RLGC ranks candidates around local pathway anchors, a metabolic repair should also be stoichiometrically active at the network level, which we audited next.

2.6 Positive-cost flux repair adds flux-carrying reactions and attributes them to gene loci

We next tested our third driver of improvement: replacing reaction-indicator MILP gap-filling with positive-cost lexicographic parsimonious flux balance analysis (pFBA) and competitive gene-locus attribution. Auditing metabolic network repairs across 108 bacterial genomes shows why negative-cost reaction-indicator MILP inflates reconstructions (Figure 5a and Supplementary Table S12): because candidate reactions above the addition threshold receive a negative objective cost, the solver adds a mean of 313.35 reactions per genome, of which 262.34 (83.72%) carry zero steady-state flux. Imposing a strictly positive addition cost over the elementally balanced universal network eliminates all zero-flux reaction padding (0.0% uncertified additions), and Stage 2B's two-step lexicographic pFBA further reduces active repairs to a minimal, flux-certified set of 5.70 mass-balanced reactions per genome while restoring 100% biomass feasibility. A positive addition cost coupled with lexicographic flux minimization therefore ensures that every added reaction actively participates in a mass-balanced metabolic route. We next asked whether attributing these flux-certified reactions competitively to candidate gene loci improves the identification of experimentally validated enzymes.

We evaluated enzyme discovery on 148 divergent bacterial enzymes across 22 host genomes whose physiological roles were established by random-barcoded transposon mutagenesis (Price-149) [54], comparing standalone predictors, METEOR v8, single-predictor GSPA, and multi-predictor GSPA (Figure 5b and Table 5). Because 55 of the 148 enzymes catalyze alternative catabolic pathways that are inactive on minimal glucose, single-medium MILP gap-filling lowers $F_{\max}$ on DeepProZyme, EnzBERT, and TopEC. Operating on a single predictor's proteome output, Stage 2B cognate-substrate probing and single-predictor GSPA improve Top-1 enzyme recovery on MAPred from 50 to 56 proteins ($F_{\max}$ from 0.5316 to 0.5625), on CLEAN from 60 to 63, and on DeepProZyme from 49 to 51 ($F_{\max}$ to 0.5757). Combining multi-predictor Stage 0 pooling with subsequent locus-level refinement produces both strong pooling gains on lower-recall baselines (where Stage 0 pooling alone provides the peak Top-1 accuracy on EnzBERT, GraphEC, and TopEC) and additive post-pooling refinement gains on the leading predictors, raising MAPred from 58 after Stage 0 pooling to 73 (49.3% Top-1 accuracy, $F_{\max} = 0.6481$), CLEAN from 63 to 66, and DeepProZyme from 54 to 59 (Table 5). Complementary predictor pooling and competitive stoichiometric locus attribution therefore




Figure 5. Stoichiometric flux certification, fitness-validated enzyme discovery, and experimental phenotype prediction. (**a**) 108-genome stoichiometric invariant audit comparing total added reactions, flux-certified additions ($|v_j| > 10^{-6}$), zero-flux reaction padding, and elementally imbalanced additions per genome across negative-cost MILP (METEOR v8), positive-cost threshold repair, and Stage 2B lexicographic pFBA (symmetric logarithmic scale). (**b**) Per-protein enzyme $F_{\max}$ on 148 RB-TnSeq fitness-validated bacterial enzymes (Price-149) across six baseline predictors, single-predictor METEOR v8, Stage 2B cognate-substrate probing, single-predictor GSPA, and multi-predictor GSPA. (**c**) Role-stratified LRO enzyme recovery ($n = 414$) across four donor-homology masking tiers (100%, $\leq 70\%$, $\leq 50\%$, and $\leq 30\%$ identity). (**d**) Experimental carbon-source growth (2,332 assays, 529 strains) and diagnostic enzyme activity (17,496 assays, 3,017 type strains) in BacDive across unsupervised reconstructions, blind Track 1 and phenotype-anchored Track 2 GSPA, genus-blocked DiASPora, and the GSPA + DiASPora hybrid.

Table 5. Refinement of 148 RB-TnSeq fitness-validated bacterial enzymes (Price-149) across 22 host genomes. Columns report correct Top-1 EC predictions (per-protein $F_{\max}$ in parentheses) for each standalone predictor, single-predictor METEOR v8, Stage 2B refinement under minimal glucose and with local-context cognate-substrate probing, full single-predictor GSPA (3 iterations), Stage 0 multi-predictor pooling, and full multi-predictor GSPA (iteration 3). The final column summarizes the Top-1 (and $F_{\max}$) change from standalone baseline to single-predictor and multi-predictor GSPA. Target-only sequence tools (DCGM and ECrecer) lack proteome-wide host background runs and are reported as standalone baselines.

Baseline Predictor	Standalone Base	+ METEOR v8	+ Stage 2B (Min. Glc.)	+ Stage 2B (Cognate Probe)	+ GSPA Single-Pred	+ Stage 0 Multi-Pool	+ GSPA Multi-Pred	Single / Multi Gain
CLEAN (2023)	60 (0.5295)	61 (0.5343)	60 (0.5300)	60 (0.5435)	63 (0.5214)	63 (0.5924)	66 (0.6085)	+3 (+0.014 probe) / +6 (+0.079)
MAPred (2025)	50 (0.5316)	52 (0.5469)	51 (0.5362)	50 (0.5356)	56 (0.5625)	58 (0.5698)	73 (0.6481)	+6 (+0.031) / +23 (+0.117)
DeepProZyme (2023)	49 (0.5574)	49 (0.5427)	49 (0.5625)	51 (0.5757)	48 (0.5620)	54 (0.5973)	59 (0.6061)	+2 (+0.018 probe) / +10 (+0.049)
EnzBERT (2023)	44 (0.4275)	46 (0.4254)	44 (0.4287)	44 (0.4310)	43 (0.4293)	62 (0.5964)	54 (0.5844)	+0.004 $F_{\max}$ / +18 **pool** (+10 **iter 3**)
GraphEC (2024)	18 (0.1404)	18 (0.1419)	18 (0.1422)	18 (0.1489)	18 (0.1558)	48 (0.5067)	48 (0.4718)	+0.015 $F_{\max}$ / +30 (+0.331)
TopEC (2025)	5 (0.0757)	5 (0.0753)	5 (0.0772)	5 (0.0811)	5 (0.0928)	46 (0.5301)	26 (0.2465)	+0.017 $F_{\max}$ / +41 **pool** (+21 **iter 3**)
DCGM (2025, Target-Only)	59 (0.5276)	n/a	n/a	n/a	n/a	n/a	n/a	Target-sequence baseline
ECrecer (2024, Target-Only)	58 (0.5282)	n/a	n/a	n/a	n/a	n/a	n/a	Target-sequence baseline

work additively on the top enzyme predictors to identify divergent, fitness-validated bacterial enzymes. Having evaluated gene-level enzyme recovery, we next examined whether systems-level pathway pooling and coupling predict experimental organismal phenotypes at scale.

2.7 Systems-level annotations predict experimental bacterial phenotypes

We next evaluated whether systems-level pathway pooling and subunit-aware modeling predict experimentally measured bacterial physiology across thousands of strains in BacDive [55, 56]. We tested blind genome-to-phenotype predictions (0% phenotype training labels) across 2,332 carbon-source utilization assays in 529 strains, 17,496 diag-

Table 6. Experimental carbon-source growth and diagnostic enzyme phenotype prediction in BacDive. Track 1 evaluates blind genome-to-phenotype prediction (0% phenotype labels) and 5-fold genus-blocked cross-validation (pooled out-of-fold metrics across unseen genera). Track 2 reveals 30% of each strain's assay rows as partial phenotypic evidence and evaluates predictions on the held-out 70%. Binary reconstruction pipelines (CarveMe, ModelSEED, and gapseq) output $\{0, 1\}$ calls whose single-point ROC-AUC equals balanced accuracy; GSPA and supervised models output continuous probabilities. Middle rows report the 10-enzyme diagnostic panel (50,075 assays across 6,313 strains), heteromeric urease (`EC 3.5.1.5`), and the 108-genome catabolic panel. Runtime is reported per genome after base reconstruction (end-to-end runtime in parentheses).

Assay Cohort & Protocol	Method / Pipeline	Strains / Assays	Precision	Recall	F_1 Score [95% CI]	Bal. Acc.	ROC-AUC	Runtime / Genome
Track 1: Carbon Growth (Blind & Genus-CV)	CarveMe v1.5.2 (Unsupervised Binary)	529/2,332	0.6844	0.4105	0.5132 [0.489, 0.537]	0.6596	0.6596	28.4 s
	ModelSEED v2.6.1 (Unsupervised Binary)	529/2,332	0.7510	0.5689	0.6474 [0.625, 0.671]	0.7392	0.7392	42.0 s
	gapseq v1.2 (Unsupervised Binary)	529/2,332	0.8071	0.7836	0.7952 [0.777, 0.812]	0.8469	0.8469	185.0 s
	GSPA Track 1 (100% Blind Continuous)	529/2,332	0.8071	0.7836	0.7952 [0.778, 0.813]	0.8469	0.8567	1.1 s (256.5 s)
	DiASPora GBDT (5-Fold Genus-Blocked CV)	529/2,332	0.8475	0.8491	0.8483 [0.833, 0.862]	0.8879	0.9519	0.05 s (255.5 s)
	GSPA + DiASPora Hybrid (5-Fold Genus-CV)	529/2,332	0.8766	0.8655	0.8710 [0.858, 0.884]	0.9035	0.9567	1.2 s (256.6 s)
Track 1: Diagnostic Enzymes (Blind & Genus-CV)	CarveMe v1.5.2 (Unsupervised Binary)	3,017/17,496	0.6808	0.3163	0.4319 [0.422, 0.442]	0.5844	0.5844	28.4 s
	gapseq v1.2 (Unsupervised Binary)	3,017/17,496	0.7366	0.8184	0.7753 [0.769, 0.781]	0.7461	0.7461	185.0 s
	GSPA Track 1 (100% Blind Continuous)	3,017/17,496	0.7366	0.8184	0.7753 [0.769, 0.781]	0.7461	0.7520	1.1 s (256.5 s)
	DiASPora GBDT (5-Fold Genus-Blocked CV)	3,017/17,496	0.8427	0.8676	0.8550 [0.850, 0.860]	0.8535	0.9061	0.05 s (255.5 s)
	GSPA + DiASPora Hybrid (5-Fold Genus-CV)	3,017/17,496	0.8712	0.8715	0.8713 [0.867, 0.876]	0.8714	0.9097	1.2 s (256.6 s)
10-EC Diagnostic Panel (50,075 Assays)	DeepProZyme Base → + METEOR v8	6,313/50,075	0.6820 → 0.6743	0.7575 → 0.7604	0.7178 → 0.7148	0.6305 → 0.6251	n/a	Over-gap-filling drop
Heteromeric Urease (`EC 3.5.1.5`)	DeepProZyme Base → + METEOR v8	5,026 strains	0.5812 → 0.5140	0.5448 → 0.5842	0.5624 → 0.5469	0.6120 → 0.5890	n/a	Isolated subunit FP
108-Genome Catabolic Assays	DeepProZyme Base → + METEOR v8	108/1,622	0.6929 → 0.6925	0.6507 → **0.6531**	0.6711 → **0.6722**	0.6716 → **0.6723**	n/a	+3 TP rescues
Track 2: Carbon Held-Out 70% (30% Revealed)	gapseq v1.2 (Unsupervised Binary)	529/1,641	0.8124	0.7730	0.7922 [0.770, 0.812]	0.8445	0.8445	185.0 s
	GSPA Track 2 (Unsupervised Bayesian Anchor)	529/1,641	0.8473	0.9308	0.8871 [0.872, 0.901]	0.9259	0.9613	1.1 s (256.5 s)
	GSPA Track 2 + DiASPora Hybrid	529/1,641	0.9163	0.9288	0.9225 [0.910, 0.935]	0.9445	0.9776	1.2 s (256.6 s)
Track 2: Enzymes Held-Out 70% (30% Revealed)	gapseq v1.2 (Unsupervised Binary)	3,017/12,316	0.7376	0.8195	0.7764 [0.769, 0.784]	0.7471	0.7471	185.0 s
	GSPA Track 2 + DiASPora Hybrid	3,017/12,316	0.8705	0.8741	0.8723 [0.867, 0.877]	0.8721	0.9102	1.2 s (256.6 s)

nostic enzyme assays in 3,017 type strains, and an extended panel of 50,075 assays for 10 diagnostic enzymes across 6,313 strains (Figure 5d and Table 6). On the 50,075-assay panel, uniform MILP gap-filling (METEOR v8) lowers overall F_1 from 0.7178 to 0.7148, driven by heteromeric enzymes such as urease (`EC 3.5.1.5`; F_1 from 0.5624 to 0.5469), where boosting an isolated UreC alpha subunit without the accessory UreA, UreB, and UreDFG proteins generates false-positive calls. By pooling calibrated pathway calls from gapseq, ModelSEED, and CarveMe with unsupervised cross-assay pathway co-occurrence coupling, blind Track 1 GSPA matches gapseq's binary F_1 on carbon growth (0.7952) and diagnostic enzymes (0.7753) while raising continuous ROC-AUC to 0.8567 on carbon growth and 0.7520 on diagnostic enzymes (Table 6). Continuous pathway coupling and subunit-complete enzyme modeling therefore yield well-calibrated physiological probabilities from genomic reconstructions alone. We next examined whether these mechanistic pathway posteriors improve out-of-genus transfer when combined with supervised phenotype classifiers and whether partial phenotypic observations from a strain refine its remaining unassayed pathways.

Under five-fold genus-blocked cross-validation, in which every test strain belongs to a taxonomic genus unseen during training, combining GSPA pathway posteriors with the DiASPora gradient-boosting classifier [57] raises out-of-genus F_1 to 0.8710 on carbon growth and 0.8713 on diagnostic enzymes, outperforming all individual metabolic reconstructions and supervised baselines across folds (Table 6 and Supplementary Table S13). In Track 2, where 30% of each strain's experimental assays are revealed as partial phenotypic evidence and the remaining 70% are held out, conditioning GSPA's pathway posteriors via an unsupervised Bayesian update lifts held-out carbon-growth F_1 from 0.7922 (gapseq) to 0.8871, and the hybrid with DiASPora reaches $F_1 = 0.9225$ (Table 6). Mechanistic pathway coupling provides domain-generalizable features for unseen genera and propagates partial experimental observations across coupled metabolic pathways. After deterministic probabilistic and stoichiometric refinement, a small set of ambiguous loci remains, which we examined in Stage 3.

2.8 Guarded language-model reasoning resolves residual ambiguity

Deterministic inference in Stages 1 and 2 resolves the vast majority of proteome annotations, leaving a small fraction of structured anomalies: unresolved pathway gaps, incomplete heteromeric complexes, uncovered essential functions, and borderline ontology conflicts. Across the 88 multi-kingdom proteomes, deterministic anomaly triggers select a mean of 1.84% of proteins per proteome (fewer than 2.5% in every domain of life), bounding the computational cost of invoking a language model. Evaluated across three random seeds on a de-identified benchmark of 120 anomalies from seven prokaryotic and eukaryotic genomes with all gene symbols masked and candidate ordering randomized, the guarded multi-turn agent reaches a mean F_1 of 0.9115 ± 0.0008 with 0.0% accepted ontology violations, whereas ablating the symbolic guardrail admits a cross-clade violation and reduces F_1 to 0.8515 (Figure 6a,b,d, Table 7, and Supplementary Table S14). Selective triggering therefore confines language-model inference to fewer than 2.5% of proteins, while deterministic verification guarantees that accepted proposals satisfy all taxonomic and anatomical axioms. We next tested whether tool-mediated reasoning can resolve multi-candidate metabolic gaps and prospective sequence targets.

When multiple candidate genes in a genomic neighborhood share similar RLGC scores for a missing reaction, resolving the true enzyme requires synthesizing domain architecture, operon structure, and pathway stoichiometry. Across 50 unique-protein ambiguous leave-reaction-out gaps in *E. coli* K-12 where Stage 1 RLGC alone ranks the true enzyme first in 34.0% of cases, Stage 3 multi-turn re-ranking lifts Top-1 recovery to a three-seed mean of




Figure 6. Evaluation of Stage 3 selective neuro-symbolic reasoning on de-identified benchmarks. (**a**) Precision, recall, F_1, and resolution accuracy across 120 de-identified anomalies (30 pathway gaps, 30 incomplete complexes, 30 borderline essential functions, and 30 borderline ontology conflicts) comparing deterministic Stages 1 and 2, posterior thresholding (≥ 0.35) with the symbolic guardrail, 5-fold cross-validated supervised classifiers on pre-tool and post-tool features, fast 1-step Stage 3, unguarded multi-turn Stage 3, and full guarded multi-turn Stage 3 (3-seed mean). (**b**) Rate of accepted taxonomic and anatomical violations across configurations. (**c**) Rank distribution of the true enzyme across 50 unique-protein ambiguous LRO metabolic gaps before and after Stage 3 re-ranking. (**d**) Concurrency-normalized wall-clock latency per anomaly across seven prokaryotic and eukaryotic species (mean 1.84% of proteome loci triggered).

Table 7. Stage 3 neuro-symbolic disambiguation compared against deterministic, threshold, and supervised baselines across 120 de-identified anomalies and 50 ambiguous metabolic gaps. All gene symbols are masked and candidate order is hash-shuffled. The 120 anomalies comprise 54 true-positive promotable gaps, 36 score-matched decoys, and 30 ontology conflicts (15 genuine cross-clade conflicts and 15 adversarial traps). Supervised classifiers are evaluated via 5-fold cross-validation using either 3 pre-tool scalars or 8 post-tool features combined with the symbolic guardrail. Latency reports concurrency-normalized wall-clock seconds per anomaly at batch concurrency 8 (raw HTTP latency in parentheses).

Resolution Method	Precision	Recall	F_1 Score	Accuracy	Prom. / Relax.	Violations Accepted	LRO-50 Hit@1 (MRR)	Latency / Tools
Stage 1 + Stage 2 Deterministic Only	0.0000	0.0000	0.0000	0.5500	0/0	0.0% (0/120)	34.0% (0.6020)	0.00 s / 0.00
Posterior Threshold (≥ 0.35) + Guardrail	0.5686	0.5370	0.5524	0.7083	29/30	0.0% (0/120)	34.0% (0.6020)	< 0.01 s / 0.00
Logistic Regression (5-Fold CV, 3 Pre-Tool Features)	0.9000	0.6429	0.7500	0.8000	36/0	2.5% (3/120)	94.0% (0.9670)	< 0.01 s / 0.00
GBDT (5-Fold CV, 3 Pre-Tool Features)	0.9535	0.7321	0.8283	0.8583	41/0	0.0% (0/120)	90.0% (0.9467)	< 0.01 s / 0.00
Stage 3 Fast 1-Step Mode (Max 2 Turns)	0.8889	0.1702	0.2857	0.6667	16/3	0.0% (0/120)	34.0% (0.6020)	1.56 s (12.5 s) / 1.92
Stage 3 Multi-Turn Without Symbolic Guardrail	0.9149	0.7963	0.8515	0.8750	43/1	0.8% (1/120)	82.0% (0.8740)	3.86 s (30.9 s) / 3.92
Stage 3 Multi-Turn + Guardrail (3-Seed Mean ± s.d.)	0.9722 ± 0.009	0.8581 ± 0.009	0.9115 ± 0.001	0.9278 ± 0.004	46.3/13.7	0.0% (0/120)	$82.7 \pm 0.9\%$ (0.879)	4.07 s (32.5 s) / 4.18
Logistic Regression (5-Fold CV, 8 Tool Features + Guardrail)	0.9583	0.8889	0.9216	0.9333	48/2	0.0% (0/120)	96.0% (0.9767)	< 0.01 s / 0.00
Random Forest (5-Fold CV, 8 Tool Features + Guardrail)	0.9286	0.9630	0.9455	0.9500	52/2	0.0% (0/120)	92.0% (0.9557)	< 0.01 s / 0.00
GBDT (5-Fold CV, 8 Tool Features + Guardrail)	0.9800	0.9074	0.9423	0.9500	49/2	0.0% (0/120)	92.0% (0.9557)	< 0.01 s / 0.00
MLP (5-Fold CV, 8 Tool Features + Guardrail)	0.9811	0.9630	0.9720	0.9750	52/2	0.0% (0/120)	92.0% (0.9557)	< 0.01 s / 0.00

$82.7 \pm 0.9\%$ ($p = 2.8 \times 10^{-7}$, exact McNemar test; Figure 6c and Table 7). On 769 prospective CAFA6 targets evaluated as isolated sequences with Stages 0, 2A, and 3 active, guarded reasoning reaches macro $F_{\max} = 0.5433$, outperforming both a multilayer perceptron trained on ESM-2 and ProstT5 3Di embeddings (0.4957) and an unguarded 14-billion-parameter agent baseline (0.5130; Supplementary Table S15). Tool-mediated evidence synthesis resolves ties among candidate enzymes after deterministic priors have narrowed the search space. Because a supervised tabular classifier can also be trained on numerical features extracted by the same tools, we compared zero-shot

guarded reasoning against cross-validated classifiers to establish the operating regime of each method.

We compared the zero-shot guarded agent against logistic regression, random forest, gradient-boosted decision tree (GBDT), and multilayer perceptron classifiers evaluated by five-fold cross-validation on either three pre-tool summary scalars or eight post-tool genomic and domain features paired with the symbolic guardrail (Table 7). On the 120 anomalies, the zero-shot agent ($F_1 = 0.9115$) outperforms all classifiers restricted to pre-tool scalars ($F_1 = 0.7500$–0.8283), while supervised classifiers trained on the eight post-tool features reach up to $F_1 = 0.9720$ on binary anomaly promotion and 96.0% Hit@1 on the 50 metabolic gaps at sub-millisecond inference latency. On borderline ontology conflicts, the zero-shot agent proposes a mean of 13.7 valid ancestral GO relaxations compared with 2 for the post-tool classifiers, which select among fixed candidate labels. Supervised post-tool classifiers are therefore preferable for high-throughput binary promotion and candidate ranking when labeled anomalies are available, whereas zero-shot guarded reasoning provides zero-label operation and multi-class ancestral GO relaxation (see Supplementary Section S6 and Supplementary Table S16 for an application to five desert extremophile isolates from the Empty Quarter).

3 Discussion

Protein function prediction has traditionally been formulated as a per-protein classification task in which each sequence or structure is scored independently [1, 20, 22]. GSPA reformulates proteome annotation as staged inference over the complete genome of an organism, and our evaluation shows that three complementary drivers account for its performance. First, calibrated pooling of sequence homology, domain profiles, protein language models, and structural similarity establishes a strong per-protein probability distribution in Stage 0. Second, whole-genome priors and deterministic cross-ontology enforcement in Stages 1 and 2A update these probabilities using organism-wide invariants over operons, core essential functions, obligate heteromeric complexes, and taxonomic and anatomical consistency. Third, stoichiometric flux modeling, competitive gene-locus attribution, and selective ontology-guarded reasoning in Stages 1, 2B, and 3 connect metabolic network requirements and multi-source genomic evidence to individual gene loci. These three layers act on complementary subsets of the proteome: per-protein pooling governs conserved proteins with strong sequence or structural similarity, while whole-genome architecture and systems-level modeling resolve twilight-zone enzymes, incomplete complexes, and cross-clade homology transfers.

Stage 0 builds directly on the multi-source ensemble paradigm introduced by GOLabeler [24] and NetGO [25, 26], which showed that integrating sequence alignment, domain signatures, motifs, network associations, and language-model embeddings outperforms individual representations in CAFA evaluations. GSPA extends this principle to genome-scale inference by calibrating raw predictor outputs via Platt scaling, grouping correlated evidence channels so that shared homology or embedding signals are counted once, and pooling across groups via Noisy-OR into an anchored log-odds prior $L^{(0)}$. Because every subsequent Stage 1 update is added to $L^{(0)}$, sequence homology, domain profiles, protein language models, and structural similarity remain the foundation of the framework. Conserved proteins retain their high-confidence Stage 0 assignments, and the recall of the Stage 0 candidate pool defines the space of candidate functions that genome-scale priors refine.

Exploiting whole-genome information in Stages 1 and 2A is timely because long-read sequencing and modern repeat-graph assemblers now routinely recover complete or near-complete chromosomes for both cultivated isolates and metagenome-assembled genomes [10–12]. Complete assemblies expose two distinct scales of genomic organization. Global organismal invariants identify which functional requirement a draft annotation violates: essentiality profiles flag missing core cellular machinery, complex stoichiometry flags obligate heteromeric complexes with missing partner subunits, and cross-ontology axioms linking GO biological processes to Uberon, PO, and CL anatomy flag processes incompatible with the organism's body plan [9, 44, 45]. Local chromosomal architecture then identifies which specific gene satisfies a missing requirement: in our leave-subunit-out experiments, global complex coherence raised subunit recall while co-directional operon synteny lifted top-ranked subunit identification from 56.0% to 93.7% (Table 2). Similarly, in multicellular animals and vascular plants, relaxing anatomically forbidden GO terms along the ontology hierarchy to their nearest valid ancestors preserved conserved tissue- and process-level biology that deletion-based filters discard (Table 3).

Our third driver advances functional inference beyond sequence homology by integrating systems biology with genomic architecture, addressing the specific mathematical limitations we identified in our earlier method METEOR [42]. METEOR coupled metabolic gap-filling with enzyme prediction through a reaction-indicator MILP that assigned negative objective costs to candidate reactions above an addition threshold and boosted all candidate genes of a selected reaction uniformly. Our 108-genome stoichiometric audit shows that negative addition costs incentivize the solver to accumulate 83.7% zero-flux reaction padding (Figure 5a), while uniform gene boosting elevates non-cognate paralogs and isolated complex subunits. GSPA resolves both issues by coupling reaction-local genomic context (RLGC) in Stage 1 with role-aware disjunctive normal form (DNF) aggregation of isozymes and required subunits, positive-cost lexicographic pFBA [48], and competitive locus responsibility against an explicit unassigned-gene hypothesis in Stage 2B. Under positive addition costs, every added reaction carries steady-state flux in an elementally balanced network, and competitive re-

sponsibility concentrates the backward log-odds update on the candidate gene supported by operon neighborhood and pathway synteny, improving gene-level discovery on RB-TnSeq fitness-validated bacterial enzymes [54] (Table 5). At the organism level, pooling calibrated metabolic reconstructions with unsupervised cross-assay pathway coupling and subunit-complete enzyme rules accurately predicts carbon-source growth and diagnostic enzyme phenotypes across unseen genera in BacDive (Table 6).

Finally, Stage 3 shows how tool-using language models and formal symbolic reasoning can be combined selectively at proteome scale. By restricting invocation to deterministic post-Stage-2 anomaly triggers that select fewer than 2.5% of proteins per proteome, GSPA restricts language-model inference to ambiguous loci while leaving conserved genes resolved by Stages 0–2 untouched. Within each triggered anomaly, the language model synthesizes heterogeneous operon, domain, pathway, and ontology evidence through read-only tools, while a deterministic symbolic guardrail verifies every proposed resolution against GO, NCBI Taxonomy, and the GO–Uberon/PO/CL bridge, mirroring neuro-symbolic verification principles in biomedical curation [46, 47]. Our comparison with five-fold cross-validated tabular classifiers delineates clear deployment regimes: supervised classifiers trained on post-tool features and paired with the symbolic guardrail provide the highest binary promotion accuracy at sub-millisecond latency when labeled anomalies exist, whereas the zero-shot guarded agent operates without anomaly training labels and performs multi-class traversal to valid ancestral GO terms.

3.1 Limitations

Our evaluation design establishes how genome-scale priors and stoichiometric refinement build upon a per-protein candidate set, and four aspects of this design define concrete next experiments. First, on well-annotated reference genomes, roughly 85% of proteins already carry high-confidence homologs in Stage 0 ($P_0 \geq 0.90$), so Stage 0 governs overall protein-centric $F_{\max}$ while Stages 1–3 concentrate their updates on twilight-zone proteins, pathway gaps, incomplete complexes, and taxonomic conflicts. Because the composite GAEF score evaluates the same structural invariants over essentiality, pathway and complex coherence, and taxonomic consistency that Stages 1 and 2 optimize, we report GAEF as an internal structural audit score paired with independent held-out GO, EC, leave-reaction-out, leave-subunit-out, Price-149, and BacDive evaluations, and we evaluate both endogenous unperturbed contradictions (1,009 eukaryotic violations) and synthetic cross-clade stress-test injections. Similarly, in leave-reaction-out evaluation, unconditional enzyme recovery is bounded by whether Stage 0 places the true gene into the candidate pool (70.7% pool coverage in *E. coli* and 52.8% in *B. subtilis*), and Track B uses released ESM-2 weights pretrained on UniRef50, making strict Track A cosine masking the primary test of de novo context recovery. In addition, the prospective sequence benchmarks (CAFA6 and Holdout-125) provide isolated target sequences without source genome coordinates, evaluating Stages 0, 2A, and 3, while the leave-subunit-out benchmark samples domain-matched decoys from non-operonic loci. Expanding Stage 0 candidate generation with remote structural fold search, evaluating protein language models trained under strict chronological splits, sampling leave-subunit-out decoys from within-operon paralogs, and linking prospective benchmark targets to their complete source genome assemblies will directly measure how increased candidate recall and full chromosomal context affect these regimes.

Second, Stage 2B models steady-state, mass-balanced metabolic flux over reactions mapped in the sanitized universal network under specified growth media, and Stage 2A enforces anatomical constraints over the 2,715 GO terms that currently carry explicit Uberon, PO, or CL logical definitions. Experimental growth and enzyme phenotypes in Price-149 and BacDive also depend on transcriptional regulation, transporter substrate specificity, and cofactor synthesis, while anatomical relaxation in unicellular fungi and non-vascular plants reflects the granularity of tissue-level ancestor annotations in reference databases. Integrating condition-specific transcriptomic expression priors and automated transporter substrate predictors into Stage 2B, alongside expanding cross-ontology logical definitions across additional GO biological processes and eukaryotic clades, will extend stoichiometric and ontological coverage to broader physiological states.

Third, Stage 3 was evaluated using a single locally hosted open-weight instruction model across three random seeds, and the 568 high-confidence pathway gap fills across 97 uncultured MAGs (as well as the extremophile pathway completions in the Empty Quarter cohort) represent ranked computational hypotheses. Benchmarking additional open-weight model families on the de-identified 120-anomaly suite, evaluating zero-shot Stage 3 against supervised post-tool classifiers under a leave-clade-out anomaly split, implementing an adaptive router that dispatches binary in-distribution promotions to the >100-fold faster supervised classifier while reserving multi-turn language-model reasoning for novel clades and ancestral GO relaxations, and validating top-ranked MAG and extremophile enzyme candidates via heterologous expression and mutant fitness assays will establish both computational generality and experimental confirmation.

4 Methods

4.1 Problem formulation and four-stage architecture

Let $\mathcal{P} = \{p_1, \ldots, p_N\}$ denote the protein-coding genes of a target genome with NCBI Taxonomy identifier τ_{tax}, and let $\mathcal{K} = \{(c_i, s_i, e_i, \zeta_i)\}_{i=1}^{N}$ denote their genomic coordinates (contig c_i, start s_i, end e_i, and strand $\zeta_i \in \{+, -\}$).

Let $\mathcal{T}$ denote the Gene Ontology (GO) directed acyclic graph under the true-path relation $t \sqsubseteq a$ (`is_a` and `part_of`), with ancestor closure $\mathrm{Anc}(t)$, $\mathcal{M}$ the universal metabolic reaction network with stoichiometric matrix $\mathbf{S}$ and reaction–EC–GO mappings, and $\mathcal{O}$ the formal taxonomic and cross-ontology anatomical axiom set. GSPA estimates a proteome-wide posterior probability matrix $\mathbf{P} \in [0,1]^{N\times|\mathcal{T}|}$ (and corresponding EC probabilities in enzyme benchmarks) conditioned jointly on $\mathcal{K}$, τ_{tax}, $\mathcal{M}$, and $\mathcal{O}$ across four sequential stages (Figure 1): Stage 0 calibrates and pools per-protein evidence channels; Stage 1 performs iterative whole-proteome Bayesian log-odds refinement; Stage 2A applies deterministic symbolic constraint enforcement with semantic ancestor relaxation while Stage 2B performs positive-cost lexicographic parsimonious flux balance analysis (pFBA) and competitive gene-locus attribution; and Stage 3 applies selective tool-mediated neuro-symbolic reasoning verified by a symbolic ontology guardrail. Supplementary Table S1 summarizes all eight benchmark suites, cohorts, Stage 0 channel configurations, and active pipeline stages.

4.2 Stage 0: multi-channel evidence calibration and correlation-grouped Noisy-OR pooling

Stage 0 ingests functional claims from either a standalone predictor (such as CLEAN [21], MAPred [58], DeepProZyme [52], EnzBERT [51], GraphEC [59], or TopEC [60]) or up to seven complementary per-protein channels $e \in \mathcal{E}_{\mathrm{ch}}$: (1) DIAMOND v2.1.8 sequence alignment against experimentally annotated Swiss-Prot [13]; (2) HMMER 3.4 profile HMM search against Pfam-A mapped via Pfam2GO [16]; (3) InterProScan v5.68-100.0 signature integration [17]; (4) ESM-2 (650M) embedding k-nearest-neighbor transfer [18]; (5) ProtT5-XL-U50 embedding k-nearest-neighbor transfer [19]; (6) neural sequence classifiers (ProteInfer [50], DeepGOWeb [61], and DeepGO-SE [20]); and (7) structural similarity transfer via FoldSeek [23] and ProstT5 3Di structural tokens [62].

Each raw claim $c = (p, f, s(c), e(c))$ with raw score $s(c) \in [0,1]$ is mapped to a calibrated probability via Platt logistic scaling, $\tilde{q}(c) = \mathrm{clip}\big(\sigma_{\log}(a_{e(c)}\, s(c) + b_{e(c)}),\ \varepsilon_{\mathrm{cal}},\ 1-\varepsilon_{\mathrm{cal}}\big)$ with $\varepsilon_{\mathrm{cal}} = 10^{-4}$, and weighted by channel reliability $r_{e(c)} \in (0,1]$ to give $q(c) = r_{e(c)}\,\tilde{q}(c)$ (channel parameters and reliabilities in Supplementary Section S1). To prevent double-counting correlated evidence sources, claims for each pair (p,f) are partitioned into correlation groups $\mathcal{G}_{\mathrm{corr}}(p,f)$ by evidence modality (homology, protein-sequence ML, structure, genomic context, genomic language models, inferred context, sequence motifs, localization, and domain-specific predictors; Supplementary Section S1). Stage 0 takes the maximum weighted probability within each group (with an optional second-opinion weight $\beta_{\mathrm{so}} = 0.0$ by default) and pools across independent groups via Noisy-OR:

$$q_g(p,f) = \mathrm{clip}\Big(\max_{c\in g}\big(r_{e(c)}\tilde{q}(c)\big) + \beta_{\mathrm{so}}\, q_{g,(2)}(p,f),\ 10^{-3},\ 0.999\Big),$$
$$P_0(p,f) = \mathrm{clip}\left(1 - \prod_{g\in\mathcal{G}_{\mathrm{corr}}(p,f)} \big(1 - q_g(p,f)\big),\ 10^{-6},\ 1-10^{-6}\right). \tag{1}$$

Stage 0 converts $P_0(p,f)$ into the anchored log-odds $L^{(0)}(p,f) = \mathrm{clip}\big(\log\frac{P_0(p,f)}{1-P_0(p,f)}, -12, 12\big)$ and propagates probabilities upward along GO `is_a` and `part_of` edges.

4.3 Stage 1: whole-proteome iterative Bayesian log-odds refinement

Stage 1 refines the proteome-wide probability matrix $\mathbf{P}^{(k)}$ over iterations $k = 1, \ldots, K_{\max}$ by adding six genome-scale prior log-odds shifts $B_m\big(p, f \mid \mathbf{P}^{(k-1)}, \mathcal{K}, \tau_{\mathrm{tax}}, \mathcal{M}, \mathcal{O}\big)$ to the fixed Stage 0 anchor $L^{(0)}(p,f)$ under Jacobi damping:

$$\tilde{L}^{(k)}(p,f) = \mathrm{clip}\Bigg(L^{(0)}(p,f) + \sum_{m=1}^{6} w_m\, B_m\big(p,f \mid \mathbf{P}^{(k-1)}, \mathcal{K}, \tau_{\mathrm{tax}}, \mathcal{M}, \mathcal{O}$$
$$L^{(k)}(p,f) = (1-\gamma_{\mathrm{iter}})L^{(k-1)}(p,f) + \gamma_{\mathrm{iter}}\tilde{L}^{(k)}(p,f), \qquad P^{(k)}(p,f) = \tag{2}$$

with unit module weights $w_m = 1.0$, damping factor $\gamma_{\mathrm{iter}} = 0.50$, and maximum iterations $K_{\max} = 6$. Iteration terminates when the mean absolute probability change across updated entries falls below $\epsilon = 5\times10^{-3}$, and reverts to the last stable iterate if the mean change increases over two consecutive steps. All 18 configurable Stage 1 and Stage 2B hyperparameters were calibrated on *E. coli* K-12 MG1655 and held frozen across all other species (Supplementary Section S2 and Supplementary Table S3). The six priors are defined as follows:

1. *Essentiality prior (B_1, $\alpha_{ess} = 1.50$):* Let $\mathcal{F}_{\mathrm{ess}}(\tau_{\mathrm{tax}})$ be the domain-specific core essential GO profile and $\mathcal{U}_{\mathrm{ess}}(\tau_{\mathrm{tax}})$ the essential categories with $\max_p P^{(0)}(p,e) \le 0.50$. For any term f in the downward subclass closure $\mathcal{D}_{\mathrm{uncov}}(\tau_{\mathrm{tax}}) = \bigcup_{e\in\mathcal{U}_{\mathrm{ess}}(\tau_{\mathrm{tax}})}\{t \in \mathcal{T} : t \sqsubseteq e\}$, candidate claims receive $B_1(p,f) = \alpha_{\mathrm{ess}}\,\mathbb{I}[f \in \mathcal{D}_{\mathrm{uncov}}(\tau_{\mathrm{tax}})]$.
2. *Pathway and heteromeric complex coherence prior (B_2, $\alpha_{coh} = 1.00$):* Missing steps of active GO `has_part` processes receive $w_{\mathrm{proc}}(f) = \max(0, 1 - c_{\mathrm{proc}})$, and missing steps of triggered reference pathways W with step coverage $c_W \in (0,1)$ receive $w_{\mathrm{path}}(f) = \max_{W\ni f}\max(0.25, 1-c_W)$. For an obligate heteromeric complex $f \in \mathcal{C}_{\mathrm{het}}$ with $N_{\mathrm{strong}}(f)$ proteins at $P^{(0)} > 0.50$ and $N_{\mathrm{cand}}(f)$ candidates at $\max(P^{(0)}, P^{(k-1)}) \ge 0.08$, candidate partners receive $w_{\mathrm{cplx}}(p,f) = 0.85$ when $N_{\mathrm{strong}} = 1 \wedge N_{\mathrm{cand}} \ge 2$ (and 0.5525 when $N_{\mathrm{strong}} \ge 2$), whereas isolated strong singletons ($N_{\mathrm{strong}} = 1 \wedge N_{\mathrm{cand}} \le 1$) receive -0.75. Therefore, $B_2(p,f) = \alpha_{\mathrm{coh}}\big(\max(w_{\mathrm{proc}}(f), w_{\mathrm{path}}(f)) + w_{\mathrm{cplx}}(p,f)\big)$.
3. *Taxonomic consistency prior (B_3, $\alpha_{cons} = 3.00$):* Active terms $\mathcal{A}^{(k-1)} = \{t \in \mathcal{T} : \max_p P^{(k-1)}(p,t) >$

0.50} are checked against direct GO taxon constraints and SAT4J unsatisfiable cores over multi-parent `is_a` and disjointness axioms. Any claim whose term or ancestor lies in the conflict set $\mathcal{V}_{\text{conf}}^{(k-1)}$ receives $B_3(p,f) = -\alpha_{\text{cons}}$.

4. *Anatomical prior (B_4, $\alpha_{anat} = 3.50$, $\alpha_{boost} = 0.65$):* Let $\mathcal{T}_{\text{forb}}^{\text{anat}}(\tau_{\text{tax}})$ denote GO terms forbidden in τ_{tax} through the GO–Uberon/PO/CL bridge (Section 4.4) and $\mathcal{U}_{\text{act}}(\tau_{\text{tax}})$ valid non-root anatomical structures supported by at least two proteins at $P^{(0)} \geq 0.50$. Claims intersecting $\mathcal{T}_{\text{forb}}^{\text{anat}}(\tau_{\text{tax}})$ receive $B_4(p,f) = -3.50$; valid claims linked to $\mathcal{U}_{\text{act}}(\tau_{\text{tax}})$ receive $+0.65$.
5. *Metabolic gap-filling and reaction-local genomic context prior (B_5, $\alpha_{gap} = 1.50$):* Candidate terms matching open metabolic network gaps receive $B_5(p,f) = \alpha_{\text{gap}}$ (1.05 for low-homology gapseq proposals). For each gap reaction R, reaction-local genomic context (RLGC) identifies pathway-neighboring GO terms at 1–2 reaction hops ($\alpha_{\text{rxn}} = 0.50$), weights chromosomal anchor genes p' with $P(p',g) \geq 0.30$ by $w_{\text{anc}}(p',R) = \max_g \alpha_{\text{rxn}}^{d(g)} P(p',g)$, and accumulates Gaussian kernel density $K_{\text{bp}}(\Delta\text{bp}) = \exp(-\Delta\text{bp}^2/(2h_{\text{bp}}^2))$ ($h_{\text{bp}} = 5{,}000$ bp):

$$D_{\text{RLGC}}(p,R) = \sum_{p':|\Delta\text{bp}(p,p')|\leq 4h_{\text{bp}}} w_{\text{anc}}(p',R)\, K_{\text{bp}}\big(\Delta\text{bp}(p,p')\big)$$

$$S_{\text{RLGC}}(p,R) = \log D_{\text{RLGC}}(p,R) + \beta_{\text{div}}\text{Div}(p) - \gamma_{\text{com}}\text{Comm}(p) - \delta_{\text{self}}\mathbb{I}[p \text{ is a gene}] + \varepsilon_{\text{str}}\mathbb{I}[\text{same strand}(p)] - \zeta_{\text{gap}}\log_{10}(\max(1,\Delta\text{bp}_{\min}(p))) \quad (3)$$

with neighbor-reaction diversity weight $\beta_{\text{div}} = 1.0$, commitment penalty $\gamma_{\text{com}} = 1.0$, self-annotation penalty $\delta_{\text{self}} = 2.0$, same-strand bonus $\varepsilon_{\text{str}} = 0.2$, and minimum-distance penalty $\zeta_{\text{gap}} = 0.1$. In LRO enzyme attribution, S_{RLGC} is combined with orthogroup co-occurrence, Pfam domain compatibility, and (in Track B) Swiss-Prot ESM-2 reaction-centroid cosine similarity.

6. *Co-directional operon prior (B_6, $\alpha_{ctx} = 1.00$, $\alpha_{gap_ctx} = 1.50$, $w_{op_cplx} = 0.95$):* Within a co-directional operon O ($|O| \geq 2$) having active pathway fraction $c_O(W) = k_O(W)/|O|$ for pathway W, a candidate (p,f) receives $B_6(p,f) = \max_{W \ni f : k_O(W) \geq 1}\big(\alpha_{\text{ctx}} c_O(W) \alpha_{\text{gap_ctx}}^{\mathbb{I}[f \in \mathcal{G}_{\text{gap}}]}\big) + w_{\text{op_cplx}}\mathbb{I}[f \in \mathcal{C}_{\text{het}} \wedge \exists p' \in O \setminus \{p\} : P^{(k-1)}(p',f) > 0.50]$.

4.4 Cross-ontology taxonomic and anatomical constraint bridge

To enforce anatomical and taxonomic consistency across domains, we extracted five programmatic constraint tables from the official OBO Foundry releases of GO, Uberon [9], Plant Ontology (PO) [44], Cell Ontology (CL) [45], and NCBI Taxonomy (Supplementary Section S3 and Supplementary Table S7): (1) 26,911 direct GO taxon constraints (13,819 `only_in_taxon` and 13,092 `never_in_taxon`) across 18,159 GO terms [8]; (2) 2,715 cross-ontology equivalence and subclass axioms linking 2,674 GO terms to 1,294 Uberon, PO, and CL classes across 30 relations (including `results_in_development_of`, `results_in_morphogenesis_of`, `results_in_formation_of`, `results_in_maturation_of`, and `occurs_in`); (3) 8,331 taxon constraints (4,767 `only_in_taxon` and 3,564 `never_in_taxon`) over 1,247 Uberon, PO, and CL anatomical classes; (4) 7,722 transitive `is_a` and `part_of` edges over 2,855 anatomical classes; and (5) 1,818 taxonomic lineage edges over 1,305 NCBI Taxonomy clades. A GO term t is entailed as anatomically forbidden in taxon τ_{tax} whenever t (or a GO ancestor of t) links to an anatomical class u whose anatomical ancestor closure carries an `only_in_taxon` or `never_in_taxon` rule incompatible with the taxonomic lineage of τ_{tax}.

4.5 Stage 2A: deterministic symbolic enforcement and semantic ancestor relaxation

After Stage 1 convergence, Stage 2A deterministically enforces all direct GO taxon constraints, cross-ontology anatomical constraints, and SAT4J logical consistency axioms. For each protein p and forbidden term t, Stage 2A traverses $\text{Anc}(t)$ (excluding the three GO aspect roots) to identify the minimal valid ancestor set $\mathcal{A}_{\text{valid}}(t,\tau_{\text{tax}})$ comprising terms both taxonomically and anatomically admissible in τ_{tax}. Reconstructing the unpenalized pre-conflict probability $P_{\text{unpen}}(p,t)$ from $L^{(0)}(p,t)$ and positive Stage 1 priors, semantic ancestor relaxation transfers this support to each minimal valid ancestor and zeroes the forbidden term:

$$P_{\text{Stage2A}}(p,a) = \max\Big(P^{(K)}(p,a),\, \gamma_{\text{relax}}\, P_{\text{unpen}}(p,t)\Big), \qquad \forall a \in \mathcal{A}_{\text{valid}}$$
$$P_{\text{Stage2A}}(p,t) = 0, \quad (4)$$

with $\gamma_{\text{relax}} = 1.0$. In addition, for any obligate heteromeric complex $f \in \mathcal{C}_{\text{het}}$ with a single protein at $P \geq 0.50$ and zero partner candidates at $P \geq 0.08$ across the genome, Stage 2A attenuates the isolated singleton to 0.45.

4.6 Stage 2B: role-aware reaction support and positive-cost lexicographic pFBA

Stage 2B operates over a sanitized universal metabolic network $\mathcal{U}_{\text{clean}}$ derived from ModelSEED [38] (47,880 reactions and 33,978 metabolites, of which 14,243 carry EC mappings and 11,826 are mass- and charge-balanced enzymatic reactions; 2,417 pool or elementally imbalanced reactions receive penalty $w_{\text{imb}}(j) = 100.0$). Each reaction j is associated with a disjunctive normal form (DNF) catalytic rule $\mathcal{M}_{\text{dnf}}(j) = \{m_1, \ldots, m_{K_j}\}$ over alternative isozyme mechanisms m, where each mechanism requires a conjunction of subunit roles $\mathcal{R}_{\text{sub}}(m) = \{u_1, \ldots, u_{D_m}\}$. Given candidate locus probabilities $s_{p,u} \in (0,1)$, Stage 2B aggregates the top $M_{\text{iso}} = 3$ candidates per role via Noisy-OR, combines required subunits via a bottleneck-gated

geometric mean at $\theta_{\text{sub}} = 0.25$, and takes the maximum across isozyme mechanisms:

$$Q(u) = 1 - \prod_{p \in \text{Top-}M_{\text{iso}}(\mathcal{C}_{\text{cand}}(u))} (1 - s_{p,u}),$$

$$S_{\text{mech}}(m) = \begin{cases} \Big(\prod_{u \in \mathcal{R}_{\text{sub}}(m)} Q(u) \Big)^{1/D_m}, & \min_u Q(u) \geq \theta_{\text{sub}}, \\ \min_{u \in \mathcal{R}_{\text{sub}}(m)} Q(u), & \text{otherwise}, \end{cases} \tag{5}$$

Reactions with $S_{\text{rxn}}(j) \geq \tau_{\text{core}} = 0.50$, together with open medium exchanges and spontaneous reactions, form the draft core $\mathcal{R}_{\text{core}}$. Candidate repair reactions $j \in \mathcal{D}_{\text{dom}} \setminus \mathcal{R}_{\text{core}}$ receive a strictly positive addition cost $c_j^{\text{add}} = \lambda_{\text{add}}(1 - S_{\text{rxn}}(j)) + w_{\text{imb}}(j) + \epsilon_{\text{par}}$ ($\lambda_{\text{add}} = 1.0$, $\epsilon_{\text{par}} = 10^{-3}$). Splitting reversible fluxes into non-negative components $v_j = v_j^+ - v_j^-$ and setting the biomass target on medium $\mathcal{E}_{\text{med}}$ to $v_{\text{bio}}^{\text{target}}(\mathcal{E}_{\text{med}}) = \max(\alpha_{\text{bio}} v_{\text{bio}}^{\max}(\mathcal{E}_{\text{med}}), 0.1)$ ($\alpha_{\text{bio}} = 0.10$), Stage 2B solves a two-step lexicographic pFBA [48] using the HiGHS linear programming solver (feasibility tolerance 10^{-9}). Step 1 minimizes the repair cost $\sum_{j \notin \mathcal{R}_{\text{core}}} c_j^{\text{add}}(v_j^+ + v_j^-)$ subject to $\mathbf{Sv} = \mathbf{0}$ and $v_{\text{bio}} \geq v_{\text{bio}}^{\text{target}}(\mathcal{E}_{\text{med}})$ (solving first over balanced reactions and admitting penalized imbalanced reactions only if infeasible) to obtain optimal repair cost $C^*(\mathcal{E}_{\text{med}})$. Step 2 minimizes total non-biomass flux $\sum_{j \neq \text{bio}}(v_j^+ + v_j^-)$ subject to $\sum_{j \notin \mathcal{R}_{\text{core}}} c_j^{\text{add}}(v_j^+ + v_j^-) \leq (1 + 10^{-6})C^*(\mathcal{E}_{\text{med}}) + 10^{-9}$. The active repair set $\mathcal{J}_{\text{rep}}(\mathcal{E}_{\text{med}}) = \{j \notin \mathcal{R}_{\text{core}} : |v_j^{\text{pFBA}}(\mathcal{E}_{\text{med}})| > 10^{-6}\}$ is evaluated by flux-variability ($1.01C^*$), reaction-knockout ($1.10C^*$), and five-replicate cost-perturbation ($\pm 5\%$) certificates to yield the flux confidence $\pi_j^{\text{FBA}} = 0.5\mathbb{I}[j \in \mathcal{R}_{\text{FVA}}] + 0.3\mathbb{I}[j \in \mathcal{R}_{\text{KO}}] + 0.2\omega_{\text{ens}}(j) \in [0, 1]$ (Supplementary Section S4 and Supplementary Table S12).

4.7 Stage 2B: competitive gene-locus responsibility and substrate probing

For each repaired reaction $j \in \mathcal{J}_{\text{rep}}$ and required role $u \in \mathcal{R}_{\text{sub}}(m_j^*)$, Stage 2B distributes locus responsibility competitively across candidate genes $p \in \mathcal{C}_{\text{cand}}(u)$ and an explicit unassigned-gene hypothesis $h_\varnothing$ with prior mass $\eta_{\text{null}} = 0.05$:

$$w_{j,u}(p) = \big(o_{p,u}\big)^{\beta_1} \text{BF}_{\text{operon}}(p, u)^{\beta_2} \exp\big(\beta_3 C_{\text{syn}}(p, u) + \beta_4 S_{\text{ESM2}}(p, u)\big),$$

$$r_{j,u}(p) = \frac{w_{j,u}(p)}{\eta_{\text{null}} + \sum_{p' \in \mathcal{C}_{\text{cand}}(u)} w_{j,u}(p')}, \tag{6}$$

where $o_{p,u} = s_{p,u}/(1 - s_{p,u})$ is the sequence odds, $\text{BF}_{\text{operon}}(p, u)$ is the RLGC operon Bayes factor, $C_{\text{syn}}(p, u)$ is the operon and pathway synteny score, $S_{\text{ESM2}}(p, u)$ is the language-model similarity score, and $\beta_1 = \beta_2 = 1.0, \beta_3 = \beta_4 = 1.5$. The backward log-odds update to locus p is $\Delta L_{\text{FBA}}(p, u) = \gamma_{\text{FBA}} \sum_{j \in \mathcal{J}_{\text{rep}}(u)} \pi_j^{\text{FBA}} r_{j,u}(p)(1 - S_{\text{rxn}}(j))$ with $\gamma_{\text{FBA}} = 2.20$, applied across damped outer iterations $\lambda_t = \lambda_0 \gamma_{\text{damp}}^{t-1}$ ($\lambda_0 = 0.70, \gamma_{\text{damp}} = 0.60$). For catabolic enzymes in Price-149, Stage 2B supplements minimal-glucose pFBA with cognate-substrate stoichiometric neighborhood probing (rewarding candidate reactions that share non-currency metabolites at 1–2 reaction hops with chromosomal neighbors within ±10 genes) and blends single-predictor and five-predictor ensemble posteriors (complete equations in Supplementary Section S4). Held-out metabolic reactions are classified by five binary recoverability indicators: structural presence and mass balance in $\mathcal{U}_{\text{clean}}$ ($\mathcal{C}_{\text{struct}}$), medium-specific flux demand ($\mathcal{C}_{\text{task}}$, partitioning core biomass Stratum A, cognate-substrate catabolic Stratum B, and secondary Stratum C), isozyme uniqueness ($\mathcal{C}_{\text{unique}}$), operon support ($\mathcal{C}_{\text{operon}}$), and multi-subunit partner presence ($\mathcal{C}_{\text{complex}}$).

$$S_{\text{rxn}}(j) = \max_{m \in \mathcal{M}_{\text{dnf}}(j)} S_{\text{mech}}(m)$$

4.8 Stage 3: selective neuro-symbolic annotation completion

Stage 3 is triggered selectively on four post-Stage-2 anomaly categories: (1) pathway gaps (pathway completeness ≥ 0.50 with candidate posteriors in $[0.05, 0.50)$; ≥ 0.65 and $[0.15, 0.49]$ in the controlled 120-anomaly benchmark); (2) incomplete obligate heteromeric complexes (one active subunit at $P \geq 0.50$ and sub-threshold partner candidates); (3) borderline uncovered essential categories; and (4) borderline taxonomic or anatomical conflicts ($P \in [0.35, 0.62]$) amenable to ancestral GO relaxation. Stage 3 evaluates at most 8 candidates per anomaly and 150 anomalies per genome.

Each anomaly is stripped of evaluation labels and gene symbols and processed in a bounded multi-turn tool-calling loop (maximum 6 turns; 2 turns in the fast mode) by a locally hosted FP8-quantized 27-billion-parameter instruction model from the Qwen open-weight family [63] served via vLLM (32,768-token context, temperature 0.0, maximum 450 output tokens per turn, evaluated across seeds 42, 43, and 44). The model invokes five deterministic read-only tools to inspect the operon neighborhood, inspect the pathway or complex gap, query protein evidence and domain signatures, check symbolic ontology constraints, and propose a structured resolution (PROMOTE, RELAX, DEMOTE, or ABSTAIN). Every proposal is verified by the deterministic symbolic guardrail against the 48,101 GO term stanzas, 1,305 NCBI Taxonomy clades, 26,911 direct GO taxon rules, and 2,715 GO–Uberon/PO/CL axioms before acceptance (tool schemas, de-identification protocol, and per-seed results in Supplementary Section S6 and Supplementary Table S14).

4.9 Experimental phenotype prediction in BacDive

We evaluated organism-level phenotype prediction across 2,332 carbon-source utilization assays in 529 ProTraits-matched bacterial strains, 17,496 diagnostic enzyme assays in 3,017 BacDive type strains, and 50,075 assays for 10 diagnostic enzymes across 6,313 strains [55, 56]. Given

binary pathway or reaction calls $g(s,y), m(s,y), c(s,y) \in \{0,1\}$ from gapseq v1.2 [40], ModelSEED v2.6.1 [38], and CarveMe v1.5.2 [39] for strain s and assay y, we form the blended matrix $B(s,y)$ ($0.55g + 0.30m + 0.15c$ for carbon; $0.55g + 0.28m + 0.17c$ for enzymes), column-center each assay $\tilde{B}_{s,y} = B(s,y) - \bar{B}_y$, and compute the row-normalized non-negative cosine coupling matrix $\Psi(y,y')$ with zero diagonal to obtain the unsupervised cross-assay coupling score $\psi(s,y) = \sum_{y'} \Psi(y,y')(B(s,y') - 0.5)$. In blind Track 1 (0% phenotype labels), binary tool calls are mapped to calibrated probabilities $p_{\text{gap}}, p_{\text{ms}}, p_{\text{cm}}$ and pooled in log-odds space:

$$P_{\text{Track1}}(s,y) = \sigma_{\log}\Big(w_{\text{gap}} \operatorname{logit}\big(p_{\text{gap}}(s,y)\big) + w_{\text{ms}} \operatorname{logit}\big(p_{\text{ms}}(s,y)\big) + w_{\text{cm}} \operatorname{logit}\big(p_{\text{cm}}(s,y)\big) + w_{\psi}\,\psi(s,y)\Big), \tag{7}$$

with weights $(0.56, 0.28, 0.16, 0.65)$ for carbon and $(0.55, 0.27, 0.18, 0.55)$ for enzymes at threshold 0.50. Supervised baselines (Traitar L1-SVM [64], BacDive Random Forest [56], and DiASPora GBDT [57]) and the GSPA + DiASPora hybrid were evaluated under five-fold genus-blocked cross-validation with all calibration, prevalence, and coupling parameters estimated strictly within training folds. In Track 2, 30% of each strain's assays are revealed to condition the Bayesian anchor $P_{\text{Track2}}(s,y)$ on revealed substrate co-occurrence or genus-level EC posteriors, and predictions are evaluated on the held-out 70% (complete parameterization in Supplementary Section S5 and Supplementary Table S13).

4.10 Benchmarks, donor-homology masking, metrics, and structural audit score

To evaluate generalization beyond sequence retrieval, donor-homology masking removes reference proteins sharing pairwise sequence identity $\text{pident}(g,r) > \eta_{\text{id}}$ ($\eta_{\text{id}} \in \{100\%, 70\%, 50\%, 30\%\}$) or belonging to the test organism's genus or family across DIAMOND databases, ESM-2 and ProtT5 neighbor indices, FoldSeek structural templates, and calibration sets. Neural enzyme predictors use their published homology-split lookup protocols with frozen released backbone weights. In leave-reaction-out benchmarks, Track A masks sequence alignments and embedding donors with cosine similarity > 0.50 without reaction centroids, while Track B masks sequence alignments and retains Swiss-Prot ESM-2 reaction centroids.

We report protein-centric CAFA $F_{\max}$, information-accretion-weighted $WF_{\max}$, pair-level micro $F_{\max}$, and minimum semantic distance $S_{\min}$ [1, 27, 65] over threshold grids $\theta \in \{0.01, \ldots, 0.99\}$ ($\{0.10, \ldots, 0.90\}$ in the eukaryotic stress test), alongside Hit@k and mean reciprocal rank (MRR) for candidate ranking, 95% bootstrap or exact Clopper–Pearson confidence intervals, and two-sided paired Wilcoxon signed-rank or exact binomial McNemar tests (Supplementary Section S5). All CPU benchmarks ran on dual-socket AMD EPYC 7763 nodes (8 threads per genome), with PLM and Stage 3 LLM inference executed on an NVIDIA A100-SXM4-80GB GPU.

The Genome-Scale Annotation Evaluation Framework (GAEF) summarizes structural proteome invariants at $\theta_{\text{eval}} = 0.50$ by combining essential function completeness C_{ess} ($w_{\text{ess}} = 0.30$), functional pathway/complex/process coherence C_{func} ($w_{\text{func}} = 0.40$), taxonomic and anatomical consistency $C_{\text{cons}} = \max(0, 1 - 0.01V(\mathcal{P}))$ ($w_{\text{cons}} = 0.30$), and functional resolution $C_{\text{res}} = \min(1, \text{meanIC}/10)$ ($w_{\text{res}} = 0.0$):

$$\text{GAEF}(\mathcal{P}) = \frac{\sum_{d \in \{\text{ess,func,cons,res}\}: C_d \neq \text{null}} w_d\, C_d}{\sum_{d: C_d \neq \text{null}} w_d}, \tag{8}$$

which equals $0.30C_{\text{ess}} + 0.40C_{\text{func}} + 0.30C_{\text{cons}}$ when all three structural dimensions are active. Because GAEF measures the structural invariants targeted by Stages 1 and 2, we report it as an internal structural audit score alongside independent held-out functional and experimental phenotype evaluations.

Data availability

All precomputed cross-ontology constraint tables and hierarchies (Supplementary Table S7), obligate heteromeric complex classifications, benchmark manifests with cryptographic checksums, frozen evaluation tables underlying Figures 1–6 and Tables 1–7, and a complete provenance map linking every manuscript value to its underlying result table are available under CC-BY-4.0 at https://github.com/bio-ontology-research-group/gspa (release v2.1.0).

Code availability

The GSPA v2.1.0 software suite, including the command-line integration and evaluation engine, the Stage 2B stoichiometric locus refiner, the Stage 3 neuro-symbolic agent and ontology guardrail, and all benchmark and figure generation scripts, is available under the GPL-3.0-or-later license at https://github.com/bio-ontology-research-group/gspa (release v2.1.0).

Acknowledgments

This work was supported by King Abdullah University of Science and Technology (KAUST) baseline funding to R.H.

Author contributions

T.A.-S., M.K., and R.H. conceived the study and designed the methodology. T.A.-S. and R.H. implemented the software, cross-ontology bridges, stoichiometric refiner, and benchmark suites. All authors analyzed the results and wrote the manuscript.

Competing interests

The authors declare no competing interests.

References

[1] Predrag Radivojac, Wyatt T. Clark, Tal Ronnen Oron, Alexandra M. Schnoes, Tobias Wittkop, Artem Sokolov, Kiley Graim, Christopher Funk, Karin Verspoor, Asa Ben-Hur, Gaurav Pandey, Jeffrey M. Yunes, Ameet S. Talwalkar, Sanna Repo, Michael L. Souza, Damiano Piovesan, Rong Zhong, Haixuan Cui, Nan Zhou, Melanie Kirac, Slobodan Vucetic, Aakrosh Ratan, Burkhard Rost, Julian Gough, Matthew Rogers, Franck Rapaport, Janet M. Thornton, Maria J. Martin, Amos Bairoch, Sean D. Mooney, Constance J. Jeffery, Iddo Friedberg, and Steven E. Brenner. A large-scale evaluation of computational protein function prediction. *Nature Methods*, 10 (3):221–227, 2013. doi: 10.1038/nmeth.2340. URL https://doi.org/10.1038/nmeth.2340.

[2] Nan Zhou, Yuxiang Jiang, Timothy R. Bergquist, Angela J. Lee, Balint Z. Kacsoh, Alex W. Crocker, Kaitlyn A. Lewis, George Georghiou, Huy N. Nguyen, Predrag Radivojac, and Iddo Friedberg. The CAFA challenge reports improved protein function prediction and new functional annotations for hundreds of genes through experimental screens. *Genome Biology*, 20 (1):244, 2019. doi: 10.1186/s13059-019-1835-8. URL https://doi.org/10.1186/s13059-019-1835-8.

[3] Paul D. Thomas, David P. Hill, Huaiyu Mi, David Osumi-Sutherland, Kimberly Van Auken, Seth Carbon, James P. Balhoff, Laurent-Philippe Albou, Benjamin Good, Pascale Gaudet, Suzanna E. Lewis, and Christopher J. Mungall. Gene Ontology Causal Activity Modeling (GO-CAM) moves beyond GO annotations to structured descriptions of biological functions and systems. *Nature Genetics*, 51(10):1429–1433, 2019. doi: 10.1038/s41588-019-0500-1. URL https://doi.org/10.1038/s41588-019-0500-1.

[4] Andrei Osterman and Ross Overbeek. Missing genes in metabolic pathways: a comparative genomics approach. *Current Opinion in Chemical Biology*, 7(2):238–251, 2003. doi: 10.1016/S1367-5931(03)00027-9. URL https://doi.org/10.1016/S1367-5931(03)00027-9.

[5] Peter D. Karp, Daniel Weaver, and Mario Latendresse. How accurate is automated gap filling of metabolic models? *BMC Systems Biology*, 12(1):59, 2018. doi: 10.1186/s12918-018-0593-7. URL https://doi.org/10.1186/s12918-018-0593-7.

[6] Madalina Giurgiu, Julian Reinhard, Barbara Brauner, Irmtraud Dunger-Kaltenbach, Gisela Fobo, Goar Frishman, Corinna Montrone, and Andreas Ruepp. Corum: the comprehensive resource of mammalian protein complexes – 2019. *Nucleic Acids Research*, 47 (D1):D559–D563, October 2018. ISSN 1362-4962. doi: 10.1093/nar/gky973. URL http://dx.doi.org/10.1093/nar/gky973.

[7] Birgit H M Meldal, Livia Perfetto, Colin Combe, Tiago Lubiana, João Vitor Ferreira Cavalcante, Hema Bye-A-Jee, Andra Waagmeester, Noemi del Toro, Anjali Shrivastava, Elisabeth Barrera, Edith Wong, Bernhard Mlecnik, Gabriela Bindea, Kalpana Panneerselvam, Egon Willighagen, Juri Rappsilber, Pablo Porras, Henning Hermjakob, and Sandra Orchard. Complex portal 2022: new curation frontiers. *Nucleic Acids Research*, 50(D1):D578–D586, October 2021. ISSN 1362-4962. doi: 10.1093/nar/gkab991. URL http://dx.doi.org/10.1093/nar/gkab991.

[8] Jennifer I. Deegan née Clark, Emily C. Dimmer, and Christopher J. Mungall. Formalization of taxon-based constraints to detect inconsistencies in annotation and ontology development. *BMC Bioinformatics*, 11: 530, 2010. doi: 10.1186/1471-2105-11-530. URL https://doi.org/10.1186/1471-2105-11-530.

[9] Christopher J. Mungall, Carlo Torniai, George V. Gkoutos, Suzanna E. Lewis, and Melissa A. Haendel. Uberon, an integrative multi-species anatomy ontology. *Genome Biology*, 13(1):R5, 2012. doi: 10.1186/gb-2012-13-1-r5. URL https://doi.org/10.1186/gb-2012-13-1-r5.

[10] Shanika L. Amarasinghe, Shian Su, Xueyi Dong, Luke Zappia, Matthew E. Ritchie, and Quentin Gouil. Opportunities and challenges in long-read sequencing data analysis. *Genome Biology*, 21(1):30, 2020. doi: 10.1186/s13059-020-1935-5.

[11] Glennis A. Logsdon, Mitchell R. Vollger, and Evan E. Eichler. Long-read human genome sequencing and its applications. *Nature Reviews Genetics*, 21(10): 597–614, 2020. doi: 10.1038/s41576-020-0236-x.

[12] Mikhail Kolmogorov, Derek M. Bickhart, Bahar Behsaz, Alexey Gurevich, Mikhail Rayko, Sung Bong Shin, Kristen Kuhn, Jeffrey Yuan, Evgeny Polevikov, Timothy P. L. Smith, and Pavel A. Pevzner. metaFlye: scalable long-read metagenome assembly using repeat graphs. *Nature Methods*, 17(11):1103–1110, 2020. doi: 10.1038/s41592-020-00971-x.

[13] Benjamin Buchfink, Klaus Reuter, and Hajk-Georg Drost. Sensitive protein alignments at tree-of-life scale using diamond. *Nature Methods*, 18(4):366–368, 2021. doi: 10.1038/s41592-021-01101-x.

[14] Wenjun Li, Kathleen R O'Neill, Daniel H Haft, Michael DiCuccio, Vyacheslav Chetvernin, Azat Badretdin, George Coulouris, Farideh Chitsaz, Myra K Derbyshire, A Scott Durkin, Noreen R Gonzales, Marc Gwadz, Christopher J Lanczycki, James S Song, Narmada Thanki, Jiyao Wang, Roxanne A Yamashita, Mingzhang Yang, Chanjuan Zheng, Aron Marchler-Bauer, and Françoise Thibaud-Nissen. Refseq: expanding the prokaryotic genome annotation pipeline reach with protein family model curation. *Nu-*

cleic Acids Research, 49(D1):D1020–D1028, December 2020. ISSN 1362-4962. doi: 10.1093/nar/gkaa1105. URL http://dx.doi.org/10.1093/nar/gkaa1105.

[15] Oliver Schwengers, Lukas Jelonek, Marius Alfred Dieckmann, Sebastian Beyvers, Jochen Blom, and Alexander Goesmann. Bakta: rapid and standardized annotation of bacterial genomes via alignment-free sequence identification: Find out more about bakta, the motivation, challenges and applications, here. *Microbial Genomics*, 7(11), November 2021. ISSN 2057-5858. doi: 10.1099/mgen.0.000685. URL http://dx.doi.org/10.1099/mgen.0.000685.

[16] Sean R. Eddy. Accelerated profile HMM searches. *PLoS Computational Biology*, 7(10):e1002195, 2011. doi: 10.1371/journal.pcbi.1002195.

[17] Philip Jones, David Binns, Hsin-Yu Chang, Matthew Fraser, Weizhong Li, Craig McAnulla, Hamish McWilliam, John Maslen, Alex Mitchell, Gift Nuka, Sebastien Pesseat, Antony F. Quinn, Amaia Sangrador-Vegas, Maxim Scheremetjew, Siew-Yit Yong, Rodrigo Lopez, and Sarah Hunter. Interproscan 5: genome-scale protein function classification. *Bioinformatics*, 30(9):1236–1240, January 2014. ISSN 1367-4803. doi: 10.1093/bioinformatics/btu031. URL http://dx.doi.org/10.1093/bioinformatics/btu031.

[18] Zeming Lin, Halil Akin, Roshan Rao, Brian Hie, Zhongkai Zhu, Wenting Lu, Nikita Smetanin, Robert Verkuil, Ori Kabeli, Yaniv Shmueli, et al. Evolutionary-scale prediction of atomic-level protein structure with a language model. *Science*, 379(6637): 1123–1130, 2023. doi: 10.1126/science.ade2574.

[19] Ahmed Elnaggar, Michael Heinzinger, Christian Dallago, Ghalia Rehawi, Yu Wang, Llion Jones, Tom Gibbs, Tamas Feher, Christoph Angerer, Martin Steinegger, Debsindhu Bhowmik, and Burkhard Rost. ProtTrans: Toward understanding the language of life through self-supervised learning. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 44(10): 7112–7127, 2022. doi: 10.1109/TPAMI.2021.3095381.

[20] Maxat Kulmanov, Francisco J. Guzmán-Vega, Paula Duek Roggli, Lydie Lane, Stefan T. Arold, and Robert Hoehndorf. Protein function prediction as approximate semantic entailment. *Nature Machine Intelligence*, 6(2):220–228, 2024. doi: 10.1038/s42256-024-00795-w. URL https://doi.org/10.1038/s42256-024-00795-w.

[21] Tianhao Yu, Haiyang Cui, Jianan Li, Yang Yang, Hui Xiong, Zixuan Cang, and Huimin Zhao. Enzyme function prediction using contrastive learning. *Science*, 379(6639):1358–1363, 2023. doi: 10.1126/science.adf2470. URL https://doi.org/10.1126/science.adf2470.

[22] Vladimir Gligorijević, P. Douglas Renfrew, Tomasz Kosciolek, Julia Koehler Leman, Daniel Berenberg, Tommi Vatanen, Chris Chandler, Ian Mavric, Thomas Hill, Kyunghyun Cho, and Richard Bonneau. Structure-based protein function prediction using graph convolutional networks. *Nature Communications*, 12:3168, 2021. doi: 10.1038/s41467-021-23303-9. URL https://doi.org/10.1038/s41467-021-23303-9.

[23] Michel van Kempen, Stephanie S. Kim, Charlotte Tumescheit, Milot Mirdita, Jeongjae Lee, Cameron L. M. Gilchrist, Johannes Söding, and Martin Steinegger. Fast and accurate protein structure search with Foldseek. *Nature Biotechnology*, 42(2):243–246, 2024. doi: 10.1038/s41587-023-01773-0.

[24] Ronghui You, Zihan Zhang, Yi Xiong, Fengzhu Sun, Hiroshi Mamitsuka, and Shanfeng Zhu. GOLabeler: improving sequence-based large-scale protein function prediction by learning to rank. *Bioinformatics*, 34(14): 2465–2473, 2018. doi: 10.1093/bioinformatics/bty130.

[25] Ronghui You, Shuwei Yao, Yi Xiong, Zhi-Ping Huang, Fengzhu Sun, Hiroshi Mamitsuka, and Shanfeng Zhu. NetGO: improving large-scale protein function prediction with graph-preserving learning and network integration. *Nucleic Acids Research*, 47(W1): W397–W402, 2019. doi: 10.1093/nar/gkz388. URL https://doi.org/10.1093/nar/gkz388.

[26] Shaojun Wang, Ronghui You, Yunjia Liu, Yi Xiong, and Shanfeng Zhu. NetGO 3.0: Protein language model improves large-scale functional annotations. *Genomics, Proteomics & Bioinformatics*, 21(2):349–358, 2023. doi: 10.1016/j.gpb.2023.04.001. URL https://doi.org/10.1016/j.gpb.2023.04.001.

[27] Yuxiang Jiang, Tal Ronnen Oron, Wyatt T. Clark, Atreya R. Bankapur, Daniel D'Andrea, Rosalba Lepore, Christopher S. Funk, Indika Kahanda, Karin M. Verspoor, Asa Ben-Hur, David Chi-Eng Koo, Dan Penfold-Brown, Dennis Shasha, Noah M. Young, Richard Bonneau, Björn Wallner, Nereo Atropo, Florence Martin, Antonio Puertas-Gallardo, Jose M. Carazo, Pedro Larrañaga, Concha Bielza, Luigi Cerulo, Alexandre Chigaev, Bruce S. Edwards, Larry A. Sklar, Emmanuel Charbonnier, François Enault, Angela Del Pozo, José M. Fernández, Michael L. Tress, Amar Bhaskar, Arne Elofsson, Shaojun Wang, Shanfeng Zhu, Ronghui You, Zhi-Ping Huang, Shanshan Dai, Shuwei Yao, Weidong Tian, S. Radha, Aakrosh Ratan, Burkhard Rost, Julian Gough, Predrag Radivojac, and Iddo Friedberg. An expanded evaluation of protein function prediction methods shows an improvement in accuracy. *Genome Biology*, 17(1):184, 2016. doi: 10.1186/s13059-016-1037-6. URL https://doi.org/10.1186/s13059-016-1037-6.

[28] Thomas Dandekar, Berend Snel, Martijn Huynen, and Peer Bork. Conservation of gene order: a fingerprint of proteins that physically interact. *Trends in Genetics*, 14(9):334–337, 1998. doi: 10.1016/S0168-9525(98)01558-3. URL https://doi.org/10.1016/S0168-9525(98)01558-3.

[29] Ross Overbeek, Michael Fonstein, Mark D'Souza, Gordon D. Pusch, and Natalia Maltsev. The use of gene clusters to infer functional coupling. *Proceedings of the National Academy of Sciences of the United States of America*, 96(6):2896–2901, 1999. doi: 10.1073/pnas.96.6.2896. URL https://doi.org/10.1073/pnas.96.6.2896.

[30] Matteo Pellegrini, Edward M. Marcotte, Michael J. Thompson, David Eisenberg, and Todd O. Yeates. Assigning protein functions by comparative genome analysis: Protein phylogenetic profiles. *Proceedings of the National Academy of Sciences of the United States of America*, 96(8):4285–4288, 1999. doi: 10.1073/pnas.96.8.4285. URL https://doi.org/10.1073/pnas.96.8.4285.

[31] Morgan N. Price, Katherine H. Huang, Eric J. Alm, and Adam P. Arkin. A novel method for accurate operon predictions in all sequenced prokaryotes. *Nucleic Acids Research*, 33(3):880–892, 2005. doi: 10.1093/nar/gki232. URL https://doi.org/10.1093/nar/gki232.

[32] Damian Szklarczyk, Rebecca Kirsch, Mikaela Koutrouli, Katerina Nastou, Falaah Mehryary, Radja Hachilif, Annika Gable, Tao Fang, Nadezhda T. Doncheva, Sampo Pyysalo, Peer Bork, Lars Juhl Jensen, and Christian von Mering. The STRING database in 2023: protein–protein association networks with increased coverage of experimental and predicted interactions. *Nucleic Acids Research*, 51 (D1):D638–D646, 2023. doi: 10.1093/nar/gkac1000. URL https://doi.org/10.1093/nar/gkac1000.

[33] Yunha Hwang, Andre L. Cornman, Elizabeth H. Kellogg, Sergey Ovchinnikov, and Peter R. Girguis. Genomic language model predicts protein co-regulation and function. *Nature Communications*, 15: 2888, 2024. doi: 10.1038/s41467-024-46947-9. URL https://doi.org/10.1038/s41467-024-46947-9.

[34] Andre Cornman, Jacob West-Roberts, Antonio Pedro Camargo, Simon Roux, Martin Beracochea, Milot Mirdita, Sergey Ovchinnikov, and Yunha Hwang. The OMG dataset: An open metagenomic corpus for mixed-modality genomic language modeling. *bioRxiv*, 2024. doi: 10.1101/2024.08.14.607850. URL https://doi.org/10.1101/2024.08.14.607850.

[35] Daulet Toibazar, Maxat Kulmanov, and Robert Hoehndorf. Context-based protein function prediction in bacterial genomes. *bioRxiv*, 2024. doi: 10.1101/2024.10.14.618363. URL https://doi.org/10.1101/2024.10.14.618363.

[36] Maciej Wiatrak, Ramon Viñas Torné, Maria Ntemourtsidou, Adam Dinan, David C. Abelson, Divya Arora, Maria Brbić, Aaron Weimann, and R. Andres Floto. A contextualised protein language model reveals the functional syntax of bacterial evolution. *bioRxiv*, 2025. doi: 10.1101/2025.07.20.665723. URL https://doi.org/10.1101/2025.07.20.665723.

[37] Michelle L. Green and Peter D. Karp. A Bayesian method for identifying missing enzymes in predicted metabolic pathway databases. *BMC Bioinformatics*, 5:76, 2004. doi: 10.1186/1471-2105-5-76. URL https://doi.org/10.1186/1471-2105-5-76.

[38] Samuel M. D. Seaver, Filipe Liu, Qizhi Zhang, Jose P. Roitberg, José P. Faria, and Christopher S. Henry. The ModelSEED biochemistry database for the integration of metabolic annotations and the reconstruction, comparison and analysis of metabolic models for plants, fungi and microbes. *Nucleic Acids Research*, 49(D1):D575–D588, 2021. doi: 10.1093/nar/gkaa746. URL https://doi.org/10.1093/nar/gkaa746.

[39] Daniel Machado, Sergej Andrejev, Melanie Tramontano, and Kiran Raosaheb Patil. Fast automated reconstruction of genome-scale metabolic models for microbial species and communities. *Nucleic Acids Research*, 46(15):7542–7553, 2018. doi: 10.1093/nar/gky537. URL https://doi.org/10.1093/nar/gky537.

[40] Johannes Zimmermann, Christoph Kaleta, and Silvio Waschina. gapseq: informed prediction of bacterial metabolic pathways and reconstruction of accurate metabolic models. *Genome Biology*, 22(1):81, 2021. doi: 10.1186/s13059-021-02295-1. URL https://doi.org/10.1186/s13059-021-02295-1.

[41] Morgan N. Price, Adam M. Deutschbauer, and Adam P. Arkin. GapMind: Automated annotation of amino acid biosynthesis. *mSystems*, 5(3):e00291–20, 2020. doi: 10.1128/mSystems.00291-20.

[42] Kexin Niu, Maxat Kulmanov, and Robert Hoehndorf. METEOR: joint genome-scale reconstruction and enzyme prediction. In *Pacific Symposium on Biocomputing 2027 (Session: Biological Molecular Function: Beyond Homology; preprint: bioRxiv 2026.02.01.701942)*, volume 32. World Scientific, 2027. doi: 10.64898/2026.02.01.701942.

[43] Christopher J. Mungall, Michael Bada, Tanya Z. Berardini, Jennifer Deegan, Amelia Ireland, Midori A. Harris, David P. Hill, and Jane Lomax. Cross-product extensions of the Gene Ontology. *Journal of Biomedical Informatics*, 44(1):80–86, 2011. doi: 10.1016/j.jbi.2010.02.002. URL https://doi.org/10.1016/j.jbi.2010.02.002.

[44] Laurel Cooper, Ramona L. Walls, Justin Elser, Maria A. Gandolfo, Dennis W. Stevenson, Barry Smith, Justin Preece, and Pankaj Jaiswal. The Plant Ontology as a tool for comparative plant anatomy and development. *International Journal of Plant Sciences*, 174(1):41–53, 2013. doi: 10.1086/668614.

[45] Alexander D. Diehl, Terrence F. Meehan, Yvonne M. Bradford, Matthew H. Brush, Wasila M. Dahdul, David S. Dougall, Yongqun He, David Osumi-Sutherland, Alan Ruttenberg, Sirarat Sarntivijai, Ceri E. Van Slyke, Nicole A. Vasilevsky, Melissa A. Haendel, Judith A. Blake, and Christopher J. Mungall.

The Cell Ontology 2016: enhanced content, modularization, and ontology interoperability. *Journal of Biomedical Semantics*, 7:44, 2016. doi: 10.1186/s13326-016-0088-0.

[46] Zhizheng Wang, Qiao Jin, Chih-Hsuan Wei, Shubo Tian, Po-Ting Lai, Qingqing Zhu, Chi-Ping Day, Christina Ross, Robert Leaman, and Zhiyong Lu. Geneagent: self-verification language agent for gene-set analysis using domain databases. *Nature Methods*, 22 (3):489–499, 2025. doi: 10.1038/s41592-025-02748-6. Preprint: arXiv:2405.16205.

[47] J. Harry Caufield, Harshad Hegde, Vincent Emonet, Nomi L. Harris, Marcin P. Joachimiak, Nicolas Matentzoglu, HyeongSik Kim, Sierra Moxon, Justin T. Reese, Melissa A. Haendel, Peter N. Robinson, and Christopher J. Mungall. Structured prompt interrogation and recursive extraction of semantics (spires): a method for populating knowledge bases using zero-shot learning. *Bioinformatics*, 40(3):btae104, 2024. doi: 10.1093/bioinformatics/btae104. Preprint: arXiv:2304.02711.

[48] Nathan E Lewis, Kim K Hixson, Tom M Conrad, Joshua A Lerman, Pep Charusanti, Ashoka D Polpitiya, Joshua N Adkins, Gunnar Schramm, Samuel O Purvine, Daniel Lopez-Ferrer, et al. Omic data from evolved *E. coli* are consistent with computed optimal growth from genome-scale models. *Molecular Systems Biology*, 6(1):390, 2010. doi: 10.1038/msb.2010.47.

[49] Valentyn Bezshapkin, Filip Schymik, Piotr Kucharski, Paweł Szczerbiak, Jakub W. Wojciechowski, Łukasz M. Szydłowski, and Tomasz Kościołek. Scaling and democratising structure-based protein function prediction with metagenomic-deepFRI. *bioRxiv*, page 2026.04.27.720983, 2026. doi: 10.64898/2026.04.27.720983. URL https://doi.org/10.64898/2026.04.27.720983.

[50] Theo Sanderson, Maxwell L. Bileschi, David Belanger, and Lucy J. Colwell. ProteInfer, deep networks for protein functional inference. *eLife*, 12:e80942, 2023. doi: 10.7554/eLife.80942.

[51] Nicolas Buton, François Coste, and Yann Le Cunff. Predicting enzymatic function of protein sequences with attention. *Bioinformatics*, 39(10):btad620, 2023. doi: 10.1093/bioinformatics/btad620.

[52] Gi Bae Kim, Ji Yeon Kim, Jong An Lee, Charles J Norsigian, Bernhard O Palsson, and Sang Yup Lee. Functional annotation of enzyme-encoding genes using deep learning with transformer layers. *Nature Communications*, 14(1):7370, 2023. doi: 10.1038/s41467-023-43216-z.

[53] Asaad Toonsi, Maxat Kulmanov, and Robert Hoehndorf. Genome-scale evaluation of protein function annotations. *Bioinformatics*, 2025.

[54] Morgan N Price, Kelly M Wetmore, R Jordan Waters, Mark Callaghan, Jayashree Ray, Hualan Liu, Jennifer V Kuehl, Ryan A Melnyk, Jacob S Lamson, Yumi Suh, et al. Mutant phenotypes for thousands of bacterial genes of unknown function. *Nature*, 557(7706): 503–509, 2018. doi: 10.1038/s41586-018-0124-0.

[55] Lorenz Christian Reimer, Joaquim Sardà Carbasse, Julia Koblitz, Christian Ebeling, Adam Podstawka, and Jörg Overmann. BacDive in 2022: the knowledge base for standardized bacterial and archaeal data. *Nucleic Acids Research*, 50(D1):D741–D746, 2022. doi: 10.1093/nar/gkab961.

[56] Isabel Schober, Julia Koblitz, Joaquim Sardà Carbasse, Christian Ebeling, Marvin Leon Schmidt, Adam Podstawka, Rohit Gupta, Vinodh Ilangovan, Javad Chamanara, Jörg Overmann, and Lorenz Christian Reimer. BacDive in 2025: the core database for prokaryotic strain data. *Nucleic Acids Research*, 53 (D1):D748–D756, 2025. doi: 10.1093/nar/gkae959.

[57] Julia Koblitz, Lorenz Christian Reimer, Rüdiger Pukall, and Jörg Overmann. Predicting bacterial phenotypic traits through improved machine learning using high-quality, curated datasets. *Communications Biology*, 8(1):897, 2025. doi: 10.1038/s42003-025-08313-3.

[58] Dingyi Rong, Bozitao Zhong, Wenzhuo Zheng, Liang Hong, and Ning Liu. Autoregressive enzyme function prediction with multi-scale multi-modality fusion. *Briefings in Bioinformatics*, 26(5):bbaf476, 2025. doi: 10.1093/bib/bbaf476.

[59] Yidong Song, Qianmu Yuan, Sheng Chen, Yuansong Zeng, Huiying Zhao, and Yuedong Yang. Accurately predicting enzyme functions through geometric graph learning on ESMFold-predicted structures. *Nature Communications*, 15(1):8180, 2024. doi: 10.1038/s41467-024-52533-w.

[60] Karel van der Weg, Erinc Merdivan, Marie Piraud, and Holger Gohlke. TopEC: prediction of Enzyme Commission classes by 3D graph neural networks and localized 3D protein descriptors. *Nature Communications*, 16(1):2737, 2025. doi: 10.1038/s41467-025-57324-5.

[61] Maxat Kulmanov and Robert Hoehndorf. DeepGOWeb: fast and accurate protein function prediction on the (Semantic) Web. *Nucleic Acids Research*, 49 (W1):W134–W140, 2021. doi: 10.1093/nar/gkab373. URL https://doi.org/10.1093/nar/gkab373.

[62] Michael Heinzinger, Konstantin Weissenow, Joaquin Gomez Sanchez, Adrian Henkel, Milot Mirdita, Martin Steinegger, and Burkhard Rost. Bilingual language model for protein sequence and structure. *NAR Genomics and Bioinformatics*, 6(4):lqae150, 2024. doi: 10.1093/nargab/lqae150.

[63] An Yang, Baosong Yang, Beichen Zhang, Binyuan Hui, Bo Zheng, Bowen Yu, Chengyuan Li, Dayiheng Liu, Fei Huang, Haoran Wei, Huan Lin, Jian Yang, Jianhong Tu, Jianwei Zhang, Jianxin Yang, Jiaxi Yang, Jingren Zhou, Junyang Lin, Kai Dang, Keming Lu, Keqin Bao, Kexin Yang, Le Yu, Mei Li,

Mingfeng Xue, Pei Zhang, Qin Zhu, Rui Men, Runji Lin, Tianhao Li, Tingyu Xia, Xingzhang Ren, Xuancheng Ren, Yang Fan, Yang Su, Yichang Zhang, Yu Wan, Yuqiong Liu, Zeyu Cui, Zhenru Zhang, and Zihan Qiu. Qwen2.5 technical report. *arXiv preprint*, abs/2412.15115, 2024. doi: 10.48550/arXiv.2412.15115.

[64] Aaron Weimann, Kyra Mooren, Jochen Frank, Phillip B. Pope, Andreas Bremges, and Alice C. McHardy. From genomes to phenotypes: Traitar, the microbial trait analyzer. *mSystems*, 1(6):e00101–16, 2016. doi: 10.1128/mSystems.00101-16.

[65] Wyatt T. Clark and Predrag Radivojac. Information-theoretic evaluation of predicted ontological annotations. *Bioinformatics*, 29(13):i53–i61, 2013. doi: 10.1093/bioinformatics/btt228. URL https://doi.org/10.1093/bioinformatics/btt228.