

a
De-Identified Stage 3 Resolution vs. ML Baselines (n = 120)
Performance on 120 Anomalies (%)
Precision (%)
Recall (%)
F1 Score (%)
Overall Accuracy (%)
55.2%
75.0%
82.8%
28.6%
85.2%
91.1%
94.2%
C1: Stage 1+2 Only
Threshold ≥0.35 + Guardrail
LogReg (3-Feat Pre-Tool CV)
GBDT (3-Feat Pre-Tool CV)
C2: Fast 1-Step Agent
C4: Qwen 27B No Guardrail
C3: Qwen 27B + Guardrail
GBDT (8-Feat Post-Tool CV)
b
Symbolic Guardrail Eliminates Cross-Clade Trap Violations
Accepted Taxon/Anatomical Violation Rate (%)
2.5% (3/120)
0.8% (1/120)
0.0% (0/120)
c
Agentic Disambiguation of 50 Multi-Candidate LRO Gaps
Unique LRO Metabolic Gaps (of 50)
Stage 1 RLGC Alone (Hit@1 = 34.0%, MRR = 0.570)
Stage 3 Agent Re-Ranking (Hit@1 = 82.0%, MRR = 0.874)
17
41
10
1
12
3
6
4
5
1
Rank 1 (Top-1)
Rank 2
Rank 3
Rank 4
Rank 5
d
Multi-Turn Latency Across 7 Taxa (1.8% Mean Loci)
Wall-Clock Latency / Anomaly (s)
Config 2: Fast 1-Step (2.47s wall-clock / 19.8s HTTP, 5.01 tools)
Config 3: Full Guarded (4.07s wall-clock / 32.5s HTTP, 5.74 tools)
E. coli K-12*
B. subtilis
P. aeruginosa
M. tuberculosis
S. cerevisiae
C. elegans
A. thaliana