



BIMM 143

Cancer Genomics & Immunoinformatics

Lecture 18

Barry Grant

UC San Diego

<http://thegrantlab.org/bimm143>

Today's Menu

Cancer Genomics

Brief review of cancer fundamentals,
What is cancer and what causes it?

Mining Cancer Genomic Data

Hands-on analysis to identify genomic changes in different cancers and identify new targets for therapy

Towards personalized cancer treatments

Recap on how the immune system normally detects cancer cells and how we can predict mutations that can be recognized by T cells

Cancer Immunoinformatics

Hands-on analysis to design personalized cancer vaccines

What is Cancer?

“Cancer is a name given to a collection of related diseases, where some of the body’s cells begin to divide without stopping and spread into surrounding tissue”

Source: <https://www.cancer.gov>

It is estimated that cancer will strike 40% of people at some point in their lifetime with frequently devastating effects.


What is Cancer?

“Cancer is a name given to a collection of related diseases, where some of the body’s cells begin to divide without stopping and spread into surrounding tissue”

Source: <https://www.cancer.gov>


Cancer is a disease of the Genome

- Caused by changes to genes that control the way our cells function, especially how they **grow and divide**.
- A major challenge in treating cancer is that every tumor is different: Each person's cancer has a unique combination of genetic changes (both "driver" & "passenger").
- As the cancer continues to grow, additional changes will occur.




Goals of Cancer Genome Research


- Identify changes in the genomes of tumors that drive cancer progression
- Identify new targets for therapy
- Select drugs based on the genomics of the tumor
- Provide early cancer detection and treatment response monitoring
- Utilize cancer specific mutations to derive neoantigen immunotherapy approaches



Finding Cancer Associated Mutations



Mutations detected: Point mutations



Mutations detected: Indels

Reference Sequence --- C T G G T G A C T A G T T ---

Mutations detected: Indels

Tumor Sequence 1: --- C T G G T G A T T ---

Reference Sequence: --- C T G G T G A **CTAG** T T ---


Deletion: CTAG deleted

Mutations detected: Indels

The diagram illustrates two mutations:

- Reference Sequence**: `- - - C T G G T G A C T A G T T - - -`
- Tumor Sequence 1**: `- - - C T G G T G A CTAG T T - - -` (CTAG deleted)
- Tumor Sequence 2**: `- - - C T G G T ATCA G A C T - - -` (ATCA inserted)

Mutations detected: Translocations



What can go wrong in cancer genomes?




Type of change	Some common technology to study changes
DNA mutations	WGS, WXS
DNA structural variations	WGS
Copy number variation (CNV)	CGH array, SNP array, WGS
DNA methylation	Methylation array, RRBS, WGBS
mRNA expression changes	mRNA expression array, RNA-seq
miRNA expression changes	miRNA expression array, miRNA-seq
Protein expression	Protein arrays, mass spectrometry





WGS = whole genome sequencing, WXS = whole exome sequencing

RRBS = reduced representation bisulfite sequencing, WGBS = whole genome bisulfite sequencing


Genomics allows us to answer the question:

How many mutations are there in typical cancers?






Identifying genes most commonly mutated in cancer



Identifying genes most commonly mutated in cancer




Three Main Types of Cancer Genes:

- **Oncogenes**, such as **Ras**, normally function to accelerate cell division and growth. They can be mutated to act like stuck gas pedals.
- **Tumor suppressor genes**, such as **p53** normal act like breaks. Mutations can cause these breaks to fail.
- **DNA repair genes**, such as **BRCA1 & 2**, normally function to fix minor damage to DNA when it replicates. When these genes are mutated, DNA damage can accumulate and lead to cancer.


Cell growth and survival genes

Many participate in signaling pathways that promote cell proliferation
(E.G. EGFR, Ras, BRAF, MEK etc.)




Cell growth and survival genes

Many participate in signaling pathways that promote cell proliferation
(E.G. EGFR, Ras, BRAF, MEK etc.)




Regulators of Cell Cycle and Cell Death



p53 Regulates Cell Division

Probably the most famous cancer gene that is mutated in about half of all tumors. Often called the '*guardian of the genome*'

- p53 normally shuts down cell division when a cell is stressed (e.g. by DNA damage)
- When DNA is damaged, p53 activates genes that stop cell growth or trigger the cell to die.
- Thus, p53 guards against changes to cells that might lead to tumor formation.
- It appears necessary to inactivate p53 to develop many forms of cancer.




Hands-on time!

https://bioboot.github.io/bimm143_F19/lectures/#18

Part 1 Only Please

Do it Yourself!



Representative H&E micrographs of rectus abdominis biopsies are displayed for two patients without cancer (left) and four patients with pancreatic cancer (right)

Next Up: Cancer Immunotherapy

Today's Menu

Cancer Genomics	Brief review of cancer fundamentals, What is cancer and what causes it?
Mining Cancer Genomic Data	Hands-on analysis to identify genomic changes in different cancers and identify new targets for therapy
Towards personalized cancer treatments	Recap on how the immune system normally detects cancer cells and how we can predict mutations that can be recognized by T cells
Cancer Immunoinformatics	Hands-on analysis to design personalized cancer vaccines

Nobel Prize 2018 in Medicine

- Identification of the molecules PD-1 and CTLA-4 that function as 'T cell brakes' (immune checkpoints)
- Blockade of PD-1 and CTLA-4 results in activation of T cells which has "fundamentally changed the outcome for certain groups of patients with advanced cancer"
- "Similar to other cancer therapies, adverse side effects are seen, which can be serious and even life threatening. They are caused by an overactive immune response leading to autoimmune reactions [...]"





© Nobel Media AB, Photo: A. Mahmoud
James P. Allison



© Nobel Media AB, Photo: A. Mahmoud
Tasaku Honjo

"for their discovery of cancer therapy by inhibition of negative immune regulation"


- Cancers genomes accumulate mutations
- Mutations in coding regions are translated in mutated protein sequences
- Mutated peptides can be presented as epitopes on **MHC** to **T cells**



- **Neoepitopes** are presumably recognized by tumor-infiltrating lymphocytes (TILs)
- **Neoepitopes** are highly tumor-specific!

Coulier et al, Nat Rev Cancer. 2014 Feb;14(2):135-46
Schumacher & Schreiber, Science. 2015 Apr 3;348(6230):69-74

DNA and RNA sequencing identifies tumor specific somatic mutations




Slide from: Bjoern Peters (LIAI)

- **Vaccination:** Introduce or boost an immune response against a specific target (**antigen**)
- Cancer cells contain non-self antigens that *could* be recognized by T cells, but the presence of cancer means this mechanism has failed, typically by the tumor suppressing immune responses
- **Checkpoint blockade treatments:** Block immune suppressive mechanisms to boost T cell immune responses against cancer cells.
- **Problem:** Checkpoint blockade is unspecific, and will also boost unwanted autoimmune responses
- **Personalized Cancer Immunotherapy:** Boost anti-tumor response with vaccine containing peptides corresponding to cancer mutations that can be recognized by T cells.


Q. How can such a vaccine be designed?

HLA Typing: Targeted sequencing of HLA locus



→ http://www.ashi-hla.org/publicationfiles/ASHI_Quarterly/25_2_2001/highthrusbt3.htm

Slide from: Bjoern Peters (LIAI)



Hands-on time!


https://bioboot.github.io/bimm143_F19/lectures/#18

Part 2: Designing a personalized cancer vaccine

Workflow:

- **Step 1:** Identify sequence regions that contain all **9-mer** peptides that are only found in the tumor
- **Step 2:** Run HLA binding prediction to identify 9-mer peptides in the sequence regions unique to the tumor to select peptides that can be **presented to T cells** in this patient
- **Step 3:** Determine if the identified peptides are specific for the tumor (i.e. are they found anywhere else?)
- **Final question:** Which peptide would you choose?

Depictions of the peptide bound MHC and T-cell receptor



Note:

- Anchor residues in the peptide bind to the allele-specific pockets of the MHC molecule.
- Certain MHC molecules (alleles) preferentially bind peptides with specific anchor residues in the 8- or 9-amino-acid peptide sequence.
- We want our tumor specific residues to be within 8 to 9-mer sequences bound by a patient HLA alleles!

Reference: <https://oncohemakey.com/how-t-cells-recognize-antigen-the-role-of-the-major-histocompatibility-complex/>

Bonus Slides (For Reference)

Slide from: Bjoern Peters (LIAI)

Calculate scoring matrix from affinities

Machine learning PSSM = Minimize the difference between predicted and measured binding affinities by varying the matrix values

N peptides with measured binding affinities	
$\log(\text{IC}_{50})$	Peptide
0.50	FQPQNGSFI
0.72	ISVANKLYM
2.37	RVYEALYYV
3.42	FQPQSGQFI
3.46	LYEKVKSQL
4.07	FKSVEFDMS
4.18	FQPQNGQFH
4.24	VLMLPVWFL
4.39	YMTLQGVVF
4.40	EDVKNAVGV
4.90	VFYEQMKRF
...	




HLA A*0201									
A	1	2	3	4	5	6	7	8	9
A	-0.3	0.8	-0.3	-0.3	-0.2	-0.3	0.0	0.0	-0.9
C	0.2	0.9	0.0	0.3	-0.5	-0.1	0.1	0.2	0.4
D	0.8	-0.9	-0.4	-0.3	0.3	0.2	0.4	0.3	0.6
E	-0.6	-0.4	0.7	-0.2	0.1	-0.4	-0.2	-0.2	-0.5
F	1.3	0.5	-0.5	0.1	-0.1	0.0	-0.3	-0.4	-0.8
G	-0.2	0.1	0.3	-0.1	0.0	0.4	0.3	-0.1	0.2
H	1.1	0.9	-0.1	0.4	0.1	0.2	0.0	0.2	0.8
I	-0.4	-0.7	-0.4	0.1	-0.1	-0.4	-0.5	0.5	-1.4
K	-0.3	0.0	1.1	0.1	0.1	0.6	0.9	0.2	0.9
L	0.0	-1.9	-0.4	-0.2	0.0	-0.2	0.0	-0.1	-1.1
M	-0.7	-1.2	-0.7	0.2	-0.6	0.0	0.0	0.0	-0.8
N	-0.1	0.3	0.1	-0.3	-0.1	-0.3	0.0	0.2	0.7
P	1.2	0.5	0.6	0.3	0.4	0.0	-0.4	-0.5	0.7
Q	0.4	-1.1	0.0	0.1	0.4	-0.2	-0.3	0.2	0.7
R	-0.2	0.9	1.0	0.3	0.1	0.4	0.7	0.0	0.9
S	-0.3	0.1	0.1	-0.4	0.1	0.3	0.2	-0.1	0.2
T	-0.2	-0.5	0.1	0.4	0.1	-0.5	0.2	0.0	-0.1
V	-0.1	-0.9	-0.1	0.2	0.0	-0.3	0.1	0.1	-1.9
W	0.0	0.7	-0.5	-0.2	-0.1	0.2	-0.3	-0.1	0.4
Y	-0.3	0.2	0.6	0.2	0.0	0.4	-0.4	-0.3	0.8

Offset: 4.3

Slide from: Bjoern Peters (LIAI)

Measuring and predicting MHC:peptide binding

Experimental Basis: MHC Binding Assay
List of peptides with allele specific binding affinity



Sequence	IC_{50}
QIVTMFEAL	3.6
LKGPDIFYK	308
NFCNLTSAF	50,000
AQSQRTRFR	38,000
CTYAGPFGM	143
CFGNTAVAK	50,000
...	

$\log(\text{IC}_{50}) \sim \text{Binding free Energy}$

low $\text{IC}_{50} \rightarrow$ high affinity

Impossible to measure all peptides

→ Predict binding peptides using machine learning


Find function F_i in F_1, F_2, F_3, \dots
 $F_i(\text{Sequence}) \approx \text{Affinity}$

Many different approaches
(ANN, SVM, HMM, LP, ...)


ORF 1	M G Q I V T M F E A L P H I T D E I N Y V I I V L I I V I T G I K A V Y N ...
ORF 2	M G L K G P D I Y K G V Y Q F K S V E F D M S H L N L T M P N A C S A N N ...
ORF 3	M H N F C N L T S A F N K K T F D H T L M S I V S S L H L S I D G N S N Y ...
ORF 4	M S A Q S O C R T F R G R V L D M F R T A F G G K Y M R S G W G W T G S D ...
ORF 5	M H C T Y A G P F G M S R I L L S Q E K T K F F T R R L A G T F T W T L S ...
ORF 6	M K C F G N T A V A K C N V N H D A E F C D M L R L I D Y N K A A L S K F ...
ORF 7	M L M R N H L I D I M G V P Y C N Y S K F W Y L E H A K T G E T S V P K C ...

Genetic and genomic approaches can identify a cancer's molecular signature to usefully stratify tumors for treatment


Stratify tumors based on molecular patterns



Stratify tumors based on molecular patterns




TCGA Pan-Cancer project



For example, breast cancer may be classified into various types based upon which proteins are expressed on the surface of the tumor cells. Breast tumors that express human epidermal growth factor 2 (HER2), estrogen receptor (ER), and progesterone receptor (PR), or are triple negative (do not express HER2, ER, or PR) behave differently and have different prognoses. Tumors that are HER2 positive are treated with medications that bind to HER2 (e.g. trastuzumab, lapatinib) and inhibit its activity. ER and PR are hormone receptors, and ER/PR positive tumors are treated with antihormonal therapies (e.g. tamoxifen and aromatase inhibitors). Triple negative tumors have the poorest prognosis and are unlikely to respond to HER2-targeted therapies or antihormonal therapies. Such cancers are usually treated very aggressively with chemotherapy.

As more has been learned about the molecular signature of various cancer subtypes, therapies that are specifically targeted to those signatures have been developed. Conventional chemotherapy acts on all rapidly dividing cells and does not distinguish between cancer cells and normal cells.

Classification of Breast Cancer



Readings to find out more...

Leading Edge
Review

The Genetic Basis for Cancer Treatment Decisions

Janet E. Dancey,^{1,2} Philippe L. Bedard,^{3,4} Nicole Onetto,³ and Thomas J. Hudson^{1,5,6,*}

¹Ontario Institute for Cancer Research, Toronto, ON M5G 0A3, Canada

²NCIC-Clinical Trials Group, Queen's University, Kingston, ON K7L 3N6, Canada

³Princess Margaret Hospital, Division of Medical Oncology and Hematology, University Health Network

⁴Department of Medicine

⁵Department of Medical Biophysics

⁶Department of Molecular Genetics

University of Toronto, Toronto, ON M5S 1A1, Canada

*Correspondence: tom.hudson@icr.on.ca


DOI 10.1016/j.cell.2012.01.014

Personalized cancer medicine is based on increased knowledge of the cancer mutation repertoire and availability of agents that target altered genes or pathways. Given advances in cancer genetics, technology, and therapeutics development, the timing is right to develop a clinical trial and research framework to move future clinical decisions from heuristic to evidence-based decisions. Although the challenges of integrating genomic testing into cancer treatment decision making are wide-ranging and complex, there is a scientific and ethical imperative to realize the benefits of personalized cancer medicine, given the overwhelming burden of cancer and the unprecedented opportunities for advancements in outcomes for patients.


Cell

Cancer immune surveillance and escape

- Mutations in cells occur frequently
- The immune system has the capacity to detect and eliminate such mutated cells, and will do so on a regular basis
- Only when mutated cells find ways to hide from- or suppress an attack by the immune system, they can grow and spread unhindered leading to clinically apparent tumors



Nobel Prize 2018 in Medicine



- Identification of the molecules PD-1 and CTLA-4 that function as 'T cell brakes' (immune checkpoints)
- Blockade of PD-1 and CTLA-4 results in activation of T cells which has "fundamentally changed the outcome for certain groups of patients with advanced cancer"
- "Similar to other cancer therapies, adverse side effects are seen, which can be serious and even life threatening. They are caused by an overactive immune response leading to autoimmune reactions [...]"

"for their discovery of cancer therapy by inhibition of negative immune regulation"

Several trials for personalized cancer vaccines are currently ongoing

Slide from: [Bjoern Peters \(LJAI\)](#)

UC San Diego Health

MEDICAL SERVICES PATIENT GUIDE FIND A PROVIDER LOCATIONS HEALTH INFO

Newsroom

Home / Newsroom / Releases / Is the Next Big Step in Cancer Therapy Personalized Vaccines?

Press Releases UC San Diego Health In The News News Features Trending Topics Podcast

Is the Next Big Step in Cancer Therapy Personalized Vaccines?

UC San Diego Health and La Jolla Institute launch clinical trial harnessing an individual's immune system in a syringe

October 12, 2016 | Yadira Galindo

After sequencing Tamara's tumor and normal tissue, the team identified mutations expressed solely by cancer cells in her body. Schoenberger and LJ's Bjoern Peters, PhD, developed a novel algorithm to select mutations that are recognized by the immune system. This algorithm was deployed to recognize the neoantigens that generated the strongest T cell response from Tamara's tissue samples. These neoantigens were then presented to Tamara's own T cells and cultured over a two-week period using 50 milliliters of her blood to develop a personalized vaccine.

Ezra Cohen, MD, UC San Diego Health physician scientist, administered the first-of-its-kind personalized cancer vaccine to Tamara Strauss, while Aaron Miller, MD, PhD, UC San Diego Health physician scientist, Tamara's mother, Iris Strauss, and Stephen Schoenberger, PhD, La Jolla Institute for Allergy and Immunology professor of immunology, look on.

Personalized Cancer Immunotherapy



After sequencing Tamara's tumor and normal tissue, the team identified mutations expressed solely by cancer cells in her body.

Schoenberger and LJ's Bjoern Peters, PhD, developed a novel algorithm to select mutations that are recognized by the immune system.

This algorithm was deployed to recognize the

neoantigens that generated the strongest T cell response from Tamara's tissue samples.

These neoantigens were then presented to Tamara's own T cells and cultured over a two-week period using 50 milliliters of her blood to develop a personalized vaccine.

14

Your Turn


Read and share your thoughts on the following class [Readings](#)

- Calling cancer's bluff with neoantigen vaccines
- Can genomics help detect early cancer and monitor treatment effectiveness?
- The increasing cost of cancer therapies

https://bioboot.github.io/bimm194_W18/readings/

1. Predict consequences of mutations

```
ACTGCCTACGTCTCACCGTCGACTTCAATCGCTTAACCGTACTCCCATGCTACTGCATCTGGGTTAACTC  
GACGTTTTTTCATGCATGTGCACCCCAATATATATGCAACTTTGTGCACCTCTGCACCGCGAGTTGCA  
CTGTCGCCCTGTGCATGTGCACTGTCTTCGCTGACTGCCTACGTCTCACCGTCGACTTCAAATCGTT  
AACCGTACTCCCATGCTACTGCATCTGGTTAACCGTTAACCGTACTGCATGTGCACCCAAATATA  
TGCAACTTTGTGCACCTCTGTACCGCGAGTTGCACGTGCCCCTGTGCATGTGCACTGTCTTCGAA
```




2. Assess the functional impact of nsSNVs

nsSNVs = non-synonymous Single Nucleotide Variant (missense)

ATC GAA GCA CGT
Met Glu Ala Gly

ATC GAC GCA CGT
Met **Asp** Ala Gly



Computational methods to assess the functional impact of nsSNVs

MutationTaster	LogRe	MutPred	SNPs&GO
CanPredict	Condel	CHASM	SNPeffect
SIFT	PolyPhen2	MutationAssessor	PMut

3. Identify cancer drivers from somatic mutations




Patient cohort



Normal cell



Cancer cell



Sequencing machines



A**T**ATGCCA
TC**A**TGTC
GGT**A**TCG
CAGC ...


Somatic mutations



Which mutations are cancer drivers?

Find signals of selection across tumors

Cancer is an evolutionary process



Yates and Campbell et al, Nat Rev Genet 2012

How to differentiate drivers from passengers?

```
ACTGCCTACGCTCACCGTCGACTTCAAATCGCTTAACCGTACTCCCATGCTACTGC
ATCTCGGGTTAACCGACGTTTTTCATGCATGTGTCACCCCAATATATATGCAACT
TTGTGCACCTCTGTACCGCGAGTTGGCAGTGTGCCCCCTGTGTCATGTGCACT
CTCTCGCTGCACTGCCAACGCTACGTCTACCGTCGACTCAATCTAACCGTACTCCC
ATGCTACTGCACTCGGGTTAACCGACGTTTTGCATGCATGTGTCACCCCAATA
TATGCAACTTTTGTCACCTCTGTACCGCGAGTTGGCAGTGTGCCCCCTGTGCA
TGTGCACTGTCTTCGAGTTTGCATGCATGTGCACTGTGACCTCTGTACGTCT
```




How to differentiate drivers from passengers?

ACTG**C**CTACGTCTACCGTCGACTTCAAATCG**C**TTAACCCGACTCCCAGTGCAGTGC
ATCTCGGGTTAACTCGACGTTTCATGCATGTGCAACCCAAATATATATGCA**A**CTT
TTGTCACCTCTGTCACGCCAGTTGGCAGTGTGCACTGTCGCCCTGTGCAATGTGCACTGT
CTC**T**CGCTGCACTGCCTACCGTCAACCGTCAACT**C**TTAACCCGACTCCC
ATGCTACTGCACTCGGGTTAACTCGACGTTTG**C**ATGCATGTGCAACCCAAATA
TATGCA**A**CTTTGTCACCTCTGTCACGCCAGTTGGCACTGTGCAACCTGTGCA
TGTGCACTGTCT**T**CGAGTTTG**C**ATGCATGTGCACTGTGCAACCTGTGCA



Find signals of positive selection across tumour re-sequenced genomes



PIK3CA is recurrently mutated in the same residue in breast tumours

Signals of positive selection

Recurrence

MuSiC-SMG / MutSigCV




○ Mutation

Identify genes mutated more frequently than background mutation rate

Mutation clustering

OncodriveCLUST



○ Mutation

<http://www.intogen.org/mutations/analysis>

IntOGen Mutations Analysis

[Download](#)

To interpret catalogs of cancer somatic mutations.

Cohort analysis

Use this if you have a list of somatic mutations for a cohort of tumors and want to identify driver mutations, genes and pathways.

[View an example](#)

[Analyse your data](#)

Single tumor analysis

Use this if you have a list of somatic mutations for a single tumor and want to rank them based on their implication in cancer development.

[View an example](#)

[Analyse your data](#)