

CHAPTER 5.8. ANTARCTIC HEXACORALS (CNIDARIA: ANTHOZOA: HEXACORALLIA). Rodríguez E., Fautin D.G. 2014.

In: De Broyer C., Koubbi P., Griffiths H.J., Raymond B., Udekem d'Acoz C. d', et al. (eds.). Biogeographic Atlas of the Southern Ocean. Scientific Committee on Antarctic Research, Cambridge, pp. 113-116.

EDITED BY:

Claude DE BROYER & Philippe KOUBBI (chief editors)

with Huw GRIFFITHS, Ben RAYMOND, Cédric d'UDEKEM d'ACOZ, Anton VAN DE PUTTE, Bruno DANIS, Bruno DAVID, Susie GRANT, Julian GUTT, Christoph HELD, Graham HOSIE, Falk HUETTMANN, Alexandra POST & Yan ROPERT-COUDERT

SCIENTIFIC COMMITTEE ON ANTARCTIC RESEARCH

THE BIOGEOGRAPHIC ATLAS OF THE SOUTHERN OCEAN

The "Biogeographic Atlas of the Southern Ocean" is a legacy of the International Polar Year 2007-2009 (www.ipy.org) and of the Census of Marine Life 2000-2010 (www.coml.org), contributed by the Census of Antarctic Marine Life (www.caml.aq) and the SCAR Marine Biodiversity Information Network (www.scarmarbin.be; www.biodiversity.aq).

The "Biogeographic Atlas" is a contribution to the SCAR programmes Ant-ECO (State of the Antarctic Ecosystem) and AnT-ERA (Antarctic Thresholds- Ecosystem Resilience and Adaptation) (www.scar.org/science-themes/ecosystems).

Edited by:

Claude De Broyer (Royal Belgian Institute of Natural Sciences, Brussels)

Philippe Koubbi (Université Pierre et Marie Curie, Paris)

Huw Griffiths (British Antarctic Survey, Cambridge)

Ben Raymond (Australian Antarctic Division, Hobart)

Cédric d'Udekem d'Acoz (Royal Belgian Institute of Natural Sciences, Brussels)

Anton Van de Putte (Royal Belgian Institute of Natural Sciences, Brussels)

Bruno Danis (Université Libre de Bruxelles, Brussels)

Bruno David (Université de Bourgogne, Dijon)
Susie Grant (British Antarctic Survey, Cambridge)
Julian Gutt (Alfred Wegener Institute, Helmoltz Centre for Polar and Marine Research, Bremerhaven)

Christoph Held (Alfred Wegener Institute, Helmoltz Centre for Polar and Marine Research, Bremerhaven)
Graham Hosie (Australian Antarctic Division, Hobart)

Falk Huettmann (University of Alaska, Fairbanks)

Alix Post (Geoscience Australia, Canberra)

Yan Ropert-Coudert (Institut Pluridisciplinaire Hubert Currien, Strasbourg)

Published by:

The Scientific Committee on Antarctic Research, Scott Polar Research Institute, Lensfield Road, Cambridge, CB2 1ER, United Kingdom (www.scar.org).

Publication funded by:

- The Census of Marine Life (Albert P. Sloan Foundation, New York)
- The TOTAL Foundation, Paris

The "Biogeographic Atlas of the Southern Ocean" shared the Cosmos Prize awarded to the Census of Marine Life by the International Osaka Expo'90 Commemorative Foundation, Tokyo, Japan.

Publication supported by:

- $The \ Belgian \ Science \ Policy (Belspo), through \ the \ Belgian \ Scientific \ Research \ Programme \ on \ the \ Antarctic \ and \ the \ "biodiversity.aq" \ network \ (SCAR-MarBIN/ANTABIF)$
- The Royal Belgian Institute of Natural Sciences (RBINS), Brussels, Belgium
 The British Antarctic Survey (BAS), Cambridge, United Kingdom
- The Université Pierre et Marie Curie (UPMC), Paris, France
- The Australian Antarctic Division, Hobart, Australia
- The Scientific Steering Committee of CAML, Michael Stoddart (CAML Administrator) and Victoria Wadley (CAML Project Manager)

Mapping coordination and design: Huw Griffiths (BAS, Cambridge) & Anton Van de Putte (RBINS, Brussels)

Editorial assistance: Henri Robert, Xavier Loréa, Charlotte Havermans, Nicole Moortgat (RBINS, Brussels)

Printed by: Altitude Design, Rue Saint Josse, 15, B-1210 Brussels, Belgium (www.altitude-design.be)

Lay out: Sigrid Camus & Amélie Blaton (Altitude Design, Brussels).

Cover design: Amélie Blaton (Altitude Design, Brussels) and the Editorial Team.

Cover pictures: amphipod crustacean (Epimeria rubrieques De Broyer & Klages, 1991), image © T. Riehl, University of Hamburg; krill (Euphausia superba Dana, 1850), image © V. Siegel, Institute of Sea Fisheries, Hamburg; fish (*Chaenocephalus* sp.), image © C. d'Udekem d'Acoz, RBINS; emperor penguin (*Aptenodytes forsteri* G.R. Gray, 1844), image © C. d'Udekem d'Acoz, RBINS; Humpback whale (*Megaptera novaeangliae* (Borowski, 1781)), image © L. Kindermann, AWI

Online dynamic version:

A dynamic online version of the Biogeographic Atlas is available on the SCAR-MarBIN / AntaBIF portal: atlas.biodiversity.aq.

Recommended citation:

De Broyer C., Koubbi P., Griffiths H.J., Raymond B., Udekem d'Acoz C. d', Van de Putte A.P., Danis B., David B., Grant S., Gutt J., Held C., Hosie G., Huettmann F., Post A., Ropert-Coudert Y. (eds.), 2014. Biogeographic Atlas of the Southern Ocean. Scientific Committee on Antarctic Research, Cambridge, XII + 498 pp.

For individual chapter:

(e.g.) Crame A., 2014. Chapter 3.1. Evolutionary Setting. In: De Broyer C., Koubbi P., Griffiths H.J., Raymond B., Udekem d'Acoz C. d', et al. (eds.). Biogeographic Atlas of the Southern Ocean. Scientific Committee on Antarctic Research, Cambridge, pp. xx-yy.

ISBN: 978-0-948277-28-3



This publication is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License



5.8. Antarctic Hexacorals (Cnidaria, Anthozoa, Hexacorallia)

Estefanía Rodríguez¹ & Daphne G. Fautin²

- ¹ Division of Invertebrate Zoology, American Museum of Natural History, New York City, NY, USA.
- ² Department of Ecology and Evolutionary Biology and Natural History Museum and Biodiversity Institute, University of Kansas, Lawrence, KS, USA.

1. Introduction

Subclass Hexacorallia (Cnidaria, Anthozoa) comprises sea anemones (members of order Actiniaria), black corals (members of order Antipatharia), tube anemones (members of order Ceriantharia), mushroom anemones (members of order Corallimorpharia), stony corals (members of order Scleractinia) and mat anemones (members of order Zoanthidea). Members of Hexacorallia, which currently contains about 4300 extant species (Daly et al. 2007), are characterised by hexamerous symmetry (although eight- or ten-part symmetry is not uncommon) and spirocysts, a type of cnida with a single-walled capsule and a tubule with tiny entangling threads. Actiniaria (~1100 species) and Scleractinia (~1600 species) are the most species-rich orders; the other four orders represent less than a quarter of the diversity (Antipatharia 235 species; Ceriantharia 100 species; Corallimorpharia 45 species; and Zoanthidea ~200 species).

Hexacorals are important components of macrobenthic communities in the Southern Ocean in terms of biomass and as part of the suspension feeder communities that dominate some habitats. The most recent taxonomic and biogeographic reviews of members of this taxon in the Southern Ocean identified 42 species of order Scleractinia (Cairns 1982) and 116 species of order Actiniaria (Dunn 1983, Fautin 1984, Rodríguez *et al.* 2007). Fautin (1984) reported two species of Corallimorpharia and described another (Fautin 2011a) that reaches the waters we inventory. Rodríguez *et al.* (2007) included six species, four of which we recognise here; they listed one that we consider a synonym, and the other was included because of differences in the geographical limits of their inventory. The other three hexacoral orders are understudied and represent perhaps only about 15 species.

Actiniaria and Scleractinia in the Southern Ocean follow the general biogeographic pattern of other invertebrates — high endemism, circumpolar species, differentiation between East and West Antarctica faunae, etc. (Cairns 1982, Rodríguez et al. 2007). Although baselines for biogeography of these two major hexacoral orders have been written, new data are available (particularly for sea anemones). No synthesis exists for any of the other orders in the Southern Ocean. Here we provide an overview of the entire subclass; our main generalisations are relatively high levels of endemism and complementarity in the distribution the six orders of hexacorals.



Photo 1 Hormathia lacunifera (Stephenson, 1918). Det. E. Rodríguez. Larsen B (Polarstern ANT-XXIII/8, st. 720-2). The elongate organisms on the column of the sea anemone are antarcturid isopods. Image: J. Gutt © AWI/Marum, University of Bremen, Germany.

2. Methods

We include bibliographic records of hexacorals (Cairns 1982, Cairns et al. 2005, Rodríguez et al. 2007, Fautin 2011b, Rodríguez & López-González 2013). The limits of our geographical coverage are those of SCAR-MarBIN for the Southern Ocean in its broad sense (e.g. Deacon 1984, Rintoul 2007). We consider the Antarctic Region (ANT) to be the zone south of the Polar Front (PF) (or Antarctic Convergence; 48°S to 63°S, average 55°S); the Sub-

Antarctic Region (SBA) to extend from the Polar Front in the south to the Sub-Tropical Front in the north (30°S to 47°S, average 43°S); and the limit of the Southern Ocean (O) starts north of the Sub-Tropical Front.

3. Biodiversity and biogeographic patterns of Antarctic and sub-Antarctic hexacorals

3.1. Actiniaria (Maps 1-5)

The sea anemone fauna of the Southern Ocean includes 116 species (Table 1), 54 in the Antarctic region (29 ANT + 23 ANT-SBA + 2 ANT-SBA-O) (Rodríguez et al. 2007, Fautin 2011b). Endemism is high at the species level (53% for the Antarctic Region, 81% for the Southern Ocean) but low at higher taxonomic levels (25% endemic genera and no endemic families). The Antarctic Peninsula area has the highest proportion of endemic species (e.g. Stephanthus antarcticus Rodríguez & López-González, 2003), several of which extend to the Weddell Sea. More than 25% of species in the Antarctic Region are circumpolar (e.g. Glyphoperidium bursa Roule, 1909 and Stomphia selaginella (Stephenson, 1918)). However, only 15% of species cross the Polar Front — that is, are shared with the Magellan Province (e.g. Isocicyonis alba (Studer, 1879), and Antholoba achates Drayton in Dana, 1846) — or extend to the islands near New Zealand. Only two Southern Ocean species are documented to extend beyond the Southern Ocean, Liponema multiporum Hertwig, 1882, and Galatheanthemum profundale Carlgren, 1956, presumably having invaded the Southern Ocean from the deep sea. More than 61% of species in the Antarctic Region are restricted to the continental shelf (<1000 m). Most eurybathic species occur in both Antarctic and Sub-Antarctic Regions.

3.2. Antipatharia (Map 6)

Seventeen species of black corals have been documented in the Southern Ocean, nine in deep waters of the Antarctic Region (5 ANT + 2 SBA + 2 ANT-SBA-O: Table 1): Abyssopathes lyriformis Opresko, 2002 (Ross Sea, 3500 m), Schizopathes affinis Brook, 1889 (north of Amundsen Sea, 4895 m), Dendrobathypathes grandis Opresko, 2002 (West Antarctica, 417–686 m), Bathypathes bifida Thompson, 1905 (Weddell Sea and Antarctic Peninsula, 2569-3018 m), B. alternata Brook, 1889 (West Antarctica, 3914-4863 m), and B. tenuis Brook, 1889 (off Amundsen Sea, 4005-4755 m) also cited extending north of the Sub-tropical Front (Indian Ocean, 4755 m). Heliopathes heterorhodzos (Cooper, 1909) (3475-4667 m) and Schizopathes crassa Brook, 1889 (2836-3040 m) have a circumpolar distribution; the latter and B. alternata also extend into the Sub-Antarctic Region. Only B. patula Brook, 1889, is distributed across the entire Southern Ocean; in the Antarctic Region, it is known from East and West Antarctica (2500-4200 m) whereas in the Sub-Antarctic Region it is known only from the eastern and southwest Pacific (Chilean fjords, 1583-1908 m; southwest Pacific islands, 604 m); in the southwest Pacific it also occurs north of the Sub-Tropical Front (4863 m). Eight other species occur in the Sub-Antarctic Region: D. isocrada Opresko, 2002 (off Auckland Island, 728 m), Stauropathes staurocrada Opresko, 2002 (off Auckland Island, 814 m), Trissopathes tristicha (van Pesch, 1914) (off Auckland Island, 728 m), Cladopathes plumosa Brook, 1889 (off Auckland and Prince Edward islands, 566-728 m), T. tetracrada Opresko, 2003 (east Pacific, 1150 m), Antipathes speciosa (Brook, 1889) (east Pacific, 320 m), and Antipathella fiordensis (Grange, 1990) (east Pacific, 15 m). An unidentified species of Cirripathes was reported off Auckland Island (728-837 m) (de Blainville 1830).

3.3. Ceriantharia (Map 7)

One species of tube anemone is recorded in the Southern Ocean; identified only to order (Arntz & Brey 2001), the specimens were collected off the Antarctic Peninsula (so in the Antarctic Region) at a depth of 150–300 m. Although this order is not very speciose (Daly *et al.* 2007), its remarkable sparsity in the Antarctic Region might be an artefact of collecting. Ceriantharians live in a tube burrowed into soft sediment; because they are able to retract rapidly into the tube when disturbed, collecting them is difficult, particularly with trawls (this conventionally used collecting gear also damages the animals).

Table 1 Species of hexacorals in each region of the Southern Ocean (SO). Numbers in brackets are species shared by ANT and SBA (therefore, number is in both columns).

Order	Antarctic region (ANT)	Sub-Antarctic region (SBA)	SBA and north (SBA-O)	All 3 regions (ANT-SBA-O)	Total (proportion endemic)
Actiniaria	29 [23]	42 [23]	20	2	116 (81%)
Antipatharia	5 [2]	8 [2]	0	2	17 (88%)
Ceriantharia	1 [0]	0 [0]	0	0	1 (?)
Corallimorpharia	0 [0]	1 [0]	2	2	5 (20%)
Scleractinia	7 [2]	7 [2]	20	6	42 (38%)
Zoanthidea	1 [0]	4 [0]	0	1	6 (83%)
TOTAL	43 [27]	62 [27]	42	13	187 (71%)



3.4. Corallimorpharia (Map 8)

Five species of mushroom anemones have been documented in the Southern Ocean, Corallimorphus profundus Moseley, 1877, C. rigidus Moseley, 1877, C. niwa Fautin, 2011a, Corynactis carnea Studer, 1879, and Corynactis chilensis Carlgren, 1941. Most specimens of Corallimorphus are from deep waters, although some records are from water shallower than 100 m in the Southern Ocean (Fautin 2011b). Corallimorphus rigidus and C. profundus are widely distributed, occurring also in the northern hemisphere, and extending south of the Polar Front; C. profundus is circumpolar, inhabiting shallower depths in the Antarctic than in the Sub-Antarctic Region (continental shelf vs. bathyal depths). Corallimorphus niwa is a sub-Antarctic species found around New Zealand. In the Southern Ocean, specimens of Corynactis have been reported to about 50 m off both coasts of South America to north of the Sub-Tropical Front.

3.5. Scleractinia (Map 9)

Of the 42 species of stony corals in the Southern Ocean (Cairns 1982, Cairns et al. 2005), four are of uncertain identity (Cairns 1982). They display three biogeographic patterns: some are confined to the Antarctic Region; some are confined to the Sub-Antarctic Region; and some are cosmopolitan or widespread, only barely reaching the Sub-Antarctic Region (Table 1). Twentysix species (20 SBA-O + 6 ANT-SBA-O; 62%) are cosmopolitan; 14 species (7 ANT + 7 SBA; 33%) are endemic to either the Antarctic or the Sub-Antarctic Region, whereas two species (<5%) are endemic to both the Antarctic and Sub-Antarctic Regions.

3.6. Zoanthidea (Map 10)

Six nominal species of mat anemones are documented in the Southern Ocean; only Parazoanthus antarcticus (Carlgren, 1927) (Antarctic Peninsula, 290 m) occurs in the Antarctic Region (Fautin 2011b). Four records of zoanthid specimens identified only as Parazoanthus sp. (Arntz & Brey 2001, 2003, 2005, Fütterer et al. 2003) are from West Antarctica (Antarctic Peninsula and eastern Weddell Sea area, 306-3925 m). Five named species are restricted to shallow waters in the Sub-Antarctic Region around South America: Parazoanthus fuegiensis Carlgren, 1899 (also Carlgren 1927; Magellan Strait), P. elongatus McMurrich, 1904 (southern Chile, 20-25 m), Epizoanthus fiordicus Sinniger & Häussermann, 2009 (southern Chile, 28 m), Mesozoanthus fossii Sinniger & Häussermann, 2009 (southern Chile, 25–29 m), and Epizoanthus patagonicus Carlgren, 1899 (Argentinean Patagonia, 20-197 m), the only species to extend north of the Sub-Tropical Front.

3.7. General remarks

Of the three orders that have been well studied, the percentage of species in the Southern Ocean relative to the total number of species worldwide is lower for Scleractinia (2.5%) than for Corallimorpharia and Actiniaria (about 11% each). We note some complementarity (Fautin 1989) in the distribution of members of the three hexacoral orders for which there are data.

Data concerning distribution of Antipatharia, Ceriantharia, and Zoanthidea are few in the Southern Ocean, many species being known from single sites. Both limited sampling and limited taxonomic expertise confound any analysis. Members of order Antipatharia are restricted to deep waters (which is typical of the taxon); species in the Sub-Antarctic Region are restricted to either the surroundings of the Auckland Island area or the east Pacific coast, and occur in shallower water than those in the Antarctic Region. That species seem to have small ranges (most of the Antarctic ones are restricted to West Antarctica, only three being circumpolar) may be an artefact of poor knowledge. Zoanthids are more or less restricted to shallow waters of the Sub-Antarctic Region; like antipatharians, they occur shallow in the Sub-Antarctic Region and deeper in the Antarctic Region. However, the only zoanthid species known from the Antarctic Region is apparently circumpolar

The distribution patterns of antipatharians and zoanthids might be related to their ecology. Many zoanthids are epizoic. The availability of substrate in the Antarctic Region is considered a key factor regulating benthic communities (Lovell & Trego 2003), which are highly structured; competition for substrate is important (Dayton 1990). Their restricted distributions could be because of their larvae: the only antipatharian species to have been studied, which live in shallow water, have lecithotrophic crawling larvae with low dispersal capacities (Grigg 1965, Miller 1998). The relatively small size of antipatharian polyps might be adapted to exploit small particles of organic matter that are available year round in "food banks" (Mincks et al. 2005), and thereby favor members of this order in deeper Antarctic waters.

Continental shelves in the Antarctic Region are dominated by solitary scleractinians and sea anemones. However, whereas about 15% of Southern Ocean scleractinian species are widespread (6 species ANT-SBA-O), only 2% (2 species ANT-SBA-O) of actiniarian species recorded in the Antarctic Region extend north of the Sub-Antarctic Region (Table 1). This difference is more remarkable in the proportion of Southern Ocean species reported only north of the Polar Front (SBA-O + ANT-SBA-O species): almost 70% of scleractinians but only 19% of actiniarians. The predominance of widely distributed or cosmopolitan species of members of Corallimorpharia resembles the pattern of Scleractinia; this is maybe because members of these two orders are closely related phylogenetically (Daly et al. 2007, Daly & Fautin 2007).

However, in contrast with Scleractinia, no corallimorpharian species seem to have diversified in the Southern Ocean — two of the five known species occur in northern latitudes and only those two barely reach the Antarctic Region.

There is no clear explanation for differences in distribution of Southern Ocean scleractinians and actiniarians. It is unlikely to be the calcified skeleton, which actiniarians lack and scleractinians have, even though the cold and deep waters of the Southern Ocean are undersaturated with calcium carbonate, because corallimorpharians, with a distribution resembling that of scleractinians, lack one as well. Dispersal could account for these differences, but, although reproductive and larval biology of Southern Ocean hexacorals has not been well studied, among actiniarians are external and internal brooders, gonochores and hermaphrodites, and broadcast spawners, so reproductive/ sexual pattern may not be important in explaining their differential success

Acknowledgments

Mercer Brugler (AMNH) is thanked for his helpful comments of the order Antipatharia. Comments from Meg Daly (Ohio State University) improved this manuscript. Huw Griffiths (BAS, Cambridge) and Anton Van de Putte (RBINS, Brussels) prepared the maps. This is CAML contribution # 106.

References

- Arntz, W.E., Brey, T. (eds.), 2001. The expedition ANTARKTIS XVII/3 (EASIZ III) of RV "Polarstern"
- to the eastern Weddell Sea in 2000. Berichte zur Polar-und Meeresforschung, 402, 1–181.

 Arntz, W.E., Brey, T. (eds.), 2003. The expedition ANTARKTIS XIX/5 (LAMPOS) of the R/V "Polarstern" in 2002. Berichte zur Polar-und Meeresforschung, **462**, 1–120.

 Arntz, W.E., Brey, T., 2005. The expedition ANTARKTIS XXI/2 (BENDEX) on RV "Polarstern" in
- 2003/2004 (eds.). Berichte zur Polar-und Meeresforschung, 503, 1–149.
 Brook, G., 1889. Report on the Antipatharia. Reports of the Scientific Results of the Voyage of the
- Challenger, Zoology, **32**, 5–222.

 Cairns, S.D., 1982. Antarctic and Subantarctic Scleractinia. Biology of the Antarctic Seas XIV, Antarctic Research Series, **34(1)**, 1–74.
- Cairns, S.D., 1989. A revision of the ahermatypic Scleractinia of the Philippine Islands and adjacent waters, Part 1: Fungiacyathidae, Micrabaciidae, Turbinoliinae, Guyniidae, and Flabellidae. Smithsonian Contributions to Zoology, 486, 1–136.
 Cairns, S.D, Häussermann, V., Försterra, G., 2005. A review of the Scleractinia (Cnidaria: Anthozoa)

- of Chile, with the description of two new species. *Zootaxa*, **1018**, 15–46.

 Carlgren, O., 1899. Zoantharien. *Hamburger Magalhaensische Sammelreise*, **4(1)**, 1–48.


 Carlgren, O., 1927. Actiniaria and Zoantharia. In: Odhner, T., (ed.). Further Zoological Results of the Swedish Antarctic Expedition 1901-1903, **2(3)**. Stockholm: Norstedt, P.A., Söner, 1–102. Carlgren, O., 1941.Corallimorpharia, Actiniaria, and Zoantharia. *Results of the Norwegian Scientific*
- Expedition to Tristan da Cunha 1937–1938, 1(8), 1–12.
 Carlgren, O., 1956. Actiniaria from depths exceeding 6000 meters. Galathea Reports, 2, 9–16
- Cooper, F. O., 1909. Antipatharia. *Transactions of the Linnean Society (London)*, 12(17), 301–321.
 Daly, M., Fautin, D.G., 2007. Corallimorpharia. In: Linnaeus Tercentenary: Progress in Invertebrate Taxonomy. *Zootaxa*, 1668, 139–140.
- Daly, M., Brugler, M.R., Cartwright, P., Collins, A.G., Dawson, M.N., Fautin, D.G., France, S.C., McFadden, C.S., Opresko, D.M., Rodriguez, E., Romano, S., Stake, J., 2007. The phylum Cnidaria: A review of phylogenetic patterns and diversity three hundred years after Linneaeus. In: Linnaeus Tercentenary: Progress in Invertebrate Taxonomy. *Zootaxa*, **1668**, 127–182.
- Dayton, P.K., 1990. Polar benthos. In: Smith, W.O. (ed.). Polar oceanography, Part B: Chemistry, biology and geology. San Diego: Academic Press, 631–685.
- Deacon, G., 1984. *The Antarctic circumpolar ocean*. Cambridge University Press, 180 pp. de Blainville, H.M., 1830. Zoophytes. Dictionnaire des Sciences naturelles. Paris: F.G. Levrault, **60**, 1-548.
- Dunn, D.F., 1983. Some Antarctic and sub-Antarctic sea anemones (Coelenterata: Ptychodactiaria and Actiniaria). Biology of the Antarctic Seas XIV, Antarctic Research Series, 39, 1–67.
- Fautin, D.G., 1984. More Antarctic and Subantarctic sea anemones (Coelenterata: Corallimorpharia and Actiniaria). Biology of the Antarctic Seas XIV, *Antarctic Research Series*, **41**, 1–42.
- Fautin, D.G., 1989. Anthozoan dominated benthic environments. Proceedings of the Sixth
- International Coral Reef Symposium, 3, 231–236. Fautin, D.G., 2011a, Corallimorphus niwa new species (Cnidaria: Anthozoa), New Zealand members of Corallimorphus, and redefinition of Corallimorphidae and its members. Zootaxa, 2775, 37-
- Fautin, D.G., 2011b, Hexacorallians of the World, Accessed through: World Register of Marine Species at http://www.marinespecies.org/aphia.php?p=taxdetails&id=290089 on 2013-09-23
- http://geoportal.kgs.ku.edu/hexacoral/anemone2/index.cfm (viewed September 2012). Fütterer, D.K., Brandt, A., Poore, G.C.B., 2003. The expeditions Antarktis-XIX/ 3 and XIX/4 of the Research Vessel Polarstern in 2002 (ANDEEP I and II: Antarctic Benthic deep-sea Biodiversity: colonisation history and recent community patterns) (eds.). Berichte zur Polar-und Meeresforschung, 470, 1–174.
- Grange, K.R., 1990, *Antipathes fiordensis*, a new species of black coral (Coelenterata: Antipatharia)

- Grange, K.R., 1990. Antipathes nordensis, a new species of black coral (Coelenterata: Antipatharia) from New Zealand. New Zealand Journal of Zoology, 17, 279–282.

 Grigg, R.W., 1965. Ecological studies of black coral in Hawaii. Pacific Science, 19, 244–260. Hertwig, R., 1882. Die Actinien der Challenger Expedition. Jena: Gustav Fischer, 119 pp. Lovell, L.L., Trego, K.D., 2003. The epibenthic megafaunal and benthic infaunal invertebrates of Port Foster, Deception Island (South Shetland Islands, Antarctica). Deep-Sea Research II, 50(11), 1799–1819.
- McMurrich, J.P., 1904. The Actiniae of the Plate Collection. Zoologische Jahrbücher, 2(6), 215–306. Marenzeller, E., 1904. Steinkorallen. Wissenschaftliche Ergebnisse der Deutschen Tiefsee-Expedition auf dem Dampfer "Valdivia" 1898–1899, 7(3), 261–318.
- Miller, K.J., 1998. Short-distance dispersal of black coral larvae: inference from spatial analysis of colony genotypes. *Marine Ecology Progress Series*, **163**, 225–233.


 Mincks, S., Smith, C., De Master, D., 2005. Persistence of labile organic matter and microbial biomass in Antarctic shelf sediments: evidence of a sediment "food bank". *Marine Ecology*
- Progress Series, 300, 3–19.

 Moseley, H.N., 1876. Preliminary report to Professor Wyville Thomson, F.R.S., director of the civilian scientific staff, on the true corals dredged by H.M.S. 'Challenger' in deep water between the dates Dec. 30th, 1870, and August 31st, 1875. Proceedings of the Royal Society of London, **24**. 544-569.
- Moseley, H.N., 1877. On new forms of *Actiniaria* dredged in the deep sea; with a description of certain pelagic surface-swimming species. *Transactions of the Linnean Society London*, ser. 2. **1(5)** 295–305
- Opresko, D.M., 2002. Revision of the Antipatharia (Cnidaria: Anthozoa). Part II. Schizopathidae.
- Zoologische Mededelingen Leiden, **76**, 411–442.
 Opresko, D.M., 2003. Revision of the Antipatharia. (Cnidaria: Anthozoa). Part III. Cladopathidae. Zoologische Mededelingen Leiden, **77**, 495–536.
- van Pesch, A.J., 1914. The Antipatharia of the Siboga Expedition. Siboga Expeditie Monographie, 17, 1–258.



Hexacorallia Maps 1–6 Antarctic and sub-Antarctic records of representative species of order Actiniaria, and records of order Antipatharia. Map 1. Examples of Antarctic endemic circumpolar actiniarians: *Glyphoperidium bursa* Roule, 1909, and *Stomphia selaginella* (Stephenson, 1918). Map 2. Example of deep-sea endemic Antarctic actiniarian: *Stephanthus antarcticus* Rodríguez & López-González, 2003. Map 3. Example of endemic circumpolar Antarctic and sub-Antarctic distributed actiniarian: *Isosicyonis alba* (Studer, 1879). Map 4. Example of cosmopolitan deep-sea species extending to Antarctica: *Galatheanthemum profundale* Carlgren, 1956. Map 5. Example of shallow-water actiniarian distributed both in Antarctica and Southern Patagonia: *Antholoba achates* Drayton in Dana, 1846. Map 6. Antarctic and sub-Antarctic records of Antipatharia.

► Cnidaria: Hexacorallia



Hexacorallia Maps 7-10 Known distribution of ceriantharians, corallimorpharians, scleractinians (selected species), and zoanthideans in the Southern Ocean. Map 7. Antarctic and sub-Antarctic records of Ceriantharia. Map 8. Antarctic and sub-Antarctic records of Corallimorpharia. Map 9. Antarctic and sub-Antarctic records of representative species of order Scleractinia: Caryophyllia antarctica Marenzeller, 1904 (endemic circumpolar Antarctic scleractinian); Flabellum apertum apertum Moseley, 1876 (sub-Antarctic endemic circumpolar scleractinian); Fungiacyathus marenzelleri (Vaughan, 1906) (cosmopolitan scleractinian extending to Antarctica). Map 10. Antarctic and sub-Antarctic records of Zoanthidea.

Rintoul, S., 2007. Antarctic Circumpolar Current. In: Riffenburgh, B. (ed.). Encyclopedia of the

Antarctic. Routledge, 234–239.
Rodríguez, E., López-González, P.J., 2003. Stephanthus antarcticus a new genus and species

of sea anemone (Actiniaria, Haloclavidae) from the South Shetland Islands, Antarctica.
Helgoland Marine Research, 57, 54–62.

Rodríguez, E., López-González, P.J., 2013. New records of Antarctic and Sub-Antarctic sea
anemones (Cnidaria, Anthozoa, Actiniaria and Corrallimorpharia) from the Weddell Sea,
Antarctic Peninsula, and Scotia Arc. Zootaxa, 3624: 1–100.

Rodríguez, E., López-González, P. J., Gili. J.M., 2007. Biogeography of Antarctic sea anemones (Anthozoa, Actiniaria): what do they tell us about the origin of the Antarctic benthic fauna?

Deep-Sea Research II, **54**, 1876–1904. Roule, L., 1909. Hexactinides. *Expédition Antarctique Française (1903–1905)*, 1–19.

Sinniger, F., Häussermann, V., 2009. Zoanthids (Cnidaria: Anthozoa: Zoanthidae) from shallow waters of the southern Chilean fjord region with the description of a new genus and two new species. Organisms Diversity and Evolution, 9, 23-36.

Squires, D.F., 1969. Distribution of selected groups of marine invertebrates in waters south of 35°S

latitude: Scleractinia. Antarctic Map Folio Series, 11, 15–18.

Stephenson, T.A., 1918. Coelenterata. I. Actiniaria. Natural History Reports on British Antarctic ("Terra Nova") Expedition 1910, 5(1), 1–68.

Studer, T., 1879. Zweite Abtheilung der Anthozoa polyactinia, welche während der Reise S.M.S. Corvette Gazelle um die Erde gesammelt wurden. Monatsberichte der könglich preussischen Akademie der Wissenschaften zu Berlin, 25, 524–550.

Thompson, J., 1905. Report on the Antipatharians, Scotia Collection. Proceedings of the Royal Physical Society of Edinburgh, 16, 76–79.

Vaughan, T.W., 1906. Madreporaria. Reports on the scientific results of the expedition to the eastern tropical Pacific, in charge of Alexander Agassiz, by the U. S. Fish Commission Steamer "Albatross," from October, 1904, to March, 1905, lieut. commander L. M. Garrett, U.S.N., commanding. *Bulletin of the Museum of Comparative Zoology* (Harvard University), **50**(3),

THE BIOGEOGRAPHIC ATLAS OF THE SOUTHERN OCEAN

Biogeographic information is of fundamental importance for discovering marine biodiversity hotspots, detecting and understanding impacts of environmental changes, predicting future distributions, monitoring biodiversity, or supporting conservation and sustainable management strategies

The recent extensive exploration and assessment of biodiversity by the Census of Antarctic Marine Life (CAML), and the intense compilation and validation efforts of Southern Ocean biogeographic data by the SCAR Marine Biodiversity Information Network (SCAR-MarBIN / OBIS) provided a unique opportunity to assess and synthesise the current knowledge on Southern

The scope of the Biogeographic Atlas of the Southern Ocean is to present a concise synopsis of the present state of knowledge of the distributional patterns of the major benthic and pelagic taxa and of the key communities, in the light of biotic and abiotic factors operating within an evolutionary framework. Each chapter has been written by the most pertinent experts in their field, relying on vastly improved occurrence datasets from recent decades, as well as on new insights provided by molecular and phylogeographic approaches, and new methods of analysis, visualisation, modelling and prediction of biogeographic distributions.

A dynamic online version of the Biogeographic Atlas will be hosted on www.biodiversity.aq.

The Census of Antarctic Marine Life (CAML)


CAML (www.caml.aq) was a 5-year project that aimed at assessing the nature, distribution and abundance of all living organisms of the Southern Ocean. In this time of environmental change, CAML provided a comprehensive baseline information on the Antarctic marine biodiversity as a sound benchmark against which future change can reliably be assessed. CAML was initiated in 2005 as the regional Antarctic project of the worldwide programme Census of Marine Life (2000-2010) and was the most important biology project of the International Polar Year 2007-2009.

The SCAR Marine Biodiversity Information Network (SCAR-MarBIN)
In close connection with CAML, SCAR-MarBIN (www.scarmarbin.be, integrated into www.biodiversity.aq) compiled and managed the historic, current and new information (i.a. generated by CAML) on Antarctic marine biodiversity by establishing and supporting a distributed system of interoperable databases, forming the Antarctic regional node of the Ocean Biogeographic Information System (OBIS, www.iobis.org), under the aegis of SCAR (Scientific Committee on Antarctic Research, www.scar.org). SCAR-MarBIN established a comprehensive register of Antarctic marine species and, with biodiversity.aq provided free access to more than 2.9 million Antarctic georeferenced biodiversity data, which allowed more than 60 million downloads.

The Editorial Team



Claude DE BROYER is a marine biologist at the Royal Belgian Institute of Natural Sciences in Brussels. His research interests cover structural and ecofunctional biodiversity and biogeography of crustaceans, and polar and deep sea benthic ecology. Active promoter of CAML and ANDEEP, he is the initiator of the SCAR Marine Biodiversity Information Network (SCAR-MarBIN). He took part to 19 polar





Philippe KOUBBI is professor at the University Pierre et Marie Curie (Paris, France) and a specialist in Antarctic fish ecology and biogeography. He is the Principal Investigator of projects supported by IPEV, the French Polar Institute. As a French representative to the CCAMLR Scientific Committee, his main input is on the proposal of Marine Protected Areas. His other field of research is on the ecoregionalisation of the high seas.



Huw GRIFFITHS is a marine Biogeographer at the British Antarctic Survey. He created and manages SOMBASE, the Southern Ocean Mollusc Database. His interests include large-scale biogeographic and ecological patterns in space and time. His focus has been on molluscs, bryozoans, sponges and pycnogonids as model groups to investigate trends at high southern latitudes.



Ben RAYMOND is a computational ecologist and exploratory data analyst, working across a variety of Southern Ocean, Antarctic, and wider research projects. His areas of interest include ecosystem modelling, regionalisation and marine protected area selection, risk assessment, animal tracking, seabird ecology, complex systems, and remote sensed data analyses.



Cédric d'UDEKEM d'ACOZ is a research scientist at the Royal Belgian Institute of Natural Sciences, Brussels. His main research interests are systematics of amphipod crustaceans, especially of polar species and taxonomy of decapod crustaceans. He took part to 2 scientific expeditions to Antarctica on board of the *Polarstern* and to several sampling campaigns in Norway and Svalbard.



Anton VAN DE PUTTE works at the Royal Belgian Institute for Natural Sciences (Brussels, Belgium). He is an expert in the ecology and evolution of Antarctic fish and is currently the Science Officer for the Antarctic Biodiveristy Portal www. biodiversity.aq. This portal provides free and open access to Antarctic Marine and terrestrial biodiversity of the Antarctic and the Southern Ocean.



Bruno DANIS is an Associate Professor at the Université Libre de Bruxelles, where his research focuses on polar biodiversity. Former coordinator of the scarmarbin. be and antabif.be projects, he is a leading member of several international committees, such as OBIS or the SCAR Expert Group on Antarctic Biodiversity Informatics. He has published papers in various fields, including ecotoxicology, physiology, biodiversity informatics, polar biodiversity or information science.



Bruno DAVID is CNRS director of research at the laboratory BIOGÉOSCIENCES, University of Burgundy. His works focus on evolution of living forms, with and more specifically on sea urchins. He authored a book and edited an extensive database on Antarctic echinoids. He is currently President of the scientific council of the Muséum National d'Histoire Naturelle (Paris), and Deputy Director at the CNRS Institute for Ecology and Environment.



Susie GRANT is a marine biogeographer at the British Antarctic Survey. Her work is focused on the design and implementation of marine protected areas, particularly through the use of biogeographic information in systematic conservation planning.



Julian GUTT is a marine ecologist at the Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Bremerhaven, and professor at the Oldenburg University, Germany. He participated in 13 scientific expeditions to the Antarctic and was twice chief scientist on board Polarstern. He is member of the SCAR committees ACCE and AnT-ERA (as chief officer). Main focii of his work are: biodiversity, ecosystem functioning and services, response of marine systems to climate change, non-invasive technologies, and outreach.



Christoph HELD is a Senior Research Scientist at the Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Bremerhaven. He is a specialis in molecular systematics and phylogeography of Antarctic crustaceans, especially



Graham HOSIE is Principal Research Scientist in zooplankton ecology at the Australian Antarctic Division. He founded the SCAR Southern Ocean Continuous Plankton Recorder Survey and is the Chief Officer of the SCAR Life Sciences Standing Scientific Group. His research interests include the ecology and biogeography of plankton species and communities, notably their response to environmental changes. He has participated in 17 marine science voyages to



Falk HUETTMANN is a 'digital naturalist' he works on three poles (Arctic, Anta and Hindu-Kush Himalaya) and elsewhere (marine, terrestrial and atmosphe He is based with the university of Alaska-Fairbank (UAF) and focuses prim on effective conservation questions engaging predictions and open access date.



Alexandra POST is a marine geoscientist, with expertise in benthic habitat mapping, sedimentology and geomorphic characterisation of the seafloor. She has worked at Geoscience Australia since 2002, with a primary focus on understanding seafloor processes and habitats on the East Antarctic margin. Most recently she has led work to understand the biophysical environment beneath the Amery Ice Shelf, and to characterise the habitats on the George V Shelf and slope following the successful CAML voyages in that region.



Yan ROPERT COUDERT spent 10 years at the Japanese National Institute of Polar Research, where he graduated as a Doctor in Polar Sciences in 2001. Since 2007, he is a permanent researcher at the CNRS in France and the director of a polar research programme (since 2011) that examines the ecological response of Adélie penguins to environmental changes. He is also the secretary of the Expert Group on Birds and Marine Mammals and of the Life Science Group of the Scientific Committee on Antarctic Research entific Committee on Antarctic Research























