

Census of Antarctic Marine Life  
SCAR-Marine Biodiversity Information Network

# BIOGEOGRAPHIC ATLAS OF THE SOUTHERN OCEAN

## ► CHAPTER 6.5. SOUTHERN OCEAN SQUID.

Rodhouse P.G.K., Griffiths H.J., Xavier J., 2014.

In: De Broyer C., Koubbi P., Griffiths H.J., Raymond B., Udekem d'Acoz C. d', et al. (eds.). Biogeographic Atlas of the Southern Ocean. Scientific Committee on Antarctic Research, Cambridge, pp. 284-289.

### EDITED BY:

Claude DE BROYER & Philippe KOUBBI (chief editors)

with Huw GRIFFITHS, Ben RAYMOND, Cédric d'UDEKEM d'ACOZ, Anton VAN DE PUTTE, Bruno DANIS, Bruno DAVID, Susie GRANT, Julian GUTT, Christoph HELD, Graham HOSIE, Falk HUETTMANN, Alexandra POST & Yan ROPERT-COUDERT



SCIENTIFIC COMMITTEE ON ANTARCTIC RESEARCH

# THE BIOGEOGRAPHIC ATLAS OF THE SOUTHERN OCEAN

The "Biogeographic Atlas of the Southern Ocean" is a legacy of the International Polar Year 2007-2009 ([www.ipy.org](http://www.ipy.org)) and of the Census of Marine Life 2000-2010 ([www.coml.org](http://www.coml.org)), contributed by the Census of Antarctic Marine Life ([www.caml.aq](http://www.caml.aq)) and the SCAR Marine Biodiversity Information Network ([www.scarmarbin.be](http://www.scarmarbin.be); [www.biodiversity.aq](http://www.biodiversity.aq)).

The "Biogeographic Atlas" is a contribution to the SCAR programmes Ant-ECO (State of the Antarctic Ecosystem) and AnT-ERA (Antarctic Thresholds- Ecosystem Resilience and Adaptation) ([www.scar.org/science-themes/ecosystems](http://www.scar.org/science-themes/ecosystems)).

## Edited by:

Claude De Broyer (Royal Belgian Institute of Natural Sciences, Brussels)  
Philippe Koubbi (Université Pierre et Marie Curie, Paris)  
Huw Griffiths (British Antarctic Survey, Cambridge)  
Ben Raymond (Australian Antarctic Division, Hobart)  
Cédric d'Udekem d'Acoz (Royal Belgian Institute of Natural Sciences, Brussels)  
Anton Van de Putte (Royal Belgian Institute of Natural Sciences, Brussels)  
Bruno Danis (Université Libre de Bruxelles, Brussels)  
Bruno David (Université de Bourgogne, Dijon)  
Susie Grant (British Antarctic Survey, Cambridge)  
Julian Gutt (Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven)  
Christoph Held (Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven)  
Graham Hosie (Australian Antarctic Division, Hobart)  
Falk Huettmann (University of Alaska, Fairbanks)  
Alix Post (Geoscience Australia, Canberra)  
Yan Ropert-Coudert (Institut Pluridisciplinaire Hubert Curien, Strasbourg)

## Published by:

The Scientific Committee on Antarctic Research, Scott Polar Research Institute, Lensfield Road, Cambridge, CB2 1ER, United Kingdom ([www.scar.org](http://www.scar.org)).

## Publication funded by:

- The Census of Marine Life (Albert P. Sloan Foundation, New York)
- The TOTAL Foundation, Paris.

The "Biogeographic Atlas of the Southern Ocean" shared the *Cosmos Prize* awarded to the Census of Marine Life by the International Osaka Expo'90 Commemorative Foundation, Tokyo, Japan.

## Publication supported by:

- The Belgian Science Policy (Belspo), through the Belgian Scientific Research Programme on the Antarctic and the "biodiversity.aq" network (SCAR-MarBIN/ANTABIF)
- The Royal Belgian Institute of Natural Sciences (RBINS), Brussels, Belgium
- The British Antarctic Survey (BAS), Cambridge, United Kingdom
- The Université Pierre et Marie Curie (UPMC), Paris, France
- The Australian Antarctic Division, Hobart, Australia
- The Scientific Steering Committee of CAML, Michael Stoddart (CAML Administrator) and Victoria Wadley (CAML Project Manager)

**Mapping coordination and design:** Huw Griffiths (BAS, Cambridge) & Anton Van de Putte (RBINS, Brussels)

**Editorial assistance:** Henri Robert, Xavier Loréa, Charlotte Havermans, Nicole Moortgat (RBINS, Brussels)

**Printed by:** Altitude Design, Rue Saint Josse, 15, B-1210 Brussels, Belgium ([www.altitude-design.be](http://www.altitude-design.be))

**Lay out:** Sigrid Camus & Amélie Blaton (Altitude Design, Brussels).

**Cover design:** Amélie Blaton (Altitude Design, Brussels) and the Editorial Team.

**Cover pictures:** amphipod crustacean (*Epimeria rubrieques* De Broyer & Klages, 1991), image © T. Riehl, University of Hamburg; krill (*Euphausia superba* Dana, 1850), image © V. Siegel, Institute of Sea Fisheries, Hamburg; fish (*Chaenocephalus* sp.), image © C. d'Udekem d'Acoz, RBINS; emperor penguin (*Aptenodytes forsteri* G.R. Gray, 1844), image © C. d'Udekem d'Acoz, RBINS; Humpback whale (*Megaptera novaeangliae* (Borowski, 1781)), image © L. Kindermann, AWI.

## Online dynamic version :

A dynamic online version of the Biogeographic Atlas is available on the SCAR-MarBIN / AntaBIF portal : [atlas.biodiversity.aq](http://atlas.biodiversity.aq).

## Recommended citation:

### For the volume:

De Broyer C., Koubbi P., Griffiths H.J., Raymond B., Udekem d'Acoz C. d', Van de Putte A.P., Danis B., David B., Grant S., Gutt J., Held C., Hosie G., Huettmann F., Post A., Ropert-Coudert Y. (eds.), 2014. Biogeographic Atlas of the Southern Ocean. Scientific Committee on Antarctic Research, Cambridge, XII + 498 pp.

### For individual chapter:

(e.g.) Crame A., 2014. Chapter 3.1. Evolutionary Setting. In: De Broyer C., Koubbi P., Griffiths H.J., Raymond B., Udekem d'Acoz C. d', *et al.* (eds.). Biogeographic Atlas of the Southern Ocean. Scientific Committee on Antarctic Research, Cambridge, pp. xx-yy.

ISBN: 978-0-948277-28-3.



This publication is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License

## 6.5. Southern Ocean Squid

**Paul G.K. Rodhouse, Huw J. Griffiths & José C. Xavier**

British Antarctic Survey, Cambridge, UK

### 1. Introduction

Data on Antarctic cephalopods have accumulated since the expeditions of *HMS Challenger* and *HMS Alert* in the 19<sup>th</sup> century and the heroic era of Antarctic exploration in the early years of the 20<sup>th</sup> century. After the 1920s there was little activity in the field until the 1960s when several nations started to send research ships to the Southern Ocean for biological research. Antarctic teuthology has since grown and matured and has been the subject of a symposium and four reviews (Clarke & Okutani 1985, Rodhouse *et al.* 1994, Filippova 2002, Collins & Rodhouse 2006).

Much of our knowledge of squid biogeography in the Antarctic and worldwide derives from the work of the Russian scientist Kir Nesis who died in 2003. His world survey of cephalopods (Abridged key to the cephalopod molluscs of the world) was published in Russian in 1982 and later in English with new material (Nesis 1987). This secondary source of information has been used here where the original papers in Russian, many by Nesis himself, are difficult to access. Xavier *et al.* (1999) published maps of all known and available records of squid captured in the Southern Ocean and, with subsequent additions, these form the basis of the maps published here.

The most sampled areas in the Southern Ocean are in the region of the larger peri-Antarctic islands, the Antarctic Peninsula and the Prydz Bay area. The least sampled region lies in the Pacific sector, roughly between 180° and 90°W.

The Southern Ocean squids (order Teuthida) are restricted to the suborder Oegopsida. The suborder Myopsida is absent from the Antarctic and Sub-Antarctic regions but one species, *Doryteuthis gahi*, occurs over the southern Patagonian Shelf. Of the octopuses (order Octopoda) both suborders Cirrata and Incirrata are represented. The nautiluses (Subclass Nautiloidea), the cuttlefish (Order Sepiida), the sepiolids (order Sepiolida) and the vampire squid (Order Vampyromorpha) are all absent.

The following southern hemisphere species, which have been included in lists of Antarctic cephalopods elsewhere, are not included here as there is little evidence that they are found south of the Sub-Tropical Front: *Promachoteuthis* sp., *Pholidoteuthis massyae*, *Octopoteuthis rugosa*, *Taningia danae*, *Histioteuthis macrohista* and *H. miranda*. A single specimen of the cranchiid squid *Taonius pavo* (Rodhouse, 1990) has also not been included.



Photo 1 *Martialia hyadesi* Rochebrune & Mabille, 1889. Image: © CP. Rodhouse.

### 2. Biogeography and depth distribution

There are some nineteen species of squid known to inhabit Antarctic and sub-Antarctic waters. Their latitudinal range can be divided into six categories from high Antarctic endemics to cosmopolitan species whose range extends into Antarctic waters. This is a loose categorisation for several reasons: the surface signatures of the fronts are often not well defined, their position vary and they do not reflect the deep structure of the water masses of the Southern Ocean. Furthermore, the interfaces between water masses probably do not form impenetrable barriers to squid dispersal.

#### 2.1. Antarctic endemics extending north to the Antarctic Polar Front (APF)

*Moroteuthis knipovitchi* Filippova, 1972 (Map 1): one of four members of the family Onychoteuthidae known in the Southern Ocean; maximum ML 450 mm; circumpolar; mesopelagic and near bottom. The species occurs north of the APF at South Georgia, Kerguelen, Crozet and Prince Edward Islands. Literature: Filippova & Yukhov (1979), Rodhouse (1988, 1989), Rodhouse *et al.* (1996), Piatkowski *et al.* (1998).

*Alluroteuthis antarcticus* Odhner, 1923 (Map 2): sole member of the family Neoteuthidae known in the Southern Ocean; maximum ML 270 mm;

circumpolar; mesopelagic. Literature: Filippova & Yukhov (1982), Rodhouse (1988, 1989); Piatkowski *et al.* (1994), Jackson *et al.* (2002).

*Mastigoteuthis psychrophila* Nesis, 1977 (Map 3): single member of the family Mastigoteuthidae known in the Southern Ocean; maximum ML 180 mm; circumpolar; mesopelagic and bathypelagic. Literature: Rodhouse (1990), Lu & Williams (1994a), Piatkowski *et al.* (1994), Rodhouse *et al.* (1996).

*Mesonychoteuthis hamiltoni* Robson, 1925 (Map 4): one of two members of the family Cranchiidae known in the Southern Ocean; maximum ML 2500 mm; circumpolar; bathypelagic and near bottom; young specimens have been recorded north of the APF. Large adults have only been caught south of the APF. They are an occasional by-catch in the South Georgia long-line fishery for Patagonian toothfish (*Dissostichus eleginoides*) and the Ross Sea fishery for Antarctic toothfish (*Dissostichus mawsoni*). Literature: Filippova & Yukhov (1979), Rodhouse & Clarke (1985), Jackson *et al.* (2002), Filippova (2002), Collins *et al.* (2010), Roberts *et al.* (2011).

#### 2.2. Antarctic endemics extending north to the Sub-Antarctic Front (SAF)

*Kondakovia longimana* Filippova, 1972 (Map 5): one of four members of the family Onychoteuthidae known in the Southern Ocean; maximum ML 1100 mm; circumpolar; epipelagic, mesopelagic, bathypelagic and near bottom. Literature: Rodhouse (1990), Lu & Williams (1994a, b), Vacchi *et al.* (1994), Lynnes & Rodhouse (2002).

*Psychroteuthis glacialis* Thiele, 1920 (Map 6): the only species of the family Psychroteuthidae described; maximum ML 440 mm; circumpolar; mesopelagic, bathypelagic, near bottom. Literature: Filippova (1972), Filippova & Yukhov (1979), Kubodera (1989), Rodhouse (1989), Piatkowski *et al.* (1990, 1994, 1998), Lu & Williams (1994a), Gröger *et al.* (2000), Anderson & Rodhouse (2002), Collins *et al.* (2004).

*Slosarczykvia circumantarctica* Lipinski, 2001 (Map 7): the only member of the family Brachioteuthidae found in Antarctic waters apart from *Brachioteuthis linkovskii* Lipinski, 2001 which is occasionally found in the Sub-Antarctic; maximum ML 90 mm; circumpolar; epipelagic, mesopelagic and bathypelagic. Literature: Kubodera (1989), Rodhouse (1989, 1990), Rodhouse *et al.* (1996), Piatkowski *et al.* (1994), Anderson & Rodhouse (2002), Collins *et al.* (2004) (referred to as notalian-Antarctic *B. riisei* by Nesis (1987) and as *B. sp.* and *B. ?picta* by other authors prior to Lipinski (2001)).

#### 2.3. Antarctic endemics extending north to the Sub-Tropical Front (STF)

*Galiteuthis glacialis* Chun, 1906 (Map 8): one of two members of the family Cranchiidae known in the Southern Ocean; maximum ML 330 mm; circumpolar; may occur north of the APF when Antarctic intermediate water moves north; bathypelagic and mesopelagic to lower epipelagic; Antarctic squid most commonly caught in research nets. Literature: Dell (1959), Filippova (1972), Lu & Mangold (1978), Kubodera & Okutani (1986), Rodhouse & Clarke (1986), Rodhouse (1989), Lu & Williams (1994a), Piatkowski & Hagen (1994), Rodhouse *et al.* (1996), Nesis *et al.* (1998), Piatkowski *et al.* (1994, 1998), Anderson & Rodhouse (2002), Jackson *et al.* (2002), Van de Putte *et al.* (2010).

*Parateuthis tunicata* Thiele, 1921 (Map 9): family *incertae sedis*: only larvae (<8 mm ML) known from deep water (>2000 m). Literature: Nesis (1987).


*Batoteuthis skolops* Young and Roper, 1968 (Map 10): the only known species of the family Batoteuthidae; circumpolar; maximum ML 95 mm; mesopelagic and bathypelagic. Literature: Filippova & Yukhov (1979), Rodhouse *et al.* (1992, 1996), Anderson & Rodhouse (2002), Jackson *et al.* (2002), Collins *et al.* (2004).

*Histioteuthis eltaninae* N. Voss, 1969 (Map 11): one of two members of the family Histioteuthidae known in the Southern Ocean; maximum ML 105 mm; circumpolar (more abundant north of the APF); mesopelagic and bathypelagic. Literature: Lu & Mangold (1978), Alexeyev (1994), Piatkowski *et al.* (1994), Rodhouse *et al.* (1996), Voss *et al.* (1998), Jackson *et al.* (2002).


#### 2.4. Sub-Antarctic (APF – STF)

*Moroteuthis ingens* Smith, 1881 (Map 12): one of four members of the family Onychoteuthidae known in the Southern Ocean; circumpolar; maximum ML 940 mm; near seabed from shelf to bathyal. Literature: Massy (1916), Filippova (1972), Filippova & Yukhov (1979), Alexeyev (1994).


*Gonatus antarcticus* Lönnberg, 1898 (Map 13): single member of the family Gonatidae described from the Southern Ocean; maximum ML 350 mm; circumpolar (extends south of the APF in the Scotia Sea); mesopelagic, bathypelagic. Literature: Kubodera & Okutani (1986), Rodhouse (1990), Rodhouse *et al.* (1996), Nesis (1999), Anderson & Rodhouse (2002).




**Map 1**  
● *Moroteuthis knipovitchi*




**Map 2**  
● *Alluroteuthis antarcticus*




**Map 3**  
● *Mastigoteuthis psychrophila*



**Map 4**  
● *Mesonychoteuthis hamiltoni*




**Map 5**  
● *Kondakovia longimana*




**Map 6**  
● *Psychroteuthis glacialis*

**Cephalopoda Decapodiformes Maps 1–6** Map 1. *Moroteuthis knipovitchi*. Map 2. *Alluroteuthis antarcticus*. Map 3. *Mastigoteuthis psychrophila*. Map 4. *Mesonychoteuthis hamiltoni*. Map 5. *Kondakovia longimana*. Map 6. *Psychroteuthis glacialis*. On all maps fronts shown from south to north are: Antarctic Polar Front, Sub-Antarctic Front and Sub-Tropical Front


► Mollusca : Cephalopoda




Map 7  
● *Slosarczykvia circumantarctica*




Map 8  
● *Batoteuthis skolops*




Map 9  
● *Galiteuthis glacialis*



Map 10  
● *Parateuthis tunicata*



Map 11  
● *Histioteuthis eltaninae*



Map 12  
● *Moroteuthis ingens*

Cephalopoda Decapodiformes Maps 7–12 Map 7. *Slosarczykvia circumantarctica*. Map 8. *Batoteuthis skolops*. Map 9. *Galiteuthis glacialis*. Map 10. *Parateuthis tunicata*. Map 11. *Histioteuthis eltaninae*. Map 12. *Moroteuthis ingens*.

*Martialia hyadesi* Rochebrune & Mabille, 1887 (Map 14): one of two members of the family Ommastrephidae from the Southern Ocean; maximum ML 500 mm; circumpolar; epipelagic, mesopelagic and near the bottom from continental shelves to near-bathyal depths. Literature: O'Sullivan *et al.* (1983), Rodhouse & Yeatman (1990), Rodhouse (1991), Piatkowski *et al.* (1991), Uozomi *et al.* (1991), Alexeyev (1994), Rodhouse *et al.* (1996), Gonzalez *et al.* (1997).

## 2.5. Southern hemisphere extending south to the APF

*Moroteuthis robsoni* Adam, 1962 (Map 15): one of four members of the family Onychoteuthidae known in the Southern Ocean; maximum ML to 750 mm; circumpolar (extends South of APF in Scotia Sea - South Georgia); pelagic and near bottom. Literature: Nesis (1987).

*Histioteuthis atlantica* Hoyle, 1885 (Map 16): one of two members of the family Histiotheuthidae known in the Southern Ocean; maximum ML 258 mm; circumpolar; mesopelagic, bathypelagic, abyssopelagic. Literature: Kubodera (1989), Alexeyev (1994), Voss *et al.* (1998), Jackson *et al.* (2002).

*Todarodes filippovae* Adam, 1975 (Map 17): one of two members of the family Ommastrephidae from the Southern Ocean; maximum ML >500 mm; circumpolar; epipelagic, mesopelagic, near seabed on continental slope. Literature: Kawakami (1976), Dunning & Brandt (1985), Dunning (1988, 1993), Roeleveld (1989), Piatkowski *et al.* (1991), Roeleveld *et al.* (1992), Alexeyev (1994), Yokawa (1994), Rodhouse (1998).

## 2.6. Cosmopolitan extending south of the APF

*Bathyteuthis abyssicola* Hoyle, 1885 (Map 18): single member of the family Bathytethidae known to extend into the Southern Ocean; maximum ML 70 mm; circumpolar; mesopelagic, epipelagic, juveniles lower epipelagic. Literature: Hoyle (1912), Odhner (1923), Roper (1969), Lu & Mangold (1978), Lu & Williams (1994a), Piatkowski *et al.* (1994, 1998). Rodhouse *et al.* (1996).

*Chiroteuthis veranyi* Ferussac, 1825 (Map 19): single member of the family Chiroteuthidae known to extend into the Southern Ocean; maximum ML 107 mm; circumpolar; bathyal, bathypelagic and mesopelagic. Literature: Alexeyev (1994), Rodhouse & Lu (1998).

## 3. Squid data from higher predator gut contents

Squid remain difficult to catch with scientific sampling gear and biogeographic knowledge of most species is patchy so conclusions about distribution are tentative. Gaps in data from nets are to some extent filled by information from studies on gut contents of higher predators. The beaks of squid are indigestible so they accumulate in predator stomachs and can be identified and used to estimate the biomass they represent (Clarke 1962). This approach has been particularly valuable in the Antarctic where an estimate has been made of total cephalopod biomass consumed by predators, especially sperm whales (Clarke 1980, 1983). Subsequent research has provided detailed information about the prey of all the major Antarctic cephalopod predators, including fish (Collins & Rodhouse 2006). The research field remains active and a project within the Census of Antarctic Marine Life Programme has resulted in the publication of an identification guide to the beaks of Southern Ocean cephalopods (Xavier & Cherel 2009). Although in this CAML Atlas predator data have been used to infer distribution of squid species in the text the distribution maps are solely based on specimens caught with scientific sampling gear or commercial fishing gear.

## 4. Inferring distribution

All the species listed here are probably circumpolar. Nevertheless, for most species there are large gaps in known distribution — there are areas that have not been sampled by nets and where predator data are scarce or absent. Productivity is low in large regions in the Antarctic Polar Current system where iron, a micronutrient, is at low concentrations despite fertile levels of macronutrients (de Baar *et al.* 1995). Cephalopods are voracious predators and have high metabolic rates so they need to forage in areas of high food supply and they are known to concentrate in the vicinity of islands, seamounts and mesoscale oceanographic features where productivity is elevated by nutrients off the land and the surrounding shelf or upwelled from the depths (Wadley 1983, Rogers 1994, Rodhouse *et al.* 1996). Analysis of remotely sensed sea surface temperature and ocean colour imagery should provide a reliable indication of the distribution of squid in the Southern Ocean.

Latitudinal distributions of all the squid species included here seem to be limited by the frontal systems of the Southern Ocean to the extent that it has been possible to classify them in terms of the frontal limits of their distribution. Nevertheless the conclusions drawn here are tentative. Sampling has been very unevenly spread and some species such as *Galiteuthis glacialis* have been observed to cross apparent frontal boundaries with incursions of large volumes of water (Nesis 1987). Other deep living species may extend northwards beyond and beneath the frontal boundaries given here. The so-called colossal squid *Mesonychoteuthis hamiltoni*, a deep living species, has been found in the gut contents of sperm whales as far north as southern Africa and Australia (Clarke 1980, Filippova 2002). It is possible that the whales had taken them further south but it is equally conceivable that the squid extends northwards in Antarctic deep water. Given the extremely limited sampling with gear suitable for catching large squid at great depth the question remains open.

## 5. Fisheries

There are commercial fisheries for squid on the Patagonian Shelf and slope bordering the Falkland Current, in the Humboldt Current and around New Zealand. Over the last 20 years there have been occasional exploratory fishing expeditions to the South Atlantic sector of the Southern Ocean to explore stocks of the ommastrephid squid *Martialia hyadesi* which is taken as a by-catch in the *Illex argentinus* fishery in the South Atlantic (Gonzalez & Rodhouse 1997, Dickson *et al.* 2004). To date no commercial fishery has developed in the Southern Ocean but catch data, combined with data from predators have been used to set precautionary measures in the event that a fishery might develop in the future (Rodhouse 1997).

## Acknowledgements


This is CAML contribution # 134.

## References


- Alexeyev, D.O., 1994. New data on the distribution and biology of squids from the Southern Pacific. *Ruthenica*, **4**, 151–166.
- Anderson, C.I.H., Rodhouse, P.G., 2002. Distribution of juvenile squid in the Scotia Sea in relation to regional oceanography. *Bulletin of Marine Science*, **71**, 97–108.
- Clarke, M.R., 1962. The identification of cephalopod 'beaks' and the relationship between beak size and total body weight. *Bulletin of the British Museum of Natural History*, **8**, 419–480.
- Clarke, M.R., 1980. Cephalopoda in the diet of sperm whales of the southern hemisphere and their bearing on sperm whale biology. *Discovery Reports*, **37**, 1–324.
- Clarke, M.R., 1983. Cephalopod biomass – estimation from predation. *Memoirs of the National Museum of Victoria*, **44**, 95–107.
- Clarke, M.R., Okutani, T., 1985. Identification key and species description for Antarctic squids. SCAR BIOMASS Handbook, **21**, 1–57.
- Clarke, M.R., 1986. A handbook for the identification of cephalopod beaks. Clarendon Press, Oxford, UK, 273 pp.
- Collins, M.A., Allcock, A.L., Belchier, M., 2004. Cephalopods of the South Georgia slope. *Journal of the Marine Biological Association of the United Kingdom*, **84**, 415–419.
- Collins, M.A., Rodhouse, P.G., 2006. Southern Ocean cephalopods. *Advances in Marine Biology*, **50**, 193–265.
- Collins, M.A., Brickle, P., Brown, J., Belchier, M., 2010. The Patagonian toothfish: biology, ecology and fishery. *Advances in Marine Biology*, **58**, 229–289. doi:10.1016/B978-0-12-381015-1.00004-6.
- de Baar, H.J.W., de Jong, J.T.M., Bakker, D.C.E., Löscher, B.M., Veth, C., Bathmann, U., Smetacek, V., 1995. Importance of iron for plankton blooms and carbon dioxide drawdown in the Southern Ocean. *Nature*, **373**, 412–415.
- Dell, R.K., 1959. Cephalopoda. *British Australian and New Zealand Antarctic Expedition*, **8**, 89–106.
- Dickson, J., Morley, S.A., Mulvey, T., 2004. New data on *Martialia hyadesi* feeding in the Scotia Sea during winter with emphasis on seasonal and annual variability. *Journal of the Marine Biological Association of the United Kingdom*, **84**, 785–788.
- Dunning, M.C., 1993. Summer populations of *Ommastrephes bartrami* (Lesueur 1821) and *Todarodes filippovae* Adam, 1975 (Cephalopoda: Ommastrephidae) from the Tasman Sea. In: Okutani, T., O'Dor, R.K., Kubodera, T. (eds.). *Recent Advances in Cephalopod Fisheries Biology*, Tokai University Press, Tokyo, pp. 97–118.
- Dunning, M., Brandt, S.R., 1985. Distribution and life history of deep-water squid of commercial interest from Australia. *Australian Journal of Marine and Freshwater Research*, **36**, 343–359.
- Filippova, J.A., 1972. New data on the squids (Cephalopoda: Oegopsida) from the Scotia Sea (Antarctic). *Malacologia*, **11**, 391–406.
- Filippova, J.A., 2002. Review of Soviet/Russian studies on squids in the Antarctic Ocean. *Bulletin of Marine Science*, **71**, 255–267.
- Filippova, J.A., Yukhov, V.L., 1979. Specific composition and distribution of cephalopod molluscs in meso- and bathypelagic Antarctic waters. *Antarktika Doklady Komissi*, **18**, 175–187.
- Filippova, J.A., Yukhov, V.L., 1982. New data on the genus *Alluroteuthis* Odhner, 1923 (Cephalopoda: Oegopsida). *Antarktika Doklady Komissi*, **21**, 157–168.
- Gonzalez, A.F., Trathan, P.N., Yau, C., Rodhouse, P.G., 1997. Interactions between oceanography, ecology and fishery biology of the ommastrephid squid *Martialia hyadesi* in the South Atlantic. *Marine Ecology Progress Series*, **152**, 205–215.
- González, A.F., Rodhouse, P.G., 1998. Fishery biology of the seven star flying squid *Martialia hyadesi* at South Georgia during winter. *Polar Biology*, **19**, 231–236.
- Gröger, J., Piatkowski, U., Heinemann, H., 2000. Beak length analysis of the Southern Ocean squid *Psychroteuthis glacialis* (Cephalopoda: Psychroteuthidae) and its use for size and biomass estimation. *Polar Biology*, **23**, 70–74.
- Hoyle, W.E., 1912. The Cephalopoda of the Scottish National Antarctic Expedition. *Transactions of the Royal Society of Edinburgh*, **48**, 272–283.
- Jackson, G.D., Finn, J., Nicol, S., 2002. Planktonic cephalopods collected off East Antarctica during the 'BROKE' survey. *Deep-Sea Research I*, **49**, 1049–1054.
- Kawakami, T., 1976. The fishing biological study on a squid *Nototodarus sloani sloani* (Gray) in the New Zealand waters. *Bulletin of the Tokai Regional Fisheries Research Laboratory*, **85**, 31–104.
- Kubodera, T., 1989. Young squids collected with a 10-foot IKPT net during the Jare-28 cruise, 1987. *Proceedings of the NIPR Symposium on Polar Biology*, **2**, 71–77.
- Kubodera, T., Okutani, T., 1986. New and rare cephalopods from the Antarctic waters. *Memoirs of the National Institute of Polar Research Special Edition*, **44**, 129–143.
- Lipinski, M.R., 2001. Preliminary description of two new species of Cephalopods (Cephalopoda: Brachioctephidae) from the South Atlantic and Antarctic waters. *Bulletin of the Sea Fisheries Institute*, **1**, 3–14.
- Lu, C.C., Mangold, K., 1978. Cephalopods of the Kerguelan Province of the Indian Ocean. In: Proceedings of the International Symposium on Marine Biogeography and Evolution in the Southern Hemisphere, Auckland, NZ, pp. 567–574.
- Lu, C.C., Williams, R., 1994a. Contribution to the biology of squid in the Prydz Bay region, Antarctica. *Antarctic Science*, **62**, 223–229.
- Lu, C.C., Williams, R., 1994b. *Kondakovia longimana* Filippova, 1972 (Cephalopoda: Onychoteuthidae) from the Indian Ocean sector of the Southern Ocean. *Antarctic Science*, **6**, 231–234.
- Lynnes, A.S., Rodhouse, P.G., 2002. A big mouthful for predators: The largest recorded specimen of *Kondakovia longimana* (Cephalopoda: Onychoteuthidae). *Bulletin of Marine Science*, **71**, 1087–1090.
- Massy, A.L., 1916. Mollusca. *British Antarctic ("Terra Nova") Expedition*, 1910. *Natural History Report, Zoology*, **2**, 141–176.
- Nesis, K.N., 1987. *Cephalopods of the world*. T.N.P publications, New York and London, 351 pp.
- Nesis, K.N., 1999. Horizontal and vertical distribution and some features of biology of the gonatid squid *Gonatus antarcticus* Lönnberg 1898 (Cephalopoda). *Ruthenica*, **9**, 129–139.
- Nesis, K.N., Nigmatullin, C.M., Nikitina, I.V., 1998. Spent females of deepwater squid *Galiteuthis glacialis* under the ice at the surface of the Weddell Sea (Antarctic). *Journal of Zoology*, **244**, 185–200.
- Odhner, N.H., 1923. Die Cephalopoden. *Further zoological results from the Swedish Antarctic Expedition 1901–1903*, **1**, 1–7.




► Mollusca : Cephalopoda




Map 13  
● *Gonatus antarcticus*




Map 14  
● *Martialia hyadesi*




Map 15  
● *Moroteuthis robsoni*



Map 16  
● *Histioteuthis atlantica*




Map 17  
● *Todarodes filippovae*



Map 18  
● *Bathyteuthis abyssicola*

Cephalopoda Decapodiformes Maps 13–18 Map 13. *Gonatus antarcticus*. Map 14. *Martialia hyadesi*. Map 15. *Moroteuthis robsoni*. Map 16. *Histioteuthis atlantica*. Map 17. *Todarodes filippovae*. Map 18. *Bathyteuthis abyssicola*.



**Map 19**  
● *Chiroteuthis veranyi*

**Cephalopoda Decapodiformes Map 19 *Chiroteuthis veranyi*.**

- O'Sullivan, D.B., Johnstone, G.W., Kerry, K.R., Imber, M.J., 1983. A mass stranding of squid *Martialia hyadesi* Rochebrunne & Mabile (Teuthoidea: Ommastrephidae) at Macquarie Island. *Papers and Proceedings of the Royal Society of Tasmania*, **117**, 161–163.
- Piatkowski, U., Hagen, W., 1994. Distribution and lipid composition of early life stages of the cranchiid squid *Galiteuthis glacialis* (Chun) in the Weddell Sea, Antarctica. *Antarctic Science*, **6**, 235–239.
- Piatkowski, U., Allcock, A.L., Hevia, M., Steimer, S., Vecchione, M., 1998. Cephalopod ecology. The expedition ANTARKTIS XIV/2 of RV *Polarstern* in 1996/1997. *Berichte zur Polarforschung*, **274**, 41–47.
- Piatkowski, U., Rodhouse, P.G., Duhamel, G., 1991. Occurrence of the cephalopod *Martialia hyadesi* (Teuthoidea, Ommastrephidae) at the Kerguelen Islands in the Indian Ocean sector of the Southern Ocean. *Polar Biology*, **11**, 273–275.
- Piatkowski, U., Rodhouse, P.G., White, M.G., Bone, D.G., Symon, C., 1994. Nekton community of the Scotia Sea as sampled by the RMT 25 during austral summer. *Marine Ecology Progress Series*, **112**, 13–28.
- Piatkowski, U., White, M.G., Dimmler, W., 1990. Micronekton of the Weddell Sea: distribution and abundance. *Berichte zur Polarforschung*, **68**, 73–81.
- Roberts, J., Xavier, J.C., Agnew, D.J., 2011. The diet of toothfish species *Dissostichus eleginoides* and *Dissostichus mawsoni* with overlapping distributions. *Journal of Fish Biology*, **79**(1), 138–154. doi:10.1111/j.1095-8649.2011.03005.x.
- Rogers, A.D., 1994. The biology of seamounts. *Advances in Marine Biology*, **30**, 305–350.
- Rodhouse, P.G., 1988. Distribution of the neoteuthid squid *Alluroteuthis antarcticus* Odhner in the Atlantic sector of the Southern Ocean. *Malacologia*, **29**, 267–274.

- Rodhouse, P.G., 1989. Pelagic cephalopods caught by nets during the Antarctic research cruises of the *Polarstern* and *Walther Herwig*, 1985–1987. *Archiv für Fischereiwissenschaft*, **39**, 111–121.
- Rodhouse, P.G., 1990. Cephalopod fauna of the Scotia Sea at South Georgia: Potential for commercial exploitation and possible consequences. In: Kerry, K.R., Hempel, G. (eds.). *Antarctic Ecosystems Ecological Change and Conservation*, Springer-Verlag, Berlin, Heidelberg, pp. 289–298.
- Rodhouse, P.G., 1991. Population structure of *Martialia hyadesi* (Cephalopoda: Ommastrephidae) at the Antarctic Polar Front and the Patagonian shelf, South Atlantic. *Bulletin of Marine Science*, **49**, 404–418.
- Rodhouse, P.G., 1997. Precautionary measures for a new *Martialia hyadesi* (Cephalopoda, Ommastrephidae) fishery in the Scotia Sea: an ecological approach. *CCAMLR Science*, **4**, 125–139.
- Rodhouse, P.G., 1998. *Todarodes filippovae* in the Southern Ocean: an appraisal for exploitation and management. In: Okutani, T. (ed.). *Large pelagic squids*, Japan Marine Fishery Resources Research Centre, Tokyo, pp. 205–217.
- Rodhouse, P.G., Clarke, M.R., 1985. Growth and distribution of young *Mesonychoteuthis hamiltoni* Robson (Mollusca: Cephalopoda): an Antarctic squid. *Vie et Milieu*, **35**, 223–230.
- Rodhouse, P.G., Clarke, M.R., 1986. Distribution of the early-life phase of the Antarctic squid *Galiteuthis glacialis* in relation to the hydrology of the Southern Ocean in the sector 15° to 30°E. *Marine Biology*, **91**, 353–357.
- Rodhouse, P.G., Prince, P.A., Trathan, P.N., Hatfield, E.M.C., Watkins, J.L., Bone, D.G., Murphy, E.J., White, M.G., 1996. Cephalopods and mesoscale oceanography at the Antarctic Polar Front - satellite tracked predators locate pelagic trophic interactions. *Marine Ecology Progress Series*, **136**, 37–50.
- Rodhouse, P.G., Lu, C.C., 1998. *Chiroteuthis veranyi* from the Atlantic sector of the Southern Ocean (Cephalopoda: Chiroteuthidae). *South African Journal of Marine Science*, **20**, 311–322.
- Rodhouse, P.G., Piatkowski, U., Lu, C.C. (eds.), 1994. Southern Ocean cephalopods: life cycles and populations. *Antarctic Science*, **6**(2), 135–291 (special issue).
- Rodhouse, P.G., Symon, C., Hatfield, E.M.C., 1992. Early life-cycle of cephalopods in relation to the major oceanographic features of the Southwest Atlantic Ocean. *Marine Ecology Progress Series*, **89**, 183–195.
- Rodhouse, P.G., Yeatman, J., 1990. Redescription of *Martialia hyadesi* Rochebrune and Mabile, 1889 (Mollusca: Cephalopoda) from the Southern Ocean. *Bulletin of the British Museum of Natural History (Zoology)*, **56**, 135–143.
- Roeleveld, M.A., 1989. The occurrence of two species of squid *Todarodes* in the Benguela system. *Suid-Afrikaanse Tydskrif vir Wetenskap*, **85**, 659–653.
- Roeleveld, M.A.C., Lipinski, M.R., Augustyn, C.J., Stewart, B.A., 1992. The distribution and abundance of cephalopods on the continental shelf of the eastern South Atlantic. *South African Journal of Marine Science*, **12**, 739–752.
- Roper, C.F.E., 1969. Systematics and zoogeography of the worldwide bathypelagic squid *Bathyteuthis* (Cephalopoda: Oegopsida). *United States National Museum Bulletin*, **291**, 1–210.
- Uozomi, Y., Forch, E.C., Okazaki, T., 1991. Distribution and morphological characters of immature *Martialia hyadesi* (Cephalopoda: Oegopsida) in New Zealand waters. *New Zealand Journal of Marine and Freshwater Research*, **25**, 275–282.
- Vacchi, M., Greco, S., Lamesa, M., 1994. *Kondakovia longimana* Filippova, 1972 (Onycotheuthidae) from Terra-Nova Bay, Ross Sea. *Antarctic Science*, **6**, 283–283.
- Van de Putte, A.P., Jackson, G.D., Pakhomov, E., Flores, H., Volckaert, F.A.M., 2010. Distribution of squid and fish in the pelagic zone of the Cosmonaut Sea and Prydz Bay region during the BROKE-West campaign. *Deep-Sea Research II*, **57**, 956–967.
- Voss, N.A., Nesis, K.N., Rodhouse, P.G., 1998. The cephalopod family Histiotheuthidae (Oegopsida): systematics, biology and biogeography. *Smithsonian Contributions to Zoology*, **586**, 293–372.
- Xavier, J.C., Rodhouse, P.G., Trathan, P.N., Wood, A.G., 1999. A geographical information system (GIS) atlas of squid distribution in the Southern Ocean. *Antarctic Science*, **11**, 61–62.
- Xavier, J.C., Cherel, Y., 2009. *Cephalopod beak guide for the Southern Ocean*. British Antarctic Survey, Cambridge, UK, 129 pp.
- Yatsu, A., 1995. The role of the slender tuna, *Allothunnus fallai*, in the pelagic ecosystems of the South Pacific Ocean. *Japanese Journal of Ichthyology*, **41**, 367–377.
- Yokawa, K., 1994. Allozyme differentiation of sixteen species of ommastrephid squid (Mollusca, Cephalopoda). *Antarctic Science*, **6**, 201–204.



# THE BIOGEOGRAPHIC ATLAS OF THE SOUTHERN OCEAN

## Scope

Biogeographic information is of fundamental importance for discovering marine biodiversity hotspots, detecting and understanding impacts of environmental changes, predicting future distributions, monitoring biodiversity, or supporting conservation and sustainable management strategies.

The recent extensive exploration and assessment of biodiversity by the Census of Antarctic Marine Life (CAML), and the intense compilation and validation efforts of Southern Ocean biogeographic data by the SCAR Marine Biodiversity Information Network (SCAR-MarBIN / OBIS) provided a unique opportunity to assess and synthesise the current knowledge on Southern Ocean biogeography.

The scope of the Biogeographic Atlas of the Southern Ocean is to present a concise synopsis of the present state of knowledge of the distributional patterns of the major benthic and pelagic taxa and of the key communities, in the light of biotic and abiotic factors operating within an evolutionary framework. Each chapter has been written by the most pertinent experts in their field, relying on vastly improved occurrence datasets from recent decades, as well as on new insights provided by molecular and phylogeographic approaches, and new methods of analysis, visualisation, modelling and prediction of biogeographic distributions.

A dynamic online version of the Biogeographic Atlas will be hosted on [www.biodiversity.aq](http://www.biodiversity.aq).

## The Census of Antarctic Marine Life (CAML)

CAML ([www.caml.aq](http://www.caml.aq)) was a 5-year project that aimed at assessing the nature, distribution and abundance of all living organisms of the Southern Ocean. In this time of environmental change, CAML provided a comprehensive baseline information on the Antarctic marine biodiversity as a sound benchmark against which future change can reliably be assessed. CAML was initiated in 2005 as the regional Antarctic project of the worldwide programme Census of Marine Life (2000-2010) and was the most important biology project of the International Polar Year 2007-2009.

## The SCAR Marine Biodiversity Information Network (SCAR-MarBIN)

In close connection with CAML, SCAR-MarBIN ([www.scarmarbin.be](http://www.scarmarbin.be), integrated into [www.biodiversity.aq](http://www.biodiversity.aq)) compiled and managed the historic, current and new information (i.a. generated by CAML) on Antarctic marine biodiversity by establishing and supporting a distributed system of interoperable databases, forming the Antarctic regional node of the Ocean Biogeographic Information System (OBIS, [www.iobis.org](http://www.iobis.org)), under the aegis of SCAR (Scientific Committee on Antarctic Research, [www.scar.org](http://www.scar.org)). SCAR-MarBIN established a comprehensive register of Antarctic marine species and, with biodiversity.aq provided free access to more than 2.9 million Antarctic georeferenced biodiversity data, which allowed more than 60 million downloads.

## The Editorial Team



**Claude DE BROUER** is a marine biologist at the Royal Belgian Institute of Natural Sciences in Brussels. His research interests cover structural and ecofunctional biodiversity and biogeography of crustaceans, and polar and deep sea benthic ecology. Active promoter of CAML and ANDEEP, he is the initiator of the SCAR Marine Biodiversity Information Network (SCAR-MarBIN). He took part to 19 polar expeditions.



**Huw GRIFFITHS** is a marine Biogeographer at the British Antarctic Survey. He created and manages SOMBASE, the Southern Ocean Mollusc Database. His interests include large-scale biogeographic and ecological patterns in space and time. His focus has been on molluscs, bryozoans, sponges and pycnogonids as model groups to investigate trends at high southern latitudes.



**Cédric d'UDEKEM d'ACOZ** is a research scientist at the Royal Belgian Institute of Natural Sciences, Brussels. His main research interests are systematics of amphipod crustaceans, especially of polar species and taxonomy of decapod crustaceans. He took part to 2 scientific expeditions to Antarctica on board of the *Polarstern* and to several sampling campaigns in Norway and Svalbard.



**Bruno DANIS** is an Associate Professor at the Université Libre de Bruxelles, where his research focuses on polar biodiversity. Former coordinator of the scarmarbin.be and antabif.be projects, he is a leading member of several international committees, such as OBIS or the SCAR Expert Group on Antarctic Biodiversity Informatics. He has published papers in various fields, including ecotoxicology, physiology, biodiversity informatics, polar biodiversity or information science.



**Susie GRANT** is a marine biogeographer at the British Antarctic Survey. Her work is focused on the design and implementation of marine protected areas, particularly through the use of biogeographic information in systematic conservation planning.



**Christoph HELD** is a Senior Research Scientist at the Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Bremerhaven. He is a specialist in molecular systematics and phylogeography of Antarctic crustaceans, especially isopods.




**Falk HUETTMANN** is a 'digital naturalist' he works on three poles (Arctic, Antarctic and Hindu-Kush Himalaya) and elsewhere (marine, terrestrial and atmosphere). He is based with the university of Alaska-Fairbank (UAF) and focuses primarily on effective conservation questions engaging predictions and open access data.



**Philippe KOUBBI** is professor at the University Pierre et Marie Curie (Paris, France) and a specialist in Antarctic fish ecology and biogeography. He is the Principal Investigator of projects supported by IPEV, the French Polar Institute. As a French representative to the CCAMLR Scientific Committee, his main input is on the proposal of Marine Protected Areas. His other field of research is on the ecoregionalisation of the high seas.



**Ben RAYMOND** is a computational ecologist and exploratory data analyst, working across a variety of Southern Ocean, Antarctic, and wider research projects. His areas of interest include ecosystem modelling, regionalisation and marine protected area selection, risk assessment, animal tracking, seabird ecology, complex systems, and remote sensed data analyses.



**Anton VAN DE PUTTE** works at the Royal Belgian Institute for Natural Sciences (Brussels, Belgium). He is an expert in the ecology and evolution of Antarctic fish and is currently the Science Officer for the Antarctic Biodiversity Portal [www.biodiversity.aq](http://www.biodiversity.aq). This portal provides free and open access to Antarctic Marine and terrestrial biodiversity of the Antarctic and the Southern Ocean.



**Bruno DAVID** is CNRS director of research at the laboratory BIOGÉOSCIENCES, University of Burgundy. His works focus on evolution of living forms, with and more specifically on sea urchins. He authored a book and edited an extensive database on Antarctic echinoids. He is currently President of the scientific council of the Muséum National d'Histoire Naturelle (Paris), and Deputy Director at the CNRS Institute for Ecology and Environment.



**Julian GUTT** is a marine ecologist at the Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Bremerhaven, and professor at the Oldenburg University, Germany. He participated in 13 scientific expeditions to the Antarctic and was twice chief scientist on board Polarstern. He is member of the SCAR committees ACCE and AnT-ERA (as chief officer). Main foci of his work are: biodiversity, ecosystem functioning and services, response of marine systems to climate change, non-invasive technologies, and outreach.



**Graham HOSIE** is Principal Research Scientist in zooplankton ecology at the Australian Antarctic Division. He founded the SCAR Southern Ocean Continuous Plankton Recorder Survey and is the Chief Officer of the SCAR Life Sciences Standing Scientific Group. His research interests include the ecology and biogeography of plankton species and communities, notably their response to environmental changes. He has participated in 17 marine science voyages to Antarctica.



**Alexandra POST** is a marine geoscientist, with expertise in benthic habitat mapping, sedimentology and geomorphic characterisation of the seafloor. She has worked at Geoscience Australia since 2002, with a primary focus on understanding seafloor processes and habitats on the East Antarctic margin. Most recently she has led work to understand the biophysical environment beneath the Amery Ice Shelf, and to characterise the habitats on the George V Shelf and slope following the successful CAML voyages in that region.



**Yan ROPERT COUDERT** spent 10 years at the Japanese National Institute of Polar Research, where he graduated as a Doctor in Polar Sciences in 2001. Since 2007, he is a permanent researcher at the CNRS in France and the director of a polar research programme (since 2011) that examines the ecological response of Adélie penguins to environmental changes. He is also the secretary of the Expert Group on Birds and Marine Mammals and of the Life Science Group of the Scientific Committee on Antarctic Research.

