

Census of Antarctic Marine Life
SCAR-Marine Biodiversity Information Network

BIOGEOGRAPHIC ATLAS OF THE SOUTHERN OCEAN

► **CHAPTER 5.28. CLASSIFICATION AND SPATIALLY EXPLICIT
ILLUSTRATION OF ANTARCTIC MACROBENTHIC ASSEMBLAGES:
A FEASIBILITY STUDY.** Gutt J., Barnes D.K.A., Lockhart S.J., 2014.

In: De Broyer C., Koubbi P., Griffiths H.J., Raymond B., Udekem d'Acoz C. d', et al. (eds.). Biogeographic Atlas of the Southern Ocean. Scientific Committee on Antarctic Research, Cambridge, pp. 229-232.

EDITED BY:

Claude DE BROYER & Philippe KOUBBI (chief editors)

with Huw GRIFFITHS, Ben RAYMOND, Cédric d'UDEKEM
d'ACQZ, Anton VAN DE PUTTE, Bruno DANIS, Bruno DAVID,
Susie GRANT, Julian GUTT, Christoph HELD, Graham HOSIE,
Falk HUETTMANN, Alexandra POST & Yan ROPERT-COUDERT



SCIENTIFIC COMMITTEE ON ANTARCTIC RESEARCH

THE BIOGEOGRAPHIC ATLAS OF THE SOUTHERN OCEAN

The “Biogeographic Atlas of the Southern Ocean” is a legacy of the International Polar Year 2007-2009 (www.ipy.org) and of the Census of Marine Life 2000-2010 (www.coml.org), contributed by the Census of Antarctic Marine Life (www.caml.aq) and the SCAR Marine Biodiversity Information Network (www.scarmarbin.be; www.biodiversity.aq).

The “Biogeographic Atlas” is a contribution to the SCAR programmes Ant-ECO (State of the Antarctic Ecosystem) and AnT-ERA (Antarctic Thresholds- Ecosystem Resilience and Adaptation) (www.scar.org/science-themes/ecosystems).

Edited by:

Claude De Broyer (Royal Belgian Institute of Natural Sciences, Brussels)
Philippe Koubbi (Université Pierre et Marie Curie, Paris)
Huw Griffiths (British Antarctic Survey, Cambridge)
Ben Raymond (Australian Antarctic Division, Hobart)
Cédric d’Udekem d’Acoz (Royal Belgian Institute of Natural Sciences, Brussels)
Anton Van de Putte (Royal Belgian Institute of Natural Sciences, Brussels)
Bruno Danis (Université Libre de Bruxelles, Brussels)
Bruno David (Université de Bourgogne, Dijon)
Susie Grant (British Antarctic Survey, Cambridge)
Julian Gutt (Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven)
Christoph Held (Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven)
Graham Hosie (Australian Antarctic Division, Hobart)
Falk Huettmann (University of Alaska, Fairbanks)
Alix Post (Geoscience Australia, Canberra)
Yan Ropert-Coudert (Institut Pluridisciplinaire Hubert Currien, Strasbourg)

Published by:

The Scientific Committee on Antarctic Research, Scott Polar Research Institute, Lensfield Road, Cambridge, CB2 1ER, United Kingdom (www.scar.org).

Publication funded by:

- The Census of Marine Life (Albert P. Sloan Foundation, New York)
- The TOTAL Foundation, Paris.

The “Biogeographic Atlas of the Southern Ocean” shared the *Cosmos Prize* awarded to the Census of Marine Life by the International Osaka Expo’90 Commemorative Foundation, Tokyo, Japan.

Publication supported by:

- The Belgian Science Policy (Belspo), through the Belgian Scientific Research Programme on the Antarctic and the “biodiversity.aq” network (SCAR-MarBIN/ANTABIF)
- The Royal Belgian Institute of Natural Sciences (RBINS), Brussels, Belgium
- The British Antarctic Survey (BAS), Cambridge, United Kingdom
- The Université Pierre et Marie Curie (UPMC), Paris, France
- The Australian Antarctic Division, Hobart, Australia
- The Scientific Steering Committee of CAML, Michael Stoddart (CAML Administrator) and Victoria Wadley (CAML Project Manager)

Mapping coordination and design: Huw Griffiths (BAS, Cambridge) & Anton Van de Putte (RBINS, Brussels)

Editorial assistance: Henri Robert, Xavier Loréa, Charlotte Havermans, Nicole Moortgat (RBINS, Brussels)

Printed by: Altitude Design, Rue Saint Josse, 15, B-1210 Brussels, Belgium (www.altitude-design.be)

Lay out: Sigrid Camus & Amélie Blaton (Altitude Design, Brussels).

Cover design: Amélie Blaton (Altitude Design, Brussels) and the Editorial Team.

Cover pictures: amphipod crustacean (*Epimeria rubriques* De Broyer & Klages, 1991), image © T. Riehl, University of Hamburg; krill (*Euphausia superba* Dana, 1850), image © V. Siegel, Institute of Sea Fisheries, Hamburg; fish (*Chaenocephalus* sp.), image © C. d’Udekem d’Acoz, RBINS; emperor penguin (*Aptenodytes forsteri* G.R. Gray, 1844), image © C. d’Udekem d’Acoz, RBINS; Humpback whale (*Megaptera novaeangliae* (Borowski, 1781)), image © L. Kindermann, AWI.

Online dynamic version :

A dynamic online version of the Biogeographic Atlas is available on the SCAR-MarBIN / AntaBIF portal : atlas.biodiversity.aq.

Recommended citation:

For the volume:

De Broyer C., Koubbi P., Griffiths H.J., Raymond B., Udekem d’Acoz C. d’, Van de Putte A.P., Danis B., David B., Grant S., Gutt J., Held C., Hosie G., Huettmann F., Post A., Ropert-Coudert Y. (eds.), 2014. Biogeographic Atlas of the Southern Ocean. Scientific Committee on Antarctic Research, Cambridge, XII + 498 pp.

For individual chapter:

(e.g.) Crame A., 2014. Chapter 3.1. Evolutionary Setting. In: De Broyer C., Koubbi P., Griffiths H.J., Raymond B., Udekem d’Acoz C. d’, et al. (eds.). Biogeographic Atlas of the Southern Ocean. Scientific Committee on Antarctic Research, Cambridge, pp. xx-yy.

ISBN: 978-0-948277-28-3.



This publication is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License

5.28. Classification and spatially explicit illustration of Antarctic macrobenthic assemblages: a feasibility study

Julian Gutt¹, David Barnes² & Susanne J. Lockhart³

¹ Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Bremerhaven, Germany

² British Antarctic Survey, Cambridge, UK

³ Southwest Fisheries Science Center, NOAA, La Jolla, CA, USA

Macrobenthos form substantial and varied assemblages on continental shelves around the Southern Ocean. It includes all benthic organisms >1mm and, thus, also the megabenthos, which is often less clearly defined. These macrobenthic assemblages cover considerable gradients in abundance, species composition (Photo 1) and in ecological functioning. They can take the form of 'forests' of sessile bioconstructors, which are mainly suspension (filter) feeders, such as sponges, corals and bryozoans, or clumps of animals such as hydroids, gorgonians, sea-squirts (ascidians) or brachiopods (Dayton 1990, Clarke & Johnston 2003). Some of these provide substratum for speciose epibiotic symbioses (see Schiaparelli chapter 5.31, this volume). Siphons of buried bivalve molluscs, polychaetes and other deposit feeders omnivorous species or grazers (often echinoderms) form low-lying shapes (see e.g. Sumida *et al.* 2008). Seeps and vents in the Southern Ocean have recently been explored in detail (see Rogers & Linse chapter 5.30, this volume). Representatives from groups such as suspension feeders, predators, deposit feeders, and grazers can each locally dominate the benthic system in its functioning and biodiversity (Kaiser *et al.* 2013). Patchiness occurs across many spatial scales in depth and geography ranging from almost circumpolar in distribution (biogeographic features) to small-scaled but high concentrations of a variety of organisms e.g. on single drop stones surrounded by near desert conditions. Abundances, biomass or proxies thereof range from almost the lowest (e.g. beneath floating ice shelves) to the highest recorded in the world, (e.g. in dense sponge communities); for review see Gutt *et al.* (2012).

Ship-based surveys of benthic structure have a long tradition, but were well underway by the late 1950s. Process orientated ecological experiments and shallow water studies have mainly been carried out from land-based research

stations. Pioneering work in the 1960–1980s by Paul Dayton and colleagues investigated benthic communities in McMurdo Sound and, consequently, this location is one of the best understood areas for Antarctic benthos. Shelf areas all around the continent and islands down to 300 m deep are known to be mechanically disturbed by ice, particularly by icebergs grounding (termed scouring) with such intensity and frequency that only rapid recolonisers and growers (pioneers) are locally abundant. Anthropogenic impacts, mainly by bottom trawling and long-line fishing, have been considerable in some areas in the past and still have a destructive effect on Antarctic benthic assemblages, but these have never been comprehensively quantified. On the other hand, a number of measures of nature conservation have recently been adopted. New sampling programmes, especially within the framework of the Census of Antarctic Marine Life (CAML), have greatly increased wider knowledge but we still have only a rudimentary understanding of community structure and the physical drivers which underlie them (Gutt *et al.* 2012). This has been recognised by the two new biology programmes of SCAR, "Antarctic Thresholds - Ecosystem Resilience and Adaptation" (AnT-ERA) and "State of the Antarctic Ecosystem" (AntEco), which focus on ecosystem functioning in a changing environment and large scale patterns shaped by long term evolution, respectively.

Below the continental shelves we know little about macrobenthic assemblages, although the continental slope and Antarctic deep-sea are increasingly being considered in survey design and sampling. However, the high species richness, large size of the Southern Ocean, and cost in time and money of deep-sea sampling means that comprehensive assemblage analyses are still very difficult and rare. In the last few years information






Photo 1 Selection of Antarctic macrobenthic shelf inhabiting assemblages. Upper left: dominated by barrel shaped hexactinellid sponges, associated fauna: demosponges, pencil sea urchin, crinoid, benthic ctenophore, bryozoans, gorgonians, ascidians, hemichordates (70.941°S 010°515'W, 260 m water depth, SE Weddell Sea, doi.pangaea.de/10.1594/PANGAEA.666982); Upper centre: assemblage with very low abundances, only one shrimp is visible but "Lebensspuren" indicate activity of other mobile organisms (54.429°S 035.684°W, 270 m water depth, South Georgia, doi.pangaea.de/10.1594/PANGAEA.220742); Upper right: assemblage dominated by one single species, the demosponge *Homaxinella* sp. Most likely this "monoculture" represents a relative early stage of recolonisation after physical disturbance due to iceberg grounding (71.107°S 011.532°W, 160 m water depth, SE Weddell Sea, doi.pangaea.de/10.1594/PANGAEA.319903); Lower left: recent iceberg scour, only few survivors are present (70.835°S 010.593°W, 275 m water depth, SE Weddell Sea, doi.pangaea.de/10.1594/PANGAEA.319904); Lower centre: predators like these king crabs (*Lithodidae*) at the Spiess Sea-mount close to Bouvet Island are rare in the Antarctic but discussed as possible threat for potential food organisms because of their durophagous feeding when invading large Antarctic shelf areas as a consequence of oceanic warming (54.733°S 000.092°W, 350 m water depth, doi.pangaea.de/10.1594/PANGAEA.691581); Lower right: assemblages dominated by deposit feeders such as ophiuroids (71.677°S 012.723°W, 235 m water depth, SE Weddell Sea, doi.pangaea.de/10.1594/PANGAEA.319888). All photographs represent approximately 1 m², with the exception of that in the lower centre representing 0.56 m².


► Macrobenthic Assemblages




Map 1
 ● SSFA
 ● SSFA-SPO
 ● SSFA-OTH
 ▲ PRED



Map 2
 ● MOIN
 ● MOIN-EPI



Map 3
 ● MOIN-INF
 ● MIX



Map 4
 ● VLAB
 ● MONO
 ● PHYCO

Macrobenthic Assemblages Maps 1–4. Illustration of macrobenthic communities on the Antarctic shelf. For definition of acronyms see Fig. 1. A first interpretation is extreme patchiness across spatial scales. Despite the general dominance of sessile suspension feeders, it appears to be regionally fractal in nature. If one zooms into the patterns they do not get less heterogeneous.

systems have begun to provide easier access to many recent and historic data sets. These are facilitating the first attempts to map the occurrence of species and biogeographic analyses of taxa. To date, this is virtually all based on presence/absence data because of the rarity of truly quantitative and comparable sampling. However, improved imagery platforms and use of equipment such as mega-cores are rapidly improving estimates of abundance and density. Such studies provide a basis for more advanced macroecological studies e.g. correlative approaches linking the environment and biota (for example see Cummings *et al.* 2010), spatial predictions based on potential habitats (for review see Gutt *et al.* 2012), projections of responses of marine life to ongoing climate change, such as warming, sea-ice loss and gain as well as acidification (see e.g. Clarke *et al.* 2007, Turner *et al.* 2008, Brandt & Gutt 2011). Community analyses and studies of ecosystem functioning require more complex data and interdisciplinary coordination. Standardisation of ecological methods and approaches remains in its infancy, and thus results from different studies are so far statistically comparable only within limited areas and taxa, or if carried out by the same research team.

To date, comparison and classification of assemblages at a circumpolar scale is only possible at the level of expert best knowledge (interpretation of data) and not by statistical methods. A first step in this direction was a coarse classification of macrobenthic assemblages, defined by taxa being dominant - predominantly in terms of biomass, volume or sea-floor cover and by a few known ecological processes (Fig. 1). Using information from the literature (references cited below) was a first attempt to assemble data from historic

**Sessile Suspension Feeders
with Associated fauna (SSFA)**

**MOBILE
deposit
feeders,
INfauna &
grazers
(MOIN)**




Figure 1 Classification of Antarctic macrobenthic assemblages changed after Gutt (2007) and Gutt *et al.* (2013b). Mobile suspension feeders such as crinoids are included in category SSFA-OTH. Mobile predators not being part of the suspension feeder assemblages (e.g., king crabs) are included in MOIN-EPI. SEEP includes visible bacterial mats and dead clams. Letters used for acronyms in capitals.

and recent surveys and depict this information in a spatially explicit way (Maps 1–4). The meta-information of this data set was also published as a data paper (Gutt *et al.* 2013a) and a first careful analysis has been carried out (Gutt *et al.* 2013b). The collection, classification, and illustration of these data provide an overview for various further purposes (e.g. nature conservation measures or identification of scientific key sites) but do not necessarily represent all existing data sets. Thus, these circumpolar approaches are open for additions including additional size fractions, e.g. meiobenthos, and could be extended to outlying shelf areas such as around sub-Antarctic Islands and the deep-sea.

A better solution than to continue with this early staging-template approach would be to develop a standardised protocol for the acquirement of assemblage or community data. It should be applicable to all Antarctic macrobenthic assemblages and would provide the basis for quantitative (statistical) circumpolar analyses. Within this approach biomass should be determined for a complete range of larger taxonomic units including the possible absence of taxa similar to that of Lockhart & Jones (2008). If error can be quantified by some ground truthing, sea-floor cover could be used as proxy for biomass when using imaging methods. With similar precautions abundances can be translated to approximate biomass if conversion factors (and associated errors) are known and size classes are determined. The ecological aspect of this concept could be strengthened if a manageable number of ecologically important species and basic ecological drivers such as biological interactions, disturbance and strength between guilds are added. If this concept is realized an *a priori* classification may not be necessary anymore, instead analyses of assemblages can be applied and adjusted to the specific scientific question. As a consequence, problems of how to define details of the *a priori* classification, e.g. the dominance of functional groups, the thresholds between assemblages and the overlap between pure faunistic composition and ecological factors would be minimised. A convention for the area covering a single sample should be discussed since large scale surveys e.g. by video transects or trawl hauls are likely to integrate across habitats (Fig. 2) in contrast to spot grab samples.

The classification and zoogeographical distribution of assemblages arising from such an advanced approach are expected to become even more complex than the information depicted in Photo 1 and Fig. 1. Nevertheless, the results presented here mark a first step and potentially important basis to search for large-scale generality. In contrast they can also be used to examine fundamental differences within Antarctic benthic systems including links between habitat and biota, to understand their functioning, quantify ecosystem services and assess direct and indirect response to environmental changes.




Figure 2 Different successive stages of recolonisation along a 946 m long ROV-video transect show how much and at which spatial scales the benthic habitat can be fractionised by the impact of grounding icebergs (South-eastern Weddell Sea, 240 m water depth). For further information and sea-bed photographs see doi.pangaea.de/10.1594/PANGAEA.198668.

Acknowledgements

Thanks are due to all who provided data for this compilation (see reference list)! And to Huw Griffiths (BAS, Cambridge) and Anton Van de Putte (RBINS, Brussels) who prepared the maps. This is CAML contribution # 126.

References

- Arnaud, P., 1974. Contribution a la bionomie marine benthique des régions antarctiques et subantarctiques. *Téthys*, **6**, 1–464.
- Azam, F., Beers, J.R., Campbell, L., Carlucci, A.F., Holm-Hansen, O., Reis, F.M.H., Karl, D.M., 1979. Occurrence and metabolic activity of organisms under the Ross Ice Shelf, Antarctica, at station J9. *Science*, **203**, 451–453.
- Bannasch, R., Feiler, K., Rauschert, M., 1984. Fortsetzung der biologischen Untersuchungen im Gebiet der sowjetischen Antarktisstation Bellingshausen. *Geodätische und geophysikalische Veröffentlichungen*, Reihe **1**, Heft **11**, 37–55.
- Barnes, D.K.A., 1995 a. Sublittoral epifaunal communities at Signy Island, Antarctica. I. The ice-foot zone. *Marine Biology*, **121**, 555–563.
- Barnes, D.K.A., 1995 b. Sublittoral epifaunal communities at Signy Island, Antarctica. II. Below the ice-foot zone. *Marine Biology*, **121**, 565–572.
- Barnes, D.K.A., Clarke, A., 1995. Epibiotic communities on sublittoral macroinvertebrates at Signy Island, Antarctica. *Journal of the Marine Biological Association of the United Kingdom*, **75**, 689–703.
- Barry, J.P., Dayton, P.K., 1988. Current patterns in McMurdo Sound, Antarctica and their relationship to local biotic communities. *Polar Biology*, **8**, 367–376.
- Barthel, D., Gutt, J., 1992. Sponge associations in the eastern Weddell Sea. *Antarctic Science*, **4**, 137–150.
- Beaman, R.J., O'Brien, P.E., 2009. Collaborative East Antarctic Marine Census (CEAMARC): Post-Survey Report, RSV Aurora Australis Voyage 3, December 2007 – January 2008. *Geoscience Australia Record 2008/05*, 61 pp.
- Bellisio, N.B., Lopez, R.B., Tomo, A.P., 1972. Distribucion vertical de la fauna bentonica en tres localidades antarticas: Bahia Esperanza, Isla Petermann y Archipelago Melchior. *Contribucion Instituto Antartico Argentino*, **142**, 1–89.
- Belyaev, G.M., Ushakov, P.V., 1957. Certain regularities in the quantitative distribution of the bottom fauna in Antarctic waters. *Doklady Akademii Nauk SSSR*, **112**, 137–140.
- Bowden, D.A., Schiaparelli, S., Clark, M.R., Rickard, G.J., 2011. A lost world? Archaic crinoid-dominated assemblages on an Antarctic seamount. *Deep-Sea Research II*, **58**, 119–127.
- Brandt, A., Gutt, J., 2011. Biodiversity of a unique environment: the Southern Ocean benthos shaped and threatened by climate change. In: Zachos, F.E., J.C. Habel, J.C. (eds). *Biodiversity hotspots*. Berlin: Springer, pp. 503–526.
- Bruchhausen, P.M., Raymond, J.A., Jacobs, S.S., DeVries, A.L., Thorndike, E.M., DeWitt, H.H., 1979. Fish, crustaceans, and the sea floor under the Ross Ice Shelf. *Science*, **203**, 449–451.
- Bullivant, J.S., 1967. Ecology of the Ross Sea benthos. *Bulletin of the New Zealand Department of Scientific and Industrial Research*, **176**, 49–78.
- Castellanos, Z.J.A. de, 1973. Estratificación del complejo bentónico de invertebrados en Puerto Paraíso (Antártida). *Contribucion del Instituto Antartico Argentino*, **164**, 4–23.
- Cattaneo-Vietti R., Chiantore, M., Albertelli, G., 1997. The population structure and ecology of the Antarctic scallop *Adamussium colbecki* (Smith, 1902) at Terra Nova Bay (Ross Sea, Antarctica). *Scientia Marina*, **61**, Supl. **2**, 15–24.
- Cattaneo-Vietti, R., Chiantore, M., Gambi, M.C., Albertelli, G., Cormaci, M., Di Geronimo, I., 2000 a. Spatial and vertical distribution of benthic littoral communities in Terra Nova Bay. In: Faranda, F.M., Guglielmo, L., Ionora, A. (eds.). *Ross Sea Ecology*. Berlin: Springer Verlag, pp. 503–514.
- Cattaneo-Vietti, R., Bavestrello, G., Cerrano, C., Gaino, E., Mazzella, L., Pansini, M., Sarà, M., 2000 b. The role of sponges in the Terra Nova Bay ecosystem. In: Faranda, F.M., Guglielmo, L., Ionora, A. (eds.). *Ross Sea Ecology*. Berlin: Springer Verlag, 539–549.
- Cerrano, C., Bavestrello, G., Calcinaï, B., Cattaneo-Vietti, R., Sarà, A., 2000. Asteroids eating sponges from Tethys Bay, East Antarctica. *Antarctic Science*, **12**, pp. 425–426.
- Chiantore, M., Cattaneo-Vietti, R., Berkman, P.A., Nigro, M., Vacchi, M., Schiaparelli, S., Albertelli, G., 2001. Antarctic scallop (*Adamussium colbecki*) spatial population variability along the Victoria Land Coast, Antarctica. *Polar Biology*, **24**, 139–143.
- Clarke, A., Johnston, N.M. 2003. Antarctic marine benthic diversity, *Oceanography and Marine Biology - An Annual Review*, **41**, 47–114.
- Clarke, A., Murphy, E.J., Meredith, M.P., King, J.C., Peck, L.S., Barnes, D.K.A., Smith, R.C. 2007. Climate change and the marine ecosystem of the western Antarctic Peninsula. *Philosophical Transactions of the Royal Society B*, **362**, 149–166.
- Cranmer, T.L., Ruhl, H.A., Baldwin, R.J., Kaumann, R.S., 2003. Spatial and temporal variation in the abundance, distribution and population structure of epibenthic megafauna in Port Foster, Deception Island. *Deep-Sea Research II*, **50**, 1821–1842.
- Cummings, V., Thrush, S., Chiantore, M., Hewitt, J., Cattaneo-Vietti, R. 2010. Macrobenthic communities of the north-western Ross Sea shelf: links to depth, sediment characteristics and latitude. *Antarctic Science*, **22**, 793–804.
- Davis, R.W., Castellini, M.A., Horning, M., Davis, M.P., 1983. Maintenance of an observation hole through the McMurdo Ice Shelf for winter oceanography. *Antarctic Journal of the United States*, **18**, 12–14.
- Dayton, P.K., 1979. Observations on growth, dispersal and population dynamics of some sponges in McMurdo Sound, Antarctica. *Colloques internationaux du Centre National de la Recherche Scientifique*, **291**, 271–282.
- Dayton, P.K., 1990. Polar benthos. In: Smith, W.O. (ed.) *Polar oceanography, Part B: Chemistry, biology, and geology*. London: Academic Press, pp. 631–685.
- Dayton, P.K., Oliver, J.S., 1977. Antarctic soft-bottom benthos in oligotrophic and eutrophic environments. *Science*, **197**, 55–58.
- Dayton, P.K., Kooyman, G.L., Barry, J.P., 1984. Benthic life under thick ice. *Antarctic Journal of the United States*, **19**, 128.
- Dayton, P.K., Robillard, G.A., Paine, R.T., 1970. Benthic faunal zonation as a result of anchor ice at McMurdo Sound Antarctica. In: Holdgate, M.W. (ed.). *Antarctic Ecology*. New York: Academic Press, pp. 244–258.
- Dayton, P.K., Robillard, G.A., Paine, R.T., Dayton, L.B., 1974. Biological accommodation in the benthic community at McMurdo Sound, Antarctica. *Ecological Monographs*, **44**, 105–128.
- Domack, E., Ishman, S., Leventer, A., Sylva, S., Willmott, V., Huber, B., 2005. A chemotrophic ecosystem found beneath Antarctic ice shelf. *EOS*, **86**(269), 271–272.
- Drescher, H.E., Hubold, G., Piatkowski, U., Plötz, J., Voß, J., 1983. Das biologische Programm der ANTARKTIS-I-Expedition mit FS "Polarstern". *Berichte zur Polarforschung*, **12**, 1–34.
- Everitt, D.A., Poore, G.C.B., Pickard, J., 1980. Marine benthos from Davis Station, east Antarctica. *Australian Journal of Freshwater Research*, **31**, 829–836.
- Gallardo, V.A., Castillo, J.G., Retamal, M.A., Yáñez, A., Moyano, H.I., Hermosilla, J.G., 1977. Quantitative studies on the soft-bottom macrobenthic animal communities of shallow Antarctic bays. In: Llano, G.A. (ed.). *Adaptations within Antarctic ecosystems*. Washington, DC: Smithsonian Institution, pp. 361–387.
- Gerdes, D., Klages, M., Arntz, W.E., Herman, R.L., Galéron, J., Hain, S. 1992. Quantitative investigations on macrobenthos communities of the southeastern Weddell Sea shelf based on multibox corer samples. *Polar Biology*, **12**, 291–301.
- Gruzov, E.N., Pushkin, A.F., 1970. Bottom communities of the upper sublittoral of Enderby Land and the South Shetlands. In: Holdgate, M.W. (ed.). *Antarctic Ecology*, Vol. 1. London: Academic Press, pp. 235–238.
- Gruzov, E.N., Propp, M.V., Pushkin, A.F., 1968. Biological associations of coastal areas of the Davis Sea (based on the observations of divers). *Soviet Antarctic Expedition Information Bulletin*, **6**(6), 523–533.
- Gutt, J. 2007. Antarctic macro-zoobenthic communities: a review and an ecological classification. *Antarctic Science*, **19**, 165–182.

- Gutt, J., Piepenburg, D., 1991. Dense aggregations of deep-sea holothurians in the southern Weddell Sea, Antarctica. *Marine Ecology Progress Series*, **68**, 277–285.
- Gutt, J., Starmans, A., 1998. Structure and biodiversity of megabenthos in the Weddell and Lazarev Seas (Antarctica): ecological role of physical parameters and biological interactions. *Polar Biology*, **20**, 229–247.
- Gutt, J., 2007. Antarctic macro-zoobenthic communities: a review and an ecological classification. *Antarctic Science*, **19**, 165–182.
- Gutt, J., Koubbi, P., Eléaume, M., 2007. Mega-epibenthic diversity off Terre Adélie (Antarctica) in relation to disturbance. *Polar Biology*, **30**, 1323–1329.
- Gutt, J., Barratt, I., Domack, E., d'Udekem d'Acoz, C., Dimmler, W., Grémare, A., Heilmayer, O., Isla, E., Janussen, D., Jorgensen, E., Kock, K.-H., Lehnert, L.S., López-González, P., Langner, S., Linse, K., Manjón-Cabeza, M.E., Meißner, M., Montiel, A., Raes, M., Robert, H., Rose, A., Sañé Schepisi, E., Saucède, T., Scheidat, M., Schenke, H.-W., Seiler, J., Smith, C., 2010. Biodiversity change after climate-induced ice-shelf collapse in the Antarctic. *Deep-Sea Research II*, **58**, 74–83.
- Gutt, J., Hosie, G., Stoddart, M. 2010 Marine Life in the Antarctic. In: McIntyre, A.D. (ed) *Life in the World's Oceans: Diversity, Distribution, and Abundance*. Oxford: Blackwell Publishing Ltd., pp. 203–220.
- Gutt, J., Zurell, D., Bracegridle, T.J., Cheung, W., Clarke, M.S., Convey, P., Danis, B., David, B., De Broyer, C., di Prisco, G., Griffiths, H., Laffont, R., Peck, L., Pierrat, B., Riddle, M.J., Saucède, T., Turner, J., Verde, C., Wang, Z., Grimm, V. 2012. Correlative and dynamic species distribution modelling for ecological predictions in the Antarctic: a cross-disciplinary concept. *Polar Research*, **31**, 11091, <http://dx.doi.org/10.3402/polar.v31i0.11091>.
- Gutt, J., Barnes, D., Lockhart, S.J., van de Putte, A. 2013a. Antarctic macrobenthic communities: a compilation of circumpolar information. *Nature Conservation*, **4**, 1–13.
- Gutt, J., Griffiths, H.J., Jones, C.D. 2013b. Circumpolar overview and spatial heterogeneity of macrobenthic communities. *Marine Biodiversity*, doi:10.1007/s12526-013-0152-9
- Hamada, E., Numanami, H., Naito, Y., Taniguchi, A., 1986. Observation on the marine benthic organisms at Syowa Station in Antarctica using a remotely operated vehicle. *Memoirs of National Institute of Polar Research (Japan)*, **40**, 289–298.
- Jazdzewski, K., Juraz, W., Kittel, W., Presler, E., Presler, P., Siciński, J., 1986. Abundance and biomass estimates of the benthic Fauna in Admiralty Bay, King George Island, South Shetland Islands. *Polar Biology*, **6**, 5–16.
- Jazdzewski, K., De Broyer, C., Pudlacz, M., Zielinski, D., 2001. Seasonal fluctuations of vagile benthos in the uppermost sublittoral of a maritime Antarctic fjord. *Polar Biology*, **24**, 910–917.
- Jones, C.D., Lockhart, S.J., 2011. Detecting Vulnerable Marine Ecosystems in the Southern Ocean using research trawls and underwater imagery. *Marine Policy*, **35**(5), 7732–7736.
- Jones, D.O.B., 2005. *Ecological controls on density, diversity and community structure of polar megabenthos*. PhD thesis, University of Southampton, 281 pp. [Unpublished.]
- Kaiser, S., Brandão, S.N., Brix, S., Barnes, D.K.A., Bowden, D.A., Brenke, N., Ingels, J., Leese, F., Schiaparelli, S., Arango, C.P., Badhe, R., Bax, N., Błażewicz-Paszkowycz, M., Brandt, A., Catarino, A.I., David, B., De Ridder, C., Dubois, P., Ellingsen K.E., Glover, A.G., Griffiths, H.J., Gutt, J., Halanych, K.M., Havermans, C., Held, C., Janussen, D., Lörz, A.-N., Pearce, D.A., Pierrat, B., Riehl, T., Rose, A., Sands, C.J., Soler-Membrives, A., Schüller, M., Strugnell, J.M., Vanreusel, A., Veit-Köhler, G., Wilson, N.G., Yasuhara, M. 2013. Pattern, process and vulnerability of Antarctic and Southern Ocean benthos - a decadal leap in knowledge and understanding. *Marine Biology*, doi 10.1007/s00227-013-2232-6
- Kirkwood, J.M., Burton, H.R., 1988. Macrobenthic species assemblages in Ellis Fjord, Vestfold Hills, Antarctica. *Marine Biology*, **97**, 445–457.
- Kohnen, H., 1984. Die Expedition ANTARKTIS-II mit FS "Polarstern" 1983/84. *Berichte zur Polarforschung*, **19**, 1–185.
- Lipps J.H., Ronan, T.E. Jr., DeLaca T.E., 1979. Life below the Ross Ice Shelf, Antarctica. *Science*, **203**, 447–449.
- Littlepage, J.L., Pearse, J.S., 1962. Biological and oceanographic observations under an Antarctic ice shelf. *Science*, **137**, 679–681.
- Lockhart, S.J., Jones, C.D., 2008. Biogeographic patterns of benthic invertebrate megafauna on shelf areas within the Southern Ocean Atlantic sector. *CCAMLR Science*, **15**, 167–192.
- Lovell, L.L., Trego, K.D., 2003. The epibenthic megafaunal and benthic infaunal invertebrates of Port Foster, Deception Island (South Shetland Islands, Antarctica). *Deep-Sea Research II*, **50**, 1799–1819.
- Lowry, J.K., 1975. Soft bottom macrobenthic community of Arthur Harbor, Antarctica. *Antarctic Research Series*, **23**, 1–19.
- Nakajima, Y., Watanabe, K., Naito, Y., 1982. Diving observations of the marine benthos at Syopwa station, Antarctica. *Memoirs of the National Institute of Polar Research (Japan)*, Special Issue, **23**, 44–54.
- Niemann, H., Fischer, D., Graffe, D., Knittel, K., Montiel, A., Heilmayer, O., Nöthen, K., Pape, T., Kasten, S., Bohrmann, G., Boetius, A., Gutt, J., 2009. Biogeochemistry of a low-activity cold seep in the Larsen B area, western Weddell Sea, Antarctica. *Biogeosciences*, **6**, 2383–2395.
- Oliver, J.S., Slattery, P.N., 1985. Effects of crustacean predators on species composition and population structure of soft-bodied infauna from McMurdo Sound, Antarctica. *Ophelia*, **24**, 155–175.
- Oliver, J.S., Watson, D.J., O'Connor, E.F., Dayton, P.K., 1976. Benthic communities of McMurdo Sound. *Antarctic Journal of the U.S.*, **11**, 58–549.
- Pabis, K., Siciński, J., Krymarys, M., 2011. Distribution patterns in the biomass of macrozoobenthic communities in Admiralty Bay (King George Island, South Shetlands, Antarctic). *Polar Biology*, **34**, 489–500.
- Piepenburg, D., Voß, J., Gutt, J., 1997. Assemblages of sea stars (Echinodermata: Asteroidea) and brittle stars (Echinodermata: Ophiuroidea) in the Weddell Sea (Antarctica) and off Northeast Greenland (Arctic): a comparison of diversity and abundance. *Polar Biology*, **17**, 305–322.
- Post, A.L., Hemer, M.A., O'Brien, P.E., Roberts, D., Craven, M., 2007. History of benthic colonisation beneath the Amery Ice Shelf, East Antarctica. *Marine Ecology Progress Series*, **344**, 29–37.
- Post, A.L., O'Brien, P.E., Beaman, R.J., Riddle, M.J., de Santis, L., 2010 a. Physical controls on deep water coral communities on the George V Land slope, East Antarctica. *Antarctic Science*, **22**, 371–378.
- Post, A.L., Beaman, R.J., O'Brien, P.E., Eléaume, M., Riddle, M.J., 2011. Community structure and benthic habitats across the George V Shelf, East Antarctica: trends through space and time. *Deep-Sea Research II*, **58**, 105–118.
- Propp, M.V., 1977. The study of bottom fauna at Haswell Island by scuba diving. In: Holdgate, M.W. (ed.). *Antarctic Ecology*, Vol. 1. London, Academic Press, pp. 239–241.
- Raguá-Gil, J.M., Gutt, J., Clarke, A., Arntz, W.E., 2004. Antarctic shallow-water mega-epibenthos: shaped by circumpolar dispersion or local conditions? *Marine Biology*, **144**, 829–839.
- Rehm, P., Hooke, R.A., Thatje, S., 2011. Macrofaunal communities on the continental shelf off Victoria Land, Ross Sea, Antarctica. *Antarctic Science*, **23**, 449–455.
- Riddle, M.J., Craven, M., Goldsworthy, P.M., Carsey, F., 2007. A diverse benthic assemblage 100 km from open water under the Amery Ice Shelf, Antarctica. *Paleoceanography*, **22**, P1204. doi:10.1029/2006PA001327.
- Rogers, A.D., Tyler, P.A., Connelly, D.P., Copley, J.T., James, R., Larter, R.D., Linse, K., Mills, R.A., Naveira Garabato, A., Pancost, R.D., Pearce, D.A., Polunin, N.V.C., German, C.R., Shank, T., Boersch-Supan, P.H., Alker, B.J., Aquilina, A., Bennett, S.A., Clarke, A., Dinley, R.J.J., Graham, A.G.C., Green, D.R.H., Hawkes, J.A., Hepburn, L., Hilario, A., Huvenne, V.A.I., Marsh, L., Ramirez-Llodra, E., Reid, W.D.K., Roterman, C.N., Sweeting, C.J., Thatje, S., Zwirgmaier, K., 2012. The Discovery of New Deep-Sea Hydrothermal Vent Communities in the Southern Ocean and Implications for Biogeography. *PLoS Biology*, **10**(1), e1001234. doi:10.1371/journal.pbio.1001234
- Sahade, R., Tatián, M., Kowalke, J., Kühne, S., Esnal, G.B., 1998. Benthic faunal associations on soft substrates at Potter Cove, King George Island, Antarctic. *Polar Biology*, **19**, 85–91.
- Saiz, J.I., García, F.J., Manjón-Cabeza, M.E., Parapar, J., Peña-Cantero, Saucède, T., Troncoso, J.S., Ramos, A., 2008. Community structure and spatial distribution of benthic fauna in the Bellingshausen Sea (West Antarctica). *Polar Biology*, **31**, 735–743 (only stations <1000 m considered).
- Siciński, J., Pabis, K., Jazdzewski, K., Konopacka, A., Błażewicz-Paszkowycz, M., 2012. Macrobenthos of two Antarctic glacial coves: a comparison with non-disturbed bottom areas. *Polar Biology*, **35**: 355–367. doi 10.1007/s00300-011-1081-3
- Smith, C.R., Grange, L.J., Honig, D.L., Naudts, L., Huber, B., Guidi, L., Domack, E., 2012. A large population of king crabs in Palmer Deep on the west Antarctic Peninsula shelf and potential invasive impacts. *Proceedings of the Royal Society B, Biological Sciences*, **279**, 1017–1026.
- Starmans, A., Gutt, J., Arntz, W.E., 1999. Mega-epibenthic communities in Arctic and Antarctic shelf areas. *Marine Biology*, **135**, 269–280.
- Teixidó, N., Garrabou, J., Arntz, W.E., 2002. Spatial pattern quantification of Antarctic benthic communities using landscape indices. *Marine Ecology Progress Series*, **242**, 1–14.
- Sumida, P.Y.G., Bernardino, A.F., Stedall, V.P., Glover, A.G., Smith, C.R., 2008. Temporal changes in benthic megafaunal abundance and composition across the West Antarctic Peninsula shelf: Results from video surveys. *Deep-Sea Research II*, **55**, 2465–2477.
- Teixidó, N., Garrabou, J., Gutt, J., Arntz, W.E., 2004. Recovery in Antarctic benthos after iceberg disturbance: trends in benthic composition, abundance, and growth forms. *Marine Ecology Progress Series*, **278**, 1–16.
- Teixidó, N., Garrabou, J., Gutt, J., Arntz, W.E., 2007. Iceberg disturbance and successional spatial patterns: the case of the shelf Antarctic benthic communities. *Ecosystems*, **10**, 142–157.
- Turner, J., Bindschadler, R., Convey, P., di Prisco, G., Fahrbach, E., Gutt, J., Hodgson, D., Mayewsky, P., Summerhayes, C. 2009. Antarctic Climate Change and the Environment. Cambridge: SCAR & Scott Polar Research Institute, 526 pp.
- Voß, J., 1988. Zoogeographie und Gemeinschaftsanalyse des Makrozoobenthos des Weddellmeeres (Antarktis). *Berichte zur Polarforschung*, **45**, 1–145.
- Watters, G., Bergström, B., Gutt, J., Petterson, J.-O., Rosenberg, J., Setran, A., Valenzuela, C., 1995. 9. Leg III: Crab and epifaunal surveys of bays, anchorages, and fjords around South Georgia. AMLR 1994/95 Field Season Report, Administrative Report LJ-95-13, National Oceanic and Atmospheric Administration, Southwest Fisheries Science Center, Antarctic Ecosystem Research Group, pp. 113–130.
- Zamorano, J.H., 1983. Zonación y Biomasa de la macrofauna bentónica en Bahía South, Archipiélago de Palmer, Antártica. *Serie científica. Instituto Antártico Chileno*, **30**, 27–38.

THE BIOGEOGRAPHIC ATLAS OF THE SOUTHERN OCEAN

Scope

Biogeographic information is of fundamental importance for discovering marine biodiversity hotspots, detecting and understanding impacts of environmental changes, predicting future distributions, monitoring biodiversity, or supporting conservation and sustainable management strategies.

The recent extensive exploration and assessment of biodiversity by the Census of Antarctic Marine Life (CAML), and the intense compilation and validation efforts of Southern Ocean biogeographic data by the SCAR Marine Biodiversity Information Network (SCAR-MarBIN / OBIS) provided a unique opportunity to assess and synthesise the current knowledge on Southern Ocean biogeography.

The scope of the Biogeographic Atlas of the Southern Ocean is to present a concise synopsis of the present state of knowledge of the distributional patterns of the major benthic and pelagic taxa and of the key communities, in the light of biotic and abiotic factors operating within an evolutionary framework. Each chapter has been written by the most pertinent experts in their field, relying on vastly improved occurrence datasets from recent decades, as well as on new insights provided by molecular and phylogeographic approaches, and new methods of analysis, visualisation, modelling and prediction of biogeographic distributions.

A dynamic online version of the Biogeographic Atlas will be hosted on www.biodiversity.aq.

The Census of Antarctic Marine Life (CAML)

CAML (www.caml.aq) was a 5-year project that aimed at assessing the nature, distribution and abundance of all living organisms of the Southern Ocean. In this time of environmental change, CAML provided a comprehensive baseline information on the Antarctic marine biodiversity as a sound benchmark against which future change can reliably be assessed. CAML was initiated in 2005 as the regional Antarctic project of the worldwide programme Census of Marine Life (2000-2010) and was the most important biology project of the International Polar Year 2007-2009.

The SCAR Marine Biodiversity Information Network (SCAR-MarBIN)

In close connection with CAML, SCAR-MarBIN (www.scarmarbin.be, integrated into www.biodiversity.aq) compiled and managed the historic, current and new information (i.a. generated by CAML) on Antarctic marine biodiversity by establishing and supporting a distributed system of interoperable databases, forming the Antarctic regional node of the Ocean Biogeographic Information System (OBIS, www.iobis.org), under the aegis of SCAR (Scientific Committee on Antarctic Research, www.scar.org). SCAR-MarBIN established a comprehensive register of Antarctic marine species and, with biodiversity.aq provided free access to more than 2.9 million Antarctic georeferenced biodiversity data, which allowed more than 60 million downloads.

The Editorial Team




Claude DE BROYER is a marine biologist at the Royal Belgian Institute of Natural Sciences in Brussels. His research interests cover structural and ecofunctional biodiversity and biogeography of crustaceans, and polar and deep sea benthic ecology. Active promoter of CAML and ANDEEP, he is the initiator of the SCAR Marine Biodiversity Information Network (SCAR-MarBIN). He took part to 19 polar expeditions.



Huw GRIFFITHS is a marine Biogeographer at the British Antarctic Survey. He created and manages SOMBASE, the Southern Ocean Mollusc Database. His interests include large-scale biogeographic and ecological patterns in space and time. His focus has been on molluscs, bryozoans, sponges and pycnogonids as model groups to investigate trends at high southern latitudes.



Cédric d'UDEKEM d'ACQZ is a research scientist at the Royal Belgian Institute of Natural Sciences, Brussels. His main research interests are systematics of amphipod crustaceans, especially of polar species and taxonomy of decapod crustaceans. He took part to 2 scientific expeditions to Antarctica on board of the *Polarstern* and to several sampling campaigns in Norway and Svalbard.



Bruno DANIS is an Associate Professor at the Université Libre de Bruxelles, where his research focuses on polar biodiversity. Former coordinator of the [scarmarbin.be](http://www.scarmarbin.be) and antibif.be projects, he is a leading member of several international committees, such as OBIS or the SCAR Expert Group on Antarctic Biodiversity Informatics. He has published papers in various fields, including ecotoxicology, physiology, biodiversity informatics, polar biodiversity or information science.



Susie GRANT is a marine biogeographer at the British Antarctic Survey. Her work is focused on the design and implementation of marine protected areas, particularly through the use of biogeographic information in systematic conservation planning.



Christoph HELD is a Senior Research Scientist at the Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven. He is a specialist in molecular systematics and phylogeography of Antarctic crustaceans, especially isopods.



Falk HUETTMANN is a 'digital naturalist' he works on three poles (Arctic, Antarctic and Hindu-Kush Himalaya) and elsewhere (marine, terrestrial and atmosphere). He is based with the university of Alaska-Fairbank (UAF) and focuses primarily on effective conservation questions engaging predictions and open access data.



Philippe KOUUBI is professor at the University Pierre et Marie Curie (Paris, France) and a specialist in Antarctic fish ecology and biogeography. He is the Principal Investigator of projects supported by IPEV, the French Polar Institute. As a French representative to the CCAMLR Scientific Committee, his main input is on the proposal of Marine Protected Areas. His other field of research is on the ecoregionalisation of the high seas.



Ben RAYMOND is a computational ecologist and exploratory data analyst, working across a variety of Southern Ocean, Antarctic, and wider research projects. His areas of interest include ecosystem modelling, regionalisation and marine protected area selection, risk assessment, animal tracking, seabird ecology, complex systems, and remote sensed data analyses.



Anton VAN DE PUTTE works at the Royal Belgian Institute for Natural Sciences (Brussels, Belgium). He is an expert in the ecology and evolution of Antarctic fish and is currently the Science Officer for the Antarctic Biodiversity Portal www.biodiversity.aq. This portal provides free and open access to Antarctic Marine and terrestrial biodiversity of the Antarctic and the Southern Ocean.



Bruno DAVID is CNRS director of research at the laboratory BIOGÉOSCIENCES, University of Burgundy. His works focus on evolution of living forms, with and more specifically on sea urchins. He authored a book and edited an extensive database on Antarctic echinoids. He is currently President of the scientific council of the Muséum National d'Histoire Naturelle (Paris), and Deputy Director at the CNRS Institute for Ecology and Environment.




Julian GUTT is a marine ecologist at the Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Bremerhaven, and professor at the Oldenburg University, Germany. He participated in 13 scientific expeditions to the Antarctic and was twice chief scientist on board *Polarstern*. He is member of the SCAR committees ACCE and AnT-ERA (as chief officer). Main foci of his work are: biodiversity, ecosystem functioning and services, response of marine systems to climate change, non-invasive technologies, and outreach.



Graham HOSIE is Principal Research Scientist in zooplankton ecology at the Australian Antarctic Division. He founded the SCAR Southern Ocean Continuous Plankton Recorder Survey and is the Chief Officer of the SCAR Life Sciences Standing Scientific Group. His research interests include the ecology and biogeography of plankton species and communities, notably their response to environmental changes. He has participated in 17 marine science voyages to Antarctica.



Alexandra POST is a marine geoscientist, with expertise in benthic habitat mapping, sedimentology and geomorphic characterisation of the seafloor. She has worked at Geoscience Australia since 2002, with a primary focus on understanding seafloor processes and habitats on the East Antarctic margin. Most recently she has led work to understand the biophysical environment beneath the Amery Ice Shelf, and to characterise the habitats on the George V Shelf and slope following the successful CAML voyages in that region.



Yan ROPERT COUDERT spent 10 years at the Japanese National Institute of Polar Research, where he graduated as a Doctor in Polar Sciences in 2001. Since 2007, he is a permanent researcher at the CNRS in France and the director of a polar research programme (since 2011) that examines the ecological response of Adélie penguins to environmental changes. He is also the secretary of the Expert Group on Birds and Marine Mammals and of the Life Science Group of the Scientific Committee on Antarctic Research.

