

5. Gondos B, Bhiraless P, Hobel CJ. Ultrastructural observations on germ cells in human fetal ovaries. *AM J OBSTET GYNECOL* 1971;110:644.
6. Streeter GL. Weight, sitting height, head size, foot length and menstrual age of the human embryo. *Carnegie Contrib Embryol* 1920;11:143.
7. Randolph M, Gondos B. Evaluation of premeiotic activity in the developing rabbit testis using a modified squash technique. *Gamete Res* 1982;6:127.
8. Byskov AG. Regulation of meiosis in mammals. *Ann Biol Anim Biochem Biophys* 1979;19:1251.
9. Hilscher B, Hilscher W. In: Byskov AG, Peters H. Development and function of reproductive organs. Amsterdam: Excerpta Medica, 1981.
10. Grinstead J, Byskov AG. Meiosis inducing and meiosis preventing substances in human male reproductive organs. *Fertil Steril* 1981;35:199.
11. Baker TG, Franchi LL. The fine structure of oogonia and oocytes in human ovaries. *J Cell Sci* 1967;2:213.

## Estrogen effects on plasma volume, arterial blood pressure, interstitial space, plasma proteins, and blood viscosity in sheep

Syuji Ueda, M.D., Vicki Fortune, B.S., Brian S. Bull, M.D.,  
Guillermo J. Valenzuela, M.D., and Lawrence D. Longo, M.D.

Loma Linda, California

In adult castrated ewes the infusion of 17 $\beta$ -estradiol for 3 weeks was associated with a 12% increase in body weight, a 20% increase in whole blood volume (mainly due to a 27% increase of plasma volume), a 13% decrease in mean arterial blood pressure, and a 40% increase in heart rate. The change in plasma volume correlated with the change in estradiol concentration ( $r = 0.72$ ). Most of the fluid was retained in the interstitial space, as represented by a 6 kg weight gain, 10% of which was in the intravascular compartment. Whole blood and plasma viscosity increased 16% and 21%, respectively, thus reversing some of the blood volume effects toward a hyperdynamic cardiovascular state. We conclude that many of the cardiovascular and hematologic changes with estrogen administration are similar to the changes observed during pregnancy, with the proposed requirement of decrease in mean arterial blood pressure as a condition for blood volume expansion. (*AM J OBSTET GYNECOL* 1986;155:195-201.)

**Key words:** Estrogen, blood volume, blood pressure, pregnancy

Human pregnancy is characterized by a 35% to 45% increase in blood volume from the nonpregnant values, mainly secondary to a 45% to 60% increase in plasma volume and a 20% to 30% increase in red cell mass.<sup>1</sup> Among other cardiovascular changes, diastolic arterial pressure decreases about 10 to 15 mm Hg and heart rate increases 10 to 15 bpm.<sup>2</sup> Also, plasma protein concentrations alter to affect both the Starling forces, and thus fluid flux across the capillaries, and blood viscosity, which is a major determinant of vascular resistance.<sup>3</sup>

During human pregnancy the concentrations of several hormones that play a role in fluid regulation increase in maternal plasma, including progesterone, de-

oxycorticosterone, aldosterone, and prolactin. A large increase also occurs in the concentration of 17 $\beta$ -estradiol and other estrogens. Experimentally it has been shown that estrogen administration to postmenopausal women, men with prostate cancer, and laboratory animals results in blood volume increases secondary to plasma volume expansion.<sup>4</sup>

In an effort to test the hypothesis that some of the cardiovascular and hematologic changes of pregnancy, such as the increase in plasma and whole blood volumes, heart rate, and decrease in mean arterial blood pressure, are a function of the circulating estrogen concentration, we infused 17 $\beta$ -estradiol into castrated ewes for 3 weeks and measured whole blood, plasma, and red cell volumes, arterial pressure, heart rate, plasma protein concentrations, and plasma and whole blood viscosities. The studies were not meant to reproduce exactly the endocrinologic milieu of pregnancy.

### Material and methods


Five adult ewes were used in the present study. Four to 7 days after they were received from a local breeder,

*From the Division of Perinatal Biology, Departments of Physiology, Obstetrics and Gynecology, and Pathology, School of Medicine, Loma Linda University.*

*Supported by United States Public Health Service Grants HD 03807 and HD 13949.*

*Received for publication September 24, 1985; revised January 27, 1986; accepted January 29, 1986.*

*Reprint requests: Lawrence D. Longo, M.D., Division of Perinatal Biology, School of Medicine, Loma Linda University, Loma Linda, CA 92350.*



**Fig. 1.** Percent change from control values for body weight, whole blood, plasma, and erythrocyte volumes, mean arterial blood pressure, and heart rate. \* =  $p < 0.05$ ; \*\* =  $p < 0.01$ .

they were oophorectomized under spinal anesthesia. We placed Tygon catheters in a hindlimb artery and vein and exteriorized them to a flank pouch. Subsequently we flushed the catheters daily with 5 ml of heparinized saline solution (50 IU of heparin per milliliter) and allowed the animals to recover from the surgical stress for 3 weeks. Then we placed the ewe in a special metabolism cart and for 3 weeks monitored their body weight, arterial blood pressure, heart rate, and blood volumes. We then continuously infused  $17\beta$ -estradiol (Sigma Chemical Co., St. Louis) at the rate of either 6.7 or 33.6 mg/kg/day for 3 weeks. Three ewes were infused in a random manner with both doses with a 4-week interval between infusions, during which time the various functions were measured. Of the other two ewes, one was infused with the lower dose, while the other received the higher dose. Thus four ewes were infused at each dose. In addition, three ewes were infused at the rate of 3.3 mg/kg/day, but no significant changes were observed in any of the parameters measured. In essence this constituted a control experiment for infusion of the vehicle. The  $17\beta$ -estradiol was dissolved in ethanol and then diluted in normal saline solution. The rate of saline infusion was 4 ml/hr, for a total of 96 ml/day; the concentration of ethanol in saline solution was 0.1%. This minimized the effects of both saline volume and ethanol on the ewe's cardiovascular system. Then at weekly intervals we determined arterial blood pressure, heart rate, plasma, erythrocyte, and whole blood volumes, blood plasma protein concentrations, and plasma viscosity.

**Blood volume measurements.** We determined the total blood volume by dilution of chromium  $^{51}$ -labeled red cells, and calculated the plasma volume indirectly from the whole blood volume and the microhematocrit.<sup>5</sup> The day of the experiment we labeled 3 ml blood with  $^{51}\text{Cr}$  using standard techniques.<sup>5</sup> We obtained a 0.5 ml blood sample for background radioactivity and then

injected the tagged erythrocytes through the venous catheter, which was flushed three times the catheter volume with heparinized saline solution. Thereafter we withdrew 0.5 ml blood samples at 10, 20, and 30 minutes through the arterial catheter. To clear the catheter, we withdrew a volume equal to six times that of the catheter before each sample was obtained and then replaced that blood. We measured the sample radioactivity in a gamma counter (Packard Instrument Co., Downers Grove, Illinois) and corrected it for hematocrit and sample weight. Then by the use of a computer program, we extrapolated the decay curve back to zero time and used this value to calculate whole blood volume. Whole blood, plasma, and erythrocyte volumes were expressed both in absolute values and per weight of the ewe.

**Arterial pressure and heart rate.** Arterial blood pressure was recorded with a pressure transducer (Gould, P23Db, Cleveland) on a polygraph (Gould, Model 200), which was connected to an on-line computer (Texas Instrument, 990/10, Houston). Following a 1-hour period of adaptation, the mean arterial pressure and heart rate were obtained for an additional hour, averaged over 1-minute intervals, and stored for later analysis.

**Plasma protein determinations.** At weekly intervals we withdrew arterial blood into a sterile, nonheparinized plastic syringe and immediately transferred it into an appropriate glass tube. We determined fibrinogen concentration by a timed conversion of fibrinogen to fibrin by topical thrombin, using a fibrometer (Fibrosystem, Cockeysville, Maryland) to detect clot formation. Each time we constructed a standard curve, using dilutions of a commercial plasma of known concentration (Data-Fi Fibrinogen Calibration Reference, Dade Diagnostics Inc., Aguada, Puerto Rico).

Total plasma protein and albumin concentrations were determined by the biuret method<sup>6</sup> and a modified bromocresol green binding method,<sup>7</sup> respectively. The

**Table I.** Cardiovascular and hematologic values in ewes before and after 3 weeks infusion with 17 $\beta$  estradiol\*

Variable	Control	Change	% change
Body weight (kg)	55.11 $\pm$ 2.14	6.44 $\dagger$ $\pm$ 1.83	11.7 $\dagger$
Whole blood volume (ml)	3,355 $\pm$ 124.5	653.8 $\dagger$ $\pm$ 125.9	19.5 $\dagger$
Plasma volume (ml)	2,346 $\pm$ 94.2	627.5 $\dagger$ $\pm$ 117.9	26.7 $\dagger$
Erythrocyte volume (ml)	1,009 $\pm$ 62.2	17.9 $\pm$ 57.6	1.8
Whole blood volume (ml/kg)	61.1 $\pm$ 1.4	4.26 $\dagger$ $\pm$ 1.03	7.0 $\dagger$
Plasma volume (ml/kg)	42.8 $\pm$ 1.7	5.58 $\dagger$ $\pm$ 2.12	13.0 $\dagger$
Erythrocyte volume (ml/kg)	18.2 $\pm$ 0.5	-1.48 $\pm$ 0.66	-8.1
Hemoglobin (gm/dl)	12.0 $\pm$ 0.6	-0.22 $\pm$ 0.39	-1.8
Hematocrit (%)	30.0 $\pm$ 1.3	-0.97 $\pm$ 1.04	-3.2
Blood pressure (mm Hg)	90.9 $\pm$ 1.9	-12.0 $\dagger$ $\pm$ 2.9	-13.2 $\dagger$
Heart rate (bpm)	65.3 $\pm$ 5.0	26.3 $\dagger$ $\pm$ 6.0	40.3 $\dagger$
Estrone (pg/ml)	4.59 $\pm$ 1.20	155.8 $\dagger$ $\pm$ 130.2	3,394 $\dagger$
17 $\beta$ -Estradiol (pg/ml)	20.0 $\pm$ 8.8	6,242 $\dagger$ $\pm$ 1,103	31,210 $\dagger$

\*Mean  $\pm$  SEM, n = 8.

$\dagger p < 0.05$ .

$\ddagger p < 0.01$ .

plasma globulin was determined by the difference of these values.

**Blood viscosity determinations.** We measured the absolute viscosity of distilled water, whole blood, and plasma at 37° C in centipoise, using a Coulter capillary viscometer (Coulter Electronics Ltd., Hialeah, Florida). Viscosity was calculated relative to distilled water.

**Estrogen assays.** Unconjugated unbound and bound serum estrone and 17 $\beta$ -estradiol were measured by radioimmunoassay by means of a modification of the method of Abraham et al.<sup>8</sup> (Radioassay Systems, Carson, California). The intra-assay and interassay coefficients of variation were 7% and 13%, respectively, for both hormones.

**Data analysis.** From the control and experimental measurements as well as paired differences between experimental and control values, we computed mean, SD, and SE of the means for each variable under consideration. We tested differences between means by a paired *t* test at the 0.05 and 0.01 levels.

## Results

**Body weight.** During the 3 weeks before estradiol infusion there was no significant change in body weight or other parameter. In contrast, during the 3 weeks of estradiol infusion the ewe's body weight increased about 12% (6.4  $\pm$  1.8 kg, mean  $\pm$  SEM) from 55.1  $\pm$  2.1 kg ( $p < 0.05$ ) at both 17 $\beta$ -estradiol doses (Fig. 1 and Table I).

**Blood volume.** Fig. 1 also shows the changes of whole blood volume, which gradually increased about 19% (654  $\pm$  126 ml) at 3 weeks, from 3355  $\pm$  125 ml at both 17 $\beta$ -estradiol doses ( $p < 0.01$ ). When calculated per kilogram of body weight, whole blood volume increased about 7% from a control value of 61.1  $\pm$  1.4 ml/kg ( $p < 0.05$ ) (Table I).

Essentially all of the whole blood volume increase could be accounted for by the rise in plasma volume,

which by the end of 3 weeks increased about 27% (628  $\pm$  118 ml) from 2346  $\pm$  94 ml ( $p < 0.01$ ). With the lower dose the rate of increase was gradual over the entire 3 weeks, while at the higher dose the increase was relatively rapid, occurring in 1 to 2 weeks. Again, when normalized to body weight, plasma volume increased 13% (5.6  $\pm$  2.1 ml/kg) from a control value of 42.8  $\pm$  1.7 ml/kg ( $p < 0.05$ ) (Table I). The whole blood and plasma volume changes returned to control values by two weeks after infusion. Erythrocyte volumes did not change significantly from the control values of 1009  $\pm$  62 and 18.2  $\pm$  0.5 ml/kg (Table I).

**Estrogen concentration.** The plasma concentrations of 17 $\beta$ -estradiol and estrone demonstrated very large increases (Table I). The mean plasma estradiol concentrations were 174  $\pm$  17, 4616  $\pm$  616, and 6991  $\pm$  875 pg/ml at the infusion rates of 3.3, 6.7, and 33.6 mg/kg/day, respectively. For estrone these values were 6.5  $\pm$  2, 179.6  $\pm$  21.2, and 239.3  $\pm$  87 pg/ml, respectively. As shown in Fig. 2, the change in plasma volume was highly correlated with the change in 17 $\beta$ -estradiol concentration ( $r = 0.72$ ,  $p < 0.001$ ).

**Arterial blood pressure and heart rate.** The mean arterial blood pressure decreased 13% (12  $\pm$  2.9 mm Hg) from 90.9  $\pm$  1.9 mm Hg ( $p < 0.01$ ) following the 3-week 17 $\beta$ -estradiol infusion. In turn, the heart rate increased about 40% (26  $\pm$  6 bpm) from a control value of 65  $\pm$  5 bpm ( $p < 0.01$ ) (Table I and Fig. 1).

**Blood plasma proteins and viscosity.** As shown in Fig. 3 and Table II, plasma fibrinogen concentration increased 86% (148  $\pm$  54 mg/dl) from 173  $\pm$  15 mg/dl. The plasma albumin concentration decreased 7% from 3.7  $\pm$  0.8 gm/dl, while the globulin increased 24% from 3.9  $\pm$  0.4 gm/dl. The albumin-to-globulin ratio decreased 31% from 0.99  $\pm$  0.13 (Fig. 3).

Plasma viscosity increased 21% from 1.58  $\pm$  0.17 centipoise ( $p < 0.01$ ), while whole blood viscosity increased 16% from 3.05  $\pm$  0.30 centipoise ( $p < 0.01$ )




Fig. 2. Change in plasma volume (L) as a function of increase in circulating  $17\beta$ -estradiol concentrations (ng/ml) following 2 and 3 weeks of estradiol infusion at 3.3 mg/kg/day ( $\square$ ),  $n = 6$ ; 6.7 mg/kg/day ( $\bullet$ ),  $n = 8$ ; and 33.6 mg/kg/day ( $\Delta$ ),  $n = 8$ .  $\blacklozenge$  represents the recovery values 3 weeks following the infusion of the two higher doses (i.e., 6.7 and 33.6 mg/kg/day,  $n = 8$ ).

(Table II and Fig. 3). As observed earlier, in three sheep infused with  $17\beta$ -estradiol at 3.3 mg/kg/day, no significant changes occurred in any of the parameters measured.

### Comment

**Hemodynamics.** The first aim of this study was to test the hypothesis that profound hematologic and cardiovascular changes, such as occur during pregnancy, can be secondary to elevated circulating estrogen concentrations. Human pregnancy is associated with some rather striking cardiovascular adaptations, such as 40% and 55% increases in whole blood and plasma volumes, respectively, several liters of fluid accumulation in the interstitial space, a 10% increase in heart rate, 30% increase in cardiac output, and decreases in both systemic vascular resistance and mean arterial blood pressure.<sup>1,9</sup>

Basically two theories have been postulated to explain these pregnancy-associated changes. Burwell et al.<sup>10, 11</sup> in the late 1930s first postulated that the placental circulation acts as an arteriovenous shunt. In experimental animals the opening of a shunt produces an immediate fall in vascular resistance, which in turn results in a drop in arterial pressure and cardiac output. Within a few seconds vasoconstriction results in compensatory blood flow redistribution. Heart rate increases, as does cardiac output, while arterial pressure gradually returns toward normal values. The decrease of renal perfusion, acting probably through the renin-angiotensin-aldosterone system, results in fluid reten-

Table II. Plasma protein and viscosity changes in ewes before and after 3 weeks' infusion of  $17\beta$ -estradiol\*

	Control	Change	% change
Fibrinogen (mg/dl)	$173 \pm 14.8$	$148 \pm 54.0^\dagger$	$85.5^\dagger$
Albumin (gm/dl)	$3.67 \pm 0.75$	$-0.25 \pm 0.09^\ddagger$	$-6.8^\ddagger$
Globulin (gm/dl)	$3.85 \pm 0.41$	$0.92 \pm 0.02^\ddagger$	$23.9^\ddagger$
Albumin/globulin ratio	$0.99 \pm 0.13$	$-0.31 \pm 0.07^\ddagger$	$-31.3^\ddagger$
Whole blood viscosity (centipoise)	$3.05 \pm 0.30$	$0.50 \pm 0.17^\ddagger$	$16.4^\ddagger$
Plasma viscosity (centipoise)	$1.58 \pm 0.17$	$0.33 \pm 0.12^\ddagger$	$20.9^\ddagger$

\*Mean value  $\pm$  SEM,  $n = 8$ .

$^\dagger p < 0.01$ .

$^\ddagger p < 0.05$ .


tion with blood volume expansion within a matter of minutes to hours.<sup>12, 13</sup>

The second theory proposes that the hormonal changes during pregnancy, chiefly large increases in circulating estrogens, increase blood volume, cardiac output, and uteroplacental blood flow.<sup>4</sup> Again, the mechanism probably involves estrogenic mediated stimulation of the renin-angiotensin system, producing an increase in aldosterone, with resultant sodium and fluid retention,<sup>14</sup> as well as direct action on the uteroplacental vessels.<sup>15, 16</sup> Estrogen administration also has been shown to decrease the number of angiotensin receptors<sup>17</sup> and to increase the amount of vasodilatory prostaglandins.<sup>18</sup> In pregnant women aldosterone secretion increases markedly, reaching a peak near the end of gestation.<sup>19</sup>

The infusion of either angiotensin II<sup>20</sup> or aldosterone<sup>21</sup> into experimental animals on a long-term basis has been shown to result in transitory fluid retention. However, the mean arterial pressure increase in turn results in increased glomerular filtration rate, so that blood volume returned to a normal value. Saline infusion demonstrates a similar effect.<sup>22</sup> Only when renal arterial pressure is maintained within normal levels with a servomechanism as control is it possible to keep the fluid volume persistently elevated.<sup>20, 21, 23</sup> Thus it appears that a prerequisite to maintaining an expanded blood volume is to maintain a relative or absolute decrease in renal arterial pressure.

During the course of gestation in sheep, whole blood and plasma volumes normally increase only about 6% and 8%, respectively.<sup>24</sup> Whole blood and plasma volumes in sheep with twins and triplets increase 9% and 12%, respectively (Ueda et al., unpublished observations). Circulating estrogen concentrations are comparably low in this species, rising from 20 to 400 pg/ml only during the last few days of gestation.<sup>25</sup>

The present findings that in castrated sheep the



**Fig. 3.** Percent change from control values for fibrinogen, albumin, globulin, albumin-to-globulin ratio, and whole blood and plasma viscosities. \* =  $p < 0.05$ ; \*\* =  $p < 0.01$ .

chronic administration of  $17\beta$ -estradiol resulted in decreased mean arterial pressure, increased heart rate, expansion of the interstitial space as reflected by the body weight, and expansion of the blood volume agree with the general mechanisms for volume expansion proposed previously.<sup>4, 20, 21</sup> Although the circulating doses of estrogen in the present study were greater than the physiologic, this should not detract from the value of the study because we were attempting to demonstrate an effect in a time length shorter than the normal duration of pregnancy.

**Plasma proteins and blood viscosity.** The second aim of the present study was to test the hypothesis that plasma protein concentrations and blood viscosity can reflect circulating estrogen concentrations. Human pregnancy is associated with marked changes in plasma protein concentrations. For instance, fibrinogen and globulin concentrations increase about 50% and 13%, respectively, while albumin decreases about 30%, so that the albumin-to-globulin ratio decreases about 36%.<sup>13, 26</sup> Also, in women using oral contraceptives the increase of fibrinogen and globulin concentrations has been alleged to result from the elevated circulating estrogen concentrations.<sup>27, 28</sup> In this study  $17\beta$ -estradiol infusion resulted in plasma protein changes similar to those observed during the course of gestation and in women on oral contraceptives (Fig. 3 and Table II).

Because both fibrinogen and globulin are known to increase both plasma and whole blood viscosity,<sup>29, 30</sup> the substantial viscosity increases observed in the present study are not surprising. Additionally, the moderate decline in albumin would enhance viscosity.<sup>29, 30</sup> These results vary somewhat from the findings reported in pregnancy. During pregnancy the changes in whole blood viscosity are more complex than those in plasma, because the hematocrit decreases as pregnancy pro-

ceeds. These hematocrit changes are superimposed on any baseline changes in plasma viscosity. Because a declining hematocrit decreases whole blood viscosity, this effect counteracts an increase resulting from plasma protein changes. Such an interaction between a rising plasma viscosity and a falling hematocrit would account for the decline in whole blood viscosity that most investigators have noted as term approaches.<sup>12, 31-34</sup>

Not all authors are, however, in agreement on whether whole blood viscosity declines during pregnancy.<sup>35</sup> At least part of this disagreement can be attributed to differences in the shear rate at which the viscosity determinations were carried out. During measurements made at low shear rates the red cells interact in such a fashion that the apparent whole blood viscosity is increased. Increases in fibrinogen enhance this aggregation and thus increase the measured viscosity of whole blood in at least two ways: The first is by directly enhancing plasma viscosity. The second is by enhancing cell-cell interaction, particularly at low shear rates. Capillary viscometers such as the one employed in this study are inherently high shear devices and as such are quite insensitive to cell-cell interactions. Because of this characteristic they respond to the increase in plasma viscosity that a heightened fibrinogen will cause but are unaffected by any viscosity increase resulting from fibrinogen-enhanced red cell aggregation.

Measurements of plasma viscosity are more straightforward, and the results obtained are essentially independent of the shear rate applied, because cell-cell interactions are excluded. Furthermore, in a series of measurements made over time there are no variations in hematocrit to complicate the interpretation. It is thus surprising that during the course of pregnancy some authors report no change or a slight increase,<sup>36</sup> others report a decrease,<sup>34</sup> and still others claim to have ob-

served an increase during early pregnancy followed by a decline at term.<sup>31</sup>

From this confusion in the literature it is clear that no experimental model, even a complex one, will succeed in reproducing such discordant results. The present studies do, however, provide strong evidence that estrogen, in isolation, produces a marked rise in fibrinogen and globulin, a moderate decrease in albumin, and an unequivocal enhancement of plasma viscosity. As such, the findings may help to sort out the effects of the hormonal changes that occur in women taking oral contraceptive medication.<sup>37</sup> They may also, if due respect is accorded the more complex hormonal interplay surrounding pregnancy, provide some understanding of both the normal physiology and the pathophysiology of this state.<sup>38-40</sup>

In conclusion, estrogen administration reproduced several of the hemodynamic changes observed in human pregnancy. A perfect reproduction of these alterations cannot be expected because of differences in species and the overall hormonal milieu.

We propose that during pregnancy increasing estrogen concentrations decrease systemic vascular resistance and mean arterial pressure, with increased glomerular filtration rate. Estrogen may also affect the lymphatic vessels and interstitial space compliance, as evidenced by the fact that most of the fluid retained is in the interstitial space. These changes result in expanded plasma and interstitial fluid volumes. A possible mechanism for these changes would be as follows: estrogen results in increases in renin, angiotensin, and aldosterone with resultant fluid retention. Estrogen administration has been shown to decrease the number of angiotension receptors and to increase the amount of vasodilatory prostaglandins in the uterine vein of castrated rabbits treated with estrogen. A combination of both factors would result in decreased tone in blood and lymphatic vessels, with decreased arterial pressure and increased heart rate. Uteroplacental vascular shunts may also contribute to these changes.

We thank P. Steven Dale, Elsie LaSalle, and James C. Westengard for technical assistance.

## REFERENCES

1. Longo LD, Hardesty JS. Maternal blood volume: measurement, hypothesis of control, and clinical considerations. *Rev Perinatol Med* 1984;5:1-28.
2. Assali NS, Brinkman CR III. Disorders of maternal circulatory and respiratory adjustments. In: Assali NS, ed. *Pathophysiology of gestation; vol 1. maternal disorders*. New York: Academic Press, 1972:269-353.
3. Guyton AC, Cowley AW Jr, Young DB, Coleman TG, Hall JE, DeClue JW. Integration and control of circulatory function. In: Guyton AC, Cowley AW Jr, eds. *Cardiovascular physiology II; vol 9*. Baltimore: University Park, 1976:341-85.
4. Longo LD. Maternal blood volume and cardiac output during pregnancy: a hypothesis of endocrinologic control. *Am J Physiol* 1983;245:R720-9.
5. Brace RA. Blood volume and its measurement in the chronically catheterized sheep fetus. *Am J Physiol* 1983; 224:H487-94.
6. Weichselbaum TE. An accurate and rapid method for the determination of proteins in small amounts of blood serum and plasma. *Am J Clin Pathol* 1946;16:40-9.
7. Rodkey FL. Direct spectrophotometric determination of albumin in human serum. *Clin Chem* 1965;11:478-87.
8. Abraham GE, Manlimos FS, Garza R. Radioimmunoassay of steroids. In: Abraham GE, ed. *Handbook of radioimmunoassay*. New York: Marcel Dekker, 1977:591.
9. Pritchard JA, MacDonald PC, Gant NF. *Williams' obstetrics*. 17th ed. Norwalk, Connecticut: Appleton-Century-Crofts, 1985:191-6.
10. Burwell CS. The influence of pregnancy on the course of heart disease. *South Med J* 1936;29:1194-7.
11. Burwell CS, Strayhorn WD, Flickinger D, Corlette MB, Bowerman EP, Kennedy JA. Circulation during pregnancy. *Arch Intern Med* 1938;62:979-1003.
12. Cohen ME, Thomson KJ. Studies on the circulation in pregnancy. I. The velocity of blood flow and related aspects of the circulation in normal pregnant women. *J Clin Invest* 1936;15:607-25.
13. Mack HC. The plasma proteins. *Clin Obstet Gynecol* 1960;3:336-48.
14. Kaulhausen H, Weyand C. Evidence for two independent effects of oestradiol benzoate on the renin-angiotensin-aldosterone system. *Acta Endocrinol* 1982;100:77-84.
15. Greiss FC Jr, Marston EL. The uterine vascular bed: effect of estrogens during ovine pregnancy. *AM J OBSTET GYNECOL* 1965;93:720-2.
16. Resnik R, Killam AP, Battaglia FC, Makowski EL, Meschia G. The stimulation of uterine blood flow by various estrogens. *Endocrinology* 1974;94:1192-6.
17. Matsuura S, Naden RP, Gant NF Jr, Parker CR Jr, Rosenfeld CR. Effect of volume expansion on pressor response to angiotensin II in pregnant ewes. *Am J Physiol* 1981;240:H908-13.
18. Antonini R, Valenzuela G, Harper MJK. Uterine blood flow and prostaglandin levels in ovariectomized rabbits. *J Reprod Fertil* 1978;54:349-56.
19. Smeaton TC, Andersen GJ, Fulton IS. Study of aldosterone levels in plasma during pregnancy. *J Clin Endocrinol Metab* 1977;44:1-7.
20. Hall JE, Granger JP, Hester RL, Coleman TG, Smith MJ Jr, Cross RB. Mechanism of escape from sodium retention during angiotensin II hypertension. *Am J Physiol* 1984;246:F627-34.
21. Hall JE, Granger JP, Smith MJ Jr, Prement AJ. Role of renal hemodynamics and arterial pressure in aldosterone "escape." *Hypertension* 1984;6(suppl 1):1183-92.
22. Young DB. Neural control of fluid volumes: volume receptors and autonomic control. In: Guyton AC, Taylor AE, Granger HJ, eds. *Circulatory physiology II: dynamics and control of the body fluids*. Philadelphia: WB Saunders, 1975:262-73.
23. Olsen ME, Hall JE, Montani JP, Guyton AC, Langford HG, Cornell JE. Mechanisms of angiotensin II natriuresis and antinatriuresis. *Am J Physiol* 1985;249:F299-307.
24. Metcalfe J, Parer JT. Cardiovascular changes during pregnancy in ewes. *Am J Physiol* 1966;210:821-5.
25. Challis JRG. Sharp increase in free circulating oestrogens immediately before parturition in sheep. *Nature* 1971;229:208.
26. Foley ME, Isherwood DM, McNicol GP. Viscosity, haematocrit, fibrinogen and plasma proteins in maternal and cord blood. *Br J Obstet Gynaecol* 1978;85:500-4.
27. Laragh JH, Sealey JE, Ledingham JGG, Newton MA. Oral contraceptives: renin, aldosterone, and high blood pressure. *JAMA* 1967;201:918-22.

28. Notelovitz M, Kitchens CS, Coone L, McKenzie L, Carter R. Low-dose oral contraceptive usage and coagulation. *AM J OBSTET GYNECOL* 1981;141:71-5.
29. Barth WF. Viscosimetry of serum in relation to the serum globulins. In: Sunderman FW, Sunderman FW Jr, eds. *Serum proteins and the dysproteinemias*. Philadelphia: JB Lippincott, 1964:102-9.
30. Harkness J. The viscosity of human blood plasma: its measurement in health and disease. *Biorheology* 1971;8:171-93.
31. Buchan PC, MacDonald HN. Rheological studies in obstetrics and gynecology. In: Lowe CDO, Barbenel JC, Forbes CD, eds. *Clinical aspects of blood viscosity and cell deformability*. New York: Springer-Verlag, 1981:175-92.
32. Dintenfass L. On changes in aggregation of red cells, blood viscosity, and plasma viscosity during normal gestation. *Clin Hemorrhheol* 1982;2:175-88.
33. Esiaschwili I. Blutgerinnung und Viskosität in der Schwangerschafts und Nachgeburtsperiode. *Zentralbl Gynakol* 1933;57:2717-21.
34. Hamilton HFH. Blood viscosity in pregnancy. *J Obstet Gynaecol Br Emp* 1950;57:530-8.
35. Heilmann L, Mattek C, Kurz E. Rheologische Veränderungen des Blutes in der normalen und pathologischen Schwangerschaft und deren Einfluss auf die Sauerstoffdiffusion. *Arch Gynecol* 1977;223:283-98.
36. Eastham RD. Plasma viscosity estimation in the improved haematological screening of antenatal patients. *J Obstet Gynaecol Br Comm* 1965;72:763-4.
37. Meade TW. Oral contraceptives, clotting factors, and thrombosis. *AM J OBSTET GYNECOL* 1982;142:758-61.
38. Hobbs JB, Oats JN, Palmer AA, et al. Whole blood viscosity in preeclampsia. *AM J OBSTET GYNECOL* 1982;142:288-92.
39. Matthews JD, Mason TW. Plasma viscosity and preeclampsia. *Lancet* 1974;2:409.
40. Thorburn J, Drummond MM, Whigham KA, et al. Blood viscosity and haemostatic factors in late pregnancy, preeclampsia and fetal growth retardation. *Br J Obstet Gynaecol* 1982;89:117-22.

---

## Pregnancy-induced hypertension: Development of a model in the pregnant sheep

Craig D. Thatcher, D.V.M., Ph.D., and James C. Keith, Jr., D.V.M., Ph.D.

Blacksburg, Virginia

Selected hemodynamic, renal, and biochemical parameters were assessed in chronically instrumented third-trimester pregnant ewes and in the same ewes after induction of pregnancy toxemia. Ewes with induced pregnancy toxemia developed hypertension, proteinuria, ketonuria, decreased glomerular filtration rate, decreased cardiac output, and decreased left uterine artery blood flow. Histological and transmission electron microscopy revealed the development of renal morphologic changes consistent with those observed in human pregnancy-induced hypertension. These studies have elucidated that pregnancy-induced hypertension can be produced experimentally in the pregnant ewe. Furthermore, the pathophysiologic features of ovine pregnancy toxemia are similar to those of human preeclampsia, and therefore the sheep provides a suitable animal model to study the human condition, which still remains a major complication of pregnancy, jeopardizing both mother and fetus. (*AM J OBSTET GYNECOL* 1986;155:201-7.)

**Key words:** Pregnancy-induced hypertension, sheep, preeclampsia, renal

Toxemia states of pregnancy are known to occur in various animal species although none to date has been

found to be a precise model for human preeclampsia. The human condition is a hypertensive disorder peculiar to pregnancy and has been associated with proteinuria,<sup>1</sup> a decrease in blood volume,<sup>2</sup> thrombocytopenia,<sup>3</sup> uteroplacental ischemia,<sup>4</sup> prostaglandin deficiency,<sup>5,6</sup> increased thromboxane A<sub>2</sub> production,<sup>7</sup> increased activity of the sympathetic nervous system<sup>8</sup> and the renin-angiotensin system,<sup>9</sup> and renal morphologic changes.<sup>10</sup>

Several investigators have compared ovine pregnancy toxemia to human pregnancy-induced hypertension. In the ewe, the toxic syndrome of pregnancy demonstrates a close resemblance to human preeclampsia with proteinuria and uteroplacental isch-

*From the Division of Veterinary Biology and Clinical Studies, Virginia-Maryland College of Veterinary Medicine, Virginia Polytechnic Institute and State University.*

*This work was supported by Virginia-Maryland College of Veterinary Medicine's New Investigator Grant Program, Virginia Polytechnic Institute and State University.*

*Presented at the Sixty-fifth Conference of Research Workers in Animal Disease, Chicago, Illinois, November 12-13, 1984.*

*Received for publication August 29, 1985; revised January 24, 1986; accepted March 7, 1986.*

*Reprint requests: Craig D. Thatcher, D.V.M., Ph.D., Division of Veterinary Biology and Clinical Studies, College of Veterinary Medicine, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061.*