

Serotonin Stimulates Adenosine 3',5'-Monophosphate Accumulation in Parathyroid Adenoma*

DONALD ZIMMERMAN,[†] HANNA E. ABBOUD,[†] LAWRENCE E. GEORGE,[‡]
ANTHONY J. EDIS, AND THOMAS P. DOUSA

Nephrology Research Laboratories, Division of Nephrology and Internal Medicine, Departments of Surgery and Physiology and Biophysics, Mayo Clinic and Foundation, Mayo Medical School, Rochester, Minnesota 55901

ABSTRACT. The effect of serotonin on cAMP accumulation in parathyroid adenoma tissue from patients with primary hyperparathyroidism was studied *in vitro*. Incubation with 10^{-5} M serotonin elicited a marked increase (of 90–150%) in cAMP content in slices of parathyroid adenoma tissue. This stimulatory effect of serotonin was already apparent after 2 min of incubation; stimulation by serotonin was dose dependent, with the highest stimulation being achieved at 10^{-4} M serotonin. The serotonin antagonists, methylsergide and cinanserin, in concentrations equimolar to serotonin completely blocked the stimu-

latory effect of serotonin on cAMP increase. The serotonin content in surgically removed parathyroid adenoma tissue, as determined by fluorometric assay, was 6.4 ± 1.2 pmol/mg wet wt ($\sim 0.8 \times 10^{-5}$ M). The present observations demonstrate that parathyroid adenoma tissue has a high content of serotonin, and serotonin stimulates cAMP accumulation in this tissue. Since cAMP acts as a mediator of parathyroid hormone (PTH) release, our results suggest that serotonin could be one of the factors regulating PTH secretion and/or contributing to PTH hypersecretion in various forms of primary hyperparathyroidism. (*J Clin Endocrinol Metab* 51: 1274, 1980)

SEROTONIN-producing neoplasms have been found in association with hyperparathyroidism in some patients with multiple endocrine neoplasia types 1 (1–11) and 2a (12). While it is not known whether there is any causal association between elevated blood serotonin levels in these patients (13, 14) and hypersecretion of parathyroid hormone (PTH), many observations *in vivo* and *in vitro* provide strong evidence that a variety of other biogenic amines, such as β -adrenergic agonists (15, 16), dopamine (17), and histamine (18), are potent stimuli of PTH secretion by parathyroid tissue. This effect of biogenic amines as well as of other stimuli of PTH secretion (15, 19, 20) appears to be mediated by increases in cAMP generation and accumulation of this cyclic nucleotide in parathyroid tissue (15, 19–21). That increased cAMP generation in this setting is due to activation of adenylate cyclase is supported by the observation that human parathyroid tissue contains adenylate cyclase which can be stimulated by β -adrenergic agonists (20, 22).

Like various other biogenic amines, serotonin has been shown in several cell types and organ systems to stimulate cAMP formation (23, 24) and/or to activate adenylate cyclase (25, 26). In view of these considerations, we addressed ourselves to the question of whether serotonin can stimulate cAMP formation and accumulation in human hyperparathyroid tissue and, by extension, whether it may potentially serve as a PTH secretagogue. To further explore the possibility of a pathogenic role for serotonin in hyperparathyroidism, we also measured serotonin levels in parathyroid adenoma tissue. The results demonstrate that serotonin is a potent agonist of cAMP formation in parathyroid adenoma tissue and that this tissue itself contains a considerable amount of serotonin.

Materials and Methods

Tissue was obtained from parathyroid adenomata of patients who underwent partial parathyroidectomy for primary hyperparathyroidism diagnosed by clinical criteria. The average age of the patients (four males and two females) was approximately 40 yr. After surgical removal of the parathyroid tissue, part of the tissue specimen was immediately chilled in ice-cold 0.9% sodium chloride solution. All subsequent preparatory steps were performed at 0–2 C. After the tissue was trimmed of adipose and connective tissue, a portion was taken for histological examination. In some patients, another part of the tissue block was removed, immediately frozen on dry ice, and stored for determination of serotonin concentration. The rest of the

Received April 21, 1980.

Address all correspondence and requests for reprints to: Thomas P. Dousa, M.D., Mayo Clinic, 901 Guggenheim Building, Rochester, Minnesota 55901.

* Presented at the 62nd Annual Meeting of The Endocrine Society, Washington, D.C., 1980. This work was supported by Research Grant AM-16105 from the NIAMDD, NIH, and the Mayo Foundation.

[†] Recipient of Renal Diseases Training Grant Fellowship AM-06173.

[‡] Student at the Mayo Medical School.

tissue was quickly transferred in cold buffer to the laboratory, and slices were prepared with a Stadie-Riggs tissue slicer. Slices of parathyroid tissue were extensively washed in ice-cold modified Krebs-Ringer solution, and the effect of serotonin on cAMP accumulation was determined using basically the design described in our previous studies of serotonin's effect on cAMP accumulation in renal tissue (24). Slices were divided into individual flasks (~5–15 mg wet wt/incubation) and incubated in 350 μ l modified Krebs-Ringer buffer (KRB) (24) at 37 C for 20 min. After the preincubation period, 50 liters of 5 mM 1-methyl-3-isobutyl xanthine and of other agents dissolved in the same buffer (KRB) were added within 6–10 sec up to a final volume of 0.5 ml. After the addition of these agents, slices were incubated for an additional 5 min, unless specified otherwise in Results. The incubation was stopped by the addition of 0.5 ml 10% trichloroacetic acid (TCA) and homogenization of the tissue. Aliquots of the TCA homogenate were taken for determination of protein. After the addition of trace amounts of [3 H]cAMP to monitor recovery of the nucleotide (27), precipitated protein was removed by centrifugation, and TCA from the supernatant was removed by subsequent extraction with ethyl ether. cAMP content in the extract was determined by RIA after acetylation with acetic anhydride and triethanol amine using the procedure described in full detail in our previous studies (24, 27). Validation of the RIA used was described in detail elsewhere (27); neither serotonin nor its antagonists interferes with cAMP determination (24).

The effects of serotonin and other agents were tested in each experiment in such a way that control and experimental tissue slices from the same block of tissue were simultaneously incubated with and without added agents. The protein and cAMP contents were determined in experimental and control homogenates (with and without drugs, respectively) at the same time using the same standards. The content of cAMP in the incubation mixture was expressed in relation to tissue protein in each incubation as picomoles of cAMP per mg tissue protein.

With use of the present experimental design, cAMP which accumulated in the tissue and that which escaped into the medium were determined together. The determination of tissue protein, RIA for cAMP, and sources of agents and biochemicals were described in detail in our previous communications (24, 27, 28).

For determination of tissue levels of serotonin, the frozen tissue was homogenized and extracted as described by Snyder *et al.* (29), except that tissues were homogenized in 4 ml ice-cold 0.1 M HCl, as employed by Bogdanski *et al.* (30), rather than in 8 ml 0.4 N perchloric acid. Serotonin content was determined by fluorometric assay and expressed as picomoles of serotonin per mg tissue. To ensure validity of measurements, in preliminary experiments, serotonin content was measured with this assay in several rat tissues (intestine, spleen, and liver). The serotonin content found in these preliminary experiments agreed with values of rat tissue serotonin reported from other laboratories (31).

Serotonin (5-hydroxytryptamine creatine sulfate) was purchased from Sigma Chemical Co. (St. Louis, MO); 1-methyl-3-isobutyl-xanthine was obtained from Aldrich Chemical Co. (Milwaukee, WI). Methysergide maleate was a gift from Sandoz

Pharmaceuticals (Hanover, NJ), and cinanserin hydrochloride was a gift from Squibb and Sons, Inc. (Princeton, NJ). All agents were dissolved before use in Krebs-Ringer phosphate buffer.

Results

Serotonin (10^{-5} M) consistently caused a marked increase in the cAMP level in incubated parathyroid adenoma tissue of examined patients (Table 1). The cAMP increase was dependent on the dose of serotonin; the highest stimulation was observed at a concentration of 10^{-4} M (Fig. 1). Serotonin caused a significant cAMP increase as early as 2 min after addition, and the effect persisted up to 30 min (Table 2). To ascertain that serotonin's action was specific for this biogenic amine, the effect of antiserotonin drugs was tested. Although methylsergide and cinanserin (10^{-5} M) alone had no effect on cAMP levels in parathyroid tissue (data not shown), these drugs completely blocked the stimulatory action of serotonin on cAMP levels (Table 3).

The average serotonin content (mean \pm SE; $n = 6$) determined in the frozen tissue samples was 6.4 ± 1.2 pmol/mg tissue. Assuming that parathyroid adenoma tissue contains about 70–80% water, as tissues do in general (32), the actual concentration of serotonin in parathyroid tissue fluid was about 0.8×10^{-5} M. This concentration of serotonin is comparable to the serotonin content of human ileum, lower than that in human spleen, and higher than that in most central nervous system structures (except the hypothalamus and pineal gland), as determined by the same method (31).

Discussion

In the present study, serotonin was found to elicit marked and dose-dependent stimulation of cAMP accumulation in parathyroid adenoma tissue. Inhibition of this stimulatory effect by specific serotonin antagonists

TABLE 1. Effect of serotonin on cAMP levels in parathyroid tissue from hyperparathyroid patients

Case no.	cAMP (pmol/mg protein)		
	Basal (no additions)	Serotonin (10^{-5} M)	$\Delta\%$ ^a
1	27.2 \pm 3.4 ^b	67.7 \pm 4.9	+149
2	12.0 \pm 0.8	28.8 \pm 2.2	+140
3	20.8 \pm 2.0	39.8 \pm 10.3	+91
4	18.6 \pm 1.0	35.6 \pm 4.7	+94
5	25.8 \pm 3.4	61.2 \pm 11.0	+137
Mean \pm SE	20.9 \pm 2.7	46.6 \pm 7.6 ^c	122.2 \pm 12 ^c

^a Percent increase from basal.

^b Each value denotes the mean \pm SE of triplicates.

^c Significantly different from basal and/or $\Delta\%$ is significant ($P < 0.005$, by paired t test).




FIG. 1. Effect of various concentrations of serotonin on cAMP accumulation in parathyroid adenoma tissue. For experimental details, see *Materials and Methods*. Each bar denotes the mean \pm SE of several incubations; the number of incubations (n) is indicated at the bottom of the bars. *, Values significantly ($P < 0.05$ or higher level of significance by t test) higher than basal value (no serotonin added).

TABLE 2. Effect of serotonin on parathyroid tissue levels of cAMP at different time intervals. (For details see *Methods*)

Time	cAMP (pmol/mg protein)		$\Delta\%$ ^a	P ^b
	Basal (no addition)	Serotonin (10 ⁻⁵ M)		
2 (5) ^c	13.7 \pm 1.4 ^d	26.9 \pm 4.1	+96	<0.01
5 (5)	17.1 \pm 3.4	44.4 \pm 9.7	+126	<0.05
15 (5)	22.6 \pm 5.3	60.0 \pm 14.0	+131	<0.05
30 (5)	20.3 \pm 4.3	44.1 \pm 3.8	+137	<0.05

^a Percent increase from basal.

^b For significance of difference of serotonin-stimulated values from the corresponding basal values during the same time interval (by t test).

^c The number of samples (incubations) is indicated in parentheses. For details see *Materials and Methods*.

^d Mean \pm SE; picomoles of cAMP per mg protein.

indicates that serotonin most likely increased cAMP levels by acting on serotonin receptors in a manner analogous to serotonin's actions on some other tissues (23–26). Since cAMP appears to be an intracellular mediator of PTH secretion (15, 19, 20), and since physiological and pharmacological stimuli which increase cAMP accumulation in parathyroid tissue invariably stimulate PTH secretion (15, 19, 20), it is reasonable to expect that

TABLE 3. Effect of methylsergide and cinanserin on stimulatory action of serotonin in parathyroid tissue

Condition	n	cAMP (pmol/mg protein)
Basal (no additions)	6	16.4 \pm 2.2 ^a
Serotonin (10 ⁻⁵ M)	6	34.3 \pm 5.3 ^b
Serotonin (10 ⁻⁵ M) + methylsergide (10 ⁻⁵ M)	6	15.9 \pm 2.2
Serotonin (10 ⁻⁵ M) + cinanserin (10 ⁻⁵ M)	5	13.3 \pm 4.9

^a Values given are the mean \pm SE. Number of observations (incubations).

^b Value significantly ($P < 0.02$, by t test) different from the basal value or values with methylsergide and cinanserin.

serotonin-stimulated cAMP accumulation will also result in PTH release from parathyroid adenoma tissue.

The present observations, therefore, point to the possibility that serotonin may play a role in the physiological regulation of PTH secretion or at least in pathological PTH hypersecretion in hyperparathyroid states. Since PTH secretion may, in part, be under neural control (33–35), it is conceivable that serotonin may serve as one of the neurotransmitters in this neural regulatory pathway (36). Under normal circumstances, circulating serum levels of serotonin were reported to be approximately 0.8×10^{-6} M (14), which is somewhat lower than the lowest concentration shown in our experiments to stimulate cAMP accumulation in parathyroid tissue. However, in the carcinoid syndrome, serum levels of serotonin may reach levels in the range of about 10^{-5} M (14), a concentration in which serotonin directly stimulates cAMP formation in parathyroid glands, as suggested in the present experiments. Thus, it should be considered a possibility that the pathologically high blood serotonin levels encountered in some cases of carcinoid syndrome may contribute to PTH hypersecretion and subsequent hyperparathyroidism which have been described in several instances (1–12).

Parathyroid adenoma tissue contains serotonin in a concentration within the range capable of stimulating cAMP accumulation. Whether these relatively high levels of serotonin result from a high rate of serotonin synthesis and storage in parathyroid tissue or, alternatively, are due to uptake of serotonin from the blood perfusing the parathyroid glands remains undetermined. It is of interest that parathyroid adenomata contain numerous argentaffin cells (37, 38), a major site of serotonin storage in various tissues (39), while normal parathyroid tissue is devoid of these cells (37, 38). It should be realized that serotonin in parathyroid adenoma tissue might be sequestered in certain cell populations or cell compartments distinct from the serotonin receptor-adenylate cyclase system in plasma membranes. It is possible that serotonin should be released from such putative

stores in response to various (including neural) stimuli before it acts on the serotonin receptor and stimulates adenylate cyclase. Alternatively, serotonin synthesized *de novo* in parathyroid adenoma tissue may have direct access to serotonin-sensitive cAMP system. However, regardless of the serotonin localization in parathyroid adenoma tissue, its presence in the range of concentrations capable of stimulating cAMP accumulation (and, by extension, PTH secretion) suggests that serotonin-stimulated elevation of cAMP may contribute to anomalously high PTH secretion in hyperparathyroid states, albeit the details of such mechanism remain to be clarified.

In conclusion, although many points remain to be clarified, the present results provide evidence that human parathyroid adenoma tissue contains a cAMP system sensitive to stimulation by serotonin, and that the tissue itself contains considerable amounts of this biogenic amine. If serotonin indeed contributes, as postulated, by increasing cAMP levels in parathyroid tissue to control PTH secretion and, especially, to produce hypersecretion in pathological situations, blockade of serotonin's action on parathyroid tissue by specific antagonists (40) may be considered as a potential pharmacological tool to inhibit PTH secretion in clinical and experimental situations.

Acknowledgments

The technical assistance of Mrs. Lise Ou and Mr. Tom Haugen and the secretarial assistance of Ms. Bonnie Becker and Mrs. Ardith Walker were much appreciated.

References

- Samaan NA, Hickey RC, Bedner TD, Ibanez ML 1975 Hyperparathyroidism and carcinoid tumor. *Ann Intern Med* 82:205
- Dry J, Pradalier A, Herman D, Leynadier F 1976 La polyadenomatous endocrinienne avec tumeur carcinoïde. *Sem Hop Paris* 52: 1691
- Wick MR, Scott RE, Li C-Y, Carney JA 1980 Carcinoid tumor of the thymus: a clinopathologic report of seven cases with a review of the literature. *Mayo Clin Proc* 55:246
- Lokich JJ, Li F 1978 Carcinoid of the thymus with hereditary hyperparathyroidism. *Ann Intern Med* 89:364
- Manes JL, Taylor HB 1973 Thymic carcinoid in familial multiple endocrine adenomatosis. *Arch Pathol* 95:252
- Rosai J, Higa E, Davie J 1972 Mediastinal endocrine neoplasm in patients with multiple endocrine adenomatosis: a previously unrecognized association. *Cancer* 29:1075
- Hill DM 1967 Pluriglandular syndrome: multiple endocrine neoplasia. *Proc R Soc Med* 60:650
- Underdahl LO, Woolner LB, Black BM 1953 Multiple endocrine adenomas: report of 8 cases in which the parathyroids, pituitary and pancreatic islets were involved. *J Clin Endocrinol Metab* 13:20
- Greenbaum D 1960 Multiple familial parathyroid adenomata. *Proc R Soc Med* 53:903
- Williams ED, Celestin CR 1962 The association of bronchial carcinoid and pluriglandular adenomatosis. *Thorax* 17:20
- Escovitz WE, Reingold IM 1961 Functioning malignant bronchial carcinoid with Cushing's syndrome and recurrent sinus arrest. *Ann Intern Med* 54:1248
- Moertel CG, Behrs OH, Woolner LB, Tyce GM 1965 Malignant carcinoid syndrome associated with non-carcinoid tumors. *N Engl J Med* 273:244
- Sjoerdsma A, Weissbach H, Underfriend S 1956 A clinical, physiologic and biochemical study of patients with malignant carcinoid (argentaffinoma). *Am J Med* 20:520
- Feldman JM 1978 Serotonin metabolism in patients with carcinoid tumors: incidence of 5-hydroxytryptophan-secreting tumors. *Gastroenterology* 75:1109
- Habener JF, Potts JT 1976 Chemistry, biosynthesis, secretion and metabolism of parathyroid hormone. In: Greep RO, Astwood EB (eds) *Handbook of Physiology*. American Physiological Society, Washington DC, p 313
- Kukreja SC, Hargis GK, Rosenthal IM, Williams GA 1973 Pheochromocytoma causing excessive parathyroid hormone production and hypercalcemia. *Ann Intern Med* 79:838
- Brown EM, Carroll RJ, Aurbach GD 1977 Dopaminergic stimulation of cyclic accumulation and parathyroid hormone release from dispersed bovine parathyroid cells. *Proc Natl Acad Sci USA* 74: 4210
- Williams GA, Longley RS, Hargis GK, Bowser EN, Kukreja SC, Johnson PA, Jackson BL, Kawahara WJ 1979 Effect of histamine on secretion of parathyroid hormone. *Clin Res* 27:704A (Abstract)
- Brown EM, Gardner DG, Windeck RA, Aurbach GD 1978 Relationship of intracellular 3',5'-adenosine monophosphate accumulation to parathyroid hormone release from dispersed bovine parathyroid cells. *Endocrinology* 106:133
- Peck WA, Klahr S 1979 Cyclic nucleotides and the control of parathyroid hormone secretion. *Adv Cyclic Nucleotide Res* 10:75
- Abboud HE, Zimmerman D, Edis AJ, Dousa TP 1980 Histamine and parathyroid adenoma: effect on cyclic AMP accumulation. *Clin Res* 28:515A
- Rodriguez HJ, Morrison A, Slatopolsky E, Klahr S 1978 Adenylate cyclase of human parathyroid gland. *J Clin Endocrinol Metab* 47: 319
- Kebabian JW 1977 Biochemical regulation and physiological significance of cyclic nucleotides in the nervous system. *Adv Cyclic Nucleotide Res* 8:421
- Shah SV, Northrup TE, Hui YSF, Dousa TP 1979 Action of serotonin (5-hydroxytryptamine) on cyclic nucleotides in glomeruli of rat renal cortex. *Kidney Int* 15:463
- Higgins WJ 1977 5-Hydroxytryptamine-induced tachyphylaxis of the molluscan heart and concomitant desensitization of adenylate cyclase. *J Cyclic Nucleotide Res* 3:293
- Enjalbert S, Bourgoin S, Hanon M, Adrien J, Bockaert J 1978 Postsynaptic serotonin-sensitive adenylate cyclase in the central nervous system. I. Development and distribution of serotonin and dopamine-sensitive adenylate cyclase in rat and guinea pig brain. *Mol Pharmacol* 14:2
- Torres VE, Northrup T, Edwards RM, Shah SV, Dousa TP 1978 Modulation of cyclic nucleotides in isolated rat glomeruli: role of histamine, carbamylcholine, parathyroid hormone and angiotensin II. *J Clin Invest* 62:1334
- Abboud HE, Shah SV, Dousa TP 1979 Effects of dexamethasone on cyclic nucleotide accumulation in glomeruli. *J Lab Clin Med* 94: 708
- Snyder SH, Axelrod J, Zweig M 1965 A sensitive and specific fluorescence assay for tissue serotonin. *Biochem Pharmacol* 14:831
- Bogdanski DF, Pletscher A, Brodie BB, Udenfriend S 1956 Identification and assay of serotonin in brain. *J Pharmacol Exp Ther* 117: 82
- Essman WB 1978 Serotonin distribution in tissues and fluids. Essman WB (ed) *Serotonin in Health and Disease*. Spectrum Publications, New York, vol 1:15.
- Giese AC 1973 *Cell Physiology*. Saunders, Philadelphia, p 53
- Altenahr E 1971 Electron microscopical evidence for innervation of chief cells in human parathyroid gland. *Experientia* 27:1077
- Yeghiayan E, Rojo-Ortega JM, Genest J 1972 Parathyroid vessel innervation: an ultrastructural study. *J Anat* 112:137
- Vora NM, Kukreja SC, Bowser EN, Jackson PA, Hargis GK, Williams GA, Effect of total sympathetic ablation on serum parathyroid hormone and calcitonin in the rat. Program of the 61st Annual Meeting of The Endocrine Society, Anaheim CA, 1979, p

83

36. Gershon MD, Dreyfus CF, Pickel VM, Jon TH, Reis DJ 1977 Serotonergic neurons in the peripheral nervous system: identification in gut by immunohistochemical localization of tryptophan hydroxylase. *Proc Natl Acad Sci USA* 74:3086
37. Ljungberg O 1972 Argentaffin cells in human thyroid and parathyroid and their relationship to c-cells and medullary carcinoma. *Acta Pathol Microbiol Scand [A]* 80:589
38. Ljungberg O, Dymling JF 1972 Pathogenesis of c-cell neoplasia in thyroid gland c-cell proliferation in a case of chronic hypercalcemia. *Acta Pathol Microbiol Scand [A]* 80:577
39. Pearse AGE 1972 *Histochemistry Theoretical and Applied*. Williams and Wilkins, Baltimore, p 1101
40. Gerattini S, Samanin R 1978 Drugs affecting serotonin: a survey. In: Essman WB (ed) *Serotonin in Health and Disease*. Spectrum Publications, New York, vol 2:247

II International Meeting of Neuroendocrinology of Barcelona

II International Meeting of Neuroendocrinology of Barcelona "*Functional Pituitary Tumours: Diagnostic and Treatment*" sponsored by the International Society of Neuroendocrinology Barcelona, January 23rd and 24th, 1981.

The scientific program will include the following topics: Neuroendocrine control of Gonadotropins and Prolactin secretion (Martini, L.—Italy); Pituitary adenomas: Pathologic aspects (Doniach, I.—UK); ACTH Secretor Tumours: Diagnostic and therapeutical aspects (Jeffcoate, W. J.—UK); Prolactin secreting tumours: Diagnostic aspects (Müller, E.—Italy); Acromegaly: Diagnostic and therapeutical aspects (Wass, J. A. H.—UK); Treatment with radiation of hypophysary tumours (López Ibáñez, P.—Spain); Surgical treatment of hypophysary tumours (Derome, P.—France).

For additional information, please address:

Dr. Isaac Levy
Servicio Endocrinología y Diabetes
Hospital Clínico
c/ Casanovas, 143
Barcelona 36 (Spain)