

for assessing such protection. Exposure of volunteers to UVA light, with and without a sun screen, is not practical in view of the time necessary to produce erythema with UVA light alone. Various alternative methods have therefore been developed—eg, photosensitisation of volunteers with a chemical such as psoralens; examination of another biological endpoint such as immediate pigment darkening; use of patients with photodermatoses that are highly sensitive to UVA; or use of an in-vitro system.<sup>4</sup>

One such system relies on a spectral measuring device and an artificial transpore membrane. By comparing transmission of UV light through the membrane with and without a sun screen, a transmission protection factor can be established at different wavelengths throughout the UVB and UVA range. The test is a cheap and convenient laboratory surrogate for in-vivo testing but it has not yet been fully validated and cannot be used for products that are alcohol-based or oil-based.<sup>9</sup>

A better method might be to use human epidermis instead of an artificial membrane. Epidermis could be obtained by raising suction blisters from surgical biopsy specimens or even from cadaveric skin, and would obviate the need for human or animal experimentation. This technique would provide doctors and manufacturers with a measure of UVA protection, and if patients or customers were confused by use of two numbers on a product, they could be told that SPF/B stands for burning and SPF/A stands for ageing, this explanation being sufficiently accurate for such a purpose. Equally, there is an urgent need for independent verification of company claims, so that doctors and patients can be satisfied of the UVB and UVA protection afforded by a prescribable formulation.<sup>10</sup>

Nevertheless the main concern for human health is the carcinogenic impact of UVB, and not the cosmetic effect of UVA. As ozone depletion quickens<sup>11</sup> the proportion of UVB to UVA will rise, so burning, snow blindness, cataracts, and skin cancer can be expected to increase, whereas UVA-induced problems will decline, not because the amount of UVA will change but because sunbathers will be driven indoors by the acute effects of UVB before much UVA exposure occurs. It has been calculated that, other things being equal, a 10% loss of total column ozone will increase the frequency of melanoma by 10–20%,<sup>12</sup> of basal cell carcinoma by 25–35%, and of squamous cell carcinoma by 50–60%.<sup>13,14</sup> There is less certainty about the relation between cataracts and UVB, but the US Environmental Protection Agency have calculated an increase of 6–10% in the prevalence of senile cataracts after a 10% loss of total column ozone.<sup>15</sup>

Whilst UVA protection is therefore an important issue, it is overshadowed by the implications for human health of ozone depletion. Nor are sun screens necessarily the best method of protection against the

deleterious effects of UV light—wearing a shirt or a hat is equally effective and in the case of cataract formation more effective than topical preparations. As Dr Robin Marks, Director of the Anti-Cancer Council of Victoria in Australia, has remarked “We are using one class of chemicals to destroy the ozone layer and another class to protect us from the consequences. Surely we can organise ourselves better than that”.

1. Anon. Shedding light on seasonal affective disorder. *Drug Ther Bull* 1991; **29**: 33–34.
2. van der Leun JC. Experimental photocarcinogenesis. In: Russell Jones R, Wigley W, eds. *Ozone depletion: health and environmental consequences*. Chichester: Wiley, 1989: 161–68.
3. Pitts D, Cullen A. Determination of infrared radiation levels for acute ocular cataractogenesis. *Arch Clin Exp Ophthalmol* 1981; **217**: 285–97.
4. Editorial. Measuring sunscreen protection against UVA. *Lancet* 1990; **ii**: 472.
5. Cole C, Forbes P, Davies R. An action spectrum for UV photocarcinogenesis. *Photochem Photobiol* 1986; **43**: 275–84.
6. Diffey BL. Tanning with ultraviolet A sunbeds. *Br Med J* 1990; **301**: 773–74.
7. Russell Jones R. Tanning with ultraviolet A sunbeds. *Br Med J* 1990; **301**: 986.
8. Hawk J. Ultraviolet A radiation: staying within the pale. *Br Med J* 1991; **302**: 1036–37.
9. Diffey BL, Robson J. A new substrate to measure sunscreen protection factors throughout the ultraviolet spectrum. *J Soc Cosmet Chem* 1989; **40**: 127–33.
10. Anon. Topical sunscreens. *Drug Ther Bull* 1990; **28**: 61–63.
11. Editorial. Ozone depletion quickens. *Lancet* 1991; **337**: 1132–33.
12. Elwood JM. Epidemiology of melanoma: its relationship to ultraviolet radiation and ozone depletion. In: Russell Jones R, Wigley W, eds. *Ozone depletion: health and environmental consequences*. Chichester: Wiley, 1989: 169–89.
13. Russell Jones R. Ozone depletion and cancer risk. *Lancet* 1987; **ii**: 443–446.
14. Russell Jones R. Consequences of human health of stratospheric ozone depletion. In: Russell Jones R, Wigley W, eds. *Ozone depletion: health and environmental consequences*. Chichester: Wiley, 1989: 207–27.
15. Environmental Protection Agency. *Assessing the risk of trace gases that can modify the atmosphere*. Washington, DC: US Environmental Protection Agency, 1987.

## Insulin resistance in puberty

Blood glucose control in diabetic children deteriorates when they reach puberty. Poor dietary compliance, emotional lability, and adolescent rebellion are undoubtedly contributing factors, but a report by Amiel and her co-workers<sup>1</sup> serves as a reminder of the physiological decrease in insulin sensitivity that occurs during puberty in normal individuals and in those with diabetes. The result is a compensatory increase in basal and stimulated insulin secretion in normal adolescents<sup>2</sup> and an increased insulin requirement in diabetic adolescents. This physiological response would be less of a puzzle during this time of change in body composition and alterations of the pituitary-gonadal axis<sup>3</sup> were it not for the return in adult life of levels of insulin sensitivity similar to those found in prepubertal children.<sup>4</sup>

Amiel and colleagues investigated the mechanism of these changes in normal prepubertal and pubertal individuals by use of the euglycaemic clamp, and found that there was an insensitivity to insulin of peripheral glucose disposal in muscle. The ability of insulin to suppress hepatic glucose production,

lipolysis, and ketogenesis was not changed by puberty in this study, but these are very insulin-sensitive processes that are often overwhelmed by higher rates of insulin infusion. That stimulation of uptake of branched chain aminoacids was also unimpaired suggests that there was no resistance to the anabolic effect of insulin.

What is the explanation? The concentrations of many circulating hormones, including androgens and growth hormone, and their binding proteins change during puberty so many possible changes coincide with altered insulin sensitivity.<sup>5,6</sup> Androgen concentrations rise throughout puberty and have a small effect on insulin sensitivity in adults.<sup>7</sup> However, there is no difference in insulin sensitivity between the sexes during puberty. In any case, adrenarche (the augmentation of adrenal cortical secretion, especially androgens, in both sexes at about 8 years) occurs well before any measurable rise in basal insulin concentrations,<sup>6</sup> and androgen concentrations remain increased into adult life, when insulin sensitivity returns to prepubertal values. The growth spurt during puberty may be more relevant. Integrated 24-hour growth hormone concentrations increase about two-fold during puberty,<sup>8</sup> and the hormone is known to have an anti-insulin effect, probably at a post-receptor level.<sup>9,10</sup> In addition, at all ages there is a correlation between serum insulin concentrations and plasma insulin-like growth factor 1 (IGF-1),<sup>6,11</sup> and IGF-1 is the main mediator of the tissue effects of growth hormone.

Is this insulin insensitivity physiologically important in puberty or is it an epiphenomenon? Insulin has well-recognised growth promoting effects and growth retardation is seen in poorly controlled diabetic children.<sup>12</sup> Fasting plasma insulin levels are closely related to height velocity in childhood.<sup>13</sup> Insulin may stimulate IGF-1 secretion directly,<sup>14</sup> but interest has tended to focus on its effects on the IGF binding proteins. The growth hormone dependent binding protein IGFBP-3 may act to stabilise the circulating pool of IGF-1, but the smaller growth hormone independent protein IGFBP-1 seems to inhibit the action of IGF-1 on tissues, fluctuates during the day, and is itself suppressed by insulin concentrations in the short term.<sup>15,16</sup> During puberty growth acceleration and sexual maturation are closely linked but by an unknown mechanism.<sup>17</sup> Integrated control of growth during childhood and puberty has been suggested—ie, insulin may mediate the relation between nutritional status and maturation via its coordinated effects on sex hormone binding globulin and IGF-1 binding proteins.<sup>18</sup> When linear growth ceases, insulin would seem to lose its role as a growth hormone and revert to a purely metabolic function, with the relation between insulin and IGFBP-1 disappearing in adult life.<sup>19</sup>

Thus it may be that the role of insulin changes during puberty, and that reduced sensitivity to the hypoglycaemic action of this hormone with a

compensatory rise in basal and stimulated insulin concentrations is permissive to its action as a growth-promoting hormone. It is also possible that a result of this decrease in insulin sensitivity is the well-recognised rise in frequency of insulin-dependent diabetes mellitus in early puberty.<sup>20</sup> Ensuring appropriate increases in insulin dose in adolescents who are dependent on an exogenous supply may be important not merely in protecting against later microvascular complications of diabetes but also in securing full growth potential.

1. Amiel SA, Caprio S, Sherwin RS, Plewe G, Haymond MW, Tamborlane WV. Insulin resistance of puberty: a defect restricted to peripheral glucose metabolism. *J Clin Endocrinol Metab* 1991; **72**: 277–82.
2. Smith CP, Archibald HR, Thomas JM, et al. Basal and stimulated insulin levels rise with advancing puberty. *Clin Endocrinol* 1988; **28**: 7–14.
3. Bloch CA, Clemons P, Sperling MA. Puberty decreases insulin sensitivity. *J Pediatr* 1987; **110**: 481–87.
4. Amiel SA, Sherwin RS, Simonson DC, Lauritano AA, Tamborlane WV. Impaired insulin action in puberty. A contributing factor to poor glycemic control in adolescents with diabetes. *N Engl J Med* 1986; **315**: 215–19.
5. Mauras N, Blizzard RM, Link K, Johnson ML, Rogol AD. Augmentation of growth hormone secretion during puberty: evidence for a pulse amplitude-modulated phenomenon. *J Clin Endocrinol Metab* 1987; **64**: 596–601.
6. Smith CP, Dunger DB, Williams AJ, et al. Relationship between insulin, insulin-like growth factor I, and dehydroepiandrosterone sulfate concentrations during childhood, puberty, and adult life. *J Clin Endocrinol Metab* 1989; **68**: 932–37.
7. Hale PJ, Wright JV, Nattrass M. Differences in insulin sensitivity between normal men and women. *Metabolism* 1985; **34**: 1133–38.
8. Hindmarsh P, Di Silvio L, Pringle PJ, Kurtz AB, Brook CGD. Changes in serum insulin concentration during puberty and their relationship to growth hormone. *Clin Endocrinol* 1988; **28**: 381–88.
9. Press M, Tamborlane WV, Sherwin RS. Importance of raised growth hormone levels on mediating the metabolic derangements of diabetes. *N Engl J Med* 1984; **310**: 810–15.
10. Rosenfeld RG, Wilson DM, Dollar LA, Bennett A, Hintz RE. Both human pituitary growth hormone and recombinant DNA-derived growth hormone cause insulin resistance at a postreceptor site. *J Clin Endocrinol Metab* 1982; **54**: 1033–38.
11. Caprio S, Plewe G, Diamond MP, et al. Increased insulin secretion in puberty: a compensatory response to reductions in insulin sensitivity. *J Pediatr* 1989; **114**: 963–67.
12. Holly JM, Amiel SA, Sandhu RR, Rees LH, Wass JA. The role of growth hormone in diabetes mellitus. *J Endocrinol* 1988; **118**: 353–64.
13. Hindmarsh PC, Matthews DR, Di Silvio L, Kurtz AB, Brook CGD. Relation between height velocity and fasting insulin concentration. *Arch Dis Child* 1988; **63**: 665–66.
14. Daughaday WH, Phillips LS, Mueller MS. The effects of insulin and growth hormone on the release of somatomedin by the isolated rat liver. *Endocrinology* 1976; **98**: 1214–19.
15. Suikkari AM, Koivisto VA, Rutanen EM, Yki-Jarvinen H, Karonen S-L, Seppala M. Insulin regulates the serum levels of low molecular weight insulin-like growth factor binding protein. *J Clin Endocrinol Metab* 1988; **66**: 266–72.
16. Holly JMP, Dunger DB, Edge JA, Smith CP, Chard T, Wass JAH. Insulin-like growth factor binding protein-1 levels in diabetic adolescents and their relationship to metabolic control. *Diab Med* 1990; **7**: 618–23.
17. Stanhope R, Pringle PJ, Brook CGD. The mechanism of the adolescent growth spurt induced by low dose pulsatile GnRH treatment. *Clin Endocrinol* 1988; **28**: 83–91.
18. Holly JMP, Smith CP, Dunger DB, et al. Relationship between the pubertal fall in sex hormone binding globulin and insulin-like growth factor binding protein-I: a synchronized approach to pubertal development? *Clin Endocrinol* 1989; **31**: 277–84.
19. Holly JMP, Smith CP, Dunger DB, et al. Levels of the small insulin-like growth factor-binding protein are strongly related to those of insulin in prepubertal and pubertal children but only weakly so after puberty. *J Endocrinol* 1989; **121**: 383–87.
20. Christau B, Kromann H, Christy M, Ortvad-Anderson O, Nerup J. Incidence of insulin dependent diabetes mellitus (0–29 years at onset) in Denmark. *Acta Med Scand* 1979; **624** (suppl): 54–60.