

AGE-ASSOCIATED INCREASED INTERLEUKIN-6 GENE EXPRESSION, LATE-LIFE DISEASES, AND FRAILTY

William B. Ershler¹ and Evan T. Keller²

¹*The Institute for the Advanced Studies in Aging and Geriatric Medicine, Washington, DC 20006; e-mail: wershler@iasia.org; and* ²*Unit for Laboratory Animal Medicine, Department of Pathology, and Institute of Gerontology, University of Michigan, Ann Arbor, Michigan 48109; e-mail: etkeller@umich.edu*

Key Words cytokines, immunodeficiency, immune dysregulation, cachexia, sarcopenia, osteoporosis, dementia

■ **Abstract** Interleukin-6 (IL-6) is a proinflammatory cytokine that is normally tightly regulated and expressed at low levels, except during infection, trauma, or other stress. Among several factors that down-regulate IL-6 gene expression are estrogen and testosterone. After menopause or andropause, IL-6 levels are elevated, even in the absence of infection, trauma, or stress. IL-6 is a potent mediator of inflammatory processes, and it has been proposed that the age-associated increase in IL-6 accounts for certain of the phenotypic changes of advanced age, particularly those that resemble chronic inflammatory disease [decreased lean body mass, osteopenia, low-grade anemia, decreased serum albumin and cholesterol, and increased inflammatory proteins such as C-reactive protein (CRP) and serum amyloid A]. Furthermore, the age-associated rise in IL-6 has been linked to lymphoproliferative disorders, multiple myeloma, osteoporosis, and Alzheimer's disease. This overview discusses the data relating IL-6 to age-associated diseases and to frailty. Like the syndrome of inappropriate antidiuretic hormone, it is possible that certain clinically important late-life changes are due to an inappropriate presence of IL-6.

INTRODUCTION

Interleukin-6 (IL-6) is a multifunctional, proinflammatory cytokine that has been implicated in the pathogenesis of several chronic diseases associated with aging, including osteoporosis, Alzheimer's disease (AD), and neoplasia. IL-6 is tightly regulated and is normally not detected in the serum of healthy young individuals unless there is trauma, infection, or other stress. Under these circumstances, IL-6 is rapidly expressed, and it contributes to a cascade of events typical of inflammation. These include leukocytosis, thrombocytosis, lymphocyte activation,

acute-phase protein synthesis [including fibrinogen, C-reactive protein (CRP), and amyloid precursor protein], and a general catabolic shift in metabolic pathways.

The regulation of the IL-6 gene has been intensely investigated in recent years. It appears that gene expression can be stimulated by a variety of transcription activators, many of which are commonly associated with acute inflammatory or proliferative states (e.g. NF- κ B, NF-IL-6, and *c-fos*). It is also apparent that, in the quiescent state, gene expression is kept in check by a complex network that involves the secondary sex steroids, including estrogen and testosterone (see below). Recently, it has also been discovered that IL-6 levels become detectable in the serum of older individuals, particularly after menopause or andropause, even without signs of illness or inflammation. It has been postulated that this increase relates to the development of frailty and the predisposition to certain diseases (1).

BIOLOGY OF INTERLEUKIN-6

The biochemical characterization of IL-6 came about a decade ago from several different laboratories including those investigating hepatic regeneration, B-cell differentiation and proliferation, myeloma (hybridoma) growth, and T-cell activation [reviewed by Van Snick (2)]. IL-6 is a 26-kDa protein produced by a wide variety of cells. Like IL-1 and tumor necrosis factor (TNF), it is a critical factor in the acute-phase inflammatory response.

The long list of synonyms (Table 1) by which IL-6 has been identified is indicative of its pleiotropic biological activity (3). The molecule was first identified during an effort to isolate and characterize the virus-induced protein inter-

TABLE 1 Synonyms for interleukin-6

B-cell stimulatory factor 2
B-cell differentiation factor
T-cell-replacing factor
Cytotoxic T-cell differentiation factor
Interferon- β_2
Interleukin hybridoma plasmacytoma factor 1
Hybridoma growth factor
26-kDa protein
Plasmacytoma growth factor
Hepatocyte-stimulating factor
Thrombopoietin 2
Macrophage granulocyte-inducing factor

feron- β from cultured human fibroblasts (4). IL-6 is now well recognized for its role in inflammation, characterized by production of a variety of hepatic proteins termed acute-phase proteins (e.g. CRP, serum amyloid A, fibrinogen, complement, and α_1 -antitrypsin) (5). In addition to its role in the acute-phase response, IL-6 is important for the development of specific immunologic responses. It induces differentiation of activated, but not resting, B cells (6–8), culminating in production of immunoglobulin (9, 10). In addition to differentiating B cells, IL-6 stimulates proliferation of thymic and peripheral T cells (11, 12), and, with IL-1 (13), induces T-cell differentiation to cytolytic T cells (14, 15) and activates natural killer cells (16). These observations emphasize the importance of IL-6 in both nonspecific and specific immune responses.

In addition to its immunologic/inflammatory role, IL-6 appears to play an important role in bone metabolism by inducing osteoclastogenesis and osteoclast activity (17, 18). Estrogen inhibits IL-6 gene expression (see below), and, in rodent models, in the absence of estrogen, IL-6 activity and bone resorption increase (19–22). The importance of this estrogen–IL-6 interaction is supported by the observation that IL-6 gene knockout mice are protected from cancellous bone loss associated with ovariectomy (23).

IL-6 also influences a number of other important cell types. For example, IL-6 stimulates the endometrial vasculature during the menstrual cycle (24, 25) and promotes spermatogenesis (26). Furthermore, IL-6 stimulates epidermal proliferation (27–29), hemostasis through induction of megakaryocytopoiesis (30–32), and neural cell differentiation and proliferation (33, 34).

THE INTERLEUKIN-6 GENE

The human IL-6 gene has been localized to chromosome 7p21 (35) and consists of 5 exons and 4 introns (36, 37). The human IL-6 molecule contains 212 amino acids, including a hydrophobic signal sequence of 28 amino acids. It includes two potential N-glycosylation sites. Murine IL-6 (chromosome 5) consists of 211 amino acids with a typical signal sequence of 24 residues. It includes no N-glycosylation site but several potential O-glycosylation sites (38).

Interleukin-6 Gene Regulation

Mechanisms of IL-6 gene regulation are incompletely understood. It was demonstrated several years ago that both protein kinase C and cAMP-dependent signal transduction pathways could trigger IL-6 gene expression (39–41). To further explain the wide range of functions of IL-6, the presence of multiple initiation sites and the preferential use of a specific initiation site in a variety of cells (37) suggest that different regulatory mechanisms may be responsible for IL-6 gene expression in different tissues [Table 2 (7, 42–55)]. In fact, sequences similar to several consensus sequences involved in gene activation are present in the 5'

TABLE 2 Factors that induce interleukin-6 promoter activity^a

Stimulus	Cell type	Transcription factors	Reference
LAM or LPS	Monocyte	NF-IL-6, NF-κB	42
HTLV-I TAX	HTLV-I-infected T cell	NF-κB	7, 43, 44
Hypoxia	Endothelial cells	NF-IL-6	45
PGE ₁ , cAMP	PU5-1.8 monocyte	AP-1, NF-IL-6, NF-κB	46
LPS	Monocyte	NF-κB	46
HIV-I TAT	B-lymphoblastoid, HeLa (cervical carcinoma)	NF-IL-6, NF-κB	47
Ionizing radiation	Fibroblast	AP-1, NF-κB	48
Mutant p53	HeLa	NF-IL-6	49
Jun, TNF-α, PKC, IL-1, db-cAMP, PMA	HepG2 (hepatocyte), HeLa	Not characterized	50
LTB4	Monocyte	NF-IL-6, NF-κB	51
LIF	Monocyte	NF-κB	52
IL-1α, LPS, cAMP	OCI-LY3 (lymphoma)	Not characterized	53
IL-1α, TNF-α	HeLa, glioblastoma, fibroblasts	NF-κB	54, 55
Forskolin	Fibroblasts	Not NF-κB	54

^aAbbreviations: cAMP, cyclic AMP; db-cAMP, dibutyl cAMP; HIV-I, human immunodeficiency virus I; HTLV-I, human T-lymphotropic virus I; IL-1α, interleukin-1α; LAM, lipoarabinomannan; LIF, leukemia inhibitory factor; LPS, lipopolysaccharide; LTB4, leukotriene B4; PGE₁, prostaglandin E₁; PKC, protein kinase C; PMA, phorbol 12-myristate 13-acetate; TNF-α, tumor necrosis factor-α.

flanking region, such as the *c-fos* serum response element, the CCAAT enhancer binding protein (C/EBPβ) domain, and the NF-κB domain (Figure 1), all of which are conserved between human and mouse genes (56). The presence of this highly conserved but complex regulatory region has been interpreted to indicate the importance of IL-6 gene regulation and its varied responses in different tissues subjected to different stimuli.

Glucocorticoids and Interleukin-6 Expression

Glucocorticoids have been shown to inhibit IL-6 production from a variety of tissues (57–59). Initial investigation into the mechanism of glucocorticoid inhibition of IL-6 expression demonstrated that dexamethasone inhibited IL-1–induced transcriptional activation of the proximal 225 base pairs of the IL-6 promoter (60). This finding prompted the examination of the interaction of the glucocorticoid receptor (GR) and the IL-6 promoter (60). GR was found to weakly bind the IL-6 promoter at several important *cis*-acting sites. A DNA-binding




Figure 1 The proximal interleukin-6 promoter. Numbers indicate distance (in base pairs) from the transcription initiation site. MRE, multiple response element; C/EBP- β , CCAAT/enhancer binding protein; RCE, retinoblastoma control element; NF- κ B, nuclear factor- κ B; SRE, serum response element.

immunoprecipitation assay using HeLa cells indicated that transfected wild-type GR cDNA was capable of binding to the -225 fragment of the IL-6 promoter (61). Additionally, mutation of the GR DNA-binding domain resulted in the loss of GR inhibition of transcriptional activation (61). Thus, it appears that corticosteroid interacts with its receptor, which then binds the IL-6 promoter and inhibits IL-6 gene expression.

However, the interaction between GR and the IL-6 promoter is weak. Thus, alternative mechanisms have been sought to account for GR inhibition of the IL-6 promoter. Ray & Prefontaine (62) questioned whether GR interacted with two transcription factors known to stimulate the IL-6 promoter, NF- κ B and NF-IL6. Using murine F9 embryonal carcinoma cells, which are devoid of endogenous NF-IL6 and NF- κ B activities, they demonstrated that dexamethasone inhibited IL-6 promoter transactivation induced by transfected p65 (a component of the NF- κ B complex) and NF-IL6 (62). In cross-precipitation assays, it was demonstrated that GR bound to p65, but not to NF-IL6. Thus, GR mediates inhibition of p65-induced activation of the IL-6 promoter through protein-protein interactions. This mechanism may occur in combination with the promoter-occlusion mechanism described above.

Still another mechanism of GR inhibition of IL-6 has recently been discovered (64, 65). Dexamethasone was shown to induce the expression of the cytoplasmic inhibitor of NF- κ B, I κ B α , and this was associated with inhibition of nuclear translocation of NF- κ B p65. Thus, it follows that GR-induced I κ B α protein synthesis promotes cytoplasmic sequestration of NF- κ B, culminating in decreased activation of the IL-6 promoter.

IL-6 induced by stress, infection, or trauma stimulates release of corticotrophin-releasing factor from the hypothalamus, which results in elevated systemic levels of corticosteroids (66, 67). This observation suggests the presence of a CNS-immune-endocrine negative-feedback loop.

Estrogen and Interleukin-6 Expression

The ability of estrogen to repress IL-6 expression was first recognized in human endometrial stromal cells (68). Additional evidence came from the observations that menopause or ovariectomy resulted in increased IL-6 serum levels (69), increased IL-6 mRNA levels in bone cells (70), and increased IL-6 secretion by mononuclear cells (71–73). Direct evidence for estrogen's ability to repress IL-6 expression is derived from studies demonstrating that estradiol inhibited bone marrow stromal-cell and osteoblast IL-6 protein and mRNA production in vitro (20, 74) and that estradiol suppressed osteoclastogenesis in murine bone marrow cultures (20) or ovariectomized mice (21) as effectively as IL-6-neutralizing antibody. These data are somewhat controversial because, in a few studies using human bone-derived cells, estrogen did not inhibit IL-6 expression (75–78). Perhaps species-specific differences account for these conflicting data. The majority of the data available implicate an estrogen-mediated repression of IL-6 expression in a variety of systems.

To define the mechanism of estrogen-induced transrepression of the IL-6 promoter, Pottratz et al (79) and Ray et al (80) performed transient transfection assays using an IL-6 promoter linked to a reporter plasmid. They found that 17 β -estradiol inhibited phorbol (an inflammatory stimulus), IL-1, or TNF-induced promoter activation in a murine bone marrow stromal-cell line, which constitutively expresses the estrogen receptor (ER). This was also observed in HeLa cells, which have very little or no endogenous ER. However, in these cells, the transfection of ER DNA was required to observe suppression of the IL-6 promoter by estradiol. Thus, 17 β -estradiol inhibits IL-6 gene transcriptional activation by an ER-dependent mechanism.

Because there is no consensus ER element within the IL-6 promoter (as indicated by gene bank analysis), it stands to reason that ER inhibits the IL-6 promoter by a mechanism other than direct binding of ER to the IL-6 promoter (80). Rather, it appears that ER directly inhibits transcription factors and thus negatively influences the IL-6 promoter. For example, Stein & Yang (82) found that ER bound both NF- κ B p65 and NF-IL-6. Furthermore, they demonstrated that the ER-transcription factor complex (whether ER-NF- κ B or ER-NF-IL-6) inhibited unbound transcription factor enhancer activity through a mechanism that did not include the induction of I κ B α . We have documented that estradiol inhibits degradation of I κ B α (83). This observation suggests that estradiol promotes sequestration of NF- κ B in the cytoplasm. Adding strength to this hypothesis is the recent observation that the NF-IL6-binding site (C/EBP β) is required for estrogen-mediated repression (84).

Androgens and Interleukin-6 Expression

The first demonstration of the inhibitory action of androgens on IL-6 protein expression was made in a murine bone marrow stromal-cell line that had been stimulated with IL-1 and TNF (85). These findings were extended in HeLa cells,

in which it was demonstrated that dihydrotestosterone (DHT) inhibited PMA-induced activation, albeit to a lesser extent than did estrogen (79). Several adrenal androgens that are not known to bind the androgen receptor (i.e. androstenedione, androstenediol, and dehydroepiandrosterone sulfate) mediate repression of the IL-6 promoter in HeLa cells transfected with the androgen receptor (85). These experiments suggest that even weakly binding androgens can mediate transrepression of IL-6 promoter activation.

We have demonstrated that DHT requires the androgen receptor to inhibit IL-6 promoter (86). DHT inhibited PMA-induced activation of the IL-6 promoter by inhibiting nuclear translocation of NF- κ B in a prostate cancer cell line, LNCaP. This was associated with maintenance of cytoplasmic I κ B α levels in the presence of PMA. In the absence of androgen, I κ B α levels in the cytoplasm decline after PMA stimulation, and the maintenance of I κ B α levels in the presence of androgen could be caused by decreased phosphorylation (degradation) or increased production.

Orchiectomy induces bone marrow IL-6 protein and mRNA expression (87). The relevance of the above observations was pointed out by Bellido et al (85) based on an orchiectomized mouse model. Although serum or bone marrow IL-6 levels were not reported, these authors found that orchiectomy resulted in increased replication of bone marrow osteoclast progenitors, which was prevented by administration of IL-6-neutralizing antibody or implantation of a slow-release form of testosterone. However, because testosterone is converted to estradiol, these results are not conclusive evidence of the action of androgen. In fact, the observation of decreased bone density in a male patient who had normal androgen levels but had a mutation resulting in a nonfunctional ER suggests that estrogen's effects on bone in men may be as important as androgen's (88). This hypothesis is supported by the observation that estradiol can inhibit the bone loss that was observed in men treated by orchiectomy for prostate cancer (89). However, the ability of estrogen to inhibit bone loss may be mediated by its transrepression of the IL-6 promoter, and thus, these observations are still consistent with increased IL-6 activity from androgen loss.

THE INTERLEUKIN-6 RECEPTOR

The human IL-6 receptor (IL-6R; also known as gp80 or the IL-6R α subunit) was first cloned from a human natural-killer-like cell line, YT (89), followed by a cloning from a human hepatoma cell line, HepG2 (90). IL-6R is an 80-kDa protein consisting of 467 amino acids. Located on chromosome 1 band q21 (91), the IL-6R gene encodes for a 5-kb mRNA containing a coding region of 1401 base pairs (89).

The structure of IL-6R has been deduced by comparative sequence analysis. A hydropathy plot revealed two major hydrophobic regions: one that encodes for the signal peptide between residues 1 and 20, and another that encodes for the

transmembrane domain in the region of residues 359–386 (89). The latter region is followed by a putative transmembrane anchoring stop codon consisting of several positively charged residues. Thus, IL-6R has a 339–amino-acid extracellular region, a 28–amino-acid transmembrane region, and an 82–amino-acid intracellular region. Intriguingly, the intracellular region does not contain any kinase domains, making it likely that this molecule is not capable of signaling activity (89).

A homology search indicated that the extracellular component of IL-6R contains a domain that shares extensive homology with the immunoglobulin superfamily (89) and two tandem fibronectin type-III motifs (92) present in a 200–amino-acid region. This region defines a domain that is found in a variety of other cytokine and growth factor receptors. It contains highly conserved components consisting of four cysteine residues in its amino-terminal region and a tryptophan-serine-X-tryptophan-serine (WSXWS) motif penultimate to the transmembrane region (92, 93). Fibronectin type-III domains are observed in cell adhesion molecules, and their presence has been interpreted to suggest that cytokine receptors evolved from an ancestral adhesive molecule (94).

As the protein structure suggests, IL-6R is not capable of inducing signal transduction directly. It is now understood that, in order to mediate signal, IL-6 first binds to gp-80, forming a low-affinity receptor complex. This complex then associates with the non-ligand-binding transmembrane glycoprotein gp-130 (95). Homodimerization of gp-130 is required for IL-6 signal transduction (96). Although it was originally believed that one unit of IL-6 and IL-6R bound to a gp-130 homodimer (96), the complex now appears to be hexameric, consisting of two molecules each of IL-6, IL-6R gp-80, and gp-130 (97). This complex provides high-affinity binding of IL-6, as opposed to the low-affinity binding of IL-6 and IL-6R gp-80 in the absence of gp-130.

IL-6R has an isoform that was first identified in human urine, the soluble IL-6R (sIL-6R) (98). In contrast to other soluble cytokine receptors (e.g. sIL-2R), which inhibit cytokine-induced signaling, sIL-6R forms a fully active hexameric IL-6:sIL-6R:gp-130 complex capable of inducing signal transduction. How sIL-6R is generated is not currently established, but both alternative splicing, resulting in loss of the transmembrane domain (99), and proteolysis of the mature cell surface IL-6R (100, 101) have been proposed.

Expression of the Interleukin-6 Receptor

IL-6R is expressed in a variety of cells including B-lymphocytes, hepatocytes, prostate cells, and osteoblasts. In general it is expressed at ~100–2000 sites/cell (102). However, up to 29,000 sites/cell have been identified in myeloma lines and Epstein-Barr virus-transformed cells (102). Except for the effects of dexamethasone, modulation of IL-6R expression by various factors has not been consistently demonstrated. Snyers et al (102) found that A23187 (a calcium

ionophore), lipopolysaccharide, prostaglandin E₁, IL-1, tumor necrosis factor (TNF), and muramyl dipeptide did not significantly alter IL-6R expression in a variety of cell lines. However, other investigators have found that IL-1 modulated IL-6R expression in various cell lines and tissues (102). Perhaps cell-specific differences in response account for these discordant results.

In contrast to these inconclusive results, dexamethasone has been repeatedly demonstrated to increase IL-6R in several tissue types, including liver primary cells and cell lines, monocyte primary cultures, myeloma cell lines, and an amniotic cell line (100). In addition to these *in vitro* observations, IL-6R mRNA expression has been documented to increase in murine bone marrow stroma postovariectomy (103).

Despite the great variety of cells that express IL-6R and its importance in many facets of physiology, the molecular mechanisms that regulate transcriptional control of the IL-6R gene have not been defined to date. Cloning and analysis of the IL-6R promoter will help in this analysis.

INTERLEUKIN-6 AND “NORMAL” AGING

A wealth of data indicate that IL-6 gene expression, as well as tissue and serum levels, increases with age [Table 3 (69, 71, 104–115)]. Early observations (104) demonstrate an age-associated rise in IL-6 in autoimmune-prone mice. This increase is considered an antecedent to the development of the lupus erythematosus-like syndrome characteristic of that strain. Subsequently, other laboratories demonstrated a similar age-associated increase in IL-6 in “normal” (i.e. non-disease-prone) strains of mice and also in rats, monkeys, and humans (Table 2).

The increase in IL-6 has not been fully explained, but it may result at least partially from coincident age-associated diseases in which elevated IL-6 has been described (see below). However, increases in circulating IL-6 have been demonstrated in apparently healthy older individuals. One proposed mechanism is the reduced influence of the normally inhibiting sex steroids on endogenous IL-6 expression (as described above). Accordingly, at the time of menopause or andropause, IL-6 gene expression is not as tightly regulated, inappropriate expression occurs (at least in some tissues), and serum levels rise. The biological importance of this rise in IL-6 has yet to be fully established but, at least for certain age-associated diseases (such as osteoporosis, neoplasia, and AD), preliminary investigations have suggested an important role. Thus, it is possible that the age-associated rise in IL-6 is of physiological consequence, rendering an individual susceptible to the myriad of processes induced by this proinflammatory signal, including lymphoproliferation, osteoclast stimulation, and a metabolic orientation towards catabolic processes.

TABLE 3 Interleukin-6 and age^a

Group (reference)	Species	Finding(s)
Tang et al (104)	MRL/lpr mice	Serum levels increase with age
Suzuki et al (105)	MRL/lpr mice	Serum soluble IL-6 receptors are elevated with age
Effros et al (106)	C3B10F ₁ mice	LPS-stimulated peritoneal macrophage IL-6 levels rose with age
Zhou et al (107)	C57BL/6J mice	Increased production of IL-6 by spleen cells from old mice
Foster et al (108)	Fisher 344 rats	LPS-stimulated peripheral MNC IL-6 production increased with age
Wei et al (109)	Humans	Serum IL-6 levels elevated in old people but no age differences in IL-1 levels
Goya et al (110)	Mice	Growth-hormone-treated old mice had higher IL-6 levels than similarly treated middle-aged or young mice
Daynes et al (71)	BALB/c and 3H/HeN mice and humans	Serum IL-6 levels rise through the lifespan
Ershler et al (111)	Rhesus monkeys	Serum IL-6 levels rise through the lifespan
	Humans	Serum IL-6 levels elevated in healthy old people
Sindermann et al (112)	Humans	IL-6 (but not interferon or IL-2) synthesis in PHA-stimulated lymphocyte cultures was greater from old compared with young healthy volunteers
McKane et al (113)	Humans	Serum IL-6 levels rise with age in healthy women
Fagiolo et al (114)	Humans	Increased IL-6 in vitro production by cells from healthy old people
Liao et al (115)	Humans	Urine IL-6 levels increase with age
Kania et al (69)	Humans	IL-6 levels increase post-menopause

^aLPS, lipopolysaccharide; MNC, mononuclear cell; PHA, phytohemagglutinin.

INTERLEUKIN-6 AND CERTAIN AGE-RELATED DISEASES

Lymphoproliferation and Lymphoproliferative Disorders

IL-6 stimulates B-cell maturation and proliferation, and its overproduction has been demonstrated in a variety of malignancies of B-cell lineage (including multiple myeloma (116–118), non-Hodgkin's lymphoma (119, 120), and chronic

lymphocytic leukemia (121). In addition, other lymphoproliferative diseases, including Castleman's disease (122, 123), Hodgkin's disease (120, 124), angioimmunoblastic lymphadenopathy with dysproteinemia (125), and certain non-B-cell lymphomas (126), are typically associated with measurably high circulating levels of this cytokine. Although this may be an important clue to the pathogenesis of these disorders, the high IL-6 level may be of additional clinical importance because it has been shown to correlate with the presence of prominent constitutional symptoms (night sweats, fever, weight loss, etc) (120), stage of disease (119, 127, 128), and survival (128).

Both experimental data and clinical observations suggest that IL-6 may be involved in the pathogenesis of some or all of these diseases. For example, it has been demonstrated in certain disorders that the neoplastic cells in culture were stimulated to proliferate in the presence of recombinant IL-6 (129). Furthermore, lymphoma and myeloma cells have been shown, both in vivo and in vitro, to produce IL-6 (130–133), and the level of IL-6 production by these neoplastic cells correlates with serum levels (133) and with survival (132).

The potential importance of IL-6 in the development of B-cell malignancies has also been demonstrated in animals, e.g. C57Bl/6 mice (Figure 2), over 50% of which will develop lymphoma by 24 months of age (134). We showed that




Figure 2 Interleukin-6 and lymphoma development. This model proposes that the rise in IL-6 level that occurs in middle-aged mice is the antecedent to a polyclonal lymphoproliferative disorder and that a second oncogenic event, such as mutation and activation of the *c-myc* oncogene, is required for lymphoma development. Ab, antibody(ies).

the appearance of IL-6 with advancing age correlated with increased lymphoid proliferation, *c-myc* expression, and ultimately lymphoma development in this strain of mice (135). Lymphomas occurred only in mice known to have prior IL-6 elevations, and mice on controlled caloric restriction [which is known to markedly reduce lymphomagenesis and prolong maximal lifespan in rodents (136)] had both lower IL-6 levels and less lymphoma development. Similarly, splenocytes from AKR mice destined to develop B-cell lymphomas have markedly elevated levels of IL-6 (and IL-1) before developing overt disease, and lymphoma occurrence can be significantly inhibited by in vivo administration of monoclonal antibody to IL-6 (137). Finally, plasma cell tumors that can readily be induced in normal mice could not be induced in IL-6-deficient mice (IL-6 knockouts generated by introduction of a germline-encoded null mutation in the IL-6 gene) (138). In those experiments, mice that were homozygous (+ / +) or heterozygous (+ / -) for the wild-type IL-6 allele yielded the typical high incidence of plasmacytoma, but mice homozygous for the IL-6 null allele (- / -) were completely resistant.

The importance of IL-6 in the pathogenesis of multiple myeloma in humans has been suspected for many years, perhaps because it was initially identified as a necessary factor for myeloma cell growth in vitro (117, 139). IL-6 levels are high in patients with multiple myeloma, and myeloma cells themselves have high levels of IL-6 receptors (140). Furthermore, serum IL-6 levels have been correlated with myeloma tumor cell mass (141) and disease severity (142), although there is some controversy about this (143).

A recent report sheds light on the importance of IL-6 dysregulation in multiple myeloma. The cell surface molecule, CD44, on plasma (and myeloma) cells mediates binding of tumor cells to stroma and regulates IL-6 production. Variants of this molecule are produced by alternative splicing of the nuclear RNA, and these variants express additional protein-binding domains, primarily in the extracellular region of the molecule. In one series, expression of specific variants (determined by immunohistochemical staining) was observed in more than one third of myeloma patients (144), and this was associated with advanced, more rapidly progressive disease and shorter survival.

Overwhelming evidence has indicated that IL-6 stimulates proliferation of tumor cells of the B-cell lineage. This conclusion has led to the development of IL-6 inhibitors as a treatment strategy. Klein et al (145, 146) treated nine multiple-myeloma patients with anti-IL-6 monoclonal antibody and found dramatic blockage of myeloma cell proliferation as well as reduction of toxicity related to overproduction of IL-6 in vivo (fever, hypercalcemia, and acute-phase proteins). The same group treated a plasma-cell leukemia patient with anti-IL-6 monoclonal antibody, again finding blockage of myeloma cell proliferation and reduction in serum calcium, serum monoclonal immunoglobulin G, and serum C-reactive protein (CRP), as well as slight improvement in performance status (147). It has also been demonstrated that other agents (125, 148), including corticosteroids (149), are capable of inhibiting IL-6 production in vitro. It has been hypothesized that

the efficacy of corticosteroids in treating patients with multiple myeloma could, at least in part, be due to their ability to inhibit IL-6 production. It is possible that other agents will be found to specifically inhibit IL-6 production or IL-6 receptor function (150), and these may also have efficacy in the treatment of multiple myeloma.

Osteoporosis

It is now clear that several cytokines and growth factors contribute to the bone-remodeling process, including IL-1, IL-6, IL-11, and TNF. IL-6 has received special attention because it is the only one of these cytokines that has consistently been demonstrated to be elevated with age. Furthermore, both IL-6 and IL-6R have been recognized as important normal contributors to the process of bone remodeling (151–153).

It is now well accepted that IL-6 stimulates bone-resorbing osteoclasts, but its effect on osteoblasts remains unclear. For example, some investigators have demonstrated that IL-6 has a moderate effect on osteoblast proliferation (154) and differentiation (155, 156), whereas others have not observed these effects (157, 158). Accordingly, it is likely that the most important effect on bone is enhanced osteoclast activity.

IL-6 and IL-6R enhance osteoclastogenesis. Parathyroid hormone, a well-recognized inducer of bone resorption, stimulates IL-6 production in bone (59, 152, 159, 160). In addition to parathyroid hormone, other substances that stimulate bone remodeling [e.g. bradykinin (161) and bone morphogenic protein-2 (162)] also stimulate IL-6 production by human osteoblast-like cells. Although occasional reports suggest only a limited role for IL-6 in bone resorption (163, 164), the majority of published findings conclude the opposite. For example, IL-6 induces differentiation of bone marrow mononuclear cells to osteoclasts in long-term human marrow cultures (17) and in cocultures of murine bone marrow and primary osteoblastic cells (18). The addition of IL-6R was required to observe this effect in vitro (18). Moreover, conditioned media from marrow cultures obtained from patients with Paget's disease (characterized by increased osteoclastogenesis) stimulated osteoclast-like cell formation in normal human marrow cultures. This was reversed by the addition of neutralizing antibody to IL-6 (165). IL-6-neutralizing antibody also blocks bone resorption induced by a variety of agents including TNF (20, 166). Along with increasing osteoclast numbers, IL-6 has been shown to stimulate bone resorption in fetal mouse metacarpal (160), calvaria (156), and bone resorption pit assays (165, 167).

The importance of IL-6 dysregulation in the development of osteoporosis is highlighted by Bellido et al (85), who demonstrated that orchietomy increased osteoclastogenesis in mice but that this increase was inhibited by androgen replacement or IL-6-neutralizing antibody. Furthermore, osteoclastogenesis was not observed in orchietomized mice deficient in IL-6 production (IL-6 knock-outs). This study must be interpreted with caution because one must assume that

the bone in IL-6-deficient mice is not representative of normal bone. However, the study provides strong evidence for the role of androgens on IL-6-mediated osteoclastogenesis. The findings also gain significance in conjunction with the earlier observation (23) that ovariectomized IL-6-knockout mice maintain bone density, unlike control mice.

Taken together, these observations provide substantial support for the hypothesis that certain proinflammatory cytokines (especially IL-1 and IL-6) are important both in the normal bone-remodeling process and in the pathogenesis of postmenopausal osteoporosis. The interesting pattern of increased late-life expression of IL-6 lends further support. It is now clear that estrogens and androgens are important inhibitors of IL-6 gene expression, and it is likely that hormone replacement therapy mediates bone preservation by effecting this inhibition. Future strategies to specifically regulate menopausally dysregulated IL-6 are the focus of great interest and research at this time.

Alzheimer's Disease

IL-6 contributes to the pathogenesis of AD. Proinflammatory cytokines (including IL-1 and IL-6) may also contribute to the development of AD (165). One theory on the pathogenesis of AD postulates that IL-6 mediates neurodegeneration by potentiating chronic inflammation in the brain (168–170). This is supported by the observation that IL-6 and certain acute-phase proteins induced by IL-6 [e.g. α 2-macroglobulin, CRP, and antichymotrypsin (171)] are found in either amyloid deposits or brain tissue extract of AD patients (172–175). Yet, with the exception of one report (176), IL-6 levels have not been found to be increased in the cerebrospinal fluid (173, 177, 178) or serum (178, 179) of AD patients, raising the likelihood that IL-6 production and activity are localized to the relevant microenvironment.

IL-6 secreted by microglial and/or neuronal cells (180, 181) may induce synthesis of certain acute-phase proteins, including amyloid precursor protein (182). Amyloid precursor protein itself can induce the expression of IL-1, but not of IL-6, by neuronal cells (181). However, IL-6 has been found in the early stage of plaque formation (diffuse plaques) (183, 184), whereas none of the other cytokines tested (including IL-1 and IL-2) were detected (175, 181, 184, 185). In fact, IL-1, which can induce IL-6 expression, was found not to be elevated in brain from AD patients, whereas IL-6 was elevated (175).

There may be some evidence that IL-6 is specifically involved in AD dementia, distinct from other forms of dementia. IL-6 secretion by peripheral blood mononuclear cells was increased in patients with AD as opposed to normal control subjects or those suffering from probable vascular dementia (186, 187), and IL-6 was found in plaques of AD patients who suffered from dementia as opposed to those who did not (184). In addition to these observations, the overexpression of IL-6 in the brain in transgenic mice has been associated with a variety of neuropathologic findings, including gliosis and selective disruption of cholinergic

transmission in the hippocampus (188–193). This latter finding is of particular interest in light of the cholinergic model of AD.

Thus, there is strong evidence that IL-6 overexpression plays an important role in the development of AD. Whether the mechanism for the IL-6 overexpression in the brain of AD patients differs from the overexpression found with advancing age is unknown (183). However, it is clear that determining this mechanism will improve understanding of the pathogenesis of AD and may lead to new treatment strategies. In this regard, it is tempting to postulate that the mechanism of the positive effects of hormone replacement therapy (194) and nonsteroidal anti-inflammatory drugs (195) is their inhibition of IL-6 expression.

INTERLEUKIN-6 AND FRAILTY

In late life, IL-6 levels rise, presumably because of a loss of the influence of the inhibitory secondary sex steroids. The predicted musculoskeletal and metabolic consequences include a pattern of changes attributed to chronic inflammatory disease or frailty. Features of frailty that may be related to increased IL-6 include decreased lean body mass, anemia, thrombocytosis, decreased bone mineral density, decreased serum cholesterol and albumin, and dementia. That such changes could be produced by IL-6 may be supported by a study of young and middle-aged rhesus monkeys that received relatively low doses of IL-6 daily for 1 month. These animals lost 10% of their body weight (primarily lean body weight as determined by dual energy X-ray absorptiometry), became osteopenic and anemic, and had significant drops in serum albumin and cholesterol and corresponding rises in CRP and alkaline phosphatase (196).

In a large series of healthy humans across the lifespan, increasing IL-6 with age was associated with functional declines typical of frailty (197). Thus, the age-associated rise in IL-6 may well explain the “chronic inflammation” appearance of some elderly individuals, even in the absence of an inflammatory focus.

SUMMARY

IL-6 is one of several proinflammatory cytokines. Under normal circumstances in young individuals, its expression is tightly regulated by the interplay of several transcription factors and hormonal factors including the secondary sex steroids and glucocorticoids. Unlike the expression of IL-1 and TNF, which are in the same class of proinflammatory cytokines, IL-6 expression increases in late life. This increase is thought to result from loss of the normally inhibiting sex steroids. It has been proposed that many of the features of frailty (particularly those that resemble chronic inflammatory disease) are the result of this age-associated dysregulation of IL-6. Prospective long-term clinical trials testing the effects of inhib-

itors of IL-6, such as estrogens, androgens, and nonsteroidal anti-inflammatory drugs, on the development of frailty will be of great interest.

ACKNOWLEDGMENTS

This work was supported by US PHS, grants #R01 AG 11970 and #R01 AG 15904.

Visit the Annual Reviews home page at www.AnualReviews.org.

LITERATURE CITED

1. Erszler WB. 1993. Interleukin-6: a cytokine for gerontologists. *J. Am. Geriatr. Soc.* 41(2):176–81
2. Van Snick J. 1990. Interleukin-6: an overview. *Annu. Rev. Immunol.* 8:253–78
3. Akira S, Taga T, Kishimoto T. 1993. Interleukin-6 in biology and medicine. *Adv. Immunol.* 54:1–78
4. Weissenbach J, Chernajovsky Y, Zeevi M, et al. 1980. Two interferon mRNA in human fibroblasts: *in vitro* translation and *Escherichia coli* cloning studies. *Proc. Natl. Acad. Sci. USA* 77:7152–56
5. Sehgal PB, Greninger G, Tosato G. 1989. Acute phase and immune responses: interleukin-6. *Ann. NY Acad. Sci.* 557:1–583
6. Hirano T, Teranishi T, Lin BH, Onoue K. 1984. Human helper T cell factor(s). IV. Demonstration of a human late-acting B cell differentiation factor acting on *Staphylococcus aureus* Cowen I-stimulated B cells. *J. Immunol.* 133:798–802
7. Muraguchi A, Hirano T, Tang B, et al. 1988. The essential role of B cell stimulatory factor 2 (BSF-2/IL-6) for the terminal differentiation of B cells. *J. Exp. Med.* 67:332–44
8. Taga T, Kawanishi Y, Hardy TT, et al. 1987. Receptors for B cell stimulatory factor 2. *J. Exp. Med.* 166:967–81
9. Ambrosino DM, Delaney NR, Shamberger RC. 1990. Human polysaccharide-specific B cells are responsive to pokeweed mitogen and IL-6. *J. Immunol.* 144:1221–26
10. Takatsuki F, Okano A, Suzuki C, et al. 1988. Human recombinant IL-6/B cell-stimulatory factor 2 augments murine antigen-specific antibody responses *in vitro* and *in vivo*. *J. Immunol.* 141:3072–77
11. Uyttenhove C, Coulie PG, Van Snick J. 1988. T cell growth and differentiation induced by interleukin HP1/IL-6, the murine hybridoma/plasmacytoma growth factor. *J. Exp. Med.* 167:1414–27
12. Lotz M, Jirik F, Kabouridis R, et al. 1988. B cell stimulating factor 2/interleukin 6 is a costimulant for human thymocytes and T lymphocytes. *J. Exp. Med.* 167:1253–58
13. Helle M, Brakenhoff JPJ, DeGroot ET, Aarden LA. 1988. Interleukin-6 is involved in interleukin-1-induced activities. *Eur. J. Immunol.* 18:957–59
14. Okada M, Kitahara N, Kishimoto S, et al. 1988. IL-6/BSF-2 functions as a killer helper factor in the *in vitro* induction of cytotoxic T cells. *J. Immunol.* 141:1543–49
15. Renauld JC, Vink A, Van Snick J. 1989. IL1 and IL6 in CTL induction-accessory signals in murine cytolytic T cell responses: dual requirement for IL1 and IL6. *J. Immunol.* 143:1894–98
16. Luger TA, Krutman J, Kirnbauer T, et al.

1989. IFN β /IL-6 augments the activity of human natural killer cells. *J. Immunol.* 143:1206–9
17. Kurihara N, Bertolini D, Suda T, et al. 1990. IL-6 stimulates osteoclast-like multinucleated cell formation in long term human marrow cultures by inducing IL-1 release. *J. Immunol.* 144:4226–30
18. Tamura T, Udagawa N, Takahashi N, et al. 1993. Soluble interleukin-6 receptor triggers osteoclast formation by interleukin-6. *Proc. Natl. Acad. Sci. USA* 90:11924–28
19. Passeri G, Girasole G, Markus T, et al. 1991. 17 β -estradiol regulates IL-6 production and osteoclast development in murine calvaria cell cultures. *J. Bone Miner. Res.* 6:S263 (Suppl. 1)
20. Girasole G, Jilka RL, Passeri G, et al. 1992. 17 β -estradiol inhibits interleukin-6 production by bone marrow-derived stromal cells and osteoblasts in vitro: a potential mechanism for the antiosteoporotic effect of estrogens. *J. Clin. Invest.* 89(3):883–91
21. Jilka RL, Hangoc C, Girasole G, et al. 1992. Increased osteoclast development after estrogen loss: mediation by interleukin-6. *Science* 257:88–91
22. Miyaara C, Kusano K, Masuzawa T, et al. 1995. Endogenous bone-resorbing factors in estrogen deficiency: cooperative effects of IL-1 and IL-6. *J. Bone Miner. Res.* 10:1365–73
23. Poli V, Balena R, Fattori E, et al. 1994. Interleukin-6 deficient mice are protected from bone loss caused by estrogen depletion. *EMBO J.* 13:1189–96
24. Tabibzadeh S, Kong QF, Babaknia A, May LT. 1995. Progressive rise in the expression of interleukin-6 in human endometrium during menstrual cycle is initiated during the implantation window. *Mol. Hum. Reprod.* 1:2793–99
25. Tabibzadeh SS, Santhanam U, Sehgal PB, May LT. 1989. Cytokine-induced production of IFN- β /IL-6 by freshly explanted human endometrial stromal cells. Modulation by estradiol-17 β . *J. Immunol.* 142(9):3134–39
26. Hakovirta H, Syed V, Jegou B, Parvinen M. 1995. Function of interleukin-6 as an inhibitor of meiotic DNA synthesis in the rat seminiferous epithelium. *Mol. Cell. Endocrinol.* 108:193–98
27. Krueger JG, Krane JF, Carter DM, Gottlieb AB. 1990. Role of growth factors, cytokines, and their receptors in the pathogenesis of psoriasis. *J. Invest. Dermatol.* 94:135–140 (Suppl.)
28. Grossman RM, Kreuger J, Yourish D, et al. 1989. Interleukin 6 is expressed in high levels in psoriatic skin and stimulates proliferation of cultured human keratinocytes. *Proc. Natl. Acad. Sci. USA* 86:6367–71
29. Yoshizaki K, Nishimoto N, Matsumoto K, et al. 1990. Interleukin-6 and expression of its receptor on epidermal keratinocytes. *Cytokines* 2:381–87
30. Navarro S, Devili N, LeCouedic JP, et al. 1991. Interleukin-6 and its receptor are expressed by human megakaryocytes: *in vitro* effects on proliferation and endoreplication. *Blood* 77:461–71
31. Sun WH, Binkley N, Bidwell DW, Ershler WB. 1993. The influence of recombinant human interleukin-6 on blood and immune parameters in middle-aged and old rhesus monkeys. *Lymphokine Cytokine Res.* 12(6):449–55
32. Hill RJ, Warren MK, Levin J. 1990. Stimulation of thrombopoiesis in mice by human recombinant interleukin-6. *J. Clin. Invest.* 85:1242–47
33. Hama T, Miyamoto M, Tsukui H, et al. 1989. Interleukin-6 as a neurotrophic factor for promoting the survival of cultured basal forebrain cholinergic neurons from postnatal rats. *Neurosci. Lett.* 104:340–44
34. Satoh T, Nakamura S, Taga T, et al. 1988. Induction of neuronal differentiation in PC12 cells by B-cell stimulatory factor 2/interleukin 6. *Mol. Cell. Biol.* 8:3546–49

35. Sehgal PB, Zilberstein A, Ruggieri MR, et al. 1986. Human chromosome 7 carries the $\beta 2$ interferon gene. *Proc. Natl. Acad. Sci. USA* 83:8957–61
36. Zilberstein A, Ruggieri R, Korn JH, Revel M. 1986. Structure and expression of cDNA and genes for human interferon b-2, a distinct species inducible by growth-stimulatory cytokines. *EMBO J.* 5:2529–37
37. Yasukawa K, Hirano T, Watanabe Y, et al. 1987. Structure and expression of human B cell stimulatory factor-2 (BSF-2/IL-6) gene. *EMBO J.* 6(10):2939–45
38. Simpson RJ, Moritz RL, Rubira MR, Van Snick J. 1988. Murine hybridoma/plasmacytoma growth factor: complete amino acid sequence and relation to human interleukin-6. *Eur. J. Biochem.* 176:187–97
39. Sehgal PB, Walther Z, Tamm I. 1987. Rapid enhancement of $\beta 2$ -interferon/B-cell differentiation factor BSF-2 gene expression in human fibroblasts by diacylglycerols and the calcium ionophore A23187. *Proc. Natl. Acad. Sci. USA* 84:3633–35
40. Zhang Y, Lin J-X, Vilcek J. 1988. Synthesis of interleukin-6 (interferon- $\beta 2$ /B cell stimulatory factor 2) in human fibroblasts is triggered by an increase in intracellular cyclic AMP. *J. Biol. Chem.* 263:6177–82
41. Zhang Y, Lin J-X, Yip YK, Vilcek J. 1988. Enhancement of cAMP levels and of protein kinase activity by tumor necrosis factor and interleukin-1 in human fibroblasts: role in the induction of interleukin-6. *Proc. Natl. Acad. Sci. USA* 85:6802–6
42. Zhang Y, Broser M, Rom WN. 1994. Activation of the interleukin 6 gene by *Mycobacterium tuberculosis* or lipopolysaccharide is mediated by nuclear factors NF-IL6 and NF-kappa B. *Proc. Natl. Acad. Sci. USA* 91(6):2225–29
43. Muraoka O, Kaisho T, Tanabe M, Hirano T. 1993. Transcriptional activation of the interleukin-6 gene by HTLV-1 p40tax through an NF-kB-like binding site. *Immunol. Lett.* 37:159–65
44. Yamashita I, Katamine S, Moriuchi R, et al. 1994. Transactivation of the human interleukin-6 gene by human T-lymphotropic virus type 1 Tax protein. *Blood* 84(5):1573–78
45. Yan SF, Tritto I, Pinsky D, et al. 1995. Induction of interleukin 6 (IL-6) by hypoxia in vascular cells. Central role of the binding site for nuclear factor-IL-6. *J. Biol. Chem.* 270(19):11463–71
46. Dendorfer U, Oettgen P, Libermann TA. 1994. Multiple regulatory elements in the interleukin-6 gene mediate induction by prostaglandins, cyclic AMP, and lipopolysaccharide. *Mol. Cell. Biol.* 14(7):4443–54
47. Scala G, Ruocco MR, Ambrosino C, et al. 1994. The expression of the interleukin 6 gene is induced by the human immunodeficiency virus 1 TAT protein. *J. Exp. Med.* 179(3):961–71
48. Brach MA, Gruss HJ, Kaisho T, et al. 1993. Ionizing radiation induces expression of interleukin 6 by human fibroblasts involving activation of nuclear factor-kappa B. *J. Biol. Chem.* 268(12):8466–72
49. Margulies L, Sehgal PB. 1993. Modulation of the human interleukin-6 promoter (IL-6) and transcription factor C/EBP β (NF-IL6) activity by p534 species. *J. Biol. Chem.* 268:15096–100
50. Janaswami PM, Kalvakolanu DV, Zhang Y, Sen GC. 1992. Transcriptional repression of interleukin-6 gene by adenoviral E1A proteins. *J. Biol. Chem.* 267(34):24886–91
51. Brach MA, de Vos S, Arnold C, et al. 1992. Leukotriene B4 transcriptionally activates interleukin-6 expression involving NK-chi B and NF-IL6. *Eur. J. Immunol.* 22(10):2705–11
52. Gruss HJ, Brach MA, Herrmann F. 1992. Involvement of nuclear factor-kappa B in induction of the interleukin-6 gene

- by leukemia inhibitory factor. *Blood* 80(10):2563–70
53. Yee CS, Messner HA, Minden MD. 1991. Regulation of interleukin-6 expression in the lymphoma cell line OCI-LY3. *J. Cell. Physiol.* 148:426–29
 54. Zhang Y, Lin J-X, Vilcek J. 1990. Interleukin-6 induction by tumor necrosis factor and interleukin-1 in human fibroblasts involves activation of a nuclear factor binding to a kappa B-like sequence. *Mol. Cell. Biol.* 10:3818–23
 55. Shimizu H, Mitomo K, Watanabe T, et al. 1990. Involvement of a NF- κ B-like transcription factor in the activation of the interleukin-6 gene by inflammatory lymphokines. *Mol. Cell. Biol.* 10:561–68
 56. Tanabe O, Akira S, Kamiya T, et al. 1988. Genomic structure of the murine IL-6 gene: high degree of conservation of potential regulatory sequences between mouse and human. *J. Immunol.* 141:3875–81
 57. Woloski BMRNJ, Smith EM, Meyer WJ III, et al. 1985. Corticotropin-releasing activity in monokines. *Science* 230: 1035–37
 58. Kohase M, Henriksen-DiStefano D, May LT, Vilcek J. 1987. Dexamethasone inhibits feedback regulation of the mitogenic activity of tumor necrosis factor, interleukin-1 and epidermal growth factor in human fibroblasts. *J. Cell. Physiol.* 132:271–78
 59. Feyen HJM, Elford P, Ki Padova FE, Trechsel U. 1989. Interleukin-6 is produced by bone and modulated by parathyroid hormone. *J. Bone Miner. Res.* 4:633–38
 60. Ray A, LaForge KS, Sehgal PB. 1990. On the mechanism for efficient repression of the interleukin-6 promoter by glucocorticoids: enhancer, TATA box, and RNA start site (Inr motif) occlusion. *Mol. Cell. Biol.* 10:5736–46
 61. Ray A, LaForge KS, Sehgal PB. 1991. Repressor to activator switch by mutations in the glucocorticoid receptor: Is direct DNA binding necessary? *Proc. Natl. Acad. Sci. USA* 88(16):7086–90
 62. Ray A, Prefontaine KE. 1994. Physical association and functional antagonism between the p65 subunit of transcription factor NF-kappa B and the glucocorticoid receptor. *Proc. Natl. Acad. Sci. USA* 91(2):752–56
 63. Deleted in proof
 64. Scheinman RI, Cogswell PC, Lofquist AK, Baldwin AS Jr. 1995. Role of transcriptional activation of I κ B α in mediation of immunosuppression by glucocorticoids. *Science* 270:283–86
 65. Auphan N, DiDonato A, Rosette C, 1995. Immunosuppression by glucocorticoids: inhibition of NF- κ B activity through induction of I κ B synthesis. *Science* 270:286–290
 66. Navarra P, Tsagarakis S, Faria M, et al. 1990. Interleukin-1 and -6 stimulate the release of corticotropin-releasing hormone from rat hypothalamus *in vitro* via eicosanoid cyclooxygenase pathway. *Endocrinology* 128:37–44
 67. Lyson K, Milenkovic K, McCann SM. 1991. The stimulatory effect of interleukin-6 on corticotropin-releasing factor and thyrotropin-releasing hormone secretion *in vitro*. *Prog. Neuroendocrinol. Immunol.* 4:161–65
 68. Tabibzadeh SS, Poubouridis D, May LT, Sehgal PB. 1989. Interleukin-6 immunoreactivity in human tumors. *Am. J. Pathol.* 135:427–33
 69. Kania DM, Binkley N, Checovich M, et al. 1995. Elevated plasma levels of interleukin-6 in postmenopausal women do not correlate with bone density. *J. Am. Geriatr. Soc.* 43(3):236–39
 70. Ralston SH. 1994. Analysis of gene expression in human bone biopsies by polymerase chain reaction: evidence for enhanced cytokine expression in postmenopausal osteoporosis. *J. Bone Miner. Res.* 9:883–90
 71. Daynes RA, Araneo BA, Ershler WB, et al. 1993. A dysregulation in the produc-

- tion of interleukin-6 with normal aging: probable linkage to the age associated depression in dehydroepiandrosterone (DHEA) and its sulfated derivative. *J. Immunol.* 150:5219–30
72. Pacifici R, Brown C, Puscheck E, et al. 1991. Effect of surgical menopause and estrogen replacement on cytokine release from human blood mononuclear cells. *Proc. Natl. Acad. Sci. USA* 88:5134–38
 73. Pioli G, Basini G, Pedtazzoni M, et al. 1992. Spontaneous release of interleukin-1 and interleukin-6 by peripheral blood mononuclear cells after oophorectomy. *Clin. Sci.* 83:503–7
 74. Kassem M, Harris SA, Spelsberg TC, Riggs BL. 1996. Estrogen inhibits interleukin-6 production and gene expression in a human osteoblastic cell line with high levels of estrogen receptor. *J. Bone Miner. Res.* 11:193–99
 75. Chaudhary LR, Spelsberg TC, Riggs BL. 1992. Production of various cytokines by normal human osteoblast-like cells in response to interleukin-1 β and tumor necrosis factor- α : lack of regulation by 17 β -estradiol. *Endocrinology* 130:2528–34
 76. Kitazawa R, Kimble RB, Vannice JL, et al. 1994. Interleukin-1 receptor antagonist and tumor necrosis factor binding protein decrease osteoclast formation and bone resorption in ovariectomized mice. *J. Clin. Invest.* 94:2397–406
 77. Rickard D, Russel G, Gowen M. 1992. Oestradiol inhibits the release of tumor necrosis factor but not interleukin-6 from adult human osteoblasts in vitro. *Osteoporosis Int.* 2:94–102
 78. Rifas L, Kenney JS, Marcelli M, et al. 1995. Production of interleukin-6 in human osteoblasts and human bone marrow stromal cells: evidence that induction by interleukin-1 and tumor necrosis factor- α is not regulated by ovarian steroids. *Endocrinology* 136:4056–67
 79. Pottratz ST, Bellido T, Mocharla H, et al. 1994. 17 β -Estradiol inhibits expression of human promoter-reporter constructs by a receptor-dependent mechanism. *J. Clin. Invest.* 93(3):944–50
 80. Ray A, Prefontaine KE, Ray P. 1994. Down-modulation of interleukin-6 gene expression by 17 beta-estradiol in the absence of high affinity DNA binding by the estrogen receptor. *J. Biol. Chem.* 269(17):12940–46
 81. Deleted in proof
 82. Stein B, Yang MX. 1995. Repression of the interleukin-6 promoter by estrogen receptor is mediated by NF- κ B and C/EBP β . *Mol. Cell. Biol.* 15(9):4971–79
 83. Sun WH, Keller ET, Stebler BS, Ershler WB. 1998. Estrogen inhibits phorbol ester-induced I κ B α transcription and protein degradation. *Biochem. Biophys. Res. Commun.* 244:691–95
 84. Galien R, Evans HF, Garcia T. 1996. Involvement of CCAAT/enhancer binding protein and nuclear factor- κ B binding sites in interleukin-6 promoter inhibition by estrogens. *Mol. Endocrinol.* 10:713–22
 85. Bellido T, Jilka RL, Boyce BF, et al. 1995. Regulation of interleukin-6, osteoclastogenesis, and bone mass by androgens. The role of the androgen receptor. *J. Clin. Inv.* 95(6):2886–95
 86. Keller ET, Chang C, Ershler WB. 1996. Inhibition of NF κ B activity through maintenance of I κ B α levels contributes to dihydrotestosterone-mediated repression of the interleukin-6 promoter. *J. Biol. Chem.* 271:26267–75
 87. Zhang J, Keller ET, Pugh TD, et al. 1998. Orchiectomy increases bone marrow interleukin-6 expression in mice. *Calcif. Tissue Int.* 62:219–26
 88. Smith EP, Boyd J, Frank GR, et al. 1994. Estrogen resistance caused by a mutation in the estrogen-receptor gene in a man. *N. Engl. J. Med.* 331:1056–61
 89. Yamasaki K, Taga T, Hirata Y, et al. 1988. Cloning and expression of the human interleukin-6 (BSF-2/IFN β 2) receptor. *Science* 241:825–28

90. Schooltink H, Stoyan T, Lenz D, et al. 1991. Structural and functional studies on the human hepatic interleukin-6 receptor. *Biochem. J.* 277:659-64
91. Kluck PMC, Wiegant J, Hansen RPM, et al. 1993. The human interleukin-6 receptor α chain gene is localized on chromosome 1 band q21. *Hum. Genet.* 90:542-44
92. Bazan JF. 1989. A novel family of growth factor receptors: a common binding domain in the growth hormone, prolactin, erythropoietin and IL-6 receptors, and the p75 IL-2 receptor β -chain. *Biochem. Biophys. Res. Commun.* 164:788-95
93. Cosman D, Lyman SD, Idzerda RL, et al. 1990. A new cytokine receptor superfamily. *Trends Biol. Sci.* 15:265-70
94. Taga T, Hibi M, Murakami M, et al. 1992. Interleukin-6 receptor and signals. *Chem. Immunol.* 1992; 51:181-204
95. Taga T, Hibi M, Hirata Y, et al. 1989. Interleukin-6 triggers the association of its receptor with a possible signal transducer, gp 130. *Cell* 58:573-81
96. Murakami M, Hibi M, Nakagawa N, et al. 1993. IL-6-induced homodimerization of gp 130 and associated activation of a tyrosine kinase. *Science* 260:1808-10
97. Ward LD, Howlett GJ, Discolo G, et al. 1994. High affinity interleukin-6 receptor is a hexameric consisting of two molecules each of interleukin-6, receptor, and gp-130. *J. Biol. Chem.* 269:23286-89
98. Novick D, Engelmann H, Wallach D, Rubinstein M. 1989. Soluble cytokine receptors are present in normal human urine. *J. Exp. Med.* 170:1409-14
99. Lust JA, Jelinek DF, Donovan KA, et al. 1994. Sequence, expression and function of an mRNA encoding of the human interleukin-6 receptor (sIL-6R). *Curr. Top. Microbiol. Immunol.* 194:199-205
100. Mullberg J, Schooltink J, Stoyan T, et al. 1993. The soluble interleukin-6 receptor is generated by shedding. *Eur. J. Immunol.* 23:473-80
101. Nakajima T, Yamamoto S, Cheng M, et al. 1992. Soluble interleukin-6 receptor is released from receptor-bearing cell line *in vitro*. *Jpn. J. Cancer Res.* 83:373-78
102. Snyers L, Fontaine V, Content J. 1989. Modulation of interleukin-6 receptors in human cells. *Ann. NY Acad. Sci.* 557:388-93
103. Lin SC, Yamate T, Taguchi Y, et al. 1997. Regulation of the gp80 and gp130 subunits of the IL-6 receptor by sex steroids in the murine bone marrow. *J. Clin. Invest.* 100:1980-90
104. Tang B, Matsuda T, Akira S, et al. 1991. Age-associated increase in interleukin 6 in MRL/lpr mice. *Int. Immunol.* 3(3): 273-78
105. Suzuki H, Yasukawa K, Saito T, et al. 1993. Serum soluble interleukin-6 receptor in MRL/lpr mice is elevated with age and mediates the interleukin-6 signal. *Eur. J. Immunol.* 23(5):1078-82
106. Effros RB, Svoboda K, Walford RL. 1993. Influence of age and caloric restriction on macrophage IL-6 and TNF production. *Lymphokine Cytokine Res.* 10(5):347-51
107. Zhou D, Chrest FJ, Adler W, et al. 1993. Increased production of TGF-beta and IL-6 by aged spleen cells. *Immunol. Lett.* 36(1):7-11
108. Foster KD, Conn CA, Kluger MJ. 1992. Fever, tumor necrosis factor, and interleukin-6 in young, mature, and aged Fischer 344 rats. *Am. J. Physiol.* 262: R211-15
109. Wei J, Xu H, Davies JL, Hemmings GP. 1992. Increase of plasma IL-6 concentrations with age in healthy subjects. *Life Sci.* 51:1953-56
110. Goya RG, Gagnerault MC, De Moraes, et al. 1992. In vivo effects of growth hormone on thymus function in aging mice. *Brain Behav. Immunol.* 6:341-54
111. Ershler WB, Sun WH, Binkley N, et al.

1993. Interleukin-6 and aging: blood levels and mononuclear cell production increase with advancing age and in vitro production is modifiable by dietary restriction. *Lymphokine Cytokine Res.* 12(4):225–30
112. Sandermann J, Kruse A, Frercks HJ, et al. 1993. Investigations of the lymphokine system in elderly individuals. *Mech. Ageing Dev.* 70(1–2):149–59
113. McKane R, Khosla S, Peterson J. 1993. Effect of age and menopause on serum interleukin-6 levels in women. *J. Bone Miner. Res.* 8:S157 (Suppl.)
114. Fagiolo U, Cossarizza A, Scala E, et al. 1993. Increased cytokine production in mononuclear cells of healthy elderly people. *Eur. J. Immunol.* 23(9):2375–78
115. Liao Z, Caucino JA, Schnipper SM, et al. 1993. Increased urinary cytokines in the elderly. *Aging Immunol. Infect. Dis.* 4:139–53
116. Kawano M, Hirano T, Matsuda T, et al. 1988. Autocrine generation and requirement of BSF-2/IL-6 for human multiple myeloma. *Nature* 332:83–85
117. Klein B, Zhang X, Jourdan M, et al. 1989. Paracrine rather than autocrine regulation of myeloma-cell growth and differentiation by interleukin-6. *Blood* 73(2):517–26
118. Asaoku H, Kaeano M, Iwato K, et al. 1988. Decrease in BSF-2/IL-6 response in advanced cases of multiple myeloma. *Blood* 72:429–32
119. Kato H, Kinoshita T, Suzuki S, et al. 1996. Elevated serum interleukin-6 (IL-6) is derived from neoplastic lymphoid cells in patients with B-cell non-Hodgkin's lymphoma: correlation with extent of IL-6 expression and serum concentration. *Br. J. Haematol.* 92(4):1014–21
120. Kurzrock R, Redman J, Cabanillas F, et al. 1993. Serum interleukin 6 levels are elevated in lymphoma patients and correlate with survival in advanced Hodgkin's disease and with B symptoms. *Cancer Res.* 53:2118–22
121. Lavabre-Bertrand T, Exbrayat C, Liautard J, et al. 1995. Detection of membrane and soluble interleukin-6 receptor, in lymphoid malignancies. *Br. J. Haematol.* 91(4):871–77
122. Yoshizaki K, Matsuda T, Nishimoto N, et al. 1989. Pathogenic significance of interleukin-6 (IL-6/BSF-2) in Castleman's disease. *Blood* 74:1360–67
123. Brandt SJ, Bodine DM, Dunbar CE, Nienhuis AW. 1990. Dysregulated interleukin 6 expression produces a syndrome resembling Castleman's disease in mice. *J. Clin. Invest.* 86:592–99
124. Gause A, Scholz R, Klein S, et al. 1991. Increased levels of circulating interleukin-6 in patients with Hodgkin's disease. *Hematol. Oncol.* 9(6):307–13
125. Yamamura M, Honda M, Yamada Y, et al. 1996. Increased levels of interleukin-6 (IL-6) in serum and spontaneous in vitro production of IL-6 by lymph node mononuclear cells of patients with angioimmunoblastic lymphadenopathy with dysproteinemia (AILD) and clinical effectiveness of cyclosporin A. *Leukemia* 10(9):1504–8
126. Kim SS, Chung SM, Choi IP, Byun KH. 1995. Expression of interleukin-6 in polymorphic reticulosis—immunohistochemical study. *J. Korean Med. Sci.* 10(5):324–28
127. Stasi R, Zinzani L, Galieni P, et al. 1995. Clinical implications of cytokine and soluble receptor measurements in patients with newly-diagnosed aggressive non-Hodgkin's lymphoma. *Eur. J. Haematol.* 54(1):9–17
128. Seymour JF, Talpaz M, Cabanillas F, et al. 1995. Serum interleukin-6 levels correlate with prognosis in diffuse large-cell lymphoma. *J. Clin. Oncol.* 13(3):575–82
129. Kanno H, Yasunaga Y, Iuchi K, et al. 1996. Interleukin-6 mediated growth enhancement of cell lines derived from pyothorax-associated lymphoma. *Lab. Invest.* 75(2):167–73
130. Freeman GJ, Freedman AS, Rabinowe

- SN, et al. 1989. Interleukin 6 gene expression in normal and neoplastic B cells. *J. Clin. Invest.* 83:1512-18
131. Yee C, Biondi A, Wang XH, et al. 1989. A possible autocrine role for interleukin-6 in two lymphoma cell lines. *Blood* 74(2):798-804
 132. Takeshita M, Sumiyoshi Y, Masuda, Y, et al. 1993. Cytokine (interleukin-1 alpha, interleukin-1 beta, tumor necrosis factor alpha, and interleukin-6)-possessing cells in lymph nodes of malignant lymphoma. *Pathol. Res. Pract.* 189(1):18-25
 133. Voorzanger N, Toutitou R, Garcia E, et al. 1996. Interleukin (IL)-10 and IL-6 are produced in vivo by non-Hodgkin's lymphoma cells and act as cooperative growth factors. *Cancer Res.* 56(23):5499-505
 134. Frith CH, Davis TM, Zolotor LA, Townsend JW. 1980. Histiocytic lymphoma in the mouse. *Leuk. Res.* 4:651-62
 135. Volk MJ, Pugh TD, Kim MJ, et al. 1993. Dietary restriction from middle age attenuates age-associated lymphoma development and IL-6 dysregulation in C57BL/6 mice. *Cancer Res.* 54:3054-63
 136. Weindruch R, Walford RL. 1988. *The Retardation of Aging and Disease by Dietary Restriction*. Springfield, IL: Thomas
 137. Peled A, Tzehoval E, Haran-Ghera N. 1995. Role of cytokines in termination of the B cell lymphoma dormant state in AKR mice. *Leukemia* 9(6):1095-101
 138. Hilbert DM, Kopf M, Mock BA, et al. 1995. Interleukin-6 is essential for in vivo development of B lineage neoplasms. *J. Exp. Med.* 182(1):243-48
 139. Zhang XG, Klein B, Bataille R. 1989. Interleukin-6 is a potent myeloma-cell growth factor in patients with aggressive multiple myeloma. *Blood* 74:11-13
 140. Gaillard JP, Bataille R, Brailly H, et al. 1993. Increased and highly stable levels of functional soluble interleukin-6 receptor in sera of patients with monoclonal gammopathy. *Eur. J. Immunol.* 23:820-24
 141. Solary E, Guiguet M, Zeller V, et al. 1992. Radioimmunoassay for the measurement of serum IL-6 and its correlation with tumor cell mass parameters in multiple myeloma. *Am. J. Hematol.* 39:163-71
 142. Bataille R, Jourdan M, Zhang XG, Klein B. 1989. Serum levels of interleukin-6, a potent myeloma cell growth factor, as a reflection of disease severity in plasma cell dyscrasias. *J. Clin. Invest.* 84:2008-11
 143. Ballester OF, Moscinski LC, Lyman GH, et al. 1994. High levels of interleukin-6 are associated with low tumor burden and low growth fraction in multiple myeloma. *Blood* 83:1903-8
 144. Stauder R, Van Driel M, Schwarzler C, et al. 1996. Different CD44 splicing patterns define prognostic subgroups in multiple myeloma. *Blood* 88(8):3101-8
 145. Klein B, Lu ZY, Gallard JP, et al. 1992. Inhibiting IL-6 in human multiple myeloma. *Curr. Top. Microbiol. Immunol.* 182:237-43
 146. Klein B, Lu ZY, Bataille R. 1992. Clinical applications of IL6 inhibitors. *Res. Immunol.* 143(7):774-76
 147. Klein B, Wijdenes J, Zhang XG, et al. 1991. Murine anti-interleukin-6 monoclonal antibody therapy for a patient with plasma cell leukemia. *Blood* 5:1198-204
 148. Sipe JD, Bartle LM, Loose LD. 1992. Modification of proinflammatory production by the antirheumatic agents Tenidap and Naproxen. A possible correlate with the clinical acute phase response. *J. Immunol.* 148:480-84
 149. Wagge A, Slupphaug G, Shalaby R. 1990. Glucocorticoids inhibit production of IL-6 from monocytes, endothelial cells and fibroblasts. *Eur. J. Immunol.* 20:2439-43
 150. Keller ET, Ershler WB. 1995. Effect of IL-6 receptor antisense oligodeoxynucleotides on interleukin-6 production in human monocytes. *J. Biol. Chem.* 270:11111-11115

- cleotides on in vitro proliferation of myeloma cells. *J. Immunol.* 154:4091–98
151. Manolagas SC, Jilka RL. 1995. Bone marrow, cytokines, and bone remodeling. Emerging insights into the pathophysiology of osteoporosis. *N. Engl. J. Med.* 332(5):305–11
 152. Lorenzo JA. 1991. The role of cytokines in the regulation of local bone resorption. *Crit. Rev. Immunol.* 11:195–213
 153. Bellido T, Girasole G, Passeri G, et al. 1994. Sex steroids regulate the expression of the gp130 transduction pathway by bone marrow and bone cells. *Bone Miner.* 25(S1):S39
 154. Fang MA, Hahn TJ. 1991. Effects of interleukin-6 on cellular function in UMR-106–01 osteoblast-like cells. *J. Bone Miner. Res.* 6:133–39
 155. Taguchi Y, Yamate T, Mocharia H, et al. 1996. Interleukin-6 induces osteoblast differentiation in uncommitted embryonic fibroblasts (EF). *J. Bone Miner. Res.* 11:S101 (Suppl.) (Abstr.)
 156. Ishimi Y, Miyaura C, Jin CH, et al. 1990. IL-6 is produced by osteoblasts and induces bone resorption. *J. Immunol.* 145:3297–303
 157. Green H, Schotland S, Sella Z, Kleeman CR. 1994. Interleukin-6 attenuates agonist-mediated calcium mobilization in murine osteoblastic cells. *J. Clin. Invest.* 93:2340–50
 158. Littlewood AJ, Aarden LA, Evans DB, et al. 1991. Human osteoblastlike cells do not respond to interleukin-6. *J. Bone Miner. Res.* 6:141–48
 159. Greenfield EM, Shaw SM, Gornik SA, Banks MA. 1995. Adenyl cyclase and interleukin 6 are downstream effectors of parathyroid hormone resulting in stimulation of bone resorption. *J. Clin. Invest.* 96:1238–44
 160. Löwik CWGM, van der Pluijm G, Bloys H, et al. 1989. Parathyroid hormone (PTH) and PTH-like protein (PLP) stimulate interleukin-6 production by osteogenic cells: a possible role of interleukin-6 in osteoclastogenesis. *Biochem. Biophys. Res. Commun.* 162:1546–52
 161. Rahman S, Bunning RAD, Dobson PRM, et al. 1992. Bradykinin stimulates the production of prostaglandin E2 and interleukin-6 in human osteoblast-like cells. *Biochim. Biophys. Acta* 1135:97–102
 162. Zheng MH, Wood DJ, Wysocki S, et al. 1994. Recombinant human bone morphogenetic protein-2 enhances expression of interleukin-6 and transforming growth factor- β 1 genes in normal human osteoblast-like cells. *J. Cell. Physiol.* 159:76–82
 163. Al-Humidan A, Ralston SH, Hughes DE, et al. 1991. Interleukin-6 does not stimulate bone resorption in neonatal mouse calvariae. *J. Bone Miner. Res.* 6:3–8
 164. Flanagan AM, Stow MD, Williams R. 1995. The effect of interleukin-6 and soluble interleukin-6 receptor protein on the bone resorptive activity of human osteoclasts generated *in vitro*. *J. Pathol.* 176:289–97
 165. Roodman GD, Kurihara N, Ohsaki Y, et al. 1992. Interleukin-6: a potential autocrine/paracrine factor in Paget's disease of bone. *J. Clin. Invest.* 89:46–52
 166. Black KS, Mundy GR, Garrett IR. 1991. Interleukin-6 causes hypercalcemia in vivo, and enhances the bone resorbing potency of interleukin-1 and tumor necrosis factor by two orders of magnitude in vitro. *J. Bone Miner. Res.* 6:S271 (Suppl.)
 167. Ohsaki Y, Takahashi S, Scarcez T, et al. 1992. Evidence for an autocrine/paracrine role for interleukin-6 in bone resorption by giant cells from giant cell tumors of bone. *Endocrinology* 131:2229–34
 168. Mrak RE, Sheng JG, Griffin WS. 1995. Glial cytokines in Alzheimer's disease review and pathogenic implications. *Hum. Pathol.* 26:816–23
 169. Bauer J, Ganter U, Strauss S, et al. 1992. The participation of interleukin-6 in the

- pathogenesis of Alzheimer's disease. *Res. Immunol.* 143(6):650–57
170. Vandenabelle P, Fiers W. 1991. Is amyloidogenesis during Alzheimer's disease due to an IL-1/IL-6-mediated "acute phase response" in the brain? *Immunol. Today* 12:217–19
 171. Ganter U, Strauss S, Jonas V, et al. 1991. Alpha-2 macroglobulin synthesis in interleukin-6 stimulated human neuronal (SH-SY5Y neuroblastoma) cells. *FEBS Lett.* 282:127–31
 172. Strauss S, Bauer J, Ganter U, et al. 1992. Detection of interleukin-6 and α 2-macroglobulin immunoreactivity in cortex and hippocampus of Alzheimer's disease patients. *Lab. Invest.* 66:223–30
 173. Bauer J, Strauss S, Schreiter-Gasser U, et al. 1991. Interleukin-6 and α 2-macroglobulin indicate an acute-phase state in Alzheimer's disease cortices. *FEBS Lett.* 285:111–14
 174. Abraham CR, Shirahama T, Potter H. 1990. α 1-Antichymotrypsin is associated solely with amyloid deposits containing the β -protein. Amyloid and cell localization of α 1-antichymotrypsin. *Neurobiol. Aging* 11:123–26
 175. Wood JA, Wood PL, Ryan R, et al. 1993. Cytokine indices in Alzheimer's temporal cortex: no changes in mature IL-1 beta or IL-1RA but increases in the associated acute phase proteins IL-6, alpha 2-macroglobulin and C-reactive protein. *Brain Res.* 629(2):245–52
 176. Blum-Degen D, Muller T, Kuhn W, et al. 1995. Interleukin-1 beta and interleukin-6 are elevated in the cerebrospinal fluid of Alzheimer's disease and de novo Parkinson's disease patients. *Neurosci. Lett.* 202(1–2):17–20
 177. Yamada K, Kono K, Umegaki H, et al. 1995. Decreased interleukin-6 level in the cerebrospinal fluid of patients with Alzheimer-type dementia. *Neurosci. Lett.* 186(2–3):219–21
 178. Chao CC, Ala TA, Hu S, et al. 1994. Serum cytokine levels in patients with Alzheimer's disease. *Clin. Diagn. Lab. Immunol.* 1(4):433–36
 179. van Duijn CM, Hofman A, Nagelkerken L. 1990. Serum levels of interleukin-6 are not elevated in patients with Alzheimer's disease. *Neurosci. Lett.* 108:350–54
 180. Cadman ED, Witte DG, Lee CM. 1994. Regulation of the release of interleukin-6 from human astrocytoma cells. *J. Neurochem.* 63(3):980–87
 181. Sebire G, Emilie D, Wallon C, et al. 1993. In vitro production of IL-6, IL-1 β , and tumor necrosis factor- α by human embryonic microglial and neural cells. *J. Immunol.* 150:1517–23
 182. Del Bo R, Angeretti N, Lucca E, et al. 1995. Reciprocal control of inflammatory cytokines, IL-1, IL-6 and beta-amyloid production in cultures. *Neurosci. Lett.* 188(1):70–74
 183. Hull M, Strauss S, Berger M, et al. 1996. The participation of interleukin-6, a stress-inducible cytokine, in the pathogenesis of Alzheimer's disease. *Behav. Brain Res.* 78(1):37–41
 184. Huell M, Strauss S, Volk B, et al. 1995. Interleukin-6 is present in early stages of plaque formation and is restricted to the brains of Alzheimer's disease patients. *Acta Neuropathol.* 89(6):544–51
 185. Cacabelos R, Franco-Maside A, Alvarez XA. 1991. Interleukin 1 in Alzheimer's disease and multi-infarct dementia: neurophysiological correlations. *Methods Exp. Clin. Pharmacol.* 13:703–8
 186. Huberman M, Sredni B, Stern L, et al. 1995. IL-2 and IL-6 secretion in dementia: correlation with type and severity of disease. *J. Neurol. Sci.* 130(2):161–64
 187. Shalit F, Sredni B, Stern L, et al. 1994. Elevated interleukin-6 secretion levels by mononuclear cells of Alzheimer's patients. *Neurosci. Lett.* 174(2):130–32
 188. Fattori E, Lazzaro D, Musiani P, et al. 1995. IL-6 expression in neurons of transgenic mice causes reactive astrogliosis and increase in ramified microglial

- cells but no neuronal damage. *Eur. J. Neurosci.* 7(12):2441–49
189. Bellinger FP, Madamba SG, Campbell IL, Siggins GR. 1995. Reduced long term potentiation in the dentate gyrus of transgenic mice with cerebral overexpression of interleukin-6. *Neurosci. Lett.* 198(2):95–98
 190. Brett FM, Mizisin AP, Powell HC, Campbell IL. 1995. Evolution of neuropathologic abnormalities associated with blood brain barrier breakdown in transgenic mice expressing interleukin-6 in astrocytes. *J. Neuropathol. Exp. Neurol.* 54(6):766–75
 191. Chiang CS, Stalder A, Samimi A, Campbell IL. 1994. Reactive gliosis as a consequence of interleukin-6 expression in the brain: studies in transgenic mice. *Dev. Neurosci.* 16(3–4):212–21
 192. Steffensen SC, Campbell IL, Henriksen SJ. 1994. Site-specific hippocampal pathophysiology due to cerebral overexpression of interleukin-6 in transgenic mice. *Brain Res.* 652(1):149–53
 193. Campbell IL, Abraham CR, Masilaah E, et al. 1993. Neurologic disease induced in transgenic mice by cerebral overexpression of interleukin-6. *Proc. Natl. Acad. Sci. USA* 90(21):10061–65
 194. Kawas C, Resnick S, Morrison A, et al. 1997. A prospective study of estrogen replacement therapy and the risk of developing Alzheimer's disease: the Baltimore Longitudinal Study of Aging. *Neurology* 48(6):1517–21
 195. Stewart WF, Kawas C, Corrada M, Metter EJ. 1997. Risk of Alzheimer's disease and duration of NSAID use. *Neurology* 48(3):626–32
 196. Binkley NC, Sun WH, Checovich MM, et al. 1994. Effects of recombinant human interleukin-6 administration on bone in rhesus monkeys. *Lymphokine Cytokine Res.* 13:221–26
 197. Cohen HJ, Pieper CF, Harris T, et al. 1997. The association of plasma IL-6 levels with functional disability in community dwelling elderly. *J. Gerontol. Med. Sci.* 52:M201–8



CONTENTS

Inferior Vena Cava Interruption: How and When?, <i>Philippe Girard, Bernard Tardy, Hervé Decousus</i>	1
Ductal Carcinoma In Situ of the Breast, <i>Melvin J. Silverstein</i>	17
Gene Therapy for Adenosine Deaminase Deficiency, <i>Robertson Parkman, Kenneth Weinberg, Gay Crooks, Jan Nolte, Neena Kapoor, Donald Kohn</i>	33
Acupuncture: An Evidence-Based Review of the Clinical Literature, <i>David J. Mayer</i>	49
Role of Telomerase in Cell Senescence and Oncogenesis, <i>Virginia Urquidi, David Tarin, Steve Goodison</i>	65
Coronary Artery Disease in the Transplanted Heart, <i>M. Weis, W. von Scheidt</i>	81
Measures of Success and Health-Related Quality of Life in Lower-Extremity Vascular Surgery, <i>Joe Feinglass, Mark Morasch, Walter J. McCarthy</i>	101
New Horizons in the Treatment of Autoimmune Diseases: Immunoablation and Stem Cell Transplantation, <i>Alberto M. Marmont</i>	115
The Surgical Treatment of Parkinson's Disease, <i>Kenneth A. Follett</i>	135
Atherogenic Lipids and Endothelial Dysfunction: Mechanisms in the Genesis of Ischemic Syndromes, <i>Mark R. Adams, Scott Kinlay, Gavin J. Blake, James L. Orford, Peter Ganz, Andrew P. Selwyn</i>	149
Management of Patients with Hereditary Hypercoagulable Disorders, <i>C. Kearon, M. Crowther, J. Hirsh</i>	169
Neurocysticercosis: Updates on Epidemiology, Pathogenesis, Diagnosis, and Management, <i>A. Clinton White Jr.</i>	187
Anti-Cytokine Therapy for Rheumatoid Arthritis, <i>R. N. Maini, P. C. Taylor</i>	207
Artificial Skin, <i>J. T. Schulz III, R. G. Tompkins, J. F. Burke</i>	231
Age-Associated Increased Interleukin-6 Gene Expression, Late-Life Diseases, and Frailty, <i>William B. Ershler, Evan T. Keller</i>	245
Streptococcal Toxic Shock Syndrome Associated with Necrotizing Fasciitis, <i>Dennis L. Stevens</i>	271
Role of Cytokines in the Pathogenesis of Inflammatory Bowel Disease, <i>Konstantinos A. Papadakis, Stephan R. Targan</i>	289
Anorexia and Bulimia Nervosa, <i>W. H. Kaye, K. L. Klump, G. K. W. Frank, M. Strober</i>	299
The Role of Protein Traffic in the Progression of Renal Diseases, <i>Piero Ruggenenti, Giuseppe Remuzzi</i>	315
Carotid Artery Dissection, <i>C. Stapf, M. S. V. Elkind, J. P. Mohr</i>	329
Bacterial Vaginosis, <i>Jack D. Sobel</i>	349
Current Concepts in Cobalamin Deficiency, <i>Ralph Carmel</i>	357
Adjuvant Therapy for Breast Cancer, <i>G. Hortobagyi</i>	377
Kidney Transplantation from Living Unrelated Donors, <i>J. Michael Cecka</i>	393
Global Epidemiology of Influenza: Past and Present, <i>N. J. Cox, K. Subbarao</i>	407
The Spectrum of Human Herpesvirus 6 Infection: From Roseola Infantum to Adult Disease, <i>Mark Young Stoeckle</i>	423

Catheter Ablation for Atrial Fibrillation, <i>Pierre Jaïs, Dipen C. Shah, Michel Haïssaguerre, Meleze Hocini, Jing Tian Peng, Jacques Clémenty</i>	431
Genetic Disorders Affecting Proteins of Iron Metabolism: Clinical Implications, <i>Sujit Sheth, Gary M. Brittenham</i>	443
Genetics of Psychiatric Disease, <i>Wade H. Berrettini</i>	465
The Marfan Syndrome, <i>Reed E. Pyeritz</i>	481
Nonsteroidal Anti-Inflammatory Drugs and Cancer Prevention, <i>John A. Baron, Robert S. Sandler</i>	511
Lymphatic Mapping and Sentinel Lymph Node Biopsy in Patients with Breast Cancer, <i>Charles E. Cox, Siddharth S. Bass, Christa R. McCann, Ni Ni K. Ku, Claudia Berman, Kara Durand, Monica Bolano, Jessica Wang, Eric Peltz, Sarah Cox, Christopher Salud, Douglas S. Reintgen, Gary H. Lyman</i>	525
The Genetics of the Amyloidoses, <i>Joel N. Buxbaum, Clement E. Tagoe</i>	543