

## ORIGINAL COMMUNICATION

# Increased lipid peroxidation during long-term intervention with high doses of n-3 fatty acids (PUFAs) following an acute myocardial infarction

H Grundt<sup>1,2\*</sup>, DWT Nilsen<sup>2</sup>, MA Mansoor<sup>1</sup> and A Nordøy<sup>3</sup>

<sup>1</sup>Department of Clinical Chemistry, Central Hospital in Rogaland, Stavanger, Norway; <sup>2</sup>Department of Medicine, Central Hospital in Rogaland, Stavanger, Norway; and <sup>3</sup>Department of Medicine, University of Tromsø, Norway

**Objective:** To assess the oxidative burden of a highly concentrated compound of n-3 PUFAs as compared to corn oil by measuring thiobarbituric acid–malondialdehyde complex (TBA–MDA) by HPLC. We also studied the influence on TBA–MDA of statins combined with n-3 PUFAs or corn oil.

**Design:** A prospective, randomised, double-blind, controlled study.

**Setting:** One hospital centre in Stavanger, Norway.

**Subjects:** A total of 300 subjects with an acute myocardial infarction (MI).

**Interventions:** Gelatine capsules, containing 850–882 mg EPA and DHA as concentrated ethylesters, or 1 g of corn oil, were ingested in a dose of two capsules twice a day for at least 1 y. Alpha-tocopherol (4 mg) was added to all capsules to protect the PUFAs against oxidation.

**Results:** After 1 y TBA–MDA increased modestly in the n-3 PUFA group ( $n=125$ ), as compared to the corn oil group ( $n=130$ ),  $P=0.027$ . Multiple linear regression analyses of fatty acids in serum total phospholipids ( $n=56$ ) on TBA–MDA measured after 12 months intervention, showed no dependency. Performing best subsets regression, serum phospholipid concentration of arachidonic acid (20:4 n-6 PUFA) was identified as a predictor of TBA–MDA at 12 months follow-up,  $P=0.004$ .

We found no impact of statins on TBA–MDA.

**Conclusion:** TBA–MDA increased modestly after long-term intervention with n-3 PUFAs compared to corn oil post-MI, suggesting biological changes induced by n-3 PUFAs, rather than simply reflecting their concentration differences. The peroxidative potential of n-3 PUFAs was not modified by statin treatment.

**Sponsorship:** Pharmacia A/S and Pronova A/S, Norway.

*European Journal of Clinical Nutrition* (2003) 57, 793–800. doi:10.1038/sj.ejcn.1601730

**Keywords:** lipid peroxidation; thiobarbituric acid–malondialdehyde complex (TBA–MDA); n-3 PUFAs; myocardial infarction

## Introduction

Despite several favourable effects of n-3 PUFAs on inflammation and atherothrombosis, concern has been raised, as n-3 PUFAs may have a potential to increase oxidative stress (Harats *et al*, 1991; Meydani *et al*, 1991), resulting in the formation of lipid peroxides. During peroxidation of PUFAs several chain reactions may modify lipids and lipoproteins

containing PUFAs (Dargel, 1992; Halliwell & Chirico, 1993). Peroxidation of lipids and lipoproteins, especially low-density lipoprotein (LDL), is thought to play an important role in the development of atherosclerotic plaques (Avogaro *et al*, 1988; Boyd *et al*, 1989; Steinberg *et al*, 1989; Rosenfeld, 1991; Wiklund *et al*, 1991).

*In vitro* studies of lipid peroxidation suggest an increased susceptibility to oxidation with increasing number of double bonds in fatty acids (Cosgrove *et al*, 1987; Liu *et al*, 1997). However, the impact of n-3 PUFAs supplementation on lipid peroxidation is controversial (Nenseter & Drevon, 1996). Several clinical studies have reported no significant change in the lipid peroxidation following increased consumption of n-3 PUFAs (Nenseter *et al*, 1992; Bittolo-Bon *et al*, 1993;

\*Correspondence: H Grundt, Department of Internal Medicine, Central Hospital in Rogaland, POB 8100, 4068 Stavanger, Norway.  
E-mail: heidi@madlalia.no

Guarantor: H Grundt.

Received 8 September 2002; revised 2 December 2002; accepted 19 December 2002

Frankel *et al*, 1994; Eritsland *et al*, 1995; Bonanome *et al*, 1996; Wander *et al*, 1996; Brude *et al*, 1997; Hansen *et al*, 1998). No enhancement of the lipid peroxidation was noted after 9 months supplementation with 4 g n-3 PUFA concentrate in the shunt occlusion trial (Eritsland *et al*, 1995). Moreover, thiobarbituric acid reactive substances (TBARS) as a global marker of peroxidation remained unchanged in healthy males given 4 g purified EPA or DHA for 5 weeks, despite an immediate increase in chylomicron peroxidation *ex vivo* (Hansen *et al*, 1998). On the other hand, Meydani (Meydani *et al*, 1991) found increased TBARS in plasma samples from women given supplements of fish oil for 12 weeks. In agreement with these results, Harats *et al* (1991) observed both an increase in LDL TBARS and an accelerated metabolism of LDL by macrophages. These divergent/inconsistent findings may be because of different experimental settings, as these studies include relatively small sample sizes, different doses and compositions of n-3 PUFAs, short intervention periods, different amounts of antioxidants supplies and different methods used to quantify the oxidative burden and different methods to estimate TBARS.

As the beneficial effects of n-3 PUFAs (Kinsella *et al*, 1990; Schmidt & Dyerberg, 1994) may be opposed by an enhanced lipid peroxidation, the main aim of this study was to prospectively assess the impact of concentrated n-3 PUFAs vs corn oil on lipid peroxidation during 12 months of treatment in 300 acute myocardial infarction (MI) patients demonstrating improvement in lipids (Nilsen *et al*, 2001). As a marker of lipid peroxidation, we determined TBA-MDA measured by HPLC.

We decided to include ultrasensitive CRP ( $\mu$ CRP) and plasma total homocysteine (p-tHcy) in the baseline evaluation of our trial subjects, as these parameters have become approved markers of cardiovascular disease (Arnesen *et al*, 1995; Ridker *et al*, 1998).

Finally, we hypothesized that additional statin therapy might favourably reduce lipid peroxidation in subjects on n-3 PUFAs, in view of a recent study demonstrating beneficial effects on serum lipids and lipoproteins by short-term combined therapy (Nordøy *et al*, 1998).

## Subjects and methods

### Subjects

This prospective study was designed as part of a randomised, parallel, double-blind study evaluating the effects on clinical outcome and serum lipids of a concentrate of n-3 PUFAs, introduced early after an acute MI (Nilsen *et al*, 2001).

A total of 300 subjects (238 men and 62 women), 28–87 y of age, with an acute MI verified by WHO criteria, were recruited at one hospital centre from September 1995 until December 1996. Written informed consent was obtained from each subject. Regular supplementation of other fish-oil products was discontinued prior to intervention. Exclusion criteria have previously been described (Nilsen *et al*, 2001). All participants were included between the 4th and the 6th

day following the acute MI. Included subjects were randomly assigned to receive gelatine capsules (Pronova A/S, Oslo, Norway), containing 850–882 mg EPA and DHA as concentrated ethylesters, or 1 g of corn oil, both administered in a dose of two capsules twice a day for at least 1 y (total 3.464 g n-3 PUFAs or 4.000 g corn oil). This dose of n-3 PUFAs has previously been shown to be therapeutically useful in lowering triglyceride levels and blood pressure (Phillipson *et al*, 1985; Eritsland *et al*, 1995; Connor & Connor, 1997). Alpha-tocopherol (4 mg) was added into all capsules to protect the unsaturated fatty acids against oxidation. The study was double-blind. Intervention was initiated immediately after inclusion and collection of baseline blood samples. Details regarding compliance have previously been reported (Nilsen *et al*, 2001).

The phospholipid fatty acid profile was analysed at baseline and after 12 months of intervention in serum from 60 subjects randomly allocated from the study participants. In all, 56 of the allocated subjects had complete data throughout the study period. The study population was compared to 30 healthy individuals from a general out-of-hospital population, matched according to age, gender, smoking habits and diet. Their diet was similar to the average Norwegian diet. These subjects were not taking any supplementation of n-3 PUFAs.

Simvastatin in a dose of 20–40 mg was initiated prior to discharge and continued beyond 12 months follow-up in 95 patients. This subgroup was compared to 79 individuals free of statins during the evaluation period of 12 months.

### Clinical examination and laboratory measurements

Clinical follow-up, including a detailed patient history, clinical examination, blood tests, electrocardiogram and capsule counting was performed at inclusion, at 6 weeks, 6 months and at 1 y. All cardiac events (cardiac death, resuscitation, recurrent MI, unstable angina pectoris, revascularisation), noncardiac death, smoking habits, ongoing medication and average fishmeals per week were recorded. The blood samples were preceded by 12 h of fasting.

Serum for analyses of TBA-MDA and ultrasensitive C-reactive protein ( $\mu$ CRP) and plasma for analyses of homocysteine (Hcy) were immediately frozen and stored at  $-80^{\circ}\text{C}$ . The analyses were performed by a technician with no knowledge of the randomisation code.

TBA-MDA was measured in duplicate in serum samples heated with thiobarbituric acid at low pH, separated and determined by HPLC-fluorescence detection, based on previously described methods by Wong *et al* (1987).

$\mu$ CRP was determined in serum samples by an ultrasensitive immunoassay, using a kit from Behring Diagnostics, Marburg, Germany.

Total plasma Hcy was determined using an HPLC assay as previously described (Mansoor *et al*, 1992).

Serum triglycerides and total cholesterol were determined by enzymatic methods with kits provided by Roche Diag-

nostics, Mannheim, Germany (GPO-PAP, CHOP-PAP). HDL-cholesterol was similarly measured after precipitation with dextran sulphate and magnesium chloride (Burstein *et al*, 1970).

Analyses of fatty acids in serum phospholipids were performed at the University of Tromsø, Norway, as previously described (Nordøy *et al*, 1998).

Troponin-T (TnT), which is a sensitive marker of minor myocardial cell injury was quantified by a cardiac-specific second-generation troponin-T ELISA assay, using a high-affinity cardiac-specific TnT isoform antibody (Roche Diagnostics, Mannheim, Germany), (Muller-Bardorff *et al*, 1997), modified by Hetland *et al* (1995).

Creatine kinase-muscle brain (CK-MB) analysis was performed on a Johnson and Johnson Vitros 700 analyser, based upon the immunoinhibition principle (Gerhardt *et al*, 1982).

The study was approved by the Regional Board of Research Ethics and the Norwegian Health Authorities, and conducted in accordance with the Helsinki Declaration of 1975 as revised in 1983.

### Statistical methods

Potential differences between intervention groups at baseline regarding quantitative variables were tested by a two-sample Student's *t*-test, or in case of non-normality by the Mann-Whitney rank-sum test. Continuously distributed variables were given as mean and standard deviation (s.d.), while variables with more skewed distributions were given as median and upper and lower quartiles.

Changes in parameter values from baseline to follow-up were calculated for each individual patient and subjected to statistical analysis. As the normality assumption was fulfilled, the *t*-test was used to evaluate whether changes in the fatty acid composition of the serum phospholipids and in TBA-MDA from baseline to 12 months were significantly different from zero, and to compare changes between treatment groups. Also, comparison of TBA-MDA between

treatment groups at 12 months was performed in a multiway ANACOVA model with adjustments for age, gender and baseline values of TBA-MDA, triglycerides, HDL-cholesterol,  $\mu$ CRP and p-tHcy.

Multiple linear regression analyses were introduced to identify concentrations of fatty acids in serum phospholipids as potential predictors of TBA-MDA. Pearson's correlation coefficient was calculated to evaluate relations between different variables.

A statistically significant level of  $P < 0.05$  was applied for all tests. Adjustment for multiple pairwise comparisons has been performed using the Bonferroni correction.

The statistical analyses were performed using the statistical package SAS version 8.2.

## Results

### Baseline data

As previously reported, 150 subjects in each group, 123 (82%) receiving n-3 PUFAs and 129 (86%) receiving corn oil, fulfilled the criteria for complete compliance after 6 weeks (Nilsen *et al*, 2001). There were no significant differences in baseline demographic or clinical characteristics, and concomitant medication at baseline and follow-up did not differ between the groups, including use of fish oil prior to inclusion. Approximately 30% in the n-3 group and 25% in the corn oil group were taking fish oil supplements prior to inclusion, comparable to a daily dose of about 1 g n-3 PUFAs. The above-mentioned baseline data have previously been described by Nilsen *et al* (2001).

The weekly consumption of fish was two meals or less in the lower quartile and three meals or more in the upper quartile. The median intake consisted of three fish meals per week. The dietary habits of the participants were essentially unchanged throughout the study. The fatty acid composition in serum phospholipids was equal in both intervention groups at baseline (Table 1). However, the content of EPA in serum phospholipids was 25% higher in the group taking

**Table 1** Concentrations of fatty acids ( $\mu$ mol/l) (mean (s.d.)) in serum total phospholipids in 56 subjects randomly allocated from the study population

Fatty acids	n-3 group (n=28)		Corn oil group (n=28)	
	Baseline	12 months	Baseline	12 months
EPA (20:5 n-3)	62.9 (34.5)	189.6 (63.2)*** <sup>c</sup>	67.4 (38.6)	97.4 (91.0)
DHA (22:6 n-3)	180.2 (52.0)	213.4 (45.3)**	207.4 (77.3)	209.9 (71.5)
LA (18:2 n-6)	796.5 (223.2)	748.6 (183.0)	828.1 (216.4)	893.4 (192.1)
AA (20:4 n-6)	264.7 (85.5)	236.0 (53.3)	291.0 (85.6)	317.5 (83.4)* <sup>b</sup>
Total n-3 PUFA	284.3 (84.0)	457.6 (96.3)*** <sup>c</sup>	320.8 (117.1)	356.6 (136.5)
Total n-6 PUFA	1189.0 (297.2)	1062.0 (215.7)	1248.8 (294.1)	1332.5 (243.3) <sup>a</sup>
Total saturated FA	2030.7 (446.6)	2023.3 (306.1)	2103.2 (386.7)	2171.3 (324.8)
Total FA	3935.5 (870.2)	3920.9 (583.6)	4136.8 (800.7)	4252.6 (789.2)

LA—linoleic acid; AA—arachidonic acid.

No significant differences between intervention groups at baseline.

Significance of difference from baseline: \*\*\* $P < 0.001$ , \*\* $P = 0.011$ , \* $P = 0.035$ .

Significance of difference between groups: <sup>c</sup> $P < 0.001$ , <sup>b</sup> $P = 0.004$ , <sup>a</sup> $P = 0.01$ .

**Table 2** Contents of fatty acids ( $\mu\text{mol/l}$ ) (mean (s.d.), and (% of total fatty acids)) in serum phospholipids at baseline and serum total cholesterol, HDL-cholesterol and triglycerides (mmol/l) in 63 subjects randomly allocated from the study population, arranged according to whether fish oil supplementation was given before inclusion or not

Fatty acids	Fish oil supplementation before inclusion	
	Yes (n=17)	No (n=46)
EPA (20:5 n-3)	81.9 (45.4)* (2.0)	59.0 (28.2) (1.5)
DHA (22:6 n-3)	217.6 (71.2) (5.3)	191.7 (64.3) (4.8)
LA (18:2 n-6)	784.4 (214.5) (19.2)	839.5 (215.6) (20.9)
AA (20:4 n-6)	278.2 (64.5) (6.8)	276.6 (88.6) (6.9)
Total n-3 PUFA	344.0 (116.0) (8.4)	294.8 (95.7) <sup>a</sup> (7.3)
Total n-6 PUFA	1188.0 (293.8) (29.1)	1244.4 (286.4) (31.0)
Total saturated FA	2084.5 (384.6) (51.1)	2079.0 (415.4) (51.8)
Total FA	4079.5 (791.7)	4017.2 (862.2) <sup>b</sup>
Total cholesterol	6.0 (1.2)	5.8 (1.1) <sup>b</sup>
HDL-cholesterol	1.1 (0.3) <sup>c</sup>	1.1 (0.3) <sup>d</sup>
Triglycerides	1.5 (0.7) <sup>c</sup>	1.7 (1.0) <sup>d</sup>

\* $P=0.047$ , otherwise no significant differences between groups.

LA—linoleic acid, AA—arachidonic acid.

<sup>a</sup>Three missing values. <sup>b</sup>Seven missing values. <sup>c</sup>One missing value.

<sup>d</sup>Nine missing values.

fish oil supplements prior to inclusion as compared to those with no fish oil supplements,  $P=0.047$  (Table 2). As previously reported (Nilsen *et al*, 2001), there were no significant differences in other lipid parameters at baseline. The serum concentrations of  $\mu\text{CRP}$  (median and 25 and 75 percentiles) were equal in the n-3 PUFAs (20.7 (10.0–49.0) mg/l), ( $n=121$ ) as compared to the corn oil group (18.8 (6.3–49.3) mg/l), ( $n=130$ ). Furthermore, no difference in baseline p-tHcy (median and 25 and 75 percentiles) was seen between the n-3 PUFAs (14.1 (11.4–17.8)  $\mu\text{mol/l}$ ) ( $n=118$ ) as compared to the corn oil group (13.7 (10.5–16.3)  $\mu\text{mol/l}$ ) ( $n=123$ ). The healthy reference population ( $n=30$ ) presented with significantly lower levels of  $\mu\text{CRP}$  (1.2 (0.6–3.0) mg/l) and p-tHcy (9.3 (7.4–11.4)  $\mu\text{mol/l}$ ) as compared to baseline values in the MI population,  $P\leq 0.001$ . Moreover, TBA-MDA concentrations at baseline did not differ between the intervention groups (Table 3) and were similar when related to age, gender, thrombolytics and smoking habits (data not shown).

In subgroups with and without statin treatment the use of thrombolytic treatment and size of infarction, as judged by TnT and CKMB levels, were similar, although the statin-treated patients were younger and had higher levels of serum cholesterol and triglycerides at inclusion,  $P<0.05$ .

Serum concentrations of TBA-MDA in the healthy reference population ( $n=30$ ) were significantly lower than the values in the coronary population,  $P=0.001$  (Table 3).

No correlation was found between baseline TBA-MDA and serum lipids, p-tHcy or  $\mu\text{CRP}$  (data not shown). However, TBA-MDA at baseline was positively correlated with the fatty acids in serum phospholipids, both unsaturated and saturated (Table 4), but this relation was disrupted during intervention.

**Table 3** TBA-MDA complex, ( $\mu\text{mol/l}$ ) (mean and (s.d.)) at baseline and 12 months after the MI, arranged according to intervention group, as compared to the healthy reference group ( $n=30$ )

n-3 Group (n=125)		Corn oil group (n=130)		Reference group (n=30)
Baseline	12 months	Baseline	12 months	—
1.95 (0.59)	3.63 <sup>a</sup> (0.82)	1.92 (0.59)	3.35 <sup>a</sup> (0.85)	1.57 (0.61) <sup>b</sup>

Statistically significant intergroup differences;  $P=0.0274$  adjusting for age, gender, baseline values for TBA-MDA, serum lipids,  $\mu\text{CRP}$  and p-tHcy (ANACOVA), and  $P=0.0270$  ( $t$ -test).

No significant difference between intervention groups at baseline.

<sup>a</sup> $P<0.001$  for intragroup difference from baseline to follow-up (Student's  $t$ -test).

<sup>b</sup>Significance of difference between study group at baseline and reference population:  $P<0.001$ .

### Changes in TBA-MDA during intervention

As previously reported (Nilsen *et al*, 2001), a highly significant improvement in serum triglycerides and HDL-cholesterol was found in the n-3 group. TBA-MDA increased after 1 y in both the n-3 PUFAs and corn oil groups,  $P<0.001$  (Table 3), but the increase was more pronounced in the n-3 group,  $P=0.0270$ . Adjusting for possible confounding factors such as age, gender, triglycerides, HDL-cholesterol, p-tHcy,  $\mu\text{CRP}$  and TBA-MDA values at baseline, a statistically significant difference between groups after 12 months of intervention was clearly evident,  $P=0.0274$ .

During 12 months of intervention the incidence of recurrent MI, defined according to the recommendations in the consensus document of The Joint European Society of Cardiology/American College of Cardiology Committee for the Redefinition of Myocardial Infarction (Alpert & Thygesen, 2000), was almost the same in the n-3 group (18.0%) as compared to the corn oil group (18.7%). Subjects with a recurrent MI over the year had similar TBA-MDA values as compared to those who did not experience a re-event.

Multiple linear regression analyses of fatty acids in serum total phospholipids ( $n=56$ ) (listed in Table 2) on TBA-MDA measured after 12 months supplementation with n-3 PUFAs or corn oil, showed no TBA-MDA dependency. Performing best subsets regression, serum phospholipid concentration of arachidonic acid (20:4 n-6 PUFA) was identified as a predictor of TBA-MDA at 12 months follow-up ( $P=0.004$ ).

Subgroup analysis of patients on statin treatment ( $n=95$ ) as compared to those without a statin ( $n=79$ ), revealed no significant intergroup differences in TBA-MDA, regardless of the background intervention with either n-3 PUFAs or corn oil (Table 5).

### Concentrations of fatty acids in serum phospholipids during intervention

In the group receiving n-3 PUFAs, the amount of EPA, DHA and the total amount of n-3 fatty acids in serum phospholipids increased significantly (Table 1), while the total

**Table 4** Correlations (Pearson's correlation coefficient; *r*) between TBA-MDA complex and fatty acids in serum total phospholipids (*n*=56) at baseline

	<i>Fatty acid</i>	18:2 n-6	20:4 n-6	Sum n-6	20:5 n-3	22:6 n-3	Sum n-3	Sum saturated
Baseline	<i>r</i>	0.449***	0.296*	0.464***	0.177	0.315*	0.293*	0.333**

\**P*<0.05. \*\**P*<0.01. \*\*\**P*<0.001.

**Table 5** TBA-MDA complex (μmol/l) (mean and (s.d.)) at baseline and 12 months after the MI, in patients on statin treatment, as compared to patients without statin treatment, analysed according to intervention with either n-3 fatty acids (PUFAs) or corn oil

Statin treatment ( <i>n</i> =95)				No statin treatment ( <i>n</i> =79)			
n-3 PUFAs ( <i>n</i> =47)		Corn oil ( <i>n</i> =48)		n-3 PUFAs ( <i>n</i> =34)		Corn oil ( <i>n</i> =45)	
Baseline	12 months	Baseline	12 months	Baseline	12 months	Baseline	12 months
2.03 (0.59)	3.55 <sup>a</sup> (0.83)	2.00 (0.57)	3.45 <sup>a</sup> (0.85)	1.86 (0.53)	3.49 <sup>a</sup> (0.78)	1.80 (0.55)	3.18 <sup>a</sup> (0.74)

No statistically significant difference between groups at baseline (*t*-test).

<sup>a</sup>*P*<0.001 for intragroup difference from baseline to follow-up (Student's *t*-test).

No statistically significant differences between any of the subgroups at 12 months (ANCOVA).

amount of n-6 fatty acids showed a trend towards an increase in the corn oil group from baseline to 12 months (ns). However, a significant increase in the amount of arachidonic acid (20:4 n-6 acid) was noted (Table 1). Moreover, the changes in the amounts of n-3 and n-6 fatty acids differed significantly between the groups during intervention (Table 1).

## Discussion

Using TBA-MDA measured by HPLC as a marker of lipid peroxidation, we found a modest increase in lipid peroxidation in subjects receiving a high-dose concentrate of n-3 PUFAs as compared to those on corn oil. The clinical implications of increased TBA-MDA levels are uncertain, but increased lipid peroxidation might contribute to the lack of clinical prognostic benefit in this study population, as previously suggested (Nilsen *et al*, 2001). Based on findings of increased TBARS and reduced vitamin E, a possible adverse effect of high doses of n-3 PUFAs on cardiac events was hypothesised in a previous study using high-dose ethylester compounds of EPA/DHA (Johansen *et al*, 1999). TBARS may serve as a marker for later restenosis after PTCA (Johansen *et al*, 2001), and several studies indicate that elevated levels of TBARS may be associated with an increased risk of cardiovascular disease (CVD) (Jayakumari *et al*, 1992; Duthie *et al*, 1994; Miwa *et al*, 1995; Cavalca *et al*, 2001). Moreover, the significantly lower serum levels of TBA-MDA in our healthy reference population (*n*=30), as compared to values in the coronary population, may support the clinical relevance of elevated TBARS levels (Diaz-Velez *et al*, 1996; Cavalca *et al*, 2001). However, we could not demonstrate any association between TBA-MDA as a marker of lipid peroxidation and cardiovascular risk factors such as p-tHcy and

μCRP, and no relations between TBA-MDA and levels of serum total cholesterol and triglycerides were found.

An intragroup increase in TBA-MDA from baseline to 12 months was observed in both intervention groups. Wander *et al* (1998) have previously shown that linoleic acid is a significant contributor to lipid peroxidation. This n-6 PUFA is present in LDL in an amount far greater than any other fatty acid. In the present study, the change in n-6 PUFAs in serum phospholipids differed significantly between the groups, but the intragroup increase in linoleic acid during 1 y of corn oil treatment did not reach statistical significance. Several studies support the notion that a high intake of n-6 PUFAs may increase lipid peroxidation (Reaven *et al*, 1991, 1993, 1994), and may explain the increase over 1 y also in our control group. Therefore, the choice of control in our study may partly mask the effect of n-3 PUFAs intervention.

The impact of n-3 PUFAs on oxidative stress of the vascular system in humans still remains an unsettled issue. Previous studies included relatively small sample sizes, different doses and compositions of n-3 PUFAs, short intervention periods, different amounts of antioxidants supplies and different methods used to quantify the oxidative burden and different methods to estimate TBARS.

The choice of dose of n-3 PUFAs in our study was based on previous studies investigating the influence of n-3 PUFAs on cardiovascular risk factors and lipid peroxidation (Phillipson *et al*, 1985; Eritsland *et al*, 1995; Connor & Connor, 1997). Later the GISSI-Prevenzione trial demonstrated beneficial effects on CVD of a lower dose (GISSI-Prevenzione Investigators, 1999). Baseline levels of total n-3 PUFAs in our study were reasonably high (Table 3), and a background diet rich in n-3 PUFAs may mask the effect of intervention. However, the baseline TBA-MDA level in our CHD population was of a similar magnitude as noted by others (Nordøy *et al*, 1998; Kharb *et al*, 2000).

It has been argued that the TBARS assay may measure the oxidation or decomposition of specific PUFAs instead of measuring the overall lipid peroxidation, as both the number of double bonds as well as the position of the first double bond from the methyl terminus play a role in the formation of the oxidative products (Yoshino *et al*, 1991; Higdon *et al*, 2000). Lack of MDA-TBA dependency on concentrations of n-3 PUFAs in serum phospholipids during treatment suggests that increased TBA-MDA may reflect actual differences in lipid peroxidation, rather than merely reflecting the differences in concentrations of n-3 PUFAs in serum phospholipids.

The TBARS assay is well accepted for determination of lipid-derived decomposition products (Armstrong & Browne, 1994), but certain methodological problems must be addressed. This biomarker estimates the circulating levels of a range of lipoperoxidation aldehydes secondary to lipid peroxidation (Janero, 1990; Armstrong & Browne, 1994), and the more acidic conditions of the method applied in the present study is likely to favour the release of TBARS precursors from the lipoproteins and protein-bound aldehydes (Lapenna *et al*, 2001). Therefore, in spite of its simplicity and high sensitivity, the TBARS test has been criticised for inadequate specificity to oxidant-driven lipid peroxidation, especially when using human plasma. Greater specificity is achieved by HPLC-separation of the MDA-TBA complex from thiobarbituric acid-reactive substances, as performed in our study, allowing for purified separation of the MDA-TBA complex from extraneous substances (Wong *et al*, 1987; Anoopkumar-Dukie *et al*, 2001). Using an external standard to which the TBA-MDA complex has been compared, provides a more accurate assessment of MDA levels and lipid peroxidation. The TBARS assay is a relatively quick and inexpensive method compared to determination of F2-isoprostanes, which is a tedious and expensive procedure. However, the source of the MDA measured still remains a pitfall, as MDA also is generated *ex vivo* by decomposition of lipid peroxides during the heating stage of the test (Gutteridge, 1986). While the interpretation of the MDA content in studies of lipid peroxidation requires caution, the isoprostanes represent stable products of lipid peroxidation formed *in situ* in cell membranes, which can be measured with great sensitivity and specificity (Meagher and FitzGerald, 2000).

The vast majority of studies investigating the oxidative potential of n-3 PUFAs include short-term supplementation periods (Harats *et al*, 1991; Nenseter *et al*, 1992; Bonanome *et al*, 1996; Wander *et al*, 1996; Hansen *et al*, 1998; Higdon *et al*, 2000). Long-term effects *in vivo* after 9 months of intervention were studied by Eritsland *et al* (1995), and susceptibility to lipid peroxidation induced *in vitro* was evaluated after 180 days of n-3 PUFAs treatment in the study by Palozza (Palozza *et al*, 1996). The present study is the first to examine the peroxidative status after 12 months of intervention with a high-dose of n-3 PUFAs.

Following thrombolytic treatment, reperfusion of ischaemic areas may be associated with reperfusion injury mediated by free-radical-induced oxidation. At the time of blood sampling 4–6 days after the acute MI, TBA-MDA levels were equal in patients who had been given thrombolytic treatment as compared to those not having received thrombolytics. In a study by Kharb *et al* (2000) TBARS were increased within the first hours after thrombolytic treatment, but this increase was no longer evident after 72 h.

Oxygen-free radicals are involved in the pathophysiology of inflammation and atherosclerosis (Dargel, 1992; Halliwell & Chirico, 1993). Treatment with statins has been shown to reduce the risk for cardiovascular events and exhibits beneficial effects on inflammation and plaque stability. In a subgroup of patients receiving simvastatin, no difference in TBA-MDA was observed between individuals on n-3 PUFAs as compared to those on corn oil. Moreover, no difference in TBA-MDA was noted between statin-treated and statin-untreated patients. However, the number of subjects in these subgroups was low, and these data should be interpreted with great care. A previous study has shown no influence on lipid peroxidation of combined treatment of simvastatin and n-3 PUFAs (Nordøy *et al*, 1998).

In conclusion, the main results of this long-term, post-infarct study show a modest increase in lipid peroxidation, measured as TBA-MDA by HPLC, in subjects on a high-dose concentrate of n-3 PUFAs as compared to a control group on corn oil. Increased TBA-MDA may reflect an actual difference in lipid peroxidation caused by n-3 PUFAs, rather than merely reflecting a change in their serum concentrations. The use of a statin did not modify the peroxidation potential of n-3 PUFAs.

#### Acknowledgements

We are grateful for the financial support by Pharmacia A/S and Pronova A/S, Norway. We acknowledge the support of Ole Kristensen, MD at the Department of Clinical Chemistry, Central Hospital in Rogaland. Finally, thanks are due to the project nurse, Solfrid Moen, Hjertelaget Research Foundation, Norway, and the steering committee of the 'OFAMI'-study: Dennis WT Nilsen, Knud Landmark, Leik Woie and Michael Abdelnoor.

#### References

- Alpert JS & Thygesen K (2000): The Joint European Society of Cardiology/American College of Cardiology Committee for the redefinition of myocardial infarction. *Eur Heart J.* **21**, 1502–1513.
- Anoopkumar-Dukie S, Walker RB & Daya S (2001): A sensitive and reliable method for the detection of lipid peroxidation in biological tissues. *J. Pharm. Pharmacol.* **53**, 263–266.
- Armstrong D & Browne R (1994): The analysis of free radicals, lipid peroxides, antioxidant enzymes and compounds related to oxidative stress as applied to the clinical chemistry laboratory. In *Free Radicals in Diagnostic Medicine: A Systems Approach to Laboratory, Technology, Clinical Correlations, and Antioxidant Therapy*, ed. D Armstrong, pp 43–58. New York: Plenum Press, XVI.

- Arnesen E, Refsum H, Bonna KH, Ueland PM, Forde OH & Nordrehaug JE (1995): Serum total homocysteine and coronary heart disease. *Int. J. Epidemiol.* **24**, 704–709.
- Avogaro P, Bittolo Bon G & Cazzolato G (1988): Presence of a modified low density lipoprotein in humans. *Arteriosclerosis* **8**, 79–87.
- Bittolo-Bon G, Cazzolato G, Alessandrini P, Soldan S, Casalino G & Avogaro P (1993): Effects of concentrated DHA and EPA supplementation on LDL peroxidation and vitamin E status in type HB hyperlipidemic patients. In *Omega-3 Fatty Acids*, eds CA Drevon, I Baksaas & HE Krokan, pp 51–58. Basel, Switzerland: Birkhauser-Verlag/Switzerland.
- Bonanome A, Biasia F, De Luca M, Munaretto G, Biffanti S, Pradella M & Pagnan A (1996): n-3 Fatty acids do not enhance LDL susceptibility to oxidation in hypertriglycerolemia hemodialyzed subjects. *Am. J. Clin. Nutr.* **63**, 261–266.
- Boyd HC, Gown AM, Wolfbauer G & Chait A (1989): Direct evidence for a protein recognized by a monoclonal antibody against oxidatively modified LDL in atherosclerotic lesions from a Watanabe heritable hyperlipidemic rabbit. *Am. J. Pathol.* **135**, 815–824.
- Brude IR, Drevon CA, Hjermann I, Seljeflot I, Lund-Katz S, Saarem K, Sandstad B, Solvoll K, Halvorsen B, Arnesen H & Nenseter MS (1997): Peroxidation of LDL from combined-hyperlipidemic male smokers supplied with  $\omega$ -3 fatty acids and antioxidants. *Arterioscler. Thromb. Vasc. Biol.* **17**, 2576–2588.
- Burstein M, Scholnick HR & Morfin R (1970): Rapid method for the isolation of lipoprotein from human serum by precipitation with polyanions. *J. Lipid. Res.* **11**, 583–595.
- Cavalca V, Cighetti G, Bamonti F, Loaldi A, Bortone L, Novembrino C, De Franceschi M, Belardinelli R & Guazzi MD (2001): Oxidative stress and homocysteine in coronary artery disease. *Clin. Chem.* **47**, 887–892.
- Connor SL & Connor WE (1997): Are fish oils beneficial in the prevention and treatment of coronary artery disease? *Am. J. Clin. Nutr.* **66** (Suppl), S1020–S1031.
- Cosgrove JP, Church DF & Pryor WA (1987): The kinetics of the autooxidation of polyunsaturated fatty acids. *Lipids* **22**, 299–304.
- Dargel R (1992): Lipid peroxidation—a common pathogenetic mechanism? *Exp. Toxicol. Pathol.* **44**, 169–181.
- Diaz-Velez CR, Garcia-Castineiras S, Mendoza-Ramos E & Hernandez-Lopez E (1996): Increased malondialdehyde in peripheral blood of patients with congestive heart failure. *Am. Heart J.* **131**, 146–152.
- Duthie GG, Beattie JA, Arthur JR, Franklin M, Morrice PC & James WPT (1994): Blood antioxidants and indices of lipid peroxidation in subjects with angina pectoris. *Nutrition* **10**, 313–316.
- Eritsland J, Arnesen H, Seljeflot I & Høstmark AT (1995): Long-term metabolic effects of n-3 polyunsaturated fatty acids in patients with coronary artery disease. *Am. J. Clin. Nutr.* **61**, 831–836.
- Frankel EN, Parks EJ, Xu R, Schneeman BO, Davis PA & German JB (1994): Effect of n-3 fatty acid-rich fish oil supplementation on the oxidation of low density lipoproteins. *Lipids* **29**, 233–236.
- Gerhardt W, Waldenström J, Horder M, Hofvendahl S, Billström R, Ljungdahl R, Berning H & Bagger P (1982): Creatine kinase and creatine kinase B-subunit activity in serum in cases of suspected myocardial infarction. *Clin. Chem.* **28**, 277–283.
- GISSI-Prevenzione Investigators (1999): Dietary supplementation with n-3 polyunsaturated fatty acids and vitamin E after myocardial infarction: results of the GISSI-Prevenzione trial. *Lancet* **354**, 447–455.
- Gutteridge JM (1986): Aspects to consider when detecting and measuring lipid peroxidation. *Free Radic. Res. Commun.* **1**, 173–184.
- Halliwel B & Chirico S (1993): Lipid peroxidation: its mechanism, measurement, and significance. *Am. J. Clin. Nutr.* **57** (Suppl), S715–S725.
- Hansen JB, Berge RK, Nordøy A & Bonna KH (1998): Lipid peroxidation of isolated chylomicrons and oxidative status in plasma after intake of highly purified eicosapentaenoic or docosahexaenoic acids. *Lipids* **33**, 1123–1129.
- Harats D, Dabach Y, Hollander G, Ben-Naim M, Schwartz R, Berry EM, Stein O & Stein Y (1991): Fish oil ingestion in smokers and nonsmokers enhances peroxidation of plasma lipoproteins. *Atherosclerosis* **90**, 127–139.
- Hetland Ø, Gøransson L & Nilsen DWT (1995): Cardiac troponin-T immunoassay on biotin-streptavidin-coated microplates, preliminary performance in acute myocardial infarction. *Scand. J. Clin. Lab. Invest.* **55**, 701–713.
- Higdon JV, Liu J, Du SH, Morrow JD, Ames BN & Wander RC (2000): Supplementation of postmenopausal women with fish oil rich in eicosapentaenoic acid and docosahexaenoic acid is not associated with greater *in vivo* lipid peroxidation compared with oils rich in oleate and linoleate as assessed by plasma malondialdehyde and F<sub>2</sub>-isoprostanes. *Am. J. Clin. Nutr.* **72**, 714–722.
- Janero DR (1990): Malondialdehyde and thiobarbituric acid-reactivity as diagnostic indices of lipid peroxidation and peroxidative tissue injury. *Free Rad. Biol. Med.* **9**, 515–540.
- Jayakumari N, Ambikakumari V, Balakrishnan KG & Subramonia IK (1992): Antioxidant status in relation to free radical production during stable and unstable angina syndromes. *Atherosclerosis* **94**, 183–190.
- Johansen O, Abdelnoor M, Brekke M, Seljeflot I, Høstmark AT & Arnesen H (2001): Predictors of restenosis after coronary angioplasty. A study on demographic and metabolic variables. *Scand. Cardiovasc. J.* **35**, 86–91.
- Johansen O, Seljeflot I, Høstmark AT & Arnesen H (1999): The effect of supplementation with omega-3 fatty acids on soluble markers of endothelial function in patients with coronary heart disease. *Arterioscler. Thromb. Vasc. Biol.* **19**, 1681–1686.
- Kharb S, Singh V, Ghalaut PS & Singh GP (2000): Oxidative stress after acute myocardial infarction: effect of thrombolytic treatment. *J. Assoc. Physicians India* **48**, 578–580.
- Kinsella JE, Lokesh B & Stone RA (1990): Dietary n-3 polyunsaturated fatty acids and amelioration of cardiovascular disease: possible mechanisms. *Am. J. Clin. Nutr.* **52**, 1–28.
- Lapenna D, Ciofani G, Pierdominico SD, Giamberardino MA & Cuccurullo F (2001): Reaction conditions affecting the relationship between thiobarbituric acid reactivity and lipid peroxides in human plasma. *Free Radic. Biol. Med.* **31**, 331–335.
- Liu J, Yeo HC, Doniger SJ & Ames BN (1997): Assay of aldehydes from lipid peroxidation: gas chromatography–mass spectrometry compared to thiobarbituric acid. *Anal. Biochem.* **245**, 161–166.
- Mansoor MA, Svardal AM & Ueland PM (1992): Determination of the *in vivo* redox status of cysteine, cysteinylglycine, homocysteine, and glutathione in human plasma. *Anal. Biochem.* **200**, 218–229.
- Meagher EA & FitzGerald GA (2000): Indices of lipid peroxidation *in vivo*: strengths and limitations. *Free Radic. Biol. Med.* **28**, 1745–1750.
- Meydani M, Natiello F, Goldin B, Free N, Woods M, Schaefer E, Blumberg JB & Gorbach SL (1991): Effect of long-term fish oil supplementation on vitamin E status and lipid peroxidation in women. *J. Nutr.* **121**, 484–491.
- Miwa K, Miyagi U & Fujita M (1995): Susceptibility of plasma low density lipoprotein to cupric ion-induced peroxidation in patients with variant angina. *J. Am. Coll. Cardiol.* **26**, 632–638.
- Muller-Bardorff M, Hallermeier K, Schroeder A, Ebert C, Borgya A, Gerhardt W, Remppis A, Zehelein J & Katus HA (1997): Improved troponin T ELISA specific for cardiac troponin T isoform: assay development and analytical and clinical validation. *Clin. Chem.* **43**, 458–466.
- Nenseter MS & Drevon CA (1996): Dietary polyunsaturates and peroxidation of low density lipoprotein. *Curr. Opin. Lipidol.* **7**, 8–13.
- Nenseter MS, Rustan AC, Lund-Katz S, Soyland E, Maelandsmo G, Phillips MC & Drevon CA (1992): Effects of dietary supplementation with n-3 polyunsaturated fatty acids on physical properties

- and metabolism of low density lipoproteins in humans. *Arterioscl. Thromb.* **12**, 369–379.
- Nilsen DWT, Albrektsen G, Landmark K, Moen S, Aarsland T & Woie L (2001): Effects of a high-dose concentrate of n-3 fatty acids or corn oil introduced early after an acute myocardial infarction on serum triacylglycerol and HDL cholesterol. *Am. J. Clin. Nutr.* **74**, 50–56.
- Nordøy A, Bonna KH, Nilsen H, Berge RK, Hansen JB & Ingebreten OC (1998): Effects of simvastatin and omega-3 fatty acids on plasma lipoproteins and lipid peroxidation in patients with combined hyperlipidaemia. *J. Int. Med.* **243**, 163–170.
- Palozza P, Sgarlata E, Luberto C, Piccioni E, Anti M, Marra G, Armelao F, Franceschelli P & Bartoli GM (1996): n-3 Fatty acids induce oxidative modifications in human erythrocytes depending on dose and duration of dietary supplementation. *Am. J. Clin. Nutr.* **64**, 297–304.
- Phillipson BE, Rothrock DW, Connor WE, Harris WS & Illingworth DR (1985): Reduction of plasma lipids, lipoproteins, and apoproteins by dietary fish oils in patients with hypertriglyceridemia. *N. Engl. J. Med.* **312**, 1210–1216.
- Reaven PD, Grasse BJ & Tribble DL (1994): Effects of linoleate-enriched and oleate-enriched diets in combination with alpha-tocopherol on the susceptibility of LDL and LDL subfractions to oxidative modification in humans. *Arterioscler. Thromb.* **14**, 557–566.
- Reaven P, Parthasarathy S, Grasse BJ, Miller E, Almazan F, Mattson FH, Khoo JC, Steinberg D & Witztum JL (1991): Feasibility of using an oleate-rich diet to reduce the susceptibility of low-density lipoprotein to oxidative modification in humans. *Am. J. Clin. Nutr.* **54**, 701–706.
- Reaven P, Parthasarathy S, Grasse BJ, Miller E, Steinberg D & Witztum JL (1993): Effects of oleate-rich and linoleate-rich diets on the susceptibility of low density lipoprotein to oxidative modification in mildly hypercholesterolemic subjects. *J. Clin. Invest.* **91**, 668–676.
- Ridker PM, Glynn RJ & Hennekens CH (1998): C-reactive protein adds to the predictive value of total and HDL cholesterol in determining risk of first myocardial infarction. *Circulation* **97**, 2007–2011.
- Rosenfeld ME (1991): Oxidized LDL affects multiple atherogenic cellular responses. *Circulation* **83**, 2137–2140.
- Schmidt EB & Dyerberg J (1994): Omega-3 fatty acids: current status in cardiovascular medicine. *Drugs* **47**, 405–424.
- Steinberg D, Parthasarathy MD, Carew TE, Khoo JC & Witztum JL (1989): Beyond cholesterol. Modification of low-density lipoprotein that increases its atherogenicity. *N. Engl. J. Med.* **320**, 915–924.
- Wander RC, Du SH, Ketchum SO & Rowe KE (1996): Effects of interaction of RRR-alpha-tocopheryl acetate and fish oil on low-density-lipoprotein oxidation in postmenopausal women with and without hormone-replacement therapy. *Am. J. Clin. Nutr.* **63**, 184–193.
- Wander RC, Du SH & Thomas DR (1998): Influence of long-chain polyunsaturated fatty acids on oxidation of low density lipoprotein. *Prostaglandins Leuk. Essent. Fatty Acids* **59**, 143–151.
- Wiklund O, Mattsson L, Bjornheden T, Camejo G & Bondjers G (1991): Uptake and degradation of low density lipoproteins in atherosclerotic rabbit aorta: role of local LDL modification. *J. Lipid. Res.* **32**, 55–62.
- Wong SHY, Knight JA, Hopfer SM, Zaharia O, Leach Jr CN & Sunderman Jr FW (1987): Lipoperoxides in plasma as measured by liquid-chromatographic separation of malondialdehyde-thiobarbituric acid adduct. *Clin. Chem.* **33**, 214–220.
- Yoshino K, Sano M, Fujita M & Tomita I (1991): Production of aliphatic aldehydes on peroxidation of various types of lipids. *Chem. Pharm. Bull.* **39**, 1788–1791.