


# Author's Accepted Manuscript

Expression of human inducible nitric oxide synthase in response to cytokines is regulated by hypoxia-inducible factor-1

Martin Lee, Christine Wang, Steven W. Jin, Mark P. Labrecque, Timothy V. Beischlag, Mark A. Brockman, Jonathan C. Choy



www.elsevier.com

PII: S0891-5849(18)32263-9  
DOI: <https://doi.org/10.1016/j.freeradbiomed.2018.10.441>  
Reference: FRB14005

To appear in: *Free Radical Biology and Medicine*

Received date: 8 May 2018  
Revised date: 19 September 2018  
Accepted date: 24 October 2018

Cite this article as: Martin Lee, Christine Wang, Steven W. Jin, Mark P. Labrecque, Timothy V. Beischlag, Mark A. Brockman and Jonathan C. Choy, Expression of human inducible nitric oxide synthase in response to cytokines is regulated by hypoxia-inducible factor-1, *Free Radical Biology and Medicine*, <https://doi.org/10.1016/j.freeradbiomed.2018.10.441>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## Expression of human inducible nitric oxide synthase in response to cytokines is regulated by hypoxia-inducible factor-1

Martin Lee<sup>1,3</sup>, Christine Wang<sup>1,3</sup>, Steven W. Jin<sup>1,2</sup>, Mark P. Labrecque<sup>2</sup>, Timothy V. Beischlag<sup>2,3</sup>, Mark A. Brockman<sup>1,2</sup>, Jonathan C. Choy<sup>1,3\*</sup>

<sup>1</sup>Department of Molecular Biology and Biochemistry, Simon Fraser University, Burnaby, British Columbia, Canada

<sup>2</sup>Faculty of Health Sciences, Simon Fraser University, Burnaby, British Columbia, Canada


<sup>3</sup>Centre for Cell Biology, Development and Disease, Simon Fraser University, British Columbia, Canada

\*To whom the correspondence should be addressed: Jonathan Choy, Department of Molecular Biology and Biochemistry, Simon Fraser University, 8888 University Drive, Burnaby, BC, Canada; Tel: 778-782-8701, Fax: 778-782-5583, E-mail: jonathan.choy@sfu.ca

### ABSTRACT

The production of nitric oxide (NO) by inducible NO synthase (iNOS) and the regulation of gene expression by hypoxia-inducible factors (HIFs) are important for many aspects of human cell biology. However, little is known about whether iNOS expression is controlled by HIFs in human cells. Stimulation of A549 human lung epithelial cells with cytokines (TNF, IL-1 and IFN $\gamma$ ) increased the nuclear accumulation of HIF-1 in normoxic conditions. Activation of HIF-1 by hypoxia or CoCl<sub>2</sub> was not sufficient to induce iNOS expression. However, pharmacological inhibition of HIF-1 reduced the induction of iNOS expression in A549 cells and primary human astrocytes. Moreover, elimination of HIF-1 $\alpha$  expression and activity by CRISPR/Cas9 gene editing significantly reduced the induction of human iNOS gene promoter, mRNA and protein expression by cytokine stimulation. Three putative hypoxic response elements (HRE) are present within the human iNOS gene promoter and elimination of an HRE at -4981 bp reduced the induction of human iNOS promoter activity in response to cytokine stimulation. These findings establish an important role for HIF-1 $\alpha$  in the induction of human iNOS gene expression in response to cytokine stimulation.

### Graphical abstract



**Abbreviations:** CoCl<sub>2</sub>, Cobalt chloride; **HIF**, Hypoxia-inducible factor; **HK-1**, Hexokinase-1; **HRE**, Hypoxia response element; **iNOS**, Inducible nitric oxide synthase; **NO**, Nitric oxide; **PHD**, prolyl hydroxylase; **sgRNA**, single-guide RNA; **VEGF-A**, Vascular endothelial growth factor-A  
**Keywords:** Nitric oxide synthase, Nitric oxide, Gene regulation, Hypoxia-inducible factor (HIF), Cytokine

## INTRODUCTION

Nitric oxide (NO) has broad ranging biological effects that control essential physiological processes and dysregulation of its production is involved in the development of many pathologies [1]. This bioactive gas is synthesized by a family of nitric oxide synthases (NOSs): neuronal (nNOS), inducible (iNOS) and endothelial (eNOS). In contrast to the largely constitutive nature of nNOS and eNOS expression, iNOS expression and activity is tightly regulated to maintain precise NO levels that have effects on immune activation, inflammation and tissue damage [2]. iNOS is normally absent in unstimulated cells and expression is highly induced in response to inflammatory stimuli, such as

cytokines and microbial components [3]. Transcriptional induction of the human iNOS promoter requires the synergistic action of several transcription factors [4]. Stimulation of cells by cytokines that activate NF- $\kappa$ B and MAPK pathways results in the activation of NF- $\kappa$ B, AP-1 and C/EBP $\beta$  transcription factors that cooperatively drive iNOS gene expression [5-8]. Activation of STAT1 by interferon- $\gamma$  (IFN $\gamma$ ) is also needed for maximal induction of this enzyme in human cells [9].

Hypoxia-inducible factors (HIFs) are central regulators of cellular responses to hypoxia and cellular metabolism [10]. HIF-1 is a heterodimer composed of HIF-1 $\alpha$  and HIF-1 $\beta$ /ARNT subunits [11]. Both subunits are actively expressed in normoxic conditions but HIF-1 $\alpha$  is rapidly degraded because of prolyl hydroxylation by prolyl hydroxylases (PHDs) that leads to its ubiquitinylation by von Hippel Lindau protein and subsequent proteasome-mediated degradation [12]. In hypoxia, PHD activity is inhibited thereby preventing HIF-1 $\alpha$  degradation and increasing its levels [13]. HIF-1 nuclear translocation and binding to HIF responsive elements (HREs) within gene promoters drives transcriptional activation of genes involved in tissue perfusion, cell survival and glycolytic metabolism that are needed for tissue protection in hypoxic conditions and regulation of inflammation [14, 15]. The induction of glycolytic genes by HIF-1 also controls the metabolic reprogramming of cells to increase glycolysis in normoxic conditions, a process termed aerobic glycolysis. Induction of aerobic glycolysis affects the regulation of immune responses and the pathogenesis of cancer [16, 17].

iNOS has been suggested to be a HIF-regulated gene. Specifically, the induction of iNOS by cytokines in murine macrophages is increased by hypoxia through an HRE-dependent mechanism [18, 19]. Cellular stresses, such as iron deprivation, also induce the HIF-dependent expression of iNOS in cooperation with IFN $\gamma$  stimulation in murine cells [20]. However, there are important differences in the mechanisms by which iNOS expression is induced in human and murine cells. Specifically, the human iNOS gene promoter is structurally different than the murine counterpart, is regulated by distinct regulatory regions, and is methylated [21-23]. These differences may underlie the disparate

expression of iNOS in mouse and human cells. Cultured human macrophages do not induce iNOS expression in response to inflammatory cytokines and the cell type-specific activity of the human iNOS promoter is different than mouse iNOS expression within tissues *in vivo* [24, 25]. iNOS expression can be induced by cytokine stimulation in certain human cell lines, such as A549 transformed lung epithelial cells, and primary cell types, such as astrocytes and bronchial lung epithelial cells [26, 27]. The synergistic action of multiple cytokines is needed for maximal induction of iNOS expression in most cell types.

We examined the role of HIF-1 in regulating the induction of human iNOS expression in normoxic conditions. The nuclear accumulation of HIF-1 $\alpha$  was increased after stimulation of human cells with cytokines. Pharmacological inhibition of HIF-1 and CRISPR/Cas9-mediated elimination of HIF-1 $\alpha$  reduced cytokine-mediated induction of human iNOS gene promoter activity, mRNA and protein expression. Further, there are 3 predicted HREs in the human iNOS gene promoter and mutational elimination of a site at -4981 bp reduced the induction of gene promoter activity by cytokines. Our findings identify HIF-1 as a cytokine-activated transcription factor that increases human iNOS expression.

## MATERIAL AND METHODS

### *Cell culture*

A549 human lung epithelial cells were purchased from ATCC and cultured in F-12K medium containing 10% FBS at 37 °C in a humidified atmosphere of 95 % air and 5 % CO<sub>2</sub>. Primary human astrocytes were purchased from Lonza, cultured in ABM medium (Lonza, Basel, Switzerland) supplemented with the AGM SingleQuot Kit (Lonza), and used between passages 4 and 12 for experiments. For all experiments using A549 cells, confluent cells were incubated in basal medium in the absence of serum for 24 h prior to cytokine treatment. To induce iNOS expression, cells were treated with a cytokine mixture of human recombinant IL-1 $\alpha$  (10 ng/ml; PeproTech, Rocky Hill, NJ),

TNF (10 ng/ml; PeproTech), and IFN $\gamma$  (100 ng/ml; Invitrogen) for the indicated times. This combination of cytokines maximally induces iNOS expression. Stimulation with individual cytokines has little effect. HIF-1 $\alpha$  expression was inhibited by incubating cells with PX-478 (20  $\mu$ M; Selleck Chemicals LLC, Houston, TX) for 16 h prior to cytokine stimulation. In some experiments cells were incubated in hypoxic conditions (1% O $_2$ ) for 24 h in a humidified CO $_2$  chamber. Chemical hypoxia was induced by treatment with 200  $\mu$ M CoCl $_2$  for 5 h. All experiments were performed at least three times unless otherwise indicated.

#### *SDS-PAGE and immunoblotting*

Cells were lysed and cell lysates immunoblotted as described previously [28]. Primary antibodies used were: mouse monoclonal anti-HIF-1 $\alpha$  (NB100-105; Novus Biologicals, Littleton, CO), rabbit monoclonal anti-HIF-1 $\alpha$  (14179S; Cell Signaling Technology, Danvers, MA), rabbit monoclonal anti-iNOS (NBP1-33780; Novus Biologicals), mouse monoclonal anti-caspase-8 (MAB4708; EMD Millipore, Temecula, CA), rabbit polyclonal anti-Histone H2A (sc-67218; Santa Cruz, Dallas, TX), and mouse monoclonal anti- $\beta$ -actin (A1978, Sigma-Aldrich, St. Louis, MO). Immunoblot images were quantified by densitometry using the National Institutes of Health ImageJ software, and protein expression data were normalized to  $\beta$ -actin levels or to appropriate loading controls for the quantification of proteins in subcellular fractions.

#### *RNA Isolation and RT-qPCR*

Total RNA was isolated from cells using a RNeasy mini kit as per the manufacturer's instructions (Qiagen) and TaqMan quantitative RT-PCR (RT-qPCR) was performed as described [29] using validated primer/probe sets for HIF-1 $\alpha$ , VEGF-A, HK1, iNOS, GAPDH and MEK1 (Applied Biosystems, Foster City, CA). The relative expression of target genes was calculated by normalization

to GAPDH. In experiments involving alterations in HIF-1 expression, target gene expression was additionally normalized to MEK1 because some studies have reported that the expression of GAPDH may be regulated by HIF-1 in response to hypoxia [30]. We did not observe any change in GAPDH levels in our experiments but included additional normalization to MEK1 as a secondary analysis. Data were acquired on an ABI 7900HT fast real-time PCR system (Applied Biosystems).

#### *Quantification of NO production*

NO production was measured by quantifying the accumulation of nitrite in culture supernatants using a NO-specific chemiluminescence analyzer as described previously (Sievers, Boulder, CO) [28].

#### *Transfection of mammalian cells*

For iNOS promoter luciferase experiments, A549 cells were transfected with 1  $\mu$ g of a human iNOS promoter-luciferase construct containing 7.2 kb of the human iNOS promoter upstream of the transcription start site (pXP2-iNOS; kindly provided by Dr. David Geller, University of Pittsburgh, Pittsburgh, PA[5]), human iNOS promoter-luciferase plasmids containing mutated HREs (pXP2-iNOS $\Delta$ HRE), or pXP2 backbone plasmids using Lipofectamine 3000 reagent as per the manufacturer's instructions (Invitrogen, Carlsbad, CA). To control for transfection efficiency, cells were also transfected with a pRL-SV40 Renilla luciferase plasmid and firefly luciferase activity normalized to the levels of Renilla luciferase. Following transfection, cells were plated into 96-well flat bottom white plates, rested for 24 h in basal F-12K media, and then used for experiments.

For CRISPR-Cas9 experiments, A549 cells were harvested by trypsinization, and  $1 \times 10^6$  cells were transfected with 1  $\mu$ g of pSpCas9(HIF-1 $\alpha$ )-2A-GFP plasmid encoding the HIF-1 $\alpha$  guide RNA or 1  $\mu$ g of the pSpCas9(BB)-2A-GFP backbone vector using Lipofectamine 3000 reagent as per the manufacturer's instructions (Invitrogen).

### *Luciferase reporter assay for iNOS promoter activity*

A549 cells were transfected with iNOS promoter luciferase plasmids and a constitutively expressed Renilla luciferase construct (pRL-SV40). Cells were then stimulated with cytokines for 8 h and luciferase activity was measured using a Dual-Glo Luciferase Assay System as per the manufacturer's instructions (Promega, Madison, WI) and as described previously [31]. Luminescence was quantified using a Tecan Infinite 200 Pro multimode reader.

The sequence of the human iNOS promoter was queried for HRE consensus sequences (5'-(A/G)CGTG-3') using the motif-based sequence analysis tool FIMO (Find individual motif occurrences). Three putative HREs were identified and each was excised from the human iNOS promoter-luciferase plasmid using overlap extension PCR [32, 33]. Briefly, PCR was performed on the wild type human iNOS promoter-luciferase construct (pXP2-iNOS) using 2 primers flanking the 7.2 kb human iNOS promoter (pXP2\_F 5'-GGTTTGTCCAAACTCATCAATGTATCTTATCATGTCTGGATCCCCAGGG-3', pXP2\_R 5'-CGAACGTGTACATCGACTG-3') and pairs of overlapping primers at each putative HRE (HRE1\_Mutagenesis primer 5'-GCAGAACAGCCAGGGGGCAGCCGCCCTCTCTG-3', HRE2\_Mutagenesis primer 5'-GTGAAGGCCCCACAGGACACTCGCCACGGGACATGC-3', HRE3\_mutagenesis primer 5'-CTTTCACTGAGGCTTCTCCTGTGTGCATCACAGACATTAAC-3'). The three mutated amplicons were then cloned into the parental pXP2-iNOS plasmid to generate three single HRE-deficient human iNOS promoter-luciferase reporter constructs (pXP2-iNOS $\Delta$ HRE1, pXP2-iNOS $\Delta$ HRE2, pXP2-iNOS $\Delta$ HRE3). Successful elimination of HREs was confirmed by Sanger sequencing with primers flanking the three sites. (iNOS\_HRE1\_F 5'-TCTGGATCCCCAGGGCCCAGC-3', iNOS\_HRE1\_R 5'-GGCAACAGTGAGACTCTGATCTC-3', iNOS\_HRE2\_F 5'-CACTTCCGTCATCTCC-3', iNOS\_HRE2\_R 5'-



CGGAGTTCCTACTGTGTGCC-3', iNOS\_HRE3\_F 5'-CCCACCTGATCCTCCTGAGT-3', iNOS\_HRE3\_R 5'-GGCCTCAAAGCACTTCGTTG-3'). A549 cells were transfected with the wild-type human iNOS promoter-luciferase construct or with the single HRE-deficient human iNOS promoter-luciferase constructs, and the luciferase reporter assay was carried out as described above.

### *CRISPR-Cas9 gene editing of HIF-1 $\alpha$*

A 20 nucleotide, CRISPR RNA (crRNA) sequence (5'-TGGTATCATATACGTGAATG-3') targeting the human HIF-1 $\alpha$  gene was designed using the online single-guide RNA (sgRNA) designer tool CRISPRko (Broad Institute). The HIF-1 $\alpha$ \_sgRNA (5'-CACCGTGGTATCATATACGTGAATG-3') (Integrated DNA Technologies) was cloned into pSpCas9(BB)-2A-GFP. A549 cells were transfected with pSpCas9(HIF-1 $\alpha$ )-2A-GFP using Lipofectamine. Following transfection, GFP-positive cells were single cell sorted and clonal populations expanded in F-12K medium containing 10% FBS at 37 °C in a humidified atmosphere of 95% air and 5% CO<sub>2</sub>.

To screen for Cas9-mediated gene editing of HIF-1 $\alpha$ , a PCR reaction was carried out using the HIF-1 $\alpha$  forward primer (5'-GGGTAAAGAACAAAACACACAGCGAAGC-3') and the HIF-1 $\alpha$ \_sgRNA reverse primer (5'-CACCGTGGTATCATATACGTGAATG-3'), the latter being complimentary to the unedited HIF-1 $\alpha$  sequence. In this scenario, the reverse primer binds to the unedited HIF-1 $\alpha$  sequence but not an edited version because it is not complimentary to the edited gene sequence. In addition to examining HIF-1 $\alpha$  gene editing by PCR for the edited region, a region of the HIF1 $\alpha$  gene was amplified by PCR using the HIF-1 $\alpha$  forward primer (5'-GTGTACCCTAACTAGCCGAGGA-3') and the HIF-1 $\alpha$  reverse primer (5'-TGGGGGCAGTAGTAGTACAGTTG-3') and the amplicon sent for Sanger sequencing (Genewiz). Further, HIF-1 $\alpha$  protein expression was determined by treating cells with 200  $\mu$ M CoCl<sub>2</sub> and immunoblotting for HIF-1 $\alpha$ .

### *Statistical analyses*

Unless otherwise indicated, quantitative data presented are the mean  $\pm$  S.D. of at least three independent experiments. Statistical comparisons between two groups were made using a Student's test and between multiple groups with analysis of variance followed by a post-hoc multiple comparisons test. Statistically significant differences were defined as  $p < 0.05$ .

## **RESULTS**

### ***Cytokines induce nuclear translocation of HIF-1 $\alpha$***

Protein levels of HIF-1 $\alpha$  are tightly regulated in normoxic conditions by post-translational modification and proteasomal degradation. However, functional levels of this transcription factor can be present in normoxic conditions [34]. We examined the expression and cellular localization of HIF-1 in unstimulated and cytokine (TNF, IL-1, IFN $\gamma$ ) stimulated cells. A549 transformed human lung epithelial cells were unstimulated or stimulated with cytokines and the expression of HIF-1 $\alpha$  was examined by immunoblotting. Basal HIF-1 $\alpha$  levels were detectable in whole cell lysates and cytokine stimulation marginally affected the total levels of this protein up to 8 h post-stimulation (Fig. 1A). The transcriptional activity of HIF-1 depends on its translocation from the cytosol into the nucleus. As such, the subcellular localization of HIF-1 $\alpha$  was assessed by fractionating cell lysates into cytosolic and nuclear components. Purity of the nuclear fraction was confirmed by high levels of histone H2A and undetectable amounts of caspase-8, and of the cytosolic fraction by high levels of caspase-8 and little histone H2A (Fig. 1B). The majority of HIF-1 $\alpha$  was present in the cytosol although there was a low level of this HIF-1 subunit in the nucleus of unstimulated cells. Cytokine stimulation increased HIF-1 $\alpha$  nuclear localization by 2 h post-stimulation and this persisted up to 8 h post-stimulation (Fig. 1B). This increase in HIF-1 $\alpha$  nuclear localization was coincident with the kinetics of iNOS induction

after cytokine stimulation, in which iNOS protein expression was apparent at 3 - 4 h post-stimulation (Fig. 1C).

### ***HIF-1 stabilization alone does not induce iNOS expression***

We first examined whether HIF-1 activation is sufficient to induce iNOS expression. A549 cells were untreated or subjected to hypoxia for 24 h, cells lysed, and the expression of HIF-1 $\alpha$  and iNOS determined by immunoblotting. Hypoxia increased HIF-1 $\alpha$  levels but did not induce iNOS expression (Fig. 2A). Also, the induction of iNOS expression by cytokines was not potentiated by hypoxia (Supplemental Figure 1). Similarly, treatment of cells with the hypoxic mimetic CoCl<sub>2</sub> increased HIF-1 $\alpha$  levels but did not induce iNOS expression (Fig. 2B). Thus, activation of HIF-1 alone is not sufficient to induce iNOS expression in human cells.

### ***Pharmacological inhibition of HIF-1 $\alpha$ reduces the induction of iNOS expression by cytokines***

Because we observed that cytokine stimulation of cells resulted in the increased nuclear localization of HIF-1 $\alpha$ , we examined whether this transcription factor is involved in the induction of iNOS expression in response to cytokine stimulation even though it was not sufficient for iNOS induction in response to hypoxia or CoCl<sub>2</sub>. A549 cells were treated with PX-478, a pharmacological inhibitor of HIF-1 $\alpha$  that prevents its expression [35, 36]. Consistent with its known effects, treatment of A549 cells with PX-478 markedly reduced HIF-1 $\alpha$  mRNA and protein expression (Fig. 3A & B). The effect of PX-478 on iNOS gene promoter activity was assessed by transfecting cells with a human iNOS promoter-luciferase plasmid, treating cells with PX-478 for 16h, and then leaving cells unstimulated or stimulating them with cytokines. The expression of luciferase was quantified 8 h later. Cytokines induced human iNOS gene promoter activity and this was significantly reduced by PX-478 (Fig. 3C). Cells were then pre-treated with PX-478 and the effect on cytokine stimulation of iNOS

mRNA and protein expression was quantified. PX-478 significantly reduced the induction of human iNOS mRNA and protein expression (Fig. 3D & E). In addition to A549 cells, the induction of iNOS expression by cytokines was also significantly reduced by PX-478 in primary human astrocytes (Fig. 4).

### ***Elimination of HIF-1 $\alpha$ reduces the induction of iNOS expression by cytokines***

The role of HIF-1 in the induction of human iNOS expression was then examined by eliminating HIF-1 $\alpha$  using CRISPR/Cas9-mediated gene editing. Cells were transfected with a plasmid encoding Cas9, a guide RNA targeting the human HIF-1 $\alpha$  gene, and eGFP. Transfected cells were isolated by single cell sorting of GFP-positive cells and clones expanded (Fig. 5A). Mutation of HIF-1 $\alpha$  in transfected clones was evaluated by PCR for the Cas9-targeted region of the HIF-1 $\alpha$  gene using primers specific for the wild-type non-edited sequence. In this scenario, primers bind to a non-edited allele but not the HIF-1 $\alpha$  gene if it has been altered by Cas9-targeting. Using this approach, two cell clones that contained edited HIF-1 $\alpha$  (HIF-1 $\alpha$ -KO) were identified and one was chosen for further experimentation (Fig. 5B; clone 2). Disruption of the HIF-1 $\alpha$  gene was confirmed by sequencing (data not shown). HIF-1 $\alpha$ -KO cells also lacked protein expression of HIF-1 $\alpha$  (Fig. 5C). The induction of HIF-regulated genes involved in the hypoxic response of cells was also assessed to determine HIF-1 function. Parental cells induced the expression of VEGF-A and HK-1 in response to the hypoxic mimetic CoCl<sub>2</sub> but the induction of these genes was completely absent in HIF-1 $\alpha$ -KO cells (Fig. 5D and Supplemental Figure 2). Finally, NF- $\kappa$ B is essential for the induction of iNOS expression in response to cytokines and its activation is sensitive to oxidative conditions in cells [37]. I $\kappa$ B $\alpha$  degradation in response to cytokine stimulation was not affected in HIF-1 $\alpha$ -KO cells, indicating that activation of the NF- $\kappa$ B pathway is not affected by the loss of HIF-1 $\alpha$  in our experiments (Supplemental Figure 3).

The induction of iNOS in HIF-1 $\alpha$ -KO cells was then examined. Parental and HIF-1 $\alpha$ -KO cells were transfected with a human iNOS promoter-luciferase plasmid and stimulated with cytokines. There was significantly less induction of human iNOS promoter activity in HIF-1 $\alpha$ -KO cells when compared to parental controls (Fig. 5E). When the expression of iNOS was examined, the induction of both iNOS mRNA and protein was reduced in HIF-1 $\alpha$ -KO cells (Fig. 5F & G and Supplemental Figure 4). In addition, cytokine-induced NO production was also reduced in HIF-1 $\alpha$ -KO cells (Fig. 5H). These data indicate that HIF-1 $\alpha$  is needed for full induction of human iNOS gene expression in response to cytokines.

***Cytokine-mediated iNOS gene promoter activity is induced by an HRE at -4981 bp***

HIF-1 mediates its transcriptional activity by binding to HREs in the gene promoters of target genes. The presence of HREs in the human iNOS promoter is poorly characterized [18]. Three putative HREs were identified at -7188, -4981 and -2220 bp upstream of the transcription start site using FIMO (Fig. 6A). To determine the role of each HRE in the induction of iNOS expression by cytokines, each putative binding element was eliminated in human iNOS promoter-luciferase plasmids using site-directed excision. A549 cells were transfected with wild type human iNOS promoter-luciferase or each mutated plasmid, left unstimulated or stimulated with cytokines, and luciferase activity quantified. Cytokine induction of the human iNOS gene promoter was significantly reduced only when the HRE at -4981 bp was eliminated (Fig. 6B). No significant effects were observed when the putative HREs at -7188 and -2220 were mutated.

**DISCUSSION**

Studying the mechanisms by which iNOS expression is regulated in human cells has implications for understanding human physiology and development of several diseases. We have shown that cytokine induction of iNOS expression in human cells involves the transcriptional activity of HIF-1. This depends on the presence of an HRE at -4981 kb of the iNOS promoter. Several studies have shown previously that iNOS expression is induced in response to hypoxia but the specific role of HIF-1 in regulating iNOS in human cells has not been determined [38-41]. To our knowledge, this is the first report of the direct regulation of human iNOS expression by HIF-1 and provides new insight into the regulation of NO production and the biological effects of HIF-1.

HIF-1 is a central regulator of cellular responses to hypoxia through the up-regulation of genes that increase tissue perfusion, alter metabolic programming, and increase cell survival. In addition to hypoxia, inflammatory cytokines also affect the actions of HIF-1. Both TNF and IL-1 have been shown to increase the DNA binding and transcriptional activity of HIF-1, and this is associated with increased levels of HIF-1 $\alpha$  in some studies but not others [42, 43]. The mechanism by which these cytokines increase HIF-1 $\alpha$  levels depends on the activation of NF- $\kappa$ B that stabilizes protein levels of this HIF-1 subunit [44, 45]. Other inflammatory stimuli that activate NF- $\kappa$ B, such as microbial products that bind toll-like receptors, also increase HIF-1 $\alpha$  expression [46]. In our studies, stimulation of A549 cells with cytokines increases nuclear accumulation of HIF-1 $\alpha$  but did not significantly increase HIF-1 $\alpha$  protein levels. This could differ from previous findings due to differences in the cell types used and/or the kinetics of HIF-1 $\alpha$  expression examined. With regard to the latter, we focused on the effect of cytokines on HIF-1 $\alpha$  levels early (within 8 h) after stimulation in order to examine processes related to the induction of gene transcription.

HIF-1 $\alpha$  expression can be inhibited using pharmacological and genetic approaches. PX-478 is a selective inhibitor of HIF-1 $\alpha$  expression and activity that has been examined as a potential therapy for cancer [47]. We observed that inhibiting HIF-1 $\alpha$  with PX-478 reduced the induction of both iNOS

mRNA and protein expression by cytokines in A549 cells and primary human astrocytes. In addition to this, our studies also eliminated HIF-1 $\alpha$  by CRISPR/Cas9-mediated gene editing. Genetic elimination of HIF-1 $\alpha$  in this manner significantly reduced the induction of iNOS promoter activity as well as mRNA and protein expression by cytokines, confirming a role for this transcription factor in the induction of iNOS gene expression. Although CRISPR/Cas9 targeting of HIF-1 $\alpha$  completely eliminated the induction of hypoxia-induced genes, there may be a small level of residual expression of HIF-1 $\alpha$  protein in HIF-1 $\alpha$ -KO cells after CoCl<sub>2</sub> exposure. A549 cells are a triploid cell line [48]. As such, editing of the HIF-1 $\alpha$  gene may have differentially inactivated HIF-1 $\alpha$  in each encoding allele by separate mechanisms such as eliminating protein expression or inactivating HIF-1 $\alpha$  function despite the persistence of limited protein expression.

Mouse models have provided information on the biology of iNOS but there are differences in the mechanisms by which human and mouse iNOS expression is induced [24, 49, 50]. The iNOS gene promoters in human and mice are different. For instance, the inducibility of the human iNOS gene promoter depends on transcription factor binding sequences dispersed throughout 8.3 kb upstream of the transcription start site while the murine iNOS gene promoter is dependent largely on a 1.6 kb region [5, 6, 51, 52]. With regard to hypoxia and HIFs, several studies have shown that hypoxia alone is sufficient to induce the expression of iNOS in murine cells [53, 54]. This effect of hypoxia is likely to be context-dependent because other studies show that hypoxia is not sufficient to induce iNOS expression [55]. There is an HRE in the murine iNOS gene promoter located between 209 - 227 bp upstream of the start site [18]. We have observed that stabilization of HIF-1 $\alpha$  alone is not sufficient to induce iNOS in human lung epithelial cells and that the HRE responsible for the induction of iNOS in response to cytokines is located at -4981 bp of the human iNOS promoter. These findings contribute to understanding how iNOS expression is specifically regulated in human cells and how this might differ from murine counterparts.

Understanding the mechanisms by which iNOS gene expression is regulated has implications for protective immunity and several pathological conditions. Production of iNOS by macrophages may be needed for the elimination of intracellular bacterial infections [56]. Recent evidence also indicates that the antimicrobial properties of macrophages depend on HIF-1 [57]. Also, HIFs increase the induction of aerobic glycolysis in macrophages, which is associated with development of a M1 macrophage phenotype that is protective against infection [58]. M1 macrophages are characterized by iNOS expression and our findings provide a potential link between HIF-1 activation and iNOS expression [59]. In addition to protective immunity, the induction of iNOS expression through a HIF-1-dependent mechanism protects against ischemic injury of the heart in murine models [60, 61]. Dysregulated activity of HIF-1 and of NO production are also involved in the development of several cancers. Expression of HIF-1 $\alpha$  in tumor cells exacerbates angiogenesis and tumor growth [62]. Increased expression of iNOS is associated with the pathogenesis of certain tumors, and dysregulated production of NO has been experimentally determined to contribute to the progression and maintenance of several types of cancer [63-66]. As such, iNOS may be a downstream target of HIF-1 that is important in tumorigenesis.

In summary, we have identified a new transcriptional regulatory mechanism of iNOS in human cells. HIF-1 $\alpha$  is needed for the maximal induction of iNOS expression in response to cytokines through amplification of human iNOS gene transcription. The findings provide insight into the role of iNOS in human biology and expand on developing knowledge of the cell biological roles of HIF in normoxia.

## ACKNOWLEDGEMENTS

We are grateful to Dr. Pascal Bernatchez for assistance with measuring nitrite levels and to Dr. David Geller for providing the human iNOS promoter-luciferase plasmid.



**Funding:** This study was funded by a grant from the Natural Sciences and Engineering Research Council of Canada (J.C.C.). JCC is the recipient of a Michael Smith Foundation for Health Research Scholar award.

## REFERENCES

1. Moncada S; Higgs EA. The discovery of nitric oxide and its role in vascular biology. *Br J Pharmacol.* 147:S193-201.;2006
2. Lowenstein CJ; Padalko E. iNOS (NOS2) at a glance. *J Cell Sci.* 117:2865-7;2004
3. Lowenstein CJ; Glatt CS; Bredt DS; Snyder SH. Cloned and expressed macrophage nitric oxide synthase contrasts with the brain enzyme. *Proc Natl Acad Sci U S A.* 89:6711-5;1992
4. Nussler AK; Di Silvio M; Billiar TR; Hoffman RA; Geller DA; Selby R; Madariaga J; Simmons RL. Stimulation of the nitric oxide synthase pathway in human hepatocytes by cytokines and endotoxin. *J Exp Med.* 176:261-4;1992
5. Taylor BS; de Vera ME; Ganster RW; Wang Q; Shapiro RA; Morris SM, Jr.; Billiar TR; Geller DA. Multiple NF-kappaB enhancer elements regulate cytokine induction of the human inducible nitric oxide synthase gene. *J Biol Chem.* 273:15148-56;1998
6. Marks-Konczalik J; Chu SC; Moss J. Cytokine-mediated transcriptional induction of the human inducible nitric oxide synthase gene requires both activator protein 1 and nuclear factor kappaB-binding sites. *J Biol Chem.* 273:22201-8;1998
7. Kristof AS; Marks-Konczalik J; Moss J. Mitogen-activated protein kinases mediate activator protein-1-dependent human inducible nitric-oxide synthase promoter activation. *J Biol Chem.* 276:8445-52. Epub 2000 Dec 8.;2001
8. Guo Z; Shao L; Feng X; Reid K; Marderstein E; Nakao A; Geller DA. A critical role for C/EBPbeta binding to the AABS promoter response element in the human iNOS gene. *Faseb J.* 17:1718-20;2003

9. Ganster RW; Taylor BS; Shao L; Geller DA. Complex regulation of human inducible nitric oxide synthase gene transcription by Stat 1 and NF-kappa B. *Proc Natl Acad Sci U S A*. 98:8638-43;2001
10. Lin N; Simon MC. Hypoxia-inducible factors: key regulators of myeloid cells during inflammation. *J Clin Invest*. 126:3661-71;2016
11. Yee Koh M; Spivak-Kroizman TR; Powis G. HIF-1 regulation: not so easy come, easy go. *Trends in biochemical sciences*. 33:526-34;2008
12. Jaakkola P; Mole DR; Tian YM; Wilson MI; Gielbert J; Gaskell SJ; von Kriegsheim A; Hebestreit HF; Mukherji M; Schofield CJ; Maxwell PH; Pugh CW; Ratcliffe PJ. Targeting of HIF-alpha to the von Hippel-Lindau ubiquitylation complex by O2-regulated prolyl hydroxylation. *Science*. 292:468-72;2001
13. Huang LE; Gu J; Schau M; Bunn HF. Regulation of hypoxia-inducible factor 1alpha is mediated by an O2-dependent degradation domain via the ubiquitin-proteasome pathway. *Proc Natl Acad Sci U S A*. 95:7987-92;1998
14. Forsythe JA; Jiang BH; Iyer NV; Agani F; Leung SW; Koos RD; Semenza GL. Activation of vascular endothelial growth factor gene transcription by hypoxia-inducible factor 1. *Mol Cell Biol*. 16:4604-13;1996
15. Semenza GL; Roth PH; Fang HM; Wang GL. Transcriptional regulation of genes encoding glycolytic enzymes by hypoxia-inducible factor 1. *J Biol Chem*. 269:23757-63;1994
16. Buck MD; Sowell RT; Kaech SM; Pearce EL. Metabolic Instruction of Immunity. *Cell*. 169:570-86;2017
17. Hanahan D; Weinberg RA. Hallmarks of cancer: the next generation. *Cell*. 144:646-74;2011
18. Melillo G; Musso T; Sica A; Taylor LS; Cox GW; Varesio L. A hypoxia-responsive element mediates a novel pathway of activation of the inducible nitric oxide synthase promoter. *J Exp Med*. 182:1683-93;1995

19. Mi Z; Rapisarda A; Taylor L; Brooks A; Creighton-Gutteridge M; Melillo G; Varesio L. Synergistic induction of HIF-1alpha transcriptional activity by hypoxia and lipopolysaccharide in macrophages. *Cell cycle (Georgetown, Tex)*. 7:232-41;2008
20. Melillo G; Taylor LS; Brooks A; Musso T; Cox GW; Varesio L. Functional requirement of the hypoxia-responsive element in the activation of the inducible nitric oxide synthase promoter by the iron chelator desferrioxamine. *J Biol Chem*. 272:12236-43;1997
21. Chartrain NA; Geller DA; Koty PP; Sitrin NF; Nussler AK; Hoffman EP; Billiar TR; Hutchinson NI; Mudgett JS. Molecular cloning, structure, and chromosomal localization of the human inducible nitric oxide synthase gene. *J Biol Chem*. 269:6765-72;1994
22. Chan GC; Fish JE; Mawji IA; Leung DD; Rachlis AC; Marsden PA. Epigenetic basis for the transcriptional hyporesponsiveness of the human inducible nitric oxide synthase gene in vascular endothelial cells. *J Immunol*. 175:3846-61;2005
23. Gross TJ; Kremens K; Powers LS; Brink B; Knutson T; Domann FE; Philibert RA; Milhem MM; Monick MM. Epigenetic silencing of the human NOS2 gene: rethinking the role of nitric oxide in human macrophage inflammatory responses. *J Immunol*. 192:2326-38;2014
24. Schneemann M; Schoedon G; Hofer S; Blau N; Guerrero L; Schaffner A. Nitric oxide synthase is not a constituent of the antimicrobial armature of human mononuclear phagocytes. *J Infect Dis*. 167:1358-63;1993
25. Yu Z; Xia X; Kone BC. Expression profile of a human inducible nitric oxide synthase promoter reporter in transgenic mice during endotoxemia. *Am J Physiol Renal Physiol*. 288:F214-20;2005
26. Robbins RA; Barnes PJ; Springall DR; Warren JB; Kwon OJ; BATTERY LD; Wilson AJ; Geller DA; Polak JM. Expression of inducible nitric oxide in human lung epithelial cells. *Biochem Biophys Res Commun*. 203:209-18;1994

27. Zhao ML; Liu JS; He D; Dickson DW; Lee SC. Inducible nitric oxide synthase expression is selectively induced in astrocytes isolated from adult human brain. *Brain Res.* 813:402-5;1998
28. Lee M; Choy JC. Positive feedback regulation of human inducible nitric-oxide synthase expression by Ras protein S-nitrosylation. *J Biol Chem.* 288:15677-86;2013
29. von Rossum A; Krall R; Escalante NK; Choy JC. Inflammatory Cytokines Determine the Susceptibility of Human CD8 T Cells to Fas-mediated Activation-induced Cell Death through Modulation of FasL and c-FLIPS Expression. *J Biol Chem.* 286:21137-44;2011
30. Graven KK; Yu Q; Pan D; Roncarati JS; Farber HW. Identification of an oxygen responsive enhancer element in the glyceraldehyde-3-phosphate dehydrogenase gene. *Biochim Biophys Acta.* 1447:208-18;1999
31. Choy JC; Wang Y; Tellides G; Pober JS. Induction of inducible NO synthase in bystander human T cells increases allogeneic responses in the vasculature. *Proc Natl Acad Sci U S A.* 104:1313-8. Epub 2007 Jan 16.;2007
32. Rahimi A; Anmole G; Soto-Nava M; Escamilla-Gomez T; Markle T; Jin SW; Lee GQ; Harrigan PR; Bangsberg DR; Martin J; Avila-Rios S; Reyes-Teran G; Brockman MA; Brumme ZL. In vitro functional assessment of natural HIV-1 group M Vpu sequences using a universal priming approach. *Journal of virological methods.* 240:32-41;2017
33. Heckman KL; Pease LR. Gene splicing and mutagenesis by PCR-driven overlap extension. *Nature protocols.* 2:924-32;2007
34. Treins C; Giorgetti-Peraldi S; Murdaca J; Semenza GL; Van Obberghen E. Insulin stimulates hypoxia-inducible factor 1 through a phosphatidylinositol 3-kinase/target of rapamycin-dependent signaling pathway. *J Biol Chem.* 277:27975-81;2002
35. Welsh S; Williams R; Kirkpatrick L; Paine-Murrieta G; Powis G. Antitumor activity and pharmacodynamic properties of PX-478, an inhibitor of hypoxia-inducible factor-1alpha. *Molecular cancer therapeutics.* 3:233-44;2004

36. Koh MY; Spivak-Kroizman T; Venturini S; Welsh S; Williams RR; Kirkpatrick DL; Powis G. Molecular mechanisms for the activity of PX-478, an antitumor inhibitor of the hypoxia-inducible factor-1alpha. *Molecular cancer therapeutics*. 7:90-100;2008
37. Haddad JJ; Olver RE; Land SC. Antioxidant/pro-oxidant equilibrium regulates HIF-1alpha and NF-kappa B redox sensitivity. Evidence for inhibition by glutathione oxidation in alveolar epithelial cells. *J Biol Chem*. 275:21130-9;2000
38. Yoshioka K; Thompson J; Miller MJ; Fisher JW. Inducible nitric oxide synthase expression and erythropoietin production in human hepatocellular carcinoma cells. *Biochem Biophys Res Commun*. 232:702-6;1997
39. Jiang ZL; Fletcher NM; Diamond MP; Abu-Soud HM; Saed GM. Hypoxia regulates iNOS expression in human normal peritoneal and adhesion fibroblasts through nuclear factor kappa B activation mechanism. *Fertility and sterility*. 91:616-21;2009
40. Kiang JG; Krishnan S; Lu X; Li Y. Inhibition of inducible nitric-oxide synthase protects human T cells from hypoxia-induced apoptosis. *Molecular pharmacology*. 73:738-47;2008
41. Nilsson I; Shibuya M; Wennstrom S. Differential activation of vascular genes by hypoxia in primary endothelial cells. *Experimental cell research*. 299:476-85;2004
42. Hellwig-Burgel T; Rutkowski K; Metzen E; Fandrey J; Jelkmann W. Interleukin-1beta and tumor necrosis factor-alpha stimulate DNA binding of hypoxia-inducible factor-1. *Blood*. 94:1561-7;1999
43. Sandau KB; Zhou J; Kietzmann T; Brune B. Regulation of the hypoxia-inducible factor 1alpha by the inflammatory mediators nitric oxide and tumor necrosis factor-alpha in contrast to desferroxamine and phenylarsine oxide. *J Biol Chem*. 276:39805-11;2001
44. Jung Y; Isaacs JS; Lee S; Trepel J; Liu ZG; Neckers L. Hypoxia-inducible factor induction by tumour necrosis factor in normoxic cells requires receptor-interacting protein-dependent nuclear factor kappa B activation. *Biochem J*. 370:1011-7;2003

45. Zhou J; Schmid T; Brune B. Tumor necrosis factor-alpha causes accumulation of a ubiquitinated form of hypoxia inducible factor-1alpha through a nuclear factor-kappaB-dependent pathway. *Mol Biol Cell*. 14:2216-25;2003
46. Ramanathan M; Pinhal-Enfield G; Hao I; Leibovich SJ. Synergistic up-regulation of vascular endothelial growth factor (VEGF) expression in macrophages by adenosine A2A receptor agonists and endotoxin involves transcriptional regulation via the hypoxia response element in the VEGF promoter. *Mol Biol Cell*. 18:14-23;2007
47. Zhao T; Ren H; Jia L; Chen J; Xin W; Yan F; Li J; Wang X; Gao S; Qian D; Huang C; Hao J. Inhibition of HIF-1alpha by PX-478 enhances the anti-tumor effect of gemcitabine by inducing immunogenic cell death in pancreatic ductal adenocarcinoma. *Oncotarget*. 6:2250-62;2015
48. Park SY; Choi HC; Chun YH; Kim H; Park SH. Characterization of chromosomal aberrations in lung cancer cell lines by cross-species color banding. *Cancer genetics and cytogenetics*. 124:62-70;2001
49. Schneemann M; Schoedon G. Species differences in macrophage NO production are important. *Nat Immunol*. 3:102.;2002
50. Pautz A; Art J; Hahn S; Nowag S; Voss C; Kleinert H. Regulation of the expression of inducible nitric oxide synthase. *Nitric Oxide*. 23:75-93;2010
51. Xie QW; Whisnant R; Nathan C. Promoter of the mouse gene encoding calcium-independent nitric oxide synthase confers inducibility by interferon gamma and bacterial lipopolysaccharide. *J Exp Med*. 177:1779-84.;1993
52. Lowenstein CJ; Alley EW; Raval P; Snowman AM; Snyder SH; Russell SW; Murphy WJ. Macrophage nitric oxide synthase gene: two upstream regions mediate induction by interferon gamma and lipopolysaccharide. *Proc Natl Acad Sci U S A*. 90:9730-4;1993
53. Jung F; Palmer LA; Zhou N; Johns RA. Hypoxic regulation of inducible nitric oxide synthase via hypoxia inducible factor-1 in cardiac myocytes. *Circ Res*. 86:319-25;2000

54. Olivares-Gonzalez L; Martinez-Fernandez de la Camara C; Hervas D; Millan JM; Rodrigo R. HIF-1alpha stabilization reduces retinal degeneration in a mouse model of retinitis pigmentosa. *FASEB J.* 2018
55. Jantsch J; Wiese M; Schodel J; Castiglione K; Glasner J; Kolbe S; Mole D; Schleicher U; Eckardt KU; Hensel M; Lang R; Bogdan C; Schnare M; Willam C. Toll-like receptor activation and hypoxia use distinct signaling pathways to stabilize hypoxia-inducible factor 1alpha (HIF1A) and result in differential HIF1A-dependent gene expression. *J Leukoc Biol.* 90:551-62;2011
56. MacMicking JD; Nathan C; Hom G; Chartrain N; Fletcher DS; Trumbauer M; Stevens K; Xie QW; Sokol K; Hutchinson N; et al. Altered responses to bacterial infection and endotoxic shock in mice lacking inducible nitric oxide synthase. *Cell.* 81:641-50;1995
57. Braverman J; Sogi KM; Benjamin D; Nomura DK; Stanley SA. HIF-1alpha Is an Essential Mediator of IFN-gamma-Dependent Immunity to Mycobacterium tuberculosis. *J Immunol.* 197:1287-97;2016
58. Palsson-McDermott EM; Curtis AM; Goel G; Lauterbach MA; Sheedy FJ; Gleeson LE; van den Bosch MW; Quinn SR; Domingo-Fernandez R; Johnston DG; Jiang JK; Israelsen WJ; Keane J; Thomas C; Clish C; Vander Heiden M; Xavier RJ; O'Neill LA. Pyruvate kinase M2 regulates Hif-1alpha activity and IL-1beta induction and is a critical determinant of the warburg effect in LPS-activated macrophages. *Cell metabolism.* 21:65-80;2015
59. Gordon S; Taylor PR. Monocyte and macrophage heterogeneity. *Nat Rev Immunol.* 5:953-64;2005
60. Natarajan R; Salloum FN; Fisher BJ; Kukreja RC; Fowler AA, 3rd. Hypoxia inducible factor-1 activation by prolyl 4-hydroxylase-2 gene silencing attenuates myocardial ischemia reperfusion injury. *Circ Res.* 98:133-40;2006

61. Belaidi E; Beguin PC; Levy P; Ribuot C; Godin-Ribuot D. Prevention of HIF-1 activation and iNOS gene targeting by low-dose cadmium results in loss of myocardial hypoxic preconditioning in the rat. *Am J Physiol Heart Circ Physiol.* 294:H901-8;2008
62. Maxwell PH; Dachs GU; Gleadle JM; Nicholls LG; Harris AL; Stratford IJ; Hankinson O; Pugh CW; Ratcliffe PJ. Hypoxia-inducible factor-1 modulates gene expression in solid tumors and influences both angiogenesis and tumor growth. *Proc Natl Acad Sci U S A.* 94:8104-9;1997
63. Lala PK; Chakraborty C. Role of nitric oxide in carcinogenesis and tumour progression. *The Lancet Oncology.* 2:149-56;2001
64. Lim KH; Ancrile BB; Kashatus DF; Counter CM. Tumour maintenance is mediated by eNOS. *Nature.* 452:646-9;2008
65. de Oliveira GA; Cheng RYS; Ridnour LA; Basudhar D; Somasundaram V; McVicar DW; Monteiro HP; Wink DA. Inducible Nitric Oxide Synthase in the Carcinogenesis of Gastrointestinal Cancers. *Antioxidants & redox signaling.* 26:1059-77;2017
66. Basudhar D; Glynn SA; Greer M; Somasundaram V; No JH; Scheiblin DA; Garrido P; Heinz WF; Ryan AE; Weiss JM; Cheng RYS; Ridnour LA; Lockett SJ; McVicar DW; Ambs S; Wink DA. Coexpression of NOS2 and COX2 accelerates tumor growth and reduces survival in estrogen receptor-negative breast cancer. *Proc Natl Acad Sci U S A.* 114:13030-5;2017

## FIGURE LEGENDS

**Figure 1. Cytokine stimulation increases HIF-1 $\alpha$  nuclear localization.** **A.** A549 cells were stimulated with cytokines (CM: TNF, IL-1, IFN $\gamma$ ), lysed at the indicated time-points, and HIF-1 $\alpha$  expression examined by immunoblot. **B.** A549 cells were untreated (UNT) or treated with cytokines, lysed at 2 h or 8 h post-stimulation, and lysates fractionated into cytosolic and nuclear components. HIF-1 $\alpha$  levels were examined by immunoblot. **C.** A549 cells were stimulated with cytokines, lysed at the indicated time-points, and iNOS protein expression examined by immunoblot. \*  $p < 0.05$ . \*\*  $p < 0.01$ .

**Figure 2. Stabilization of HIF-1 $\alpha$  alone is not sufficient to induce human iNOS expression.** A549 cells were **A.** subjected to hypoxia for 24 h, **B.** treated with the hypoxic mimetic CoCl $_2$  for 5 h, or




stimulated with cytokines (CM) for 5 h. Cells were lysed and the expression of HIF-1 $\alpha$  and iNOS determined by immunoblot. Data shown is representative of 3 independent experiments.

**Figure 3. Pharmacological inhibition of HIF-1 $\alpha$  reduces cytokine-mediated induction of iNOS expression in A549 cells.** **A & B.** A549 cells were treated with the HIF-1 $\alpha$  inhibitor PX-478 overnight and left untreated (UNT) or stimulated with cytokines (CM). **A.** HIF-1 $\alpha$  mRNA expression was quantified by RT-qPCR 2 h post-stimulation. **B.** HIF-1 $\alpha$  protein expression was determined by immunoblot at 6 h post-stimulation. **C.** Cells were transfected with an empty control plasmid (pXP2) or a human iNOS promoter-luciferase plasmid (pXP2-iNOS). Cells were untreated or treated with PX-478 prior to stimulation with cytokines. The induction of luciferase activity was quantified 8 h post-stimulation. **D & E.** Cells were untreated or treated with PX-478 and stimulated with cytokines. **D.** iNOS mRNA expression was quantified by RT-qPCR 2h post-stimulation. **E.** iNOS protein expression was examined by immunoblot at 6h post-stimulation \*  $p < 0.05$ , \*\*\*  $p < 0.001$ .


**Figure 4. Pharmacological inhibition of HIF-1 $\alpha$  reduces cytokine-mediated induction of iNOS expression in primary human astrocytes.** Primary human astrocytes were untreated or treated with PX-478 prior to stimulation with cytokines (CM). Cells were lysed after 6 h and iNOS expression determined by immunoblot. \*\*  $p < 0.01$

**Figure 5. CRISPR/Cas9-mediated gene editing of HIF-1 $\alpha$  reduces cytokine-mediated induction of iNOS expression.** **A.** Schematic of CRISPR/Cas9-mediated gene editing of HIF-1 $\alpha$ . **B.** DNA was isolated from cell clones that had been transfected with a plasmid not encoding a sgRNA or with a plasmid containing a HIF-1 $\alpha$  sgRNA. PCR using primers complimentary to the sgRNA-targeted sequence in HIF-1 $\alpha$  was performed to detect the presence of non-edited alleles. **C.** Parental A549 cells (WT) and a clone containing edited alleles of HIF-1 $\alpha$  (HIF-1 $\alpha$ -KO; clone 2) were stimulated with CoCl<sub>2</sub> and HIF-1 $\alpha$  expression determined by Western blot. Western blot shown is representative of 2 independent experiments. **D.** Parental A549 cells and HIF-1 $\alpha$ -KO cells were treated with CoCl<sub>2</sub> overnight and the mRNA expression of HIF-regulated genes VEGF-A and HK-1 quantified by RT-qPCR. Graph is the mean  $\pm$  SD of triplicate measurements in one of two replicate experiments. **E.** Parental A549 cells and HIF-1 $\alpha$ -KO cells were transfected with an empty control plasmid (pXP2) or a human iNOS promoter luciferase plasmid (pXP2-iNOS). Cells were stimulated with cytokines (CM) and luciferase activity quantified. **F-H.** Parental A549 cells and HIF-1 $\alpha$ -KO cells were stimulated with cytokines. **F.** iNOS mRNA expression was quantified by RT-qPCR. **G.** iNOS protein expression determined by immunoblot. **H.** NO production was determined by quantification of nitrite levels in the supernatant. \*  $p < 0.05$ , \*\*\*  $p < 0.001$ .

**Figure 6. An HRE at -4981 bp of the human iNOS promoter induces iNOS promoter activity in response to cytokines.** **A.** Schematic representation of the location of putative HREs in the human iNOS promoter and the human iNOS promoter luciferase plasmids that were generated to eliminate each of the HREs. **B.** A549 cells were transfected with plasmids encoding human iNOS promoter-luciferase or the iNOS promoters in which each of the putative HREs were eliminated. Cells were stimulated with cytokines (CM) and luciferase activity quantified. \*  $p < 0.05$ .




**Figure 1. Cytokine stimulation increases HIF-1α nuclear localization.** **A.** A549 cells were stimulated with cytokines (CM: TNF, IL-1, IFN $\gamma$ ), lysed at the indicated time-points, and HIF-1α expression examined by immunoblot. **B.** A549 cells were untreated (UNT) or treated with cytokines, lysed at 2 h or 8 h post-stimulation, and lysates fractionated into cytosolic and nuclear components. HIF-1α levels were examined by immunoblot. **C.** A549 cells were stimulated with cytokines, lysed at the indicated time-points, and iNOS protein expression examined by immunoblot. \*  $p < 0.05$ . \*\*  $p < 0.01$ .




**Figure 2. Stabilization of HIF-1 $\alpha$  alone is not sufficient to induce human iNOS expression.** A549 cells were subjected to **A.** hypoxia for 24 h, **B.** treated with the hypoxic mimetic CoCl<sub>2</sub> for 5 h, or stimulated with cytokines (CM) for 5 h. Cells were lysed and the expression of HIF-1 $\alpha$  and iNOS determined by immunoblot. Results are representative of 3 independent experiments.


A.




B.




C.




D.





E.





**Figure 4. Pharmacological inhibition of HIF-1 $\alpha$  reduces cytokine-mediated induction of iNOS expression in primary human astrocytes.** Primary human astrocytes were untreated or treated with PX-478 prior to stimulation with cytokines (CM). Cells were lysed after 6 h and iNOS expression determined by immunoblot. \*\* p < 0.01





**Figure 6. An HRE at -4981 bp of the human iNOS promoter induces iNOS promoter activity in response to cytokines.** **A.** Schematic representation of the location of putative HREs in the human iNOS promoter and the human iNOS promoter luciferase plasmids that were generated to eliminate each of the HREs. **B.** A549 cells were transfected with plasmids encoding human iNOS promoter-luciferase or the iNOS promoters in which each of the putative HREs were eliminated. Cells were stimulated with cytokines (CM) and luciferase activity quantified. \*  $p < 0.05$ .

**Highlights**

- Cytokines induce nuclear translocation of HIF-1 $\alpha$  in normoxia
- HIF-1 is needed for maximal induction of human iNOS expression
- There is a functional hypoxia response element at -4981 bp of the human iNOS gene promoter

Accepted manuscript