

Şinasi Özsoylu

Department of Pediatrics,
Hematology Unit, Faculty of Medicine,
Hacettepe University and
Hacettepe Children's Hospital,
Ankara, Turkey

I have read with interest the paper by Bartal et al. [1] in a recent issue of the Journal. The authors stated that 'lipid peroxidation was much higher in iron-deficient cells than in control cells'. However, the malonyldialdehyde (MDA) levels were given as nanomoles per gram hemoglobin (Hb). In iron deficiency anemia, more cells are required for 1 g Hb than in normal cells. MDA is not related to Hb but to the cell membrane. Therefore, MDA levels should be expressed per red blood cell rather than per gram Hb.

When I determined catalase activity in iron deficiency anemia [2], which was the first demonstration of an iron-containing enzyme deficiency, it could be shown when ex-

pressed per red cell but not with Hb level, as later stated by Acharya et al. [3].

It is surprising to see that some studies are not appreciated in the literature despite priority and several reinforcements [2, 4, 5].

Accepted:
March 24, 1994

Prof. Şinasi Özsoylu, MD
Department of Paediatrics, Faculty of Medicine
Hacettepe University
Hacettepe Children's Hospital
TR-06100 Ankara (Turkey)

© 1994 S. Karger AG, Basel
0001-5792/94/0913-0170
\$ 8.00/0

M. Bartal, D. Mazor
A. Dvilansky, N. Meyerstein

Department of Physiology,
Faculty of Health Sciences,
Ben-Gurion University of the Negev,
Beer-Sheva, Israel

We do agree that malonyldialdehyde (MDA) reflects lipid peroxidation and is not related to hemoglobin (Hb). In their detailed paper about auto-oxidation of human red blood cell lipids, Stocks and Dormandy [1] outlined the TBA reaction and calculated MDA in nanomoles per gram Hb. Later, Acharya et al. [2] emphasized that iron deficiency anemia cells were more sensitive to

oxidation when MDA levels were expressed per gram Hb. Our paper supports these findings. Unfortunately, we could not quote Prof. Özsoylu's paper in the *Turkish Journal of Pediatrics* (1964), as this journal is not available in our library.

Accepted:
April 24, 1994

Dr. M. Bartal
Department of Physiology, Faculty of Health Sciences
Ben-Gurion University of the Negev
POB 653
IL-84105 Beer-Sheva (Israel)

© 1994 S. Karger AG, Basel
0001-5792/94/0913-0170
\$ 8.00/0

Lipid Peroxidation in Iron Deficiency Anemia

References

- 1 Bartal M, Mazor D, Dvilansky A, Meyerstein N: Iron deficiency anemia: Recovery from in vitro oxidative stress. *Acta Haematol* 1993;90: 94-98.
- 2 Özsoylu Ş: Decreased activity of iron-containing enzyme catalase in human iron deficiency anemia. *Turk J Pediatr* 1964;6:200-210.
- 3 Acharya J, Punchard NA, Taylor JA, Thompson RPH, Pearson TC: Red cell lipid peroxidation and antioxidant enzymes in iron deficiency. *Eur J Haematol* 1991;47:287-291.
- 4 Özsoylu Ş: Detection of iron containing enzyme deficiency anemia. *J Pediatr* 1973;82:168.
- 5 Özsoylu Ş: Note to the paper. *Acta Haematol* 1986;76:181.

Reply

References

- 1 Stocks J, Dormandy TL: The autooxidation of human red cell lipids induced by hydrogen peroxidase. *Br J Haematol* 1971;20:95.
- 2 Acharya J, Punchard NA, Taylor JA, Thompson RPH, Pearson TC: Red cell lipid peroxidation and antioxidant enzymes in iron deficiency. *Eur J Haematol* 1991;47:287.