

# Bone Loss in Rats with Aldosteronism

ALIYE L. RUNYAN, VIKRAM S. CHHOKAR, MD, YAO SUN, MD, PhD;  
SYAMAL K. BHATTACHARYA, PhD; JOHN W. RUNYAN, MD; KARL T. WEBER, MD

**ABSTRACT:** *Objective:* We hypothesized that aldosteronism is accompanied by hypercalciuria and hypermagnesuria that lead to bone loss, which could be rescued by hydrochlorothiazide and spironolactone. *Methods:* We monitored 24-hour urinary  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  excretion; plasma ionized  $[\text{Ca}^{2+}]_o$  and  $[\text{Mg}^{2+}]_o$  and plasma  $\text{K}^+$ ; and bone mineral density of the femur. The following groups ( $n = 5$  in each group) were studied: age- and gender-matched, untreated controls; controls + 4 weeks hydrochlorothiazide; 4 weeks aldosterone/salt treatment (ALDOST,  $0.75 \mu\text{g/h}$  and dietary 1% NaCl/0.4% KCl); 4 weeks ALDOST + hydrochlorothiazide (50 mg/kg in prepared food); and 4 weeks ALDOST + hydrochlorothiazide + spironolactone (200 mg/kg day in divided doses by twice-daily gavage). *Results:* ALDOST increased ( $P < 0.05$ ) urinary  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  excretion four- and twofold, respectively; hydrochlorothiazide co-treatment attenuated ( $P < 0.05$ )  $\text{Ca}^{2+}$  excretion in controls and during ALDOST without affecting augmented  $\text{Mg}^{2+}$  excretion whereas hydrochlorothiazide + spironolactone normalized  $\text{Ca}^{2+}$  and reduced  $\text{Mg}^{2+}$  excretion ( $P < 0.05$ ). Compared with controls, plasma  $[\text{Ca}^{2+}]_o$  at 4 weeks of ALDOST was reduced ( $0.89 \pm 0.02$  versus  $0.83 \pm 0.03 \text{ mmol/L}$ ;  $P < 0.05$ ) but remained no different from levels in controls with hydrochlorothiazide and

hydrochlorothiazide + spironolactone ( $0.88 \pm 0.04$  and  $0.97 \pm 0.03 \text{ mmol/L}$ , respectively). Plasma  $[\text{Mg}^{2+}]_o$  fell ( $P < 0.05$ ) with ALDOST + hydrochlorothiazide ( $0.23 \pm 0.01$  versus  $0.34 \pm 0.01 \text{ mmol/L}$ ) and was prevented with spironolactone co-treatment ( $0.33 \pm 0.01 \text{ mmol/dL}$ ). Hypokalemia ( $2.9 \pm 0.2 \text{ mmol/L}$ ) occurred in rats with ALDOST + hydrochlorothiazide but not with spironolactone co-treatment. At 4 weeks of ALDOST, plasma parathyroid hormone was increased ( $30 \pm 4$  versus  $11 \pm 3 \text{ pg/mL}$ ;  $P < 0.05$ ) and bone mineral density was reduced ( $0.153 \pm 0.006$  versus  $0.170 \pm 0.002 \text{ g/cm}^2$ ;  $P < 0.05$ ). Co-treatments with either hydrochlorothiazide or hydrochlorothiazide + spironolactone each prevented bone loss. *Conclusions:* Hypercalciuria and hypermagnesuria accompany aldosteronism and account for a decline in their plasma ionized concentrations and secondary hyperparathyroidism with bone resorption. Attenuation of bone loss in aldosteronism can be achieved with hydrochlorothiazide; however, mono- and divalent cation homeostasis, together with bone integrity, are each preserved with the combination hydrochlorothiazide + spironolactone. **KEY INDEXING TERMS:** Aldosterone; Calcium; Magnesium; Parathyroid hormone; Bone loss. [Am J Med Sci 2005;330(1):1–7.]

**C**ongestive heart failure (CHF), a clinical syndrome whose origins are rooted in neurohormonal activation that includes effector hormones of the circulating renin-angiotensin-aldosterone system, is accompanied by a systemic illness. Pathophysiologic features include oxi/nitrosative stress in blood and such diverse tissues as skin, skeletal mus-

cle, immune cells, and heart<sup>1–5</sup> and a proinflammatory phenotype with elevated plasma levels of chemokines and cytokines and a catabolic state with loss of soft tissues and bone that eventuate in a wasting syndrome termed *cardiac cachexia*.<sup>6–12</sup> Pathogenic mechanisms involved in the appearance of this illness are under investigation.

An animal model of aldosteronism has been used to represent one aspect of the CHF neurohormonal profile. Uninephrectomized rats receive 1% NaCl in drinking water and aldosterone (ALDO) by minipump to inappropriately (relative to dietary  $\text{Na}^+$  intake) raise its plasma levels to those seen in cases of human CHF. Referred to as *aldosterone/salt treatment (ALDOST)*, this regimen is accompanied by a proinflammatory phenotype induced by the marked and persistent excretion of  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  in both urine and feces and which adversely influences extra- and intracellular concentrations of these divalent cations.<sup>13–18</sup> Pathophysiologic responses accompany the negative balance of  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$ . They include parathyroid

From the Divisions of Cardiovascular Diseases (ALR, VSC, YS, KTW) and Endocrinology (JWR), Department of Medicine and the Department of Surgery (SKB), University of Tennessee Health Science Center, Memphis, Tennessee.

Submitted for publication October 19, 2004; accepted for publication March 13, 2005.

Presented at the 2005 Southern Regional Meeting for the Society for Clinical Investigation.

This work was supported in part by NIH/NHLBI grant R01-HL73043 (KTW).

Correspondence: Karl T. Weber, MD, Division of Cardiovascular Diseases, University of Tennessee Health Science Center, 920 Madison Ave., Suite 300, Memphis, TN 38163. (E-mail: kwtweber@utmem.edu).

hormone-mediated loss of bone mineral density (and bone strength) and intracellular  $\text{Ca}^{2+}$  loading with induction of oxi/nitrosative stress in diverse cells and an immunostimulatory state with activation of peripheral blood mononuclear cells that invade the coronary and systemic circulations to create a proinflammatory vascular phenotype.<sup>13-22</sup>

Herein we focused on the bone loss seen with aldosteronism,<sup>13</sup> which far exceeds that seen solely with the hypercalciuria accompanying a high-NaCl diet alone.<sup>23</sup> We hypothesized that pharmacologic intervention could attenuate the urinary loss of these divalent cations seen with ALDOST and thereby would prevent bone loss. To uncouple hypercalciuria and hypermagnesuria during ALDOST, we selected hydrochlorothiazide, to promote urinary  $\text{Ca}^{2+}$  resorption and compared this intervention to the combination of hydrochlorothiazide and spironolactone, an ALDO receptor antagonist, to reduce both  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  excretion. Age- and gender-matched, unoperated, untreated rats served as controls. Accordingly, we monitored 24-hour urinary  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  excretion, plasma ionized  $[\text{Ca}^{2+}]_o$  and  $[\text{Mg}^{2+}]_o$ , plasma  $\text{K}^+$ , and femur mineral density.

## Methods

### Animal Model

Male, 8- to 12-week-old Sprague-Dawley rats (Harlan Sprague Dawley, Indianapolis, IN) were used in this study, approved by the institution's Animal Care and Use Committee. There were five groups with five rats in each group, unless otherwise specified. Age- and gender-matched, unoperated, untreated rats served as controls. As previously reported, ALDOST consists of uninephrectomized rats who receive ALDO (0.75  $\mu\text{g}/\text{h}$ ) by implanted minipump (Alzet, Cupertino, CA) together with a 1% NaCl/0.4% KCl drinking water and standard laboratory chow (Harlan Tekland 2215 Rodent Diet) containing 1.13%  $\text{Ca}^{2+}$ .<sup>24</sup> Separate groups of animals received ALDOST plus hydrochlorothiazide or hydrochlorothiazide alone (treated controls). Hydrochlorothiazide was mixed into powdered standard chow (50 mg/kg body weight) and water was added. The mixture was placed in a pan and cut into small squares and air dried. We did not monitor daily food intake. Yet another group received ALDOST and hydrochlorothiazide plus spironolactone (200 mg/kg/day) in divided doses by twice-daily gavage. At 4 weeks of treatment, the animals were anesthetized and killed, and blood and femur samples were harvested.

### Urinary $\text{Ca}^{2+}$ and $\text{Mg}^{2+}$ Excretion

On the day of the metabolic study, food was withheld but water with 1% NaCl was provided. Animals were "bathed" in distilled water to remove any feces or food that could contaminate collected urine. Ani-

mals were then placed in a cleaned, minerally decontaminated and distilled-deionized, water-rinsed metabolic cage. Urine was collected over 24 hours and kept frozen for  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  assay. After each use, cages were manually cleaned with deionized water; all nonmetallic parts were washed with diluted hydrochloric acid (3N), rinsed three times with deionized water, and finally rinsed twice with distilled-deionized water for future use as previously reported.<sup>13,14</sup>

Urinary  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  concentrations were determined as reported elsewhere<sup>13,14</sup> using an atomic absorption spectrophotometer. Urinary  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  excretion rates were calculated from the product of their concentration ( $\mu\text{g}/\text{mL}$ ) by the 24-hour urine volume ( $\text{mL}/24$  hours) and expressed as  $\mu\text{g}/24$  hours.

### Plasma Ionized $[\text{Ca}^{2+}]_o$ and $[\text{Mg}^{2+}]_o$ and Plasma $\text{K}^+$

The concentrations of plasma ionized  $[\text{Ca}^{2+}]_o$  and  $[\text{Mg}^{2+}]_o$  and  $\text{K}^+$  were determined by the direct ion-selective electrode technique using a Nova 8 Analyzer (Nova Biomedical, Waltham, MA) and expressed in  $\text{mmol}/\text{L}$ .

### Parathyroid Hormone

Plasma parathyroid hormone (PTH) was measured by the intact PTH immunoassay (IRMA) using a commercial kit (Nichols Institute Diagnostics, San Clemente, CA). IRMA is a two-site immunoradiometric assay for the measurement of the biologically intact 84 amino acid chain of PTH molecule. Blood (2 mL) was collected from the rat heart into a chilled EDTA tube and immediately centrifuged ( $1600 \times g$ ) for 15 minutes. Plasma was then separated and kept at  $-80^\circ \text{C}$ . For IRMA, each plasma sample (200  $\mu\text{L}$ ) was added to the tube containing 100  $\mu\text{L}$  of the  $^{125}\text{I}$ -PTH antibody solution and PTH antibody-coated beads and incubated for 24 hours. Beads were then washed twice with washing solution and each test tube was counted with a gamma counter for 1 minute. A standard curve was generated using prepared intact PTH standards and plasma PTH values and expressed as  $\text{pg}/\text{mL}$  plasma.

### Bone Mineral Density

Bone mineral density was determined for excised, manually cleaned femurs by peripheral dual-energy x-ray absorptiometry using GE Lunar PIXImus2 (GE Healthcare, Fairfield, CT). Quality control and calibration were carried out within 24 hours of each scanning period. This method has been validated for rat long bones.<sup>25</sup> We have previously reported on the equivalence of this noninvasive assessment of tibia and femur bone mineral density with their total concentrations of  $\text{Mg}^{2+}$  and  $\text{Ca}^{2+}$  determined by atomic absorption spectrophotometry.<sup>13</sup>

### Statistical Analysis

Values are presented as mean  $\pm$  SEM. Data were analyzed using analysis of variance. Significant differences between individual means were determined using the post hoc Bonferroni multiple comparisons test. Significance was assigned to  $P < 0.05$ .

## Results

### Urinary $\text{Ca}^{2+}$ and $\text{Mg}^{2+}$ Excretion

We ascertained that the 24-hour urinary  $\text{Ca}^{2+}$  excretion rate in unoperated controls was  $895.7 \pm 138.2 \mu\text{g}/24$  hours. Treatment of control rats with hydrochlorothiazide led to an expected reduction in urinary  $\text{Ca}^{2+}$  excretion (Table 1). Four weeks of ALDOST raised urinary  $\text{Ca}^{2+}$  excretion more than fourfold. Co-treatment with hydrochlorothiazide reduced the 24-hour urinary  $\text{Ca}^{2+}$  excretion seen with ALDOST, whereas the addition of spironolactone to the hydrochlorothiazide regimen further reduced ( $P < 0.05$ ) urinary  $\text{Ca}^{2+}$  excretion to levels seen in untreated controls.

Urinary  $\text{Mg}^{2+}$  excretion in unoperated, untreated controls was  $969.6 \pm 137.2 \mu\text{g}/24$  hours. This was significantly increased by hydrochlorothiazide treatment (see Table 1). At 4 weeks of ALDOST, 24-hour urinary  $\text{Mg}^{2+}$  excretion was markedly increased ( $P < 0.05$ ) above control values, as was the case with ALDOST + hydrochlorothiazide. The addition of spironolactone to the ALDOST + hydrochlorothiazide regimen significantly reduced urinary  $\text{Mg}^{2+}$  excretion compared with ALDOST alone, whereas it did not normalize urinary  $\text{Mg}^{2+}$  excretion to levels seen in controls.

### Body Weight

Body weights for 12-week-old male controls treated with hydrochlorothiazide for 4 weeks were not significantly different from one another ( $310 \pm 6$  versus  $286 \pm 6$  mg, respectively). At 4 weeks of ALDOST, however, body weight was significantly less than either of these controls ( $224 \pm 11$ ), in keeping with our previous findings, in which ALDOST is accompanied by reduced weight gain.<sup>17</sup> Co-treatment with either hydrochlorothiazide or hydrochlorothiazide + spironolactone prevented this

failure to gain weight ( $288 \pm 9$  and  $276 \pm 4$  mg, respectively), which we also previously reported for spironolactone co-treatment.<sup>17</sup>

### Plasma Ionized $[\text{Ca}^{2+}]_o$ and $[\text{Mg}^{2+}]_o$

At week 4 of ALDOST, plasma  $[\text{Ca}^{2+}]_o$  was reduced ( $P < 0.05$ ) compared with control values ( $0.83 \pm 0.03$  versus  $0.89 \pm 0.02$  mmol/L, respectively). However, ALDOST + hydrochlorothiazide prevented this fall in ionized  $\text{Ca}^{2+}$  ( $0.88 \pm 0.04$  mmol/L) and co-treatment with hydrochlorothiazide + spironolactone was accompanied by a rise in  $[\text{Ca}^{2+}]_o$  to  $0.97 \pm 0.03$  mmol/L that was greater ( $P < 0.05$ ) than ALDOST values, but not greater than the values in the five control rats in this study or the 19 historical controls in this laboratory ( $0.97 \pm 0.01$ ;  $0.88$ – $1.09$  mmol/L).

Plasma ionized  $[\text{Mg}^{2+}]_o$  was reduced at 4 weeks of ALDOST, but did not reach statistical significance ( $0.31 \pm 0.02$  mmol/L) versus control values ( $0.34 \pm 0.01$  mmol/L). At 4 weeks of ALDOST + hydrochlorothiazide, plasma  $[\text{Mg}^{2+}]_o$  was significantly less than with ALDOST alone ( $0.23 \pm 0.01$  mmol/L), and this was totally rescued by ALDOST + hydrochlorothiazide + spironolactone ( $0.33 \pm 0.01$  mmol/L).

### Plasma $\text{K}^+$

In separate studies, plasma  $\text{K}^+$  concentrations were determined in 9-, 10- and 12-week-old untreated control rats. These values were found to be  $4.4 \pm 0.1$ ,  $3.8 \pm 0.1$ , and  $4.3 \pm 0.2$  mmol/L, respectively, and ranged between 3.4 and 5.1 mmol/L. A value of less than 3.4 mmol/L was therefore chosen to define hypokalemia. At week 4 of hydrochlorothiazide treatment, none of the control rats developed hypokalemia ( $3.6 \pm 0.1$  mmol/L), as was also the case for those receiving ALDOST alone ( $4.0 \pm 0.3$  mmol/L). Four of five rats with ALDOST + hydrochlorothiazide had hypokalemia. Plasma  $\text{K}^+$  fell below 3.0 mmol/L in 2 animals receiving ALDOST + hydrochlorothiazide. However, none of the rats receiving ALDOST + hydrochlorothiazide + spironolactone had hypokalemia. Plasma  $\text{K}^+$  did not exceed 5.1 mmol/L in any animal in any of the treatment groups, including those with spironolactone co-treatment. At week 4 of ALDOST + hydrochlorothiazide

**Table 1.** Urinary  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  Excretion at Week 4 of Treatment

	$\text{Ca}^{2+}$ g/24 h (n)	$\text{Mg}^{2+}$ g/24 h (n)
Control	$895.7 \pm 138.2$ (5)	$969.6 \pm 137.2$ (5)
Control + hydrochlorothiazide	$152.4 \pm 14.0^a$ (5)	$2245.8 \pm 556.1^a$ (5)
ALDOST	$4968.5 \pm 1077.8^a$ (4)	$3856.3 \pm 439.6^a$ (5)
ALDOST + hydrochlorothiazide	$2066.5 \pm 846.8$ (5)	$3181.7 \pm 1065.7$ (5)
ALDOST + hydrochlorothiazide + spironolactone	$854.0 \pm 152.4^b$ (5)	$1887.4 \pm 707.6^b$ (5)

Values mean  $\pm$  SEM.

<sup>a</sup>  $P < 0.05$  vs. control.

<sup>b</sup>  $P < 0.05$  vs. ALDOST.



treatment, plasma  $K^+$  level was  $2.9 \pm 0.2$  and was significantly ( $P < 0.001$ ) less than the  $4.3 \pm 0.2$  mmol/L seen with ALDOST + hydrochlorothiazide + spironolactone.

#### Parathyroid Hormone

At 4 weeks of ALDOST, plasma PTH levels were increased ( $P < 0.05$ ) compared with control values ( $30 \pm 4$  versus  $11 \pm 3$  pg/mL).

#### Bone Mineral Density

Bone mineral density for the femur of 12-week-old, male, unoperated, untreated control rats was  $0.170 \pm 0.005$  g/cm<sup>2</sup> (Figure 1). At 4 weeks of ALDOST, bone mineral density was significantly ( $P < 0.05$ ) reduced to  $0.153 \pm 0.014$  g/cm<sup>2</sup>. This reduction in bone mineral density was rescued by co-treating rats with ALDOST using either hydrochlorothiazide alone or hydrochlorothiazide plus spironolactone. Four weeks' treatment of controls with hydrochlorothiazide did not result in a fall in bone mineral density.

### Discussion


This study led to several major findings. First, and confirmatory to recent reports from our laboratory,<sup>13,14</sup> is the hypercalciuria and hypermagnesuria induced by long-term ALDOST. Short-term studies in rodents<sup>26-29</sup> have identified the hypercalciuria seen with mineralocorticoid excess using either ALDO or deoxycorticosterone (DOC), together with dietary NaCl. An elevation in urinary  $Ca^{2+}$  excretion is also seen in humans during short-term administration of a mineralocorticoid, which is inappropriate relative to dietary sodium intake.<sup>30,31</sup> In primary aldosteronism, basal hypercalciuria is accentuated by dietary  $Na^+$  loading<sup>31</sup> and is normalized by spironolactone or surgical removal of diseased adrenal tissue.<sup>32</sup> Hypermagnesuria is also

seen in patients with primary aldosteronism and resolves after adrenal surgery or in response to spironolactone treatment.<sup>33</sup>

The stimulus for the hypercalciuria and hypermagnesuria that accompany ALDOST is not completely understood. Elevations in arterial pressure, which occur in response to chronic mineralocorticoid excess, and the mineralocorticoid hormone per se, in the absence of inappropriate dietary  $Na^+$  intake, have long been discounted.<sup>26,27</sup> Additionally, polydipsia and metabolic alkalosis, which accompany chronic mineralocorticoid/salt treatment, have likewise been eliminated.<sup>29</sup> Dietary  $Na^+$  loading alone (e.g., an 8% NaCl diet) leads to a suppression of plasma ALDO and is accompanied by hypercalciuria.<sup>34-38</sup> Prevailing opinion suggests the mechanism responsible for the hypercalciuria seen during ALDOST is likely related to an expansion of extravascular fluid volume including intravascular and interstitial compartments. This results in decreased proximal tubular  $Na^+$  and  $Ca^{2+}$  resorption and their increased distal delivery. The mineralocorticoid hormone promotes distal  $Na^+$  resorption without an effect on  $Ca^{2+}$ , whereby  $Ca^{2+}$  excretion is increased.<sup>26-28,31</sup> Furthermore, nitric oxide-mediated enhancement in medullary blood flow may be facilitative.<sup>39-41</sup>

The sustained increment in urinary  $Ca^{2+}$  and  $Mg^{2+}$  excretion that accompanies 4 weeks of ALDOST led to a fall in plasma ionized  $[Ca^{2+}]_o$  and  $[Mg^{2+}]_o$ , which was statistically significant for  $[Ca^{2+}]_o$ . Co-treatment with hydrochlorothiazide or hydrochlorothiazide + spironolactone each prevented the fall in  $[Ca^{2+}]_o$ . However, the fall in  $[Mg^{2+}]_o$  was accentuated by ALDOST + hydrochlorothiazide but could be rescued with spironolactone co-treatment. The fall in  $[Ca^{2+}]_o$  and  $[Mg^{2+}]_o$  provide well-known stimuli to the parathyroid glands to release PTH. Plasma levels of PTH were significantly increased at week 4 of ALDOST. Increased plasma PTH levels would be expected to increase  $Ca^{2+}$  and  $Mg^{2+}$  resorption from bone and  $Ca^{2+}$  absorption from kidneys. PTH levels are increased in rats treated with DOC/salt.<sup>29,34,42</sup> Secondary hyperparathyroidism has been observed in patients with primary hyperaldosteronism,<sup>43</sup> in whom reduced serum ionized  $Ca^{2+}$  and elevated PTH levels are normalized by spironolactone or adrenal surgery.<sup>32,43</sup> In addition to bone stores, gastrointestinal  $Ca^{2+}$  absorption would be augmented by  $1,25(OH)_2D_3$  formed in kidneys and driven by PTH. Increased duodenal absorption of  $Ca^{2+}$  by this active vitamin D metabolite accompanies the hypercalciuria associated with DOC/salt treatment.<sup>29</sup>

Our second major finding is the loss of bone mineral density that accompanies 4 weeks of ALDOST. The compensatory responses provided by PTH to preserve plasma ionized  $Ca^{2+}$  and  $Mg^{2+}$  were likely responsible for the fall in femur bone mineral den-



**Figure 1.** Bone mineral density (BMD) was monitored in rat femur for the various treatment groups and in age- and gender-matched controls (C). CT4, control + 4 weeks hydrochlorothiazide; A4, 4 weeks ALDOST; AT4, 4 weeks ALDOST + hydrochlorothiazide; ATS4, 4 weeks ALDOST + hydrochlorothiazide + spironolactone.

sity at week 4 of ALDOST. Bone loss in the rat model of mineralocorticoid/salt excess has been suggested by others<sup>42,44,45</sup> based on increased urinary hydroxyproline, a marker of bone resorption. The secondary aldosteronism that appears in patients with CHF is accompanied by reductions in bone mineral density of moderate to marked severity, together with elevated levels of intact PTH or its C-terminal PTH-related peptide.<sup>7,8,10-12,46-49</sup> The influence of furosemide, a potent loop diuretic commonly used in these patients that promotes urinary  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  excretion, may exacerbate bone loss and secondary hyperparathyroidism.

An expected reduction in urinary  $\text{Ca}^{2+}$  excretion was found with hydrochlorothiazide alone in control rats. This was also the case when hydrochlorothiazide was given during ALDOST and in combination with spironolactone during ALDOST. This represents our third major finding and one not previously reported. In rescuing urinary  $\text{Ca}^{2+}$  excretion, each regimen prevented bone loss. Hydrochlorothiazide has demonstrated pharmacologic actions within the distal tubule that include enhanced  $\text{Ca}^{2+}$  resorption while raising urinary  $\text{Mg}^{2+}$  excretion.<sup>50-52</sup> It is this property that reverses secondary hyperparathyroidism and the hypercalciuria associated with vitamin D administration.<sup>53,54</sup> We found hydrochlorothiazide to reduce ALDOST-induced hypercalciuria on average by 70%, which, over the course of 4 weeks of ALDOST, prevented the reduction in bone mineral density.

Aldosterone increases thiazide receptor density principally in the renal cortex, and this, in turn, has led to a greater natriuresis than that observed in thiazide-treated controls and in situations in which the magnitude of the natriuresis correlated with receptor density.<sup>55-58</sup> We did not measure urinary  $\text{Na}^+$  excretion in this study. However, relative to its influence on urinary  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  excretion, we found hydrochlorothiazide responses in control rats and in ALDOST rats to be comparable. The use of spironolactone in combination with hydrochlorothiazide provided an additive effect in further reducing urinary  $\text{Ca}^{2+}$  excretion and would be in keeping with its antagonism of ALDO receptors present within more distal segments of the nephron. The combination of hydrochlorothiazide + spironolactone in the management of CHF has not been systematically examined in recent times and may prove as effective a diuretic regimen as furosemide alone without the marked hypercalciuria and hypermagnesuria that accompany this loop diuretic.

The reduction in urinary  $\text{Ca}^{2+}$  excretion provided by hydrochlorothiazide, however, occurred at the expense of increased urinary  $\text{Mg}^{2+}$  and  $\text{K}^+$  excretion, which can adversely influence their intracellular levels in diverse tissues.<sup>59-64</sup> Hypokalemia appeared in a substantial number of rats receiving ALDOST + hydrochlorothiazide despite a diet sup-

plemented with 0.4% KCl and in some cases was marked ( $<3.0$  mmol/L). The addition of spironolactone to hydrochlorothiazide prevented the appearance of hypokalemia in rats with ALDOST, as is the case in diuretic-treated individuals.<sup>65,66</sup> An elevation in plasma  $\text{K}^+$  levels above 5.1 mmol/L was not observed in any of the experimental groups, each of whom had normal renal function.

Our findings suggested that the hypercalciuria and hypermagnesuria of aldosteronism can be attenuated by hydrochlorothiazide and hydrochlorothiazide + spironolactone, and this will prevent the loss of bone mineral density. This conclusion would imply that bone strength would also be preserved.<sup>13,14</sup> In preventing osteopenia, these interventions may prevent bone fractures, an important health care problem, particularly among elderly populations, in whom heart failure is so prevalent. In the setting of aldosteronism, such as appears with CHF, the appearance of hydrochlorothiazide-induced hypokalemia can be offset by spironolactone co-administration. Our study suggested that the combination of hydrochlorothiazide + spironolactone will better preserve both mono- and divalent cation homeostasis and bone integrity in aldosteronism. The addition of spironolactone to an angiotensin-converting enzyme inhibitor and loop diuretic in the overall management of CHF has proven effective in reducing mortality and morbidity events, including ventricular arrhythmias and the risk of sudden cardiac death.<sup>67-69</sup> The renal and gastrointestinal losses of  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  that accompany aldosteronism are considerable and are measured in milligram quantities in feces and microgram levels in urine. A marked reduction in bone mineral density is found at and beyond week 4 of ALDOST.<sup>13</sup> A high-NaCl diet alone and a loop diuretic will each induce hypercalciuria and hypermagnesuria. In the setting of aldosteronism, these interventions could lead to an exaggerated loss of these divalent cations and to adverse responses, including parathyroid hormone-mediated bone loss and  $\text{Ca}^{2+}$  loading of diverse tissues (e.g., myocardium) and cells (e.g., platelets and lymphocytes).<sup>70</sup> The judicious use of a thiazide diuretic, in combination with spironolactone, may protect against such adverse outcomes.

## References

1. Miyamoto M, Kishimoto C, Shioji K, et al. Cutaneous arteriolar thioredoxin expression in patients with heart failure. *Circ J* 2003;67:116-8.
2. Tsutsui H, Ide T, Hayashidani S, et al. Enhanced generation of reactive oxygen species in the limb skeletal muscles from a murine infarct model of heart failure. *Circulation* 2001;104:134-6.
3. Cesselli D, Jakoniuk I, Barlucchi L, et al. Oxidative stress-mediated cardiac cell death is a major determinant of ventricular dysfunction and failure in dog dilated cardiomyopathy. *Circ Res* 2001;89:279-86.

4. **Singh N, Dhalla AK, Seneviratne C, et al.** Oxidative stress and heart failure. *Mol Cell Biochem* 1995;147:77–81.
5. **Hill MF, Singal PK.** Antioxidant and oxidative stress changes during heart failure subsequent to myocardial infarction in rats. *Am J Pathol* 1996;148:291–300.
6. **Mann DL, Young JB.** Basic mechanisms in congestive heart failure: recognizing the role of proinflammatory cytokines. *Chest* 1994;105:897–904.
7. **Muchmore JS, Cooper DK, Ye Y, et al.** Prevention of loss of vertebral bone density in heart transplant patients. *J Heart Lung Transplant* 1992;11:959–63.
8. **Van Cleemput J, Daenen W, Nijs J, et al.** Timing and quantification of bone loss in cardiac transplant recipients. *Transpl Int* 1995;8:196–200.
9. **Anker SD, Chua TP, Ponikowski P, et al.** Hormonal changes and catabolic/anabolic imbalance in chronic heart failure and their importance for cardiac cachexia. *Circulation* 1997;96:526–34.
10. **Anker SD, Clark AL, Teixeira MM, et al.** Loss of bone mineral in patients with cachexia due to chronic heart failure. *Am J Cardiol* 1999;83:612–5.
11. **Lee AH, Mull RL, Keenan GF, et al.** Osteoporosis and bone morbidity in cardiac transplant recipients. *Am J Med* 1994;96:35–41.
12. **Shane E, Mancini D, Aaronson K, et al.** Bone mass, vitamin D deficiency, and hyperparathyroidism in congestive heart failure. *Am J Med* 1997;103:197–207.
13. **Chhokar VS, Sun Y, Bhattacharya SK.** Loss of bone minerals and strength in rats with aldosteronism. *Am J Physiol Heart Circ Physiol* 2004;287:H2023–6.
14. **Chhokar VS, Sun Y, Bhattacharya SK, et al.** Hyperparathyroidism and the calcium paradox of aldosteronism. *Circulation* 2005;111:871–8.
15. **Sun Y, Zhang J, Lu L, et al.** Aldosterone-induced inflammation in the rat heart: role of oxidative stress. *Am J Pathol* 2002;161:1773–81.
16. **Gerling IC, Sun Y, Ahokas RA, et al.** Aldosteronism: an immunostimulatory state precedes the proinflammatory/fibrogenic cardiac phenotype. *Am J Physiol Heart Circ Physiol* 2003;285:H813–21.
17. **Ahokas RA, Warrington KJ, Gerling IC, et al.** Aldosteronism and peripheral blood mononuclear cell activation: a neuroendocrine-immune interface. *Circ Res* 2003;93:e124–e35.
18. **Ahokas RA, Sun Y, Bhattacharya SK, et al.** Aldosteronism and a proinflammatory vascular phenotype: role of  $Mg^{2+}$ ,  $Ca^{2+}$  and  $H_2O_2$  in peripheral blood mononuclear cells. *Circulation* 2005;111:51–7.
19. **Pu Q, Neves MF, Virdis A, et al.** Endothelin antagonism on aldosterone-induced oxidative stress and vascular remodeling. *Hypertension* 2003;42:49–55.
20. **Virdis A, Neves MF, Amiri F, et al.** Spironolactone improves angiotensin-induced vascular changes and oxidative stress. *Hypertension* 2002;40:504–10.
21. **Park JB, Schiffrin EL.** Cardiac and vascular fibrosis and hypertrophy in aldosterone-infused rats: role of endothelin-1. *Am J Hypertens* 2002;15:164–9.
22. **Rocha R, Rudolph AE, Friedrich GE, et al.** Aldosterone induces a vascular inflammatory phenotype in the rat heart. *Am J Physiol* 2002;283:H1802–10.
23. **Lalande A, Roux C, Graulet AM, et al.** The diuretic indapamide increases bone mass and decreases bone resorption in spontaneously hypertensive rats supplemented with sodium. *J Bone Min Res* 1998;13:1444–50.
24. **Brilla CG, Pick R, Tan LB, et al.** Remodeling of the rat right and left ventricle in experimental hypertension. *Circ Res* 1990;67:1355–64.
25. **Nagy TR, Prince CW, Li J.** Validation of peripheral dual-energy X-ray absorptiometry for the measurement of bone mineral in intact and excised long bones of rats. *J Bone Min Res* 2001;16:1682–7.
26. **Suki WN, Schwetzmans RS, Rector FC Jr, et al.** Effect of chronic mineralocorticoid administration on calcium excretion in the rat. *Am J Physiol* 1968;215:71–4.
27. **Massry SG, Coburn JW, Chapman LW, et al.** The effect of long-term desoxycorticosterone acetate administration on the renal excretion of calcium and magnesium. *J Lab Clin Med* 1968;71:212–9.
28. **Gehr MK, Goldberg M.** Hypercalciuria of mineralocorticoid escape: clearance and micropuncture studies in the rat. *Am J Physiol* 1986;251:F879–88.
29. **Zikos D, Langman C, Gaftor U, et al.** Chronic DOCA treatment increases Ca absorption: role of hypercalciuria and vitamin D. *Am J Physiol* 1986;251:E279–84.
30. **Cappuccio FP, Markandu ND, MacGregor GA.** Renal handling of calcium and phosphate during mineralocorticoid administration in normal subjects. *Nephron* 1988;48:280–3.
31. **Rastegar A, Agus Z, Connor TB, et al.** Renal handling of calcium and phosphate during mineralocorticoid "escape" in man. *Kidney Int* 1972;2:279–86.
32. **Rossi E, Sani C, Perazzoli F, et al.** Alterations of calcium metabolism and of parathyroid function in primary aldosteronism, and their reversal by spironolactone or by surgical removal of aldosterone-producing adenomas. *Am J Hypertens* 1995;8:884–93.
33. **Horton R, Biglieri EG.** Effect of aldosterone on the metabolism of magnesium. *J Clin Endocrinol Metab* 1962;22:1187–92.
34. **Wright GL, Rankin GO.** Concentrations of ionic and total calcium in plasma of four models of hypertension. *Am J Physiol* 1982;243:H365–70.
35. **Chan AY, Poon P, Chan EL, et al.** The effect of high sodium intake on bone mineral content in rats fed a normal calcium or a low calcium diet. *Osteoporos Int* 1993;3:341–4.
36. **Chan EL, Swaminathan R.** Effect of different amounts of sodium intake for 4 months on calcium metabolism in normal and oophorectomized rats. *J Bone Min Res* 1993;8:1185–9.
37. **Lemann J Jr, Pleuss, JA, Hornick L, et al.** Dietary NaCl-restriction prevents the calciuria of KCl-deprivation and blunts the calciuria of  $KHCO_3$ -deprivation in healthy adults. *Kidney Int* 1995;47:899–906.
38. **Sellmeyer DE, Schloetter M, Sebastian A.** Potassium citrate prevents increased urine calcium excretion and bone resorption induced by a high sodium chloride diet. *J Clin Endocrinol Metab* 2002;87:2008–12.
39. **Salazar FJ, Llinas MT, Gonzalez JD, et al.** Role of prostaglandins and nitric oxide in mediating renal response to volume expansion. *Am J Physiol* 1995;268:R1442–R48.
40. **Zou AP, Cowley AW Jr.** Role of nitric oxide in the control of renal function and salt sensitivity. *Curr Hypertens Rep* 1999;1:178–86.
41. **Granger JP, Alberola AM, Salazar FJ, et al.** Control of renal hemodynamics during intrarenal and systemic blockade of nitric oxide synthesis in conscious dogs. *J Cardiovasc Pharmacol* 1992;20(Suppl 12):S160–2.
42. **Berthelot A, Pernot F, Gairard A.** Increase in urinary excretion of phosphates, hydroxyproline, and cAMP during acetate desoxycorticosterone + saline treatment in the rat. *Arch Int Pharmacodyn Ther* 1978;235:196–203.
43. **Resnick LM, Laragh JH.** Calcium metabolism and parathyroid function in primary aldosteronism. *Am J Med* 1985;78:385–90.



44. **Berthelot A, Desplan C, Pernot F, et al.** I. PTH and U. cAMP during mineralocorticoid hypertension in the rat. *Horm Metab Res* 1978;10:83–4.
45. **Palmieri GM, Hawrylko J.** Effects of aldosterone on the urinary excretion of total and non-dialyzable hydroxyproline. *Horm Metab Res* 1977;9:507–9.
46. **Stefenelli T, Pacher R, Woloszczuk W, et al.** Parathyroid hormone and calcium behavior in advanced congestive heart failure. *Zschr Kardiol* 1992;81:121–5.
47. **Schmid C, Kiowski W.** Hyperparathyroidism in congestive heart failure. *Am J Med* 1998;104:508–9.
48. **Ogino K, Ogura K, Kinugasa Y, et al.** Parathyroid hormone-related protein is produced in the myocardium and increased in patients with congestive heart failure. *J Clin Endocrinol Metab* 2002;87:4722–7.
49. **Zittermann A, Schleithoff SS, Tenderich G, et al.** Low vitamin D status: a contributing factor in the pathogenesis of congestive heart failure? *J Am Coll Cardiol* 2003;41:105–12.
50. **Puschett JB.** Sites and mechanisms of action of diuretics in the kidney. *J Clin Pharmacol* 1981;21:564–74.
51. **Krause U, Zielke A, Schmidt-Gayk H, et al.** Direct tubular effect on calcium retention by hydrochlorothiazide. *J Endocrinol Invest* 1989;12:531–5.
52. **Stefikova K, Spustova V, Dzurik R.** Acute effect of hydrochlorothiazide on renal calcium and magnesium handling in postmenopausal women. *Physiol Res* 1999;48:327–30.
53. **Favus MJ, Coe FL, Kathpalia SC, et al.** Effects of chlorothiazide on 1,25-dihydroxyvitamin D<sub>3</sub>, parathyroid hormone, and intestinal calcium absorption in the rat. *Am J Physiol* 1982;242:G575–81.
54. **Alon U, Wellons MD, Chan JC.** Reversal of vitamin-D<sub>2</sub>-induced hypercalciuria by chlorothiazide. *Pediatr Res* 1983;17:117–9.
55. **Beaumont K, Vaughn DA, Fanestil DD.** Thiazide diuretic drug receptors in rat kidney: identification with [<sup>3</sup>H]metolazone. *Proc Natl Acad Sci U S A* 1988;85:2311–4.
56. **Chen Z, Vaughn DA, Blakely P, et al.** Adrenocortical steroids increase renal thiazide diuretic receptor density and response. *J Am Soc Nephrol* 1994;5:1361–8.
57. **Kim GH, Masilamani S, Turner R, et al.** The thiazide-sensitive Na-Cl cotransporter is an aldosterone-induced protein. *Proc Natl Acad Sci U S A* 1998;95:14552–7.
58. **Abdallah JG, Schrier RW, Edelstein C, et al.** Loop diuretic infusion increases thiazide-sensitive Na<sup>+</sup>/Cl<sup>-</sup>-cotransporter abundance: role of aldosterone. *J Am Soc Nephrol* 2001;12:1335–41.
59. **Okusa MD, Velazquez H, Ellison DH, et al.** Luminal calcium regulates potassium transport by the renal distal tubule. *Am J Physiol*. 1990;258(2 Pt 2):F423–8.
60. **Dyckner T, Wester PO.** Renal excretion of electrolytes in patients on long-term diuretic therapy for arterial hypertension and/or congestive heart failure. *Acta Med Scand* 1985;218:443–8.
61. **Dyckner T, Wester PO, Widman L.** Amiloride prevents thiazide-induced intracellular potassium and magnesium losses. *Acta Med Scand* 1988;224:25–30.
62. **Ryan MP.** Magnesium and potassium-sparing diuretics. *Magnesium* 1986;5:282–92.
63. **Dorup I, Skajaa K, Clausen T, et al.** Reduced concentrations of potassium, magnesium, and sodium-potassium pumps in human skeletal muscle during treatment with diuretics. *Br Med J (Clin Res Ed)* 1988;296:455–8.
64. **Hollifield JW.** Thiazide treatment of hypertension: effects of thiazide diuretics on serum potassium, magnesium, and ventricular ectopy. *Am J Med* 1986;80:8–12.
65. **Murdoch DL, Forrest G, Davies DL, et al.** A comparison of the potassium and magnesium-sparing properties of amiloride and spironolactone in diuretic-treated normal subjects. *Br J Clin Pharmacol* 1993;35:373–8.
66. **Dyckner T, Wester PO.** Intracellular magnesium loss after diuretic administration. *Drugs* 1984;28(Suppl 1):161–6.
67. **Ramires FJ, Mansur A, Coelho O, et al.** Effect of spironolactone on ventricular arrhythmias in congestive heart failure secondary to idiopathic dilated or to ischemic cardiomyopathy. *Am J Cardiol* 2000;85:1207–11.
68. **Pitt B, Zannad F, Remme WJ, et al.** The effect of spironolactone on morbidity and mortality in patients with severe heart failure: Randomized Aldactone Evaluation Study Investigators. *N Engl J Med* 1999;341:709–17.
69. **Domanski M, Norman J, Pitt B, et al.** Diuretic use, progressive heart failure, and death in patients in the Studies Of Left Ventricular Dysfunction (SOLVD). *J Am Coll Cardiol* 2003;42:705–8.
70. **Weber KT.** Furosemide in the long-term management of heart failure. The good, the bad and the uncertain. *J Am Coll Cardiol* 2004;44:1308–10.