
REVIEW/HYPOTHESIS

Nutrition, hormones, and breast cancer: Is insulin the missing link?

Rudolf Kaaks

(Received 17 October 1995; accepted in revised form 5 August 1996)

Breast cancer incidence rates are high in societies with a Western lifestyle characterized by low levels of physical activity, and by an energy-dense diet rich in total and saturated fat and refined carbohydrates. Epidemiologic studies, so far mostly on postmenopausal women, have shown that breast cancer risk is increased in hyperandrogenic women, with decreased levels of plasma sex-hormone binding globulin, and with increased levels of testosterone and of free estrogens. This paper describes the role of hyperinsulinemia as a physiologic link between nutritional lifestyle factors, obesity, and the development of a hyperandrogenic endocrine profile, and reviews evidence that may or may not support the theory that chronic hyperinsulinemia is an underlying cause of breast cancer. An hypothesis is presented, stipulating that breast cancer risk is increased not only in hyperandrogenic postmenopausal women, but also in premenopausal women with mild hyperandrogenism and normal (ovulatory) menstrual cycles. The author suggests further investigation as to whether there is a positive association between risk of breast cancer before menopause and subclinical forms of the polycystic ovary syndrome (PCOS), and to what extent diet and physical activity during childhood, by modulating the degree of insulin resistance during adolescence, may or may not be determinants of a PCO-like hyperandrogenic endocrine profile persisting into adulthood. *Cancer Causes and Control* 1996, 7, 605-625

Key words: Breast cancer, insulin, insulin-like growth factors, nutrition, obesity, polycystic ovary syndrome, puberty, steroid hormones.

1. Introduction

Breast cancer incidence rates are up to five, or even 10 times higher in western Europe and North America than in rural Africa or Asia.¹ Studies on changing cancer incidence patterns in migrant populations,^{2,3} and on time trends in breast cancer incidence rates over time in Western or 'Westernizing' countries⁴⁻⁶ suggest that these differences in breast cancer incidence are determined largely by environmental factors, probably diet and nutrition-related lifestyle. In countries with high breast cancer incidence rates, dietary intake patterns generally are characterized by high intake levels of total and saturated fat and of refined carbohydrates.^{7,8} In these high-

risk countries, the average age at menarche tends to be lower,⁹ and women reach a higher adult body stature,^{10,11} which suggests that factors determining breast cancer risk act early in life. Additional suggestions that environmental influences may act at an early age come from studies on changes over time in age-specific breast cancer incidence rates, which appear to occur cohort-wise (*i.e.*, by birth year).^{5,6,12} The cohort-wise changes in incidence rates also suggest that similar environmental factors determine the risk of developing breast cancer before or after menopause. Effects of nutrition may be related particularly to energy balance. A high level of physical activity during

Dr Kaaks is with the International Agency for Research on Cancer, Lyon, France. Address correspondence to Dr Kaaks, International Agency for Research on Cancer, 150 cours Albert-Thomas, 69372 Lyon Cedex 08, France.

adolescence has been found to be protective against breast cancer.¹³⁻¹⁵ In animal studies, caloric restriction has been shown to reduce considerably the development of mammary tumors.¹⁶

Case-control studies and prospective cohort studies have confirmed the existence of associations of the risk of developing breast cancer – either before or after menopause – with age at menarche¹⁷ and body stature.¹⁸ In addition, these studies show a relation between overweight, as body mass index (wt/ht²) (BMI), and breast cancer risk, but only for breast cancers occurring after menopause. Besides the BMI, a high ratio of waist circumference divided by hip circumference (waist-to-hip ratio, or WHR) is associated with increased risk of postmenopausal breast cancer, suggesting that intra-abdominal fat stores in particular are a risk factor.¹⁹⁻²³ Interestingly, however, there is no strong evidence for a positive association between a high BMI and breast cancer before menopause; on the contrary, results from a recent meta-analysis indicate that this association may even be weakly negative.²⁴

Results from case-control and cohort studies on the relation between breast cancer risk and dietary composition are less consistent. Reviews and meta-analyses of case-control studies suggest that the risk of developing breast cancer may be increased by high intake levels of either total or, more specifically, saturated fat, whereas risk may be decreased by high consumption levels of vegetables, fruits, or dietary fiber.^{18, 25-27} These results, however, have not been confirmed clearly by prospective cohort studies thus far.^{18,27} One hypothesis, that has been put forward to explain the lack of clear associations between breast cancer risk and dietary intake patterns in mid-life, is that nutritional factors may have their strongest impact before or around puberty.^{18,25}

In terms of physiologic mechanisms, nutritional factors are thought to modify breast cancer risk by influencing levels of endogenous hormones. Nutrition-induced differences in hormonal metabolism also would explain associations between breast cancer risk and age at menarche, adult body stature, and obesity. Until recently, however, the understanding of the link between nutrition and endocrine profiles predisposing to the development of breast cancer has been rather limited.

Section 2 of this review briefly describes two main hypotheses on endogenous steroid hormones in relation to breast cancer, for which evidence has accumulated from epidemiologic studies. Sections 3 and 4 discuss the relations between hyperinsulinemia and steroid hormonal profiles that seem to be associated with increased breast cancer risk, and how plasma insulin levels may be co-determined by dietary composition and physical activity. Sections 5 and 6 cover the transient insulin resistance generally observed during puberty, and its possible role

in the development of the polycystic ovary syndrome (PCOS) as a form of 'functional ovarian hyperandrogenism' (FOH). Section 7 ('Discussion and synthesis') is an overall discussion of observations that do, or do not, support the hypothesis that breast cancer risk generally is increased in women with a hyperandrogenic endocrine profile, which may develop mainly as a consequence of chronic hyperinsulinemia.

2. Endogenous steroid hormones and breast cancer: free estrogens and ovarian androgen excess

Early hypotheses on associations between endogenous hormones and development of breast cancer focused primarily on the effects of ovarian hormones, and especially estrogens, because it had been recognized very early that:

- i. estrogens can induce mammary tumors in rodents;²⁸
- ii. the risk of developing breast cancer risk is increased among women with early menarche²⁹ or late menopause,³⁰ and age-specific incidence rates rise less steeply after the age of natural menopause when ovarian estrogen production has ceased;³¹ and
- iii. breast cancer risk decreases after artificial menopause (ovariectomy).³²

These observations initially led to a search for evidence of an excess of endogenous estrogen production in women with breast cancer. The 'estrogen excess hypothesis' evolved into the 'free estrogen hypothesis' (reviewed by Zumoff³³) when, in the early 1980s, Siiteri *et al*³⁴ reported that only plasma levels of free estradiol, unbound to sex-hormone binding globulin (SHBG), were elevated in women with breast cancer, even though total estradiol concentrations did not differ. It is now a well-established concept that only the fraction of estrogens unbound to SHBG is biologically available to receptors in target cells.^{35,36} Since the 1960s, a large number of case-control studies on endogenous steroid hormone levels have been conducted, the results of which have been reviewed by Key and Pike,³⁷ and by Bernstein and Ross.³⁸ As discussed in these reviews, there is substantial evidence that high plasma levels of available estrogen are associated with increased risk of breast cancer, at least among postmenopausal women (on whom most studies have focused thus far). This conclusion receives further support from the New York Women's Health Study³⁹ on 130 cases of breast cancer detected during an average of five and one-half years of follow-up of a cohort of about 7,000 postmenopausal women, with a relative risk (RR) of about 4.0 between the extreme quartiles of the plasma concentrations of free

estradiol. A similar RR of about 4.0 for high *cf* low levels of free estradiol was observed recently in a prospective study of 71 breast cases in a cohort of 3,375 postmenopausal women (the 'Breast Cancer Serum Bank' study).⁴⁰

In parallel with the earliest studies leading to the estrogen-excess hypothesis, Grattarola^{41,42} observed that a substantial subset of breast cancer patients had chronic anovulation, if premenopausal, and excessive urinary excretion of testosterone irrespective of menopausal status. Histologically, resected ovaries of these patients showed hyperplasia of interstitial cells, as already reported by Sommers.^{43,44} These histologic changes in the ovary are similar to those seen in women with polycystic ovary syndrome (PCOS), a condition characterized by enlarged ovaries containing an increased number of follicles, and by clinical symptoms related to hyperandrogenism (irregularity of menstrual cycles, anovulation, hirsutism).⁴⁵⁻⁴⁷ Subsequent studies by Secreto *et al* confirmed Grattarola's findings, and showed an elevated urinary excretion of androgens,^{48,49} as well as elevated plasma concentrations of testosterone in both premenopausal^{50,51} and postmenopausal^{52,53} patients with breast cancer. These observations eventually led to the formulation of the 'ovarian androgen excess' hypothesis.³³ Besides the studies by Secreto *et al*, increased plasma levels of testosterone have been observed in several other studies comparing breast cancer cases with controls.⁵⁴⁻⁵⁷ Although in two early prospective studies, breast cancer risk was not associated significantly with total plasma testosterone levels, either before⁵⁸ or after menopause,^{58,59} significantly increased risks were observed in the Italian 'ORDET' study⁶⁰ (24 women with breast cancer compared with 88 selected controls within a cohort of 4,053 postmenopausal women; RR = 7.0 for the highest *cf* lowest tertile), and in the Breast Cancer Serum Bank Study⁴⁰ (RR = 6.2 for the highest *cf* lowest quartile). In the New York Women's Health Study, breast cancer risk also was increased significantly in postmenopausal women with high plasma testosterone levels (Toniolo, New York University, NY, USA; personal communication, June 1996).

3. Free estrogens and ovarian androgen excess: relation to obesity and insulin resistance

Studies on both pre- and postmenopausal women show that obesity is associated with increased plasma concentrations of testosterone,⁶¹⁻⁶⁶ and decreased concentrations of sex-hormone binding globulin (SHBG).^{61,64-70} Due to the rise in testosterone production and the drop in SHBG levels, and because testosterone has a higher affinity than estrogens for SHBG,^{71,72} free estrogen concentrations increase. In addition, obesity often is associated with insulin resistance, a nutritional-metabolic state charac-

terized by less than normal sensitivity of target tissues (liver, muscle, adipose tissue) to the physiologic effects of insulin (stimulation of glucose uptake, inhibition of lipolysis). To compensate for the loss of insulin sensitivity, larger amounts of insulin are secreted by the pancreatic β -cells to maintain normal plasma concentrations of glucose. Thus, even though there may be no clear signs of glucose intolerance (*i.e.*, the pancreas may still secrete sufficient amounts of insulin to maintain plasma glucose concentrations within the normal homeostatic range),⁷³ fasting plasma concentrations of insulin and postprandial secretions of insulin by the pancreas are increased.^{61-65,67-70,73,74} These hormonal imbalances (androgen excess, low SHBG levels, and insulin resistance) are related in particular to a high WHR as a measure of 'central' or 'intra-abdominal' obesity^{66,75-77} and, importantly, this relation between hormonal imbalances and WHR also is observed in normal-weight women,^{62,63} or in overweight women after adjustment for BMI.^{66,78}

Until the late 1980s, the predominant view was that insulin resistance arises subsequent to an overproduction of ovarian androgens, in particular testosterone,^{77,79} or to an imbalance between androgens from either ovarian, or adrenal origin.^{80,81} Direct mechanisms by which androgens may affect insulin sensitivity are the subject of much debate (see discussions in Buffington *et al*⁸¹ and Poretsky⁸²). Nevertheless, it seems clear that androgens may favor a more centripetal body fat distribution pattern,^{75,83-85} or increase the mobilization of free fatty acids from adipose tissue,^{86,87} which relates to development of insulin resistance after long-term overeating (see section 4). On the other hand, there is substantial evidence for an inverse causal relation, whereby high insulin levels potentiate the luteinizing hormone (LH)-stimulated production of ovarian steroids, and particularly of testosterone.⁸⁸⁻⁹²

The decrease in plasma levels of SHBG in obese women originally was thought also to be the result of high plasma concentrations of testosterone, or by a high ratio of testosterone to estradiol.^{71,72} Again, however, there is now at least equally compelling evidence that insulin is the central regulating factor.^{93,94} Insulin has been shown to inhibit the production of SHBG in liver cells,^{93,95-97} and relatively strong negative correlations have been found between plasma concentrations of insulin and SHBG in obese women, or in women with PCOS.⁹⁸⁻¹⁰²

Insulin, and insulin-like growth factors

Insulin-like growth factor I (IGF-I) is a structural homologue of insulin,¹⁰³ and has long been known to be a mediator of growth hormone signal, stimulating anabolic processes required for physical growth during puberty and adolescence, as well as during adult life. Many tissue types have been shown to have receptors for IGF-I.¹⁰⁴⁻¹⁰⁷

These receptors also show considerable homology with those for insulin,¹⁰⁸ and insulin and IGF-I both have some capacity to cross-bind to their mutual receptors.¹⁰⁹

A major part of the IGF-I circulating in blood plasma is released by the liver, where its production is stimulated primarily by growth hormone (GH). In addition, IGF-I is known to be produced within many of its target tissues,^{105,107,110} where it is involved in the paracrine control of cell proliferation.^{104,107} Like insulin, IGF-I has a gonadotropic action, potentiating the production of steroid hormones – in particular androgens – in the ovaries^{90,111} and, like insulin, it inhibits the hepatic production of SHBG.^{96,97}

Plasma concentrations of IGF-I are several hundred-fold higher than those of insulin, but most of the circulating IGF-I (± 95 percent) is bound to a number of different IGF-binding proteins (IGFBP) and therefore is not directly available to target tissues.^{112,113} The plasma concentrations of IGF-I, and of its most abundant binding protein, IGFBP-3, are regulated primarily by growth hormone (GH) activity,¹¹³ as a function of nutritional status, in particular an adequate dietary supply of energy and protein.^{107,113-115} Fasting,¹¹⁶ weight loss,¹¹⁷ or an increase in energy expenditure due to physical activity¹¹⁸ can induce a drop in plasma IGF-I levels within days, while these levels can be restored quickly by nutritional repletion.^{107,113,115}

A smaller fraction of IGF-I is bound to a smaller binding protein, IGFBP-1. This binding protein shows marked circadian variation that parallels the natural secretory pattern of insulin during the day, and plasma levels of the binding protein were shown to decrease immediately after infusion of either glucose or insulin.¹¹⁹⁻¹²³ Moreover, in women with obesity,⁶⁵ or with PCOS,^{100,124} and in diabetic adolescents,¹²⁵ fasting insulin concentrations have been found to be correlated negatively with IGFBP-1 levels. More definite evidence that plasma levels of IGFBP-1 are down-regulated by insulin, rather than by glucose, has come from a study by Suikkari *et al*¹²⁶ where human volunteers were given an infusion of insulin while blood glucose was clamped at euglycemic levels. In addition, insulin has been shown to inhibit IGFBP-1 production in liver (hepatoma) cells.^{97,127-129}

The decrease in IGFBP-1 concentrations is believed to correspond to an increased biological activity of IGF-I at the level of target organs which, among a wide variety of body tissues, includes the ovaries and the breast. Thus, through a down-regulation of IGFBP-1, insulin may amplify its effects on steroid hormone metabolism by increasing the availability of IGF-I within the ovaries, and the effects of insulin on steroid hormone metabolism even may be mediated entirely by an increased physiologic activity of IGF-I⁹⁰ (see Figure 1). The mechanisms underlying this increased availability, however, are not




Figure 1. The role of insulin resistance in the development of a hyperandrogenic endocrine profile.

yet fully understood. Possibly, a decrease in IGFBP-1 levels results in a higher concentration of unbound IGF-I, which then can diffuse freely towards target cells. There also is some evidence, however, that insulin stimulates the active transport of the IGFBP-1-IGF-1 across the endothelial surface of blood vessels.¹³⁰

Like IGF-I itself, IGF-binding proteins are produced locally within most of its target tissues,^{104,107,113} and IGFBP-1 (as well as other IGF-binding proteins) is synthesized by ovarian granulosa cells,^{89,131,132} and it has been detected in follicular fluid.¹²⁴ Thus, another postulated mechanism by which high-plasma insulin levels may produce hyperandrogenism is a decrease in the local production of IGFBP-1 within the ovarian tissues, thereby increasing the local biological activity of IGF-I.^{90,133}

4. Insulin resistance and hyperinsulinemia: relations with diet and physical activity

Some severe, relatively infrequent forms of insulin resistance are known to be caused by a genetic defect in the insulin receptor ('type A' syndrome), or by the production of auto-antibodies against the insulin receptor ('type B' syndrome).¹³⁴ The more common forms of insulin resistance usually are related to obesity, or to diseases associated with obesity such as non-insulin-dependent diabetes mellitus (NIDDM) or PCOS.^{134,135} Nevertheless, subgroups of young adults with reduced insulin sensitivity in the absence of overweight also have been identified,¹³⁶⁻¹³⁹ and, in a survey in the United States, it has been estimated that about 20 percent of the normal-weight men and women, with normal glucose tolerance, had decreased insulin sensitivity comparable to that of individuals with obesity or non-insulin-dependent

diabetes mellitus (NIDDM).¹⁴⁰ As in overweight individuals, however, reduced insulin sensitivity in normal-weight individuals is associated with an increased WHR.^{62,63}

A pathogenic model of insulin resistance

Mechanistic models to explain the nutrition-related pathogenesis of insulin resistance emphasize the role of excess oxidation of free (non-esterified) fatty acids (FFA).^{73,86,87,141-144} Excessive oxidation of FFA by liver or muscle cells leads to a biochemical inhibition of enzymes involved in the intracellular glycolytic cascade (Krebs cycle), and in intracellular glucose storage in the form of glycogen.^{141,145-149} In response to these lowered enzyme activities, the rate of glucose transport into the cell decreases. Variability in the rate of insulin-stimulated glucose storage,¹⁵⁰ and in the glycogen synthesizing capacity of muscle cells,¹⁵¹ has been shown to account for a large proportion of between-person variations in insulin sensitivity impairment in normal subjects, and in NIDDM patients, respectively. In skeletal muscle, insulin action also has been shown to be related inversely to the size of intracellular stores of triglycerides, which can be increased by a high-fat diet.^{152,153}

The starting point in the development of insulin resistance induced by overnutrition is thought to be an increase in the amount of intra-abdominal fat.^{86,87,154} Intra-abdominal fat stores are mobilized more easily than fat stored in femoral-gluteal region, because of their sensitivity to lipolytic stimuli (e.g., by adrenaline, through β 3-adrenergic receptors¹⁵⁵), and because of a relative insensitivity to the antilipolytic effects of insulin.¹⁵⁶ The release of FFA from visceral fat into the portal blood may inhibit glucose uptake and metabolism by the liver and impair hepatic glycogen storage.¹⁴¹ The mobilization of fat from body fat deposits also will increase FFA levels in peripheral blood, with similar effects on the insulin sensitivity of skeletal muscles. In addition, increased oxidation of FFA by the liver is associated with a stimulation of gluconeogenesis.¹⁵⁷⁻¹⁶² The reduced uptake of glucose by the liver and skeletal muscle, combined with an increased hepatic gluconeogenesis, results in raised fasting plasma-glucose levels which, in turn, lead to chronically elevated (fasting) plasma-insulin concentrations. There is experimental evidence that insulin can down-regulate the concentration of its own receptors on the membranes of target cells,^{161,162} and the degree of hyper-insulinemia in insulin-resistant individuals correlates with the reduction in the number of insulin receptors in liver, muscle, or adipose tissue cells.^{163,164} The resulting reduction of insulin sensitivity may be aggravated by additional, post-receptor defects in the response to insulin binding.¹³⁴ As discussed in the previous sections, an increase in insulin levels reduces plasma concentrations of SHBG, and may increase concentrations of testosterone (Figure 1). The

resulting increase in androgenicity, combined with high plasma insulin levels, favors a centripetal type of body fat distribution^{75,83-85,165} and, in interaction with GH, at the same time may stimulate the mobilization of FFA from body fat deposits.^{86,87} Thus, a cycle is established in which a decrease in insulin sensitivity, initially associated with relatively mild intra-abdominal adiposity, through an increase in free androgen levels causes a more androgenic body fat distribution which, in turn, will exacerbate the insulin resistance (Figure 2).




Figure 2. Mechanisms of nutritionally-induced insulin resistance.

It is well-established that genetic factors play an important role in determining the predisposition to overweight,^{85,166} and in determining body fat distribution and insulin resistance.^{85,167} Nevertheless, the development of central obesity and related insulin resistance in susceptible individuals generally requires interaction with an environmental precipitant in the form of overnutrition. There is much evidence to suggest that overnutrition may be caused by a Western type of diet, rich in total and saturated fat and rapidly digestible carbohydrates, and by a lack of physical activity. The requirement of an environmental precipitant, even in the presence of a very strong genetic predisposition to develop obesity, insulin resistance, and NIDDM, is well-illustrated by studies on Australian aboriginals, and on Pima Indians, who develop these metabolic disorders at much higher rates after adopting a Western type of lifestyle.^{144,168}

Dietary fat, obesity, and insulin resistance

According to a well-accepted theoretical model by Flatt,¹⁶⁹ supported by detailed metabolic and biochemical studies, energy balance under unrestricted food intake is achieved by a separate regulation of carbohydrate, fat, and protein balances.^{166,169-172} To cover minimal carbohydrate requirements for energy metabolism, and because body stores of carbohydrates are very limited, a higher energy intake

is required on a high-fat diet than on a low-fat diet. It has been shown that a positive fat balance does not enhance fat oxidation acutely^{173,174} but rather through a slower, progressive increase in fat stores. Fat oxidation then is raised due to the increasing availability of free fatty acids in blood plasma, the increase in intracellular fat stores, and the development of insulin resistance. The development of insulin resistance thus may be regarded as a compensatory mechanism by which fat oxidation is increased to an amount commensurate with dietary fat intake, and by which the gain in body weight eventually will flatten off to reach a stable body mass.^{169,175,176} Thus, a primary dietary factor leading to obesity and insulin resistance would be a high percentage of energy from fat. The latter is confirmed by many,¹⁷⁷⁻¹⁸² though not all,^{183,184} clinical studies in which experimental high-fat diets had adverse effects on glucose and insulin metabolism. Cross-sectional epidemiologic studies on free-living populations^{176,185-191} also indicate that a diet with a high percentage of energy from fat favors the development of overweight. Similar cross-sectional studies have shown an association between a high intake of total or saturated fat and reduced insulin sensitivity¹⁹²⁻¹⁹⁵ or glucose intolerance.¹⁹⁶⁻¹⁹⁸

While high total fat intake generally has been found to be associated with a reduction in insulin sensitivity, it appears that this effect may be modified by the type of fat consumed. In rats fed high-fat diets, the development of insulin resistance was prevented by increasing the ratio of poly-unsaturated to saturated fatty acids (P/S ratio),¹⁹⁹ or by increasing the amount of long-chain n-3 fatty acids from fish oil.^{200,201} Similarly, insulin sensitivity correlated positively with the P/S ratio of phospholipids in serum,²⁰² muscle cells,²⁰³ and red blood-cell membranes²⁰⁴ of human subjects. In a study of obese pubertal children,²⁰⁵ peak insulinemia and the insulinemic area resulting from a standard oral glucose-tolerance test were associated positively with concentrations of saturated, as well as mono-unsaturated fatty acids, and negatively with concentrations of poly-unsaturated 20:5 n-3 fatty acids, measured in total plasma, and in circulating and erythrocyte phospholipids. In the Zutphen Elderly Study,²⁰⁶ insulin levels during a glucose tolerance test correlated positively with the estimated intakes of saturated, and negatively with the intakes of polyunsaturated fats. In an Italian population of almost 5,000 men and women, a negative correlation was found between consumption levels of olive oil (particularly rich in mono-unsaturated fatty acids) and fasting blood glucose concentrations.²⁰⁷ One suggested explanation for the effects due to fat composition is that high intake of polyunsaturated fatty acids tends to lower plasma levels of (VLDL-) triglycerides,^{208,209} thereby reducing fat available for storage in muscle cells. Another possible explanation, supported

by evidence from *in vitro* experiments,²⁰⁹⁻²¹¹ is that poly-unsaturated fatty acids may improve insulin action by modifying the fluidity of cellular membranes.

Postprandial glycemic and insulinemic responses to foods rich in carbohydrates

The consumption of carbohydrates induces an immediate, postprandial rise in plasma glucose concentrations, which, in turn, stimulates the release of insulin from the pancreas. The glycemic and insulinemic responses to food intake (both measured as the integrated areas under the curves of concentrations against time) are correlated highly for a given individual. The glycemic response depends on the physical form of a food, as well as on the specific types of carbohydrate it contains.^{212,213} For directly available sugars (mono- and disaccharides), the effect on the rise of blood glucose levels is a function primarily of the proportions of fructose (with low glycemic response) and of glucose (with high response). For starches, the glycemic response is determined by the morphology of the starch granules, and by the degree of branching of the polysaccharide chains. Of the two types of starch that are hydrolyzed enzymatically in the human gut – amylopectin and amylose – the former has more branched polysaccharide chains that are more accessible to the enzyme amylase and hence are digested more easily. Industrial processing (e.g., gelatinization) often aims at increasing the degree of branching of starch polysaccharide chains to make foods more palatable, and thereby also enhance glycemic and insulinemic responses. Similarly, cooking and processing with heat also may produce a greater digestibility of starches, and increase the glycemic response to foods.^{214,215}

Water-soluble dietary fibers from fruits, vegetables, or legumes – consisting of pectines and hemicelluloses – can form a gel-like matrix in the small intestine which retains nutrients and attenuates the glycemic and insulinemic responses.^{216,217} The addition of water-insoluble types of fiber from cereals to flour products such as bread or spaghetti does not have such attenuating effects on the glycemic response.²¹⁸ Nevertheless, the disruption of the physical structure of grains and cereals by grinding and milling (breaking cell walls) has been shown to increase the digestibility of cereal starches.^{219,220}

Besides determining plasma insulin responses immediately after a meal, there is some evidence that high consumption of sugars or foods with high glycemic response, in the long term, may lead to a decrease in insulin sensitivity of target tissues. In several epidemiologic studies, reduced insulin sensitivity, or the presence or development of glucose intolerance, was correlated positively with high intakes of sugar,^{206,221,222} and negatively with high intakes of dietary fiber.^{296,206,222,223} One suggested mechanism to explain these results is that, in the long

term, high postprandial insulinemic responses may down-regulate the levels of insulin receptors in target cells. Other epidemiologic studies,²²⁴ however, have provided conflicting results on the relation between insulin resistance and sugar intake. Also, from a recent review of epidemiologic and metabolic studies, there appears to be little evidence for a positive association between high-sugar diets and obesity;²²⁵ on the contrary, epidemiologic data quite consistently show an inverse correlation between obesity and percent energy derived from both simple and complex carbohydrates (both correlated with a lower percent of energy from fat) but a positive association with fat intake.

Physical activity and insulin resistance

It is well-recognized that a high level of physical activity helps to prevent the development of overweight. Even independently of influencing body mass, however, physical activity is associated positively with measures of insulin sensitivity, and negatively with the risk of developing glucose intolerance.^{206,226-231}

5. Insulin resistance, and steroid hormone profiles during puberty and adolescence

The metabolic-hormonal pattern observed in obese adult women, or in women with PCOS, shows interesting parallels with that during puberty, a period of physical development which also is marked by reduced plasma levels of SHBG,²³²⁻²³⁷ and IGFBP-1,^{232,238,239} as well as by insulin resistance.^{232,238,240-246} During early adulthood, insulin sensitivity normally returns to pre-pubertal levels. Plasma concentrations of IGF-I also show a steady increase from birth to puberty and, under the influence of increased growth hormone secretion, peak at about 13 years of age.^{239,244,247-250} After reaching their peak, IGF-I concentrations progressively decrease more than 2.5-fold towards the third decade of life, and even further in old age.²⁵¹ In studies comparing children and adults in different age groups, Holly and co-workers^{232,238} found that the pubertal decreases in both SHBG and IGFBP-1 levels were correlated with the rise in plasma insulin concentrations. Thus, as in obese adult women, the drop in plasma levels of SHBG and IGFBP-1 seems to be due to direct downregulation by insulin, which thus appears to be implicated in the physiologic mechanisms of sexual maturation and growth.^{232,238}

Like insulin, plasma concentrations of growth hormone reach peak levels during puberty (resulting in an increase also of IGF-I and IGFBP-3), and the reduction in insulin sensitivity correlates with the onset of pulsatile growth hormone secretion.²⁵² The transient decrease in insulin sensitivity during adolescence appears to be due largely to the lipolytic actions of growth hormone, as

opposed to those of insulin, leading to increased oxidation of fats instead of carbohydrates.²⁵³⁻²⁵⁹ Nevertheless, various observations suggest that the degree of insulin resistance in adolescent children is modulated also by type of diet, and by the level of physical activity. First, plasma levels of insulin in pubertal and adolescent children show associations similar to those among adults with BMI or WHR^{260,261} and, in obese children, insulin sensitivity improves after weight loss induced by calorie restriction.²⁶²⁻²⁶⁴ Second, the increase over time in the BMI of adolescent children was related inversely to level of physical activity during leisure time, and was reduced by a decrease in daily fat intake.^{265,266} Finally, as mentioned earlier,²⁰⁵ indices of insulin sensitivity were found to be associated with the fatty acid composition of total plasma lipids of obese pubertal children.

6. Insulin resistance, hyperinsulinemia, and the polycystic ovary syndrome

As mentioned in the introduction, Grattarola *et al*⁴¹ found that breast cancer patients often had ovaries with similar characteristics to those of women with PCOS: an overall increase in size, an increased amount of stromal tissue and hyperplasia of the thecal-interstitial cells where androgens are produced, and an increased number of follicles.^{46,47,133} Usually, PCOS is diagnosed only when women seek medical advice for problems of hirsutism, oligomenorrhea, or infertility. These women generally also have increased plasma concentrations of androgens, particularly testosterone and Δ^4 -androstenedione. Further, women with PCOS generally are insulin resistant, and often have increased plasma levels of LH, or a high ratio of LH relative to follicle stimulating hormone (FSH).

About 30 to 50 percent of women with clinical symptoms of PCOS are overweight, often even severely so, but PCOS also is observed in normal-weight women.^{46,47} In studies where body fat distribution has been measured, PCOS was associated with an increased WHR.²⁶⁷⁻²⁶⁹ Normal-weight PCOS patients are insulin-resistant compared with normal-weight women without clinical signs of ovarian disease.²⁷⁰⁻²⁷⁶ The combined presence of PCOS and obesity, however, is associated with a much stronger degree of insulin resistance than is observed, on average, in women with obesity alone, or in normal-weight women with PCOS.^{99,272,276} In contrast, plasma concentrations of testosterone and LH can be elevated equally in obese, or non-obese women with PCOS.^{45-47,99,270,271} Since plasma levels of SHBG and IGFBP-1 correlate negatively with fasting insulin concentrations, the levels of SHBG and IGFBP-1 generally are decreased more in obese, than in non-obese PCOS patients.^{47,98,99,101,124,273,277,278} Probably because of their

lower plasma levels of SHBG levels, and hence a further increase in free testosterone concentrations, oligomenorrhea – a major clinical manifestation of PCOS – is more frequent in the more insulin resistant women.^{100,102,273,275,277,279} In some studies,^{275,276} where PCOS was defined by the combined presence of hyperandrogenism and chronic anovulation, insulin resistance was found to be of equal magnitude in obese and non-obese patients.

Depending on the diagnostic criteria used, PCOS has been estimated to be present in about five to eight percent of premenopausal women.^{46,47} Using ultrasound (echographic) detection methods, however, the prevalence of an ovarian morphology typical of PCOS (increased ovarian volume, increased number of follicles) was shown to be as high as 20 percent among pubertal girls,²⁸⁰ as well as among adult women.^{46,47,281-283} In a considerable proportion of these relatively asymptomatic women with PCO-like ovarian morphology, plasma levels of testosterone and LH were elevated,^{47,281,283} but on average their BMI was similar to that of women with normal ovarian morphology and function.^{46,283}

PCOS has been described as the result of an 'exaggerated puberty,'²⁸⁴ as it resembles physiological adolescent anovulation in a number of ways:

- i. both are characterized by a similar 'polycystic' ovarian morphology;
- ii. the pattern of gonadotropin secretion – hyper-response of LH to exogenous luteinizing hormone-releasing hormone (LHRH), and elevated plasma levels of LH – resembles that seen in puberty; and
- iii. the 17 α -hydroxylase and 17-20-desmolase activities of the enzyme cytochrome P450c17 – an enzyme which is involved in ovarian and adrenal androgen production, and of whose susceptibility to stimulation by adreno- corticotrope hormone (ACTH) increases sharply at adrenarche, an early event in sexual maturation – are excessively accentuated.²⁸⁵

The origin of PCOS is called 'functional,' as it has no single identifiable organic cause, but appears to result from a complex dysregulation of hormonal metabolism.²⁸⁶ PCOS also is being increasingly referred to as a form of 'functional ovarian hyperandrogenism' (FOH). Most authors now concur that chronic overstimulation of the ovarian steroidogenesis by insulin and/or IGF-I is a central feature of the hormonal dysregulations leading to the syndrome (Figure 3).^{90,133,284,286-288} The hyperandrogenism associated with PCOS may be due particularly to a hyperfunction of the enzyme P450c17 in the ovarian thecal cells.²⁸⁹ The expression of this enzyme has been shown to be under the direct influence of insulin and IGF-I, in interaction with luteinizing hormone.^{290,291} The




Figure 3. Endocrine mechanisms related to the development of polycystic ovary disease (PCOD).

intra-ovarian androgen excess, possibly accentuated by a down-regulation of plasma SHBG levels due to hyperinsulinemia, is believed to be the direct cause of follicular atresia, and of hyperplasia of thecal-interstitial cells. The combination of hyperplasia of thecal-interstitial cells (which produce testosterone), and atresia of follicular cells (which convert testosterone into estrogens), may explain why overstimulation of the ovarian steroid production gradually leads to an excess of androgens, but not of estradiol. The peripheral aromatization of androgens in adipose tissue leads to increased levels of estrone, which may inhibit slightly the secretion of follicle stimulating hormone (FSH), but which may increase the secretion of LH by sensitizing the pituitary to the effects of LHRH.^{133,284,286} Further, an increase in bioactive IGF-I may augment the release of LHRH.²⁹² The rise in plasma concentrations of LH, combined with the lower levels of FSH, further aggravates the ovarian androgen excess and interstitial hyperplasia, and leads to the formation of multiple follicles arrested in their course of maturation.

The onset of clinical symptoms of PCOS is usually at about the time of menarche, seldom in adulthood. Even in women developing type II diabetes with insulin resistance and hyperinsulinemia, the onset of PCOS at adult age has not been reported.¹³³ This contrast, plus the fact that PCOS seems a continuation of a physiologic state of anovulation often seen during puberty (but which normally evolves after adolescence), have led several authors to postulate that the critical period for the development of PCOS may be during the peripubertal years^{90,133,284} (although some believe that development of the syndrome may have its earliest origins *in utero*²⁹³). The endocrine-metabolic profile during puberty – characterized by increased plasma levels of GH and total IGF-I, as well as insulin resistance and reduced levels of SHBG and IGFBP-1 – would particularly enhance the

gonadotropic action (of LH and insulin/IGF-I) on the ovary. In many girls, the PCO-like symptoms and changes in ovarian morphology disappear after adolescence, when plasma levels of insulin and of available IGF-I decrease progressively.²⁹⁴ In girls susceptible to becoming hyperandrogenic, however, a cycle of insulin resistance leading to ovarian androgen excess, thecal-interstitial hyperplasia and follicular atresia, and of androgen excess contributing to the maintenance of the insulin resistant state may persist into adulthood.^{90,133,284} Such persistence of a hyperandrogenic endocrine profile was demonstrated by a longitudinal study in Finland,²⁹⁵ showing positive correlations between serum androgen concentrations during adolescence, and 12 years later.

Studies of familial clustering of PCOS show that genetic predisposition may contribute to the development of the syndrome.²⁹⁶ In a family study by Carey *et al*,²⁹⁷ polycystic ovaries and male baldness were demonstrated to be the female and male phenotypes of a condition that seemed to be due to a single gene defect. Further work²⁹⁸ showed that the expression of PCOS/male baldness was associated significantly (odds ratio of 3.57) with a single base change in the 5' promotor region of P450c17, which would create an additional promotor site for the gene, and thereby lead to its increased expression. The single base change was excluded, however, as a primary defect responsible for the familial clustering. Dunaif²⁹⁹ proposed that two types of genetic predisposition may be involved in the pathogenesis of PCOS: a 'PCO gene' conferring a specific susceptibility to developing the reproductive abnormalities of PCOS (e.g., by hypersensitivity of ovarian androgen production to stimulation by insulin/IGF-I and LH), and a genetic predisposition to developing a minimum level of insulin resistance, needed for the PCO syndrome to be expressed.

It is still unknown why some women with PCOS become obese and, on average, more insulin resistant, whereas others stay comparatively lean. One can speculate that, due to variations in genetic susceptibility (e.g., the 'PCO-gene'; see above), different magnitudes of insulin resistance, augmented or nonaugmented by obesity, may be required for the ovarian androgen production to be overexpressed. Further, some forms of insulin resistance observed frequently in women with PCOS may be determined genetically and be relatively independent of body weight.²⁷⁶ Alternatively, on the basis of observations that plasma levels of GH were significantly lower in obese than in non-obese women with PCOS,^{300,301} Insler^{302,303} has postulated that, in obese girls or women, hyperinsulinemia may be the more predominant derangement causing hyperfunction of androgen producing enzymes in the ovary, whereas in normal-weight women the more predominant derangement would be an increased activity of GH (particularly during

puberty). Besides raising plasma and tissue concentrations of IGF-I, which directly stimulates ovarian androgen production, an increased GH activity contributes to the development of insulin resistance because of its lipolytic action. In addition, especially in interaction with androgens, GH favors the growth of lean body mass but limits the size of body fat deposits.

Given the postulated central role for hyperinsulinemia in the etiology of PCOS, it seems reasonable to suppose that lifestyle factors (diet, physical activity) that modulate plasma insulin levels also may influence the development or persistence of the syndrome. In studies with adult PCOS patients,^{92,304-307} and with obese adolescent girls,³⁰⁸ weight loss induced by a calorie-restricted diet indeed resulted in a significant fall of plasma levels of insulin, total and free testosterone, free estradiol and the ratio of LH/FSH, and in a significant rise in SHBG levels. This suggests that obesity, by adding to the relative degree of hyperinsulinemia, at least may contribute to the maintenance of the PCO syndrome. Longitudinal studies are needed, however, to investigate whether diet and physical activity, especially during peripubertal years and adolescence, may or may not have long-lasting effects on the development of a hyperandrogenic endocrine profile persisting into adulthood, and whether such effects are similar in girls who do, or who do not, develop obesity.

7. Discussion and synthesis

This paper reviews the possible role of insulin in the relation between nutrition-related lifestyle factors and the development of a hormonal-metabolic profile associated with an increased risk of breast cancer in women.

The current view of carcinogenesis is that it results from an imbalance between proliferation, differentiation, and programmed death (apoptosis) of cells, with clonal selection and expansion of colonies of preneoplastic and, ultimately, neoplastic cells that have become progressively independent of normal growth control.³⁰⁹⁻³¹³ A high rate of cell proliferation is believed to be a key factor in increasing the risk of cancer development, as cell division is necessary for the accumulation of genetic mutations related to neoplastic transformation and clonal selection of transformed cells, as well as for the growth of established tumors. A hormonal environment that inhibits full differentiation of cells may increase breast cancer risk by augmenting the pool of cells remaining responsive to proliferative stimuli.

Steroid hormones, and peptide growth factors have been shown to be crucial in the paracrine and endocrine intercellular communication processes controlling the differentiation³¹⁴ and proliferation of human breast cells. Regarding the effect of steroid hormones on the proliferation of epithelial cells, evidence suggests that

progestogens may be the predominant mitogen for normal epithelium, and that estrogens (especially 17- β -estradiol) may potentiate this effect via induction of increased progesterone receptor numbers. On the other hand, in neoplastic epithelium, estrogens appear to be the more predominant mitogen.^{311,312} Based on clinical observations that proliferation rates of normal breast epithelium are highest during the luteal phase of the menstrual cycle, Henderson *et al*^{313,315} have postulated that tumor development is promoted most strongly at a simultaneous exposure to estrogens plus progesterone. Pike *et al*^{37,316,317} have argued that this 'estrogen-plus-progestogen' hypothesis also might explain the finding that, in premenopausal women, extreme obesity seems to be associated with a reduction in breast cancer risk, even though obesity increases the risk of endometrial cancer (which is believed to be due to an exposure to estrogens 'unopposed' by progestogens).³¹⁸

As for peptide hormones and growth factors, numerous studies during the last five to 10 years (reviewed in Macaulay³¹⁹ and Yee *et al*³²⁰) have shown that IGF-I, especially in interaction with estrogens, is a potent inducer of mitosis in breast cancer cells. During the 1970s, experiments with animals and breast cancer cell lines, reviewed by Hilf³²¹ had demonstrated already that insulin also might promote the growth of breast tumors. The high degree of homology between insulin and IGF-I (at that time known as somatomedin C) and between their receptors, however, had not yet been established, and effects on tumor growth originally attributed to the binding of insulin to its specific receptor actually may have been mediated by an increased availability of IGF-I (in animal experiments), or by the cross-binding of insulin (often used at quite high concentrations in studies with cell cultures) to the IGF-I receptor.

In line with the experimental and clinical evidence relating estrogens – probably in interaction with progesterone and IGF-I – to increased growth rates of breast tumors, epidemiologic observations (reviewed in section 2) show that women developing breast cancer tend to have a hyperandrogenic hormonal profile characterized by higher plasma levels of testosterone, lower plasma levels of SHBG and, hence, higher levels of free estradiol. As described in sections 3 and 6, hyperinsulinemia appears to play a key role in the development of this endocrine profile, either in association with simple obesity, or as a form of functional ovarian hyperandrogenism. One therefore may hypothesize that chronic elevation of plasma insulin levels increases the risk of developing breast cancer. This 'insulin hypothesis' fits with the positive association between obesity and breast cancer observed in postmenopausal women, as recently discussed also by Stoll³²² and with results from ecologic studies showing that breast cancer incidence rates are high in

countries with a more sedentary lifestyle, and with a diet rich in saturated (animal) fats and refined carbohydrates. The hypothesis has received its first direct support from a case-control study by Bruning *et al*,³²³ in which both pre- and postmenopausal breast cancer patients were shown to have increased serum levels of C-peptide, a marker of pancreatic insulin secretion.

Although the above considerations lend considerable support to the insulin hypothesis, there are also several complicating observations.

First, in apparent contrast with the insulin hypothesis, case-control and cohort studies have not provided consistent evidence for an association between breast cancer risk and diabetes.³²⁴⁻³³⁰ The interpretation of these studies, however, is problematic. In some women with non-insulin dependent (Type 2) diabetes, residual pancreatic β -cell activity can be sufficient for them to have fasting plasma insulin levels as high as insulin-resistant, but glucose-tolerant, women; and, it has been shown that these women may have a similar steroid hormonal profile, marked by low plasma SHBG levels and high free steroid concentrations, to non-diabetic women with similar BMI.³³¹ Women with insulin treated (Type 1 or Type 2) diabetes, however, in whom the endogenous production of insulin is reduced substantially or even practically nonexistent, can have a steroid hormone profile either very similar to, or different from, that of insulin resistant but non-diabetic women. For instance, in a study of postmenopausal women with insulin-treated Type 2 diabetes, total levels of estrogens and androgens were increased, but a simultaneous rise in SHBG kept the free fractions of these sex hormones within normal limits.³³² In another study of adult women with IDDM, by Djursing *et al*,³³³ diabetics with regular menstrual cycles had significantly higher serum concentrations of androgens than controls with a normal cycle, whereas amenorrheic patients had lower serum concentrations of SHBG, LH, total androgens, and free and total estrogens. Prelevic *et al*³³⁴ further distinguished between women with IDDM who did (C-peptide positives), or did not (C-peptide negatives) have some residual pancreatic insulin secretion. All C-peptide negatives with amenorrhea had decreased LH levels, a low LH/FSH ratio, and low levels of testosterone, as observed in the study by Djursin *et al*.³³³ On the contrary, C-peptide positives with amenorrhea had the classic hormonal profile of PCOS (increased serum testosterone and LH/FSH ratio, decreased SHBG), and was associated with a history of oligomenorrhea and excess weight before the onset of diabetes. In view of this heterogeneity of steroid hormonal profiles in diabetic women, and in the absence of further information about the type and severity of diabetes, it seems difficult to predict whether diabetes will generally augment or decrease breast cancer risk.

In several studies, acromegaly – a pathologic condition characterized not only by severe insulin resistance but also by high plasma levels of IGF-I – was shown to be associated with increased occurrence of cancer.³³⁵⁻³³⁹ Although most of these studies were too small to estimate the relative incidence rates of cancer at specific sites, one study showed an excess of breast cancer among females. A number of additional studies provided evidence for an increased occurrence of colon cancer, and more particularly of colonic polyps, a relatively frequent preneoplastic lesion.³⁴⁰⁻³⁴⁶

Second, in contrast to international correlation studies, individual-level studies, especially those of a prospective cohort design, have failed to show a clear association between high intake levels of (saturated) fat and breast cancer risk. This seems surprising, since physiologic studies of energy metabolism and cross-sectional epidemiologic analyses indicate that a high intake of (especially saturated) fat is one of the dietary factors associated most clearly with development of obesity and insulin resistance (see section 4). The lack of association between level of fat intake and breast cancer risk in individual-level studies has been attributed to methodologic problems, such as the difficulty of measuring quite small differences in dietary intake patterns of individuals all living in the same, geographically limited area.³⁴⁷ A complimentary explanation could be that, in populations with a predominantly Western lifestyle and generalized conditions of overnutrition, in most individuals the development of chronic hyperinsulinemia and related endocrine dysregulations may depend more on genetic factors determining their physiologic response to overnutrition, than on comparatively small differences in their habitual diets. In addition, the development of breast cancer through nutritionally induced endocrine dysregulations may be determined predominantly during pre- or peripubertal years, while dietary habits during middle life may no longer reflect those of childhood.^{18,25} In this respect, breast cancer seems to differ from colon cancer – although international variations in incidence rates very much parallel those of breast cancer and the development of colon cancer has also recently been postulated to be related to chronic hyperinsulinemia^{348,349} – as colon cancer risk in prospective studies appears associated more clearly to aspects of habitual diet (*i.e.*, measured relatively shortly before the diagnosis of the disease).

A third confusing observation is the lack of positive association between obesity and the risk of breast cancer before menopause. This is particularly puzzling, as several case-control studies have shown that not only post-, but also premenopausal women who develop breast cancer have a more hyperandrogenic endocrine profile⁴⁹⁻⁵¹ and higher plasma insulin levels³²³ than disease-free controls. Further, cross-sectional epidemiologic studies of disease-

free, premenopausal women do show a positive association between BMI and insulin resistance.^{61,63,74,78} These puzzling observations lead to the following questions:

- i. How can obesity be unassociated, or possibly even associated negatively, with risk of breast cancer in premenopausal women? and
- ii. How can a hyperandrogenic endocrine profile with increased plasma levels of free estrogens, and with some degree of insulin resistance, develop in the absence of obesity?

An answer to the first question may be that, especially in hyperandrogenic women, overweight is associated with more severe degrees of insulin resistance and, before menopause, with oligomenorrhea. Henderson^{313,315} and Pike *et al*^{37,316,317} have postulated that women with menstrual irregularities spend a smaller cumulative lifetime in the luteal phase of their cycles, and therefore have a lower lifetime exposure to the combination of estradiol (produced during both the follicular and luteal phases of the cycle) and progesterone (produced during the luteal phase only). Moreover, during anovulatory cycles, the maturation of the dominant follicle is incomplete and the ovarian production of progesterone is subnormal. Thus, assuming that the development of breast tumors is promoted most strongly by a combined exposure to estradiol and progesterone, hyperandrogenic women who are also obese may have a relative protection against breast cancer before menopause. On the other hand, following the same estrogen-plus-progestogen hypothesis, hyperandrogenic women with regular menstrual cycles (in general less obese, and less severely insulin resistant) would be expected to have a relative increase in breast cancer risk, due to greater cumulative exposure to (free) estrogens plus normal levels of progesterone.

As for the second question, an answer may be found by further investigating the mechanisms of development of FOH (see section 6). About 30 to 50 percent of women seeking medical advice for symptoms typical of PCOS are obese, the others having a normal body weight. Moreover, studies based on echographic screening have shown that the prevalence of milder, subclinical forms of PCOS may be considerably higher than that of the clinically manifest syndrome,^{45,281-283} and that, on average, women with milder forms of PCOS have a BMI similar to that of women with normal ovarian appearance and function.^{282,283} It may be that particularly the latter category of relatively asymptomatic women has increased risk of breast cancer before menopause.

In summary, I surmise that the risk of developing breast cancer generally is increased in hyperandrogenic women, because of a higher exposure to estrogens unbound to SHBG. Chronic hyperinsulinemia appears

to be a central physiologic link between dietary composition, nutritional status (size of triglyceride stores in adipose tissue and skeletal muscle; relative contributions of the oxidation of fats and carbohydrates to energy metabolism), and development of the hyperandrogenic endocrine profile. In young women, hyperandrogenism may develop more often as a form of FOH, due to a genetically determined sensitivity of the ovaries to the stimulating effects of insulin/IGF-I and LH on androgen production, or because of a genetic predisposition to develop a higher-than-average degree of insulin resistance (with or without obesity) at a young age. In these women, hyperandrogenism may frequently also come to expression in the form of an ovarian morphology typical of PCOS. With age, high plasma androgen levels are probably associated increasingly with a high BMI, due to the rising prevalence of obesity in women less strongly predisposed to develop obesity and hyperandrogenism earlier in life.

Following the estrogen-plus-progestogen hypothesis, the increase in breast cancer risk due to a rise in free estrogens is amplified by the presence of progesterone. Therefore, premenopausal women with more severe forms of insulin resistance and hyperandrogenism, and with chronic anovulation, may turn out to have a relative protection against breast cancer compared with the general population (at least before menopause). The risk of breast cancer is expected to be increased, however, in hyperandrogenic women who maintain regular menstrual cycles. After menopause, when the ovarian production of progesterone and estradiol has ceased, and when breast cancer risk in all women is determined more predominantly by available levels of estrogens, formed in adipose tissue by the aromatization of androgens,^{350,351} breast cancer risk would be expected to be increased in all hyperandrogenic women, especially when they are obese, and even when they are more severely insulin resistant.

Further prospective cohort studies are needed to confirm further the presence of the hypothesized relations between breast cancer risk and plasma hormone levels (testosterone, SHBG, total and free estradiol, insulin), not only after, but also before menopause. Additionally, a case-control study should be undertaken to investigate whether the incidence of breast cancer before menopause is increased in women with normal cycles and subclinical forms of PCOS (e.g., diagnosed by echographic examination of the ovaries, combined with measurements of plasma hormone levels).

Thus far, only two epidemiologic studies appear to have been conducted to investigate the possible relation between PCOS and increased breast cancer risk. In a cohort study on 1,270 women with previously diagnosed PCOS,³⁵² there were three, four and five cases with pre-, peri-, and postmenopausal breast cancer, respectively. The

five postmenopausal cases were in significant excess compared with the number of (1.4) cases expected, as computed from average breast cancer rates in the corresponding general population. In the second, a multi-center study of 4,730 cases and 4,688 geographically matched controls,³⁵³ self-reports of PCOS (diagnosed previously by a physician) were against all expectations associated negatively with premenopausal breast cancer risk. Both studies focused on women with rather severe manifestations of the PCO syndrome, however, who sought medical advice because of complaints of hirsutism, menstrual irregularity, or infertility. The negative association between PCOS and breast cancer risk observed in the second study therefore may be explained by reduced exposure to progesterone, because of chronic anovulation, and thus would not contradict the hypothesis of an excess breast cancer risk in hyperandrogenic, but normally cycling premenopausal women.

The observation of an increased prevalence of milder forms of PCOS in premenopausal breast cancer patients may give additional support to the idea that breast cancer risk is determined largely by metabolic dysregulation occurring early in life. Further studies then might evaluate whether the development of PCOS is related to early sexual maturation (as has been suggested recently by some studies on sexually precocious girls^{354,355}) a well-established risk factor for breast cancer. Finally, it would then be of interest to investigate whether puberty is indeed the critical period for development of the syndrome (also its subclinical forms), and to evaluate to what extent diet and physical activity during peripubertal years and adolescence, by influencing the degree of insulin resistance, may or may not be determinants for the development of a PCO-like hyperandrogenic profile persisting into adulthood.

Acknowledgements — I thank Dr Franco Berrino, Istituto Nazionale dei Tumori, Milano (Italy) and Dr Elio Riboli, IARC, Lyon, for helpful comments, and the three anonymous reviewers for their constructive criticisms.

References

1. Parkin DM, Pisani P, Ferlay J. Estimates of the worldwide incidence of eighteen major cancers in 1985. *Int J Cancer* 1993; 54: 594-606.
2. Dunn JE. Breast cancer among American Japanese in the San Francisco Bay area. *Natl Cancer Inst Monogr* 1977; 47: 157-60.
3. Trichopoulos D, Yen S, Brown J, Cole P, MacMahon B. The effect of Westernization on urine estrogens, frequency of ovulation, and breast cancer risk. (A study of ethnic Chinese in the Orient and the USA.) *Cancer* 1984; 53: 187-92.
4. Kato I, Tominaga S, Kuroishi T. Relationship between westernization of dietary habits and mortality from breast and ovarian cancers in Japan. *Gann* 1987; 78: 349-57.

5. Bjarnason O, Day NE, Snaedal G, Tulinius H. The effect of year of birth on the breast cancer age curve in Iceland. *Int J Cancer* 1974; 13: 689-96.
6. Wakai K, Suzuki S, Ohno Y, Kawamura T, Tamakoshi A, Aoki R. Epidemiology of breast cancer in Japan. *Int J Epidemiol* 1995; 24: 285-91.
7. Armstrong B, Doll R. Environmental factors and cancer incidence and mortality in different countries, with special reference to dietary practices. *Int J Cancer* 1975; 15: 617-31.
8. Hems G. The contributions of diet and childbearing to breast-cancer rates. *Br J Cancer* 1978; 37: 974-82.
9. Eveleth PB, Tanner JM. *Worldwide Variation In Human Growth*. Cambridge, UK: Cambridge University Press, 1976.
10. Micozzi MS. Nutrition, body size, and breast cancer. *Year-book Phys Anthropol* 1985; 28: 175-206.
11. Albanes D, Taylor P. International differences in body height and weight and their relationship to cancer incidence. *Nutr Cancer* 1990; 14: 69-77.
12. Tarone RE, Chu KC. Implications of birth cohort patterns in interpreting trends in breast cancer rates. *JNCI* 1992; 84: 1402-10.
13. Frisch RE, Wyshak G, Albright NL, et al. Lower lifetime occurrence of breast cancer and cancers of the reproductive system among former college athletes. *Am J Clin Nutr* 1987; 45: 328-35.
14. Bernstein L, Henderson BE, Hanish R, Sullivan-Halley J, Ross RK. Physical exercise and reduced breast cancer risk in young women. *JNCI* 1994; 86: 1403-8.
15. Mittendorf R, Longnecker MP, Newcomer PA, et al. Strenuous physical activity in young adulthood and risk of breast cancer (United States). *Cancer Causes Control* 1995; 6: 347-53.
16. Welsh CW. Dietary fat, calories, and mammary gland tumorigenesis. In: Jacobs MM, ed. *Exercise, Calories, Fat and Cancer*. New York, NY (USA): Plenum Press 1992: 203-22.
17. Kelsey JL, Gammon MD, John EM. Reproductive factors and breast cancer. *Epidemiol Rev* 1993; 15: 36-47.
18. Hunter DJ, Willett WC. Diet, body size, and breast cancer. *Epidemiol Rev* 1993; 15: 110-32.
19. Folsom AR, Kaye SA, Prineas RJ, Potter JD, Gapstur SM, Wallace RB. Increased incidence of carcinoma of the breast associated with abdominal adiposity in postmenopausal women. *Am J Epidemiol* 1990; 131: 794-803.
20. Bruning PF, Bonfrère JMG, Hart AAM, et al. Body measurements, estrogen availability and the risk of human breast cancer: a case-control study. *Int J Cancer* 1992; 51: 14-9.
21. Schapira DV, Kumar NB, Lyman GH, Cox CE. Abdominal obesity and breast cancer risk. *Ann Intern Med* 1990; 112: 182-6.
22. Sellers TA, Kushi LH, Potter JD, et al. Effect of family history, body fat distribution and reproductive factors on the risk of postmenopausal breast cancer. *N Engl J Med* 1992; 326: 1323-9.
23. den Tonkelaar I, Seidell JC, Collette H. Body fat distribution in relation to breast cancer in women participating in the DOM-project. *Breast Cancer Res Treat* 1995; 34: 55-61.
24. Ursin G, Longnecker MP, Haile RW, Greenland S. A meta-analysis of body mass index and risk of premenopausal breast cancer. *Epidemiology* 1995; 6: 137-41.
25. Byers T. Nutritional risk factors for breast cancer. *Cancer* 1994; 74: 288-95.
26. Howe GR, Hirohata T, Bishop TG, et al. Dietary factors and risk of breast cancer: combined analysis of 12 case-control studies. *JNCI* 1990; 82: 561-9.
27. Boyd NF, Noffel MM, Lockwood GA, Trichler DL. A meta-analysis of dietary fat and breast cancer risk. *Br J Cancer* 1993; 68: 627-36.
28. MacKenzie I. The production of mammary cancer in rats using oestrogens. *Br J Cancer* 1955; 9: 284-99.
29. Staszewski J. Age at menarche and breast cancer. *JNCI* 1971; 47: 935-40.
30. Trichopoulos D, MacMahon B, Cole P. Menopause and breast cancer risk. *JNCI* 1972; 48: 605-13.
31. Clemmesen J. *Statistical Studies in Malignant Neoplasms. Volume 1*. Copenhagen, Denmark: Munksgaard, 1965.
32. Lilienfeld AM. The relationship of cancer of the female breast to artificial menopause and marital status. *Cancer* 1956; 9: 927-34.
33. Zumoff B. Hormonal profiles in women with breast cancer. *Anticancer Res* 1988; 8: 627-36.
34. Siiteri PK, Hammond GL, Nisker JA. Increased availability of serum estrogens in breast cancer: a new hypothesis. In: Pike MC, Siiteri PK, Welsch CW, eds. *Hormones and Breast Cancer*. Cold Spring Harbor, NY (USA): Cold Spring Harbor Laboratory, 1981; Banbury Report No. 8: 87-106.
35. Partridge WM. Serum availability of sex steroid hormones. *Clin Endocrinol Metab* 1986; 15: 259-78.
36. Mendel CM. The free hormone hypothesis: a physiologically based mathematical model. *Endocr Rev* 1989; 10: 232-74.
37. Key TJA, Pike MC. The role of oestrogens and progestagens in the epidemiology and prevention of breast cancer. *Eur J Cancer Clin Oncol* 1988; 24: 29-43.
38. Bernstein L, Ross RK. Endogenous hormones and breast cancer risk. *Epidemiol Rev* 1993; 15: 48-65.
39. Toniolo PG, Levitz M, Zeleniuch-Jacquotte A, et al. A prospective study of endogenous estrogens and breast cancer in postmenopausal women. *JNCI* 1995; 87: 190-7.
40. Dorgan JF, Longcope C, Stephenson HE, et al. Relation of prediagnostic serum estrogen and androgen levels to breast cancer risk. *Cancer Epidemiol Biomark Prev* 1996; 5: 533-9.
41. Grattarola R. Androgens in breast cancer. I. Atypical endometrial hyperplasia and breast cancer in married premenopausal women. *Am J Obstet Gynaecol* 1973; 116: 324-8.
42. Grattarola R, Secreto G, Recchione C, Castellini W. Androgens and breast cancer. II. Endometrial adenocarcinoma and breast cancer in married postmenopausal women. *Am J Obstet Gynaecol* 1974; 118: 173-8.
43. Sommers SC, Theloh HA. Ovarian stromal hyperplasia in breast cancer. *Arch Pathol* 1952; 53: 160-7.
44. Sommers SC. Endocrine abnormalities in women with breast cancer. *Lab Invest* 1955; 4: 160-74.
45. Adams J, Polson DW, Franks S. Prevalence of polycystic ovaries in women with anovulation and idiopathic hirsutism. *Br Med J* 1986; 293: 355-9.
46. Conway GS, Honour JW, Jacobs HS. Heterogeneity of the polycystic ovary syndrome: clinical, endocrine, and ultrasound features in 556 patients. *Clin Endocrinol* 1989; 30: 459-70.
47. Franks S. Polycystic ovary syndrome: a changing perspective. *Clinical Endocrinol* 1989; 31: 87-120.
48. Secreto G, Fariselli G, Bandieramonte G, Recchione C, Dati V, DiPietro S. Androgen excretion in women with a family history of breast cancer or with epithelial hyperplasia or cancer of the breast. *Eur J Cancer Clin Oncol* 1983; 19: 5-10.
49. Secreto G, Toniolo P, Berrino F, et al. Increased androgenic activity and breast cancer risk in premenopausal women. *Cancer Res* 1984; 44: 5902-5.
50. Secreto G, Recchione C, Fariselli G, DiPietro S. High testosterone and low progesterone levels in premenopausal

- patients with hyperplasia and cancer of the breast. *Cancer Res* 1984; 44: 841-4.
51. Secreto G, Toniolo P, Pisani P, *et al.* Androgens and breast cancer in premenopausal women. *Cancer Res* 1989; 49: 471-6.
 52. Secreto G, Recchione C, Cavalleri M, Miraglia M, Dati V. Circulating levels of testosterone, 17 β -oestradiol, luteinizing hormone and prolactin in postmenopausal breast cancer patients. *Br J Cancer* 1983; 47: 269-75.
 53. Secreto G, Toniolo P, Berrino F, *et al.* Serum and urinary androgens and risk of breast in postmenopausal women. *Cancer Res* 1991; 51: 2572-6.
 54. Malarkey WB, Schroeder LL, Stevens VC, James AG, Lanese RR. Twenty-four-hour preoperative endocrine profiles in women with benign and malignant breast disease. *Cancer Res* 1977; 37: 4655-9.
 55. McFadyen IJ, Forrest APM, Prescott RJ, *et al.* Circulating hormone levels in women with breast cancer. *Lancet* 1976; i: 1100-2.
 56. Adami HO, Johansson EDB, Vegelius J, Victor A. Serum concentrations of estrone, androstenedione, testosterone and sex-hormone-binding globulin in postmenopausal women with breast cancer and in age-matched controls. *Uppsala J Med Sci* 1979; 84: 259-74.
 57. Hill P, Garbaczewski L, Kasumi F. Plasma testosterone and breast cancer. *Eur J Cancer Clin Oncol* 1985; 21: 1256-66.
 58. Wysowski DK, Comstock GW, Helsing KJ, Lau HL. Sex hormone levels in serum in relation to the development of breast cancer. *Am J Epidemiol* 1987; 125: 791-9.
 59. Garland CF, Friedlander NJ, Barrett-Connor E, Khaw KT. Sex hormones and postmenopausal breast cancer: a prospective study in an adult community. *Am J Epidemiol* 1992; 135: 1220-30.
 60. Berrino F, Muti P, Micheli A, *et al.* Serum sex hormone levels after menopause and subsequent breast cancer. *JNCI* 1996; 88: 291-6.
 61. Evans D, Hoffmann RG, Kalkhoff RK, Kissebah AH. Relationship of androgenic activity to body fat topography, fat cell morphology, and metabolite aberrations in premenopausal women. *J Clin Endocrinol Metab* 1983; 57: 304-10.
 62. Seidell JC, Cigolini M, Deurenberg P, Oosterlee A, Doornbos G. Fat distribution, androgens, and metabolism in nonobese women. *Am J Clin Nutr* 1989; 50: 269-73.
 63. Seidell JC, Cigolini M, Chazewska J, *et al.* Androgenicity in relation to body fat distribution and metabolism in 38-year old women - the European fat distribution study. *J Clin Epidemiol* 1990; 43: 21-34.
 64. Krotkiewski M, Seidell JC, Björntorp P. Glucose tolerance and hyperinsulinemia in obese women: role of adipose tissue distribution, muscle fiber characteristics and androgens. *J Intern Med* 1990; 228: 385-92.
 65. Weaver JU, Holly JMP, Kopelman PG, *et al.* Decreased sex hormone binding globulin (SHBG) and insulin-like growth factor binding protein (IGFBP-1) in extreme obesity. *Clin Endocrinol* 1990; 33: 415-22.
 66. Kaye SA, Folsom AR, Soler JT, Prineas RJ, Potter JD. Associations of body mass and fat distribution with sex hormone concentrations in postmenopausal women. *Int J Epidemiol* 1991; 20: 151-6.
 67. Peiris AN, Sothman MS, Aiman MS, Kissebah AH. The relationship of insulin to sex hormone-binding globulin: role of adiposity. *Fertil Steril* 1989; 52: 69-72.
 68. Soler JT, Folsom AR, Kaye SA, Prineas RJ. Associations of abdominal adiposity, fasting insulin, sex hormone binding globulin, and estrone with lipids and lipoproteins in post-menopausal women. *Atherosclerosis* 1989; 79: 21-7.
 69. Haffner SM, Dunn JF, Katz MS. Relationship of sex-hormone-binding globulin to lipid, lipoprotein, glucose, and insulin concentrations in postmenopausal women. *Metabolism* 1992; 41: 278-84.
 70. Preziosi P, Barrett-Connor E, Papoz L, *et al.* Interrelation between plasma sex hormone-binding globulin and plasma insulin in healthy adult women: the Telecom study. *J Clin Endocrinol Metab* 1993; 76: 283-7.
 71. Anderson DC. Sex-hormone-binding globulin. *Clin Endocrinol* 1974; 3: 69-96.
 72. Rosner W. The functions of corticosteroid-binding globulin and sex-hormone-binding globulin: recent advances. *Endocr Rev* 1990; 11: 80-91.
 73. Reaven GM. Role of insulin resistance in human disease. *Diabetes* 1988; 37: 1595-607.
 74. Peiris AN, Sothman MS, Hennes MI, *et al.* Relative contribution of obesity and body fat distribution to alterations in glucose insulin homeostasis: predictive values of selected indices in premenopausal women. *Am J Clin Nutr* 1989; 49: 758-64.
 75. Kirschner MA, Samojlik E, Drejka M, Szmal E, Schneider G, Ertel N. Androgen-estrogen metabolism in women with upper body versus lower body obesity. *J Clin Endocrinol Metab* 1990; 70: 473: 9.
 76. Kissebah AH, Vydelingum N, Murray, *et al.* Relation of body fat to metabolic complications of obesity. *J Clin Endocrinol Metab* 1982; 54: 254-60.
 77. Evans DJ, Hoffman RG, Kalkhoff RK, Kissebah AH. Relationship of body fat topography to insulin sensitivity and metabolic profiles in premenopausal women. *Metabolism* 1984; 33: 68-75.
 78. Amemiya T, Kawahara R, Yoshino M, Komori K, Hirata Y. Population study on the adipose tissue distribution in Japanese women born in 1948. *Diabetes Res Clin Prac* 1990; 10: S71-6.
 79. Evans DJ, Barth JH, Burke CW. Body fat topography in women with androgen excess. *Int J Obes* 1988; 12: 157-62.
 80. Schriock ED, Buffington CK, Hubert GD, *et al.* Divergent correlations of circulating dehydroepiandrosterone sulfate and testosterone with insulin levels and insulin receptor binding. *J Clin Endocrinol Metab* 1988; 66: 1329-31.
 81. Buffington CK, Givens JR, Kitabchi AE. Opposing actions of dehydroepiandrosterone and testosterone on insulin sensitivity. *Diabetes* 1991; 40: 693-700.
 82. Poretsky L. On the paradox of insulin-induced hyperandrogenism in insulin-resistant states. *Endocr Rev* 1991; 12: 3-13.
 83. Kissebah AH, Peiris A, Evans DJ. Mechanisms associating body fat distribution with the abnormal metabolic profile in obesity. *Rec Adv Obes Res* 1987; 5: 54-9.
 84. Leibel RL, Edens NK, Fried SK. Physiologic basis for the control of body fat distribution in humans. *Annu Rev Nutr* 1989; 9: 417-43.
 85. Bouchard C, Desprès JP, Mauriège P. Genetic and non-genetic determinants of regional fat distribution. *Endocr Rev* 1993; 14: 72-93.
 86. Björntorp P. Insulin resistance and physical exercise in obesity. In: Oomura Y *et al.*, eds. *Progress in Obesity Research*. Washington, DC: John Libbey & Company 1990: 259-68.
 87. Björntorp P. Fatty acids, hyperinsulinaemia, and insulin resistance: which comes first? *Curr Opin Lipidol* 1994; 5: 166-74.
 88. Hernandez ER, Resnick CE, Holtzclaw D, Payne DW, Adashi EY. Insulin as a regulator of androgen biosynthesis by cultured rat ovarian cells: cellular and pharmacological hormonal actions. *Endocrinology* 1988; 122: 2034-43.

89. Adashi EY, Resnick CE, D'Ercole AJ, Svoboda ME, van Wyk JJ. Insulin-like growth factors as intraovarian regulators of granulosa cell growth and function. *Endocrinol Rev* 1985; 6: 400-20.
90. Poretsky L, Kalin MF. The gonadotropic function of insulin. *Endocrinol Rev* 1987; 8: 132-41.
91. Nestler JE, Strauss JF. Insulin as an effector of human ovarian and adrenal steroid metabolism. *Endocrinol Metab Clin North Am* 1991; 20: 807-23.
92. Kiddy DS, Hamilton-Fairley D, Seppälä M, et al. Diet-induced changes in sex hormone binding globulin and free testosterone in women with normal or polycystic ovaries: correlation with serum insulin and insulin-like growth factor-I. *Clin Endocrinol* 1989; 31: 757-63.
93. von Scholtz B, Carlström K. On the regulation of sex-hormone-binding globulin – a challenge of an old dogma and outlines of an alternative mechanism. *J Steroid Biochem* 1989; 2: 327-34.
94. Toscano V, Balducci P, Bianchi P, Mangiantini A, Sciarra F. Steroidal and non-steroidal factors in plasma sex hormone binding globulin regulation. *J Steroid Biochem Mol Biol* 1992; 43: 431-7.
95. Plymate SR, Matej LA, Jones RE, Friedl KE. Inhibition of sex hormone-binding globulin production in the human hepatoma (Hep G2) cell line by insulin and prolactin. *J Clin Endocrinol Metab* 1988; 67: 460-4.
96. Plymate SR, Hoop RC, Jones RE, Matej LA. Regulation of sex hormone-binding globulin production by growth factors. *Metabolism* 1990; 39: 967-70.
97. Singh A, Hamilton-Fairley D, Koistinen R, et al. Effect of insulin-like growth factor-type I (IGF-I) and insulin on the secretion of sex-hormone binding globulin and IGF-I binding protein (IBP-I) by human hepatoma cells. *J Endocrinol* 1990; 124: R1-3.
98. Holte J, Bergh T, Gennarelli G, Wide L. The independent effects of polycystic ovary syndrome and obesity on serum concentrations of gonadotrophins and sex steroids in premenopausal women. *Clin Endocrinol* 1994; 41: 473-81.
99. Rajkhowa M, Bicknell J, Jones M, Clayton RN. Insulin sensitivity in women with polycystic ovary syndrome: relationship to hyperandrogenemia. *Fertil Steril* 1994; 61: 605-12.
100. Robinson S, Kiddy D, Gelding SV, et al. The relationship of insulin insensitivity to menstrual pattern in women with hyperandrogenism and polycystic ovaries. *Clin Endocrinol* 1993; 39: 351-5.
101. Franks S, Kiddy DS, Hamilton-Fairley D, Bush A, Sharp PS, Reed MJ. The role of nutrition and insulin in the regulation of sex hormone binding globulin. *J Steroid Biochem Mol Biol* 1991; 39: 835-8.
102. Sharp PS, Kiddy DS, Reed MJ, Anyaoku V, Johnston DG, Franks S. Correlation of plasma insulin and insulin-like growth factor-I with indices of androgen transport and metabolism in women with polycystic ovary syndrome. *Clin Endocrinol* 1991; 35: 253-7.
103. Van den Brande JL. Structure of the human insulin-like growth factors: relationship to function. In: Schofield PN. *The Insulin-like Growth Factors; Structure and Biological Functions*. Oxford, UK: Oxford University Press, 1992: 12-44.
104. Holly JMP, Wass JAH. Insulin-like growth factors: autocrine, paracrine or endocrine? New perspectives of the somatomedin hypothesis in the light of recent developments. *J Endocrinol* 1989; 122: 611-8.
105. Clemmons DR. Structural and functional analysis of insulin-like growth factors. *Br Med Bull* 1989; 45: 465-80.
106. Humbel RE. Insulin-like growth factors I and II. *Eur J Biochem* 1990; 190: 445-62.
107. Bang P, Hall K. Insulin-like growth factors as endocrine and paracrine hormones. In: Schofield PN. *The Insulin-like Growth Factors; Structure and Biological Functions*. Oxford, UK: Oxford University Press, 1992: 151-77.
108. Moxham C, Jacobs S. Insulin-like growth factor receptors. In: Schofield PN. *The Insulin-like Growth Factors; Structure and Biological Functions*. Oxford, UK: Oxford University Press, 1992: 80-109.
109. Kahn CR, White MF. The insulin receptor and molecular mechanisms of insulin action. *J Clin Invest* 1988; 82: 1151-6.
110. D'Ercole AJ, Stiles AD, Underwood LE. Tissue concentrations of somatomedin C: Further evidence for multiple sites of synthesis and paracrine or autocrine mechanisms of action. *Proc Natl Acad Sci USA* 1984; 81: 935-9.
111. Cara JF, Rosenfield RL. Insulin-like growth factor I and insulin potentiate luteinizing hormone-induced androgen synthesis by rat ovarian thecal-interstitial cells. *Endocrinology* 1988; 123: 733-9.
112. Baxter RC, Martin JL. Binding proteins for the insulin-like growth factors: structure, regulation and function. *Prog Growth Factor Res* 1989; 1: 49-68.
113. Clemmons DR, Underwood LE. Nutritional regulation of IGF-I and IGF binding proteins. *Annu Rev Nutr* 1991; 11: 393-412.
114. Isley WL, Underwood LE, Clemmons DR. Changes in plasma somatomedin-C in response to ingestion of diets with variable protein and energy content. *J Parenteral Enteral Nutr* 1984; 8: 407-11.
115. Straus DS. Nutritional regulation of hormones and growth factors that control mammalian growth. *FASEB J* 1994; 8: 6-12.
116. Clemmons DR, Klibanski A, Underwood LE, et al. Reduction of plasma immunoreactive somatomedin C during fasting in humans. *J Clin Endocrinol Metab* 1981; 53: 1247-50.
117. Minuto F, Barreca A, Adami GF, et al. Insulin-like growth factor-I in human malnutrition: relationship with some body composition and nutritional parameters. *J Parenteral Enteral Nutr* 1989; 13: 392-6.
118. Smith AT, Clemmons DR, Underwood LE, Ben-Ezra V, McMurray R. The effect of exercise on plasma somatomedin-C/insulin-like growth factor I concentrations. *Metabolism* 1987; 36: 533-7.
119. Baxter RC, Cowell CT. Diurnal rhythm of the growth hormone-independent binding protein for insulin-like growth factors in human plasma. *J Clin Endocrinol Metab* 1987; 65: 432-40.
120. Conover CA, Lee PD, Kanaley JA, Clarkson JT, Jensen MD. Insulin regulation of insulin-like growth factor binding protein-1 in obese and nonobese humans. *J Clin Endocrinol Metab* 1992; 74: 1355-60.
121. Holly JM, Biddlecombe RA, Dunger DB, et al. Circadian variation of GH-independent IGF-binding protein in diabetes mellitus and its relationship to insulin. A new role for insulin? *Clin Endocrinol Oxf* 1988; 29: 667-75.
122. Brismar K, Gutniak M, Pova G, Werner S, Hall K. Insulin regulates the 35 kDa IGF binding protein in patients with diabetes mellitus. *J Endocrinol Invest* 1988; 11: 599-602.
123. Brismar K, Grill V, Efendic S, Hall K. The insulin-like growth factor binding protein-1 in low and high insulin responders before and during dexamethasone treatment. *Metabolism* 1991; 40: 728-32.
124. Pekonen F, Laatikainen T, Buyalos R, Rutanen EM. Decreased 34K insulin-like growth factor binding protein in polycystic ovarian disease. *Fertil Steril* 1989; 51: 972-5.
125. Holly JM, Eden JA, Alaghband-Zadeh J, et al. Insulin-like growth factor binding proteins in follicular fluid from normal

- dominant and cohort follicles, polycystic and multicystic ovaries. *Clin Endocrinol Oxf* 1990; 33: 53-64.
126. Suikkari AM, Koivisto VA, Rutanen EM, Yki-Järvinen H, Karonen SL, Seppälä M. Insulin regulates the serum levels of low molecular weight insulin-like growth factor binding protein. *J Clin Endocrinol Metab* 1988; 66: 266-72.
 127. Cotterill AM, Cowell CT, Silink M. Insulin and variation in glucose levels modify the secretion rates of the growth hormone-independent insulin-like growth factor binding protein-1 in the human hepatoblastoma cell line Hep G2. *J Endocrinol* 1989; 123: R17-20.
 128. Conover CA, Lee PD. Insulin regulation of insulin-like growth factor-binding protein production in cultured HepG2 cells. *J Clin Endocrinol Metab* 1990; 70: 1062-7.
 129. Powell DR, Suwanichkul A, Cabbage ML, DePaolis LA, Snuggs MB, Lee PDK. Insulin inhibits transcription of the human gene for insulin-like growth factor-binding protein-1. *J Biol Chem* 1991; 266: 18868-76.
 130. Bar RS, Boes M, Clemmons D, et al. Insulin differentially alters transcapillary movement of intravascular IGFBP-1, IGFBP-2 and endothelial cell IGF-binding proteins in rat heart. *Endocrinology* 1990; 127: 497-9.
 131. Adashi EY, Resnick CE, Hurwitz A, et al. Insulin-like growth factors: the ovarian connection. *Hum Reprod* 1991; 6: 1213-9.
 132. Koistinen R, Suikkari AM, Tiitinen A, Kontula K, Seppälä M. Human granulosa cells contain insulin-like growth factor binding protein (IGFBP-1) mRNA. *Clin Endocrinol Oxf* 1990; 32: 635-40.
 133. Yen SSC. Chronic anovulation caused by peripheral endocrine disorders. In: Yen SCC, Jaffe RB, eds. *Reproductive Endocrinology: Physiology, Pathophysiology, and Clinical Management*. Philadelphia, PA (USA): Saunders Company, 1991: 576-629.
 134. Moller DE, Flier JS. Insulin resistance – mechanisms, syndromes, and implications. *N Engl J Med* 1991; 26: 938-48.
 135. Dunaif A. Hyperandrogenic anovulation (PCOS): a unique disorder of insulin action associated with an increased risk of non-insulin-dependent diabetes mellitus. *Am J Medicine* 1995; 98(Suppl 1A): 33S-39S.
 136. Zavaroni I, Bonara E, Pagliara M, et al. Risk factors for coronary artery disease in healthy persons with hyperinsulinemia and normal glucose tolerance. *N Engl J Med* 1989; 320: 702-6.
 137. Ferranini E, Buzzigoli G, Bonadonna R, et al. Insulin resistance in essential hypertension. *N Engl J Med* 1987; 317: 350-7.
 138. Beatty OL, Harper B, Sheridan B, Atkinson AB, Bell PM. Insulin resistance in offspring of hypertensive parents. *Br Med J* 1993; 307: 92-6.
 139. Allemann Y, Horber FF, Colombo M, et al. Insulin sensitivity and body fat distribution in normotensive offspring of hypertensive parents. *Lancet* 1993; 341: 327-31.
 140. Hollenbeck C, Reaven GM. Variations in insulin-stimulated glucose uptake in healthy individuals with normal glucose tolerance. *J Clin Endocrinol Metab* 1987; 64: 1169-73.
 141. Randle PJ, Garland PB, Hales CN, Newsholme EA. The glucose fatty-acid cycle, its role in insulin sensitivity and the metabolic disturbances of diabetes mellitus. *Lancet* 1963; i: 785-9.
 142. Clark MG, Rattigan S, Clark DG. Obesity with insulin resistance: experimental results. *Lancet* 1983; ii: 1236-40.
 143. DeFronzo RA. The triumvirate: β -cell, muscle, liver – a collusion responsible for NIDDM. *Diabetes* 1988; 37: 667-87.
 144. O'Dea K. Westernization, insulin resistance and diabetes in Australian Aborigines. *Med J Aust* 1991; 155: 258-64.
 145. Taylor SI, Mukherjee C, Jungas RL. Studies on the mechanism of activation of adipose tissue pyruvate dehydrogenase by insulin. *J Biol Chem* 1973; 248: 73-81.
 146. Taylor SI, Mukherjee C, Jungas RL. Regulation of pyruvate dehydrogenase in isolated rat liver mitochondria. *J Biol Chem* 1975; 250: 2028-35.
 147. Jeanrenaud B, Halimi S, van de Werve G. Neuro-endocrine disorders seen as triggers of the triad: obesity – insulin resistance – abnormal glucose tolerance. *Diabetes Metab Rev* 1985; 1: 261-92.
 148. Wititsuwannakul D, Kim K. Mechanism of palmitoyl co-enzyme A inhibition on liver glycogen synthase. *J Biol Chem* 1977; 252: 7812-7.
 149. Ebeling P, Koivisto VA. Non-esterified fatty acids regulate lipid and glucose oxidation and glycogen synthesis in healthy man. *Diabetologia* 1994; 37: 202-9.
 150. Lillioja S, Mott DM, Zawadzki JK, Young AA, Abbott WG, Bogardus C. Glucose storage is a major determinant of in vivo 'insulin resistance' in subjects with normal glucose tolerance. *J Clin Endocrinol Metab* 1986; 62: 922-7.
 151. Shulman GI, Rothman DL, Jue T, Stein P, DeFronzo RA, Shulman RG. Quantitation of muscle glycogen synthesis in normal subjects and subjects with non-insulin-dependent diabetes by ^{13}C nuclear magnetic resonance spectroscopy. *N Eng J Med* 1990; 322: 223-8.
 152. Kraegen EW, Clark PW, Jenkins AB, Dalley EA, Chilholm DJ, Storlien LH. Development of muscle insulin resistance after liver insulin resistance in high-fat-fed rats. *Diabetes* 1991; 40: 1397-403.
 153. Kiens B, Essen-Gustavsson B, Gad P, Lithell H. Lipoprotein lipase activity and intramuscular triglyceride stores after long-term high-fat and high-carbohydrate diets in physically trained men. *Clin Physiol* 1987; 7: 1-9.
 154. Björntorp P. Metabolic abnormalities in visceral obesity. *Ann Med* 1992; 24: 3-5.
 155. Lonnqvist F, Thome A, Nilzell K, Hoffstedt J, Arner P. A pathogenic role of visceral fat beta 3-adrenoreceptors in obesity. *J Clin Invest* 1995; 95: 1109-16.
 156. Mårin P, Andersson B, Ottoson M, et al. The morphology and metabolism of intraabdominal adipose tissue in men. *Metabolism* 1992; 41: 1242-8.
 157. Ferranini E, Barrett EJ, Bevilacqua S. Effect of fatty acids on glucose production and utilization in man. *J Clin Invest* 1983; 72: 1737-47.
 158. Golay A, Swislocki ALM, Chen YDI, Reaven GM. Relationships between plasma free fatty acid concentration, endogenous glucose production, and fasting hyperglycemia in normal and non-insulin-dependent diabetic individuals. *Metabolism* 1987; 7: 692-6.
 159. Williamson JR, Kreisberg RA, Felts PW. Mechanism for the stimulation of gluconeogenesis in perfused rat liver. *Proc Natl Acad Sci USA* 1966; 56: 247-54.
 160. Williamson JR, Browning ET, Olson M. Interrelations between fatty acid oxidation and the control of gluconeogenesis in perfused rat liver. *Adv Enzyme Regul* 1968; 6: 67-100.
 161. Bar RS, Harrison LC, Muggeo M, Gorden P, Kahn CR, Roth J. Regulation of insulin receptors in normal and abnormal physiology in humans. *Adv Intern Med* 1972; 24: 23-52.
 162. Gavin JR, Roth J, Neville DM, de Meyts P, Buell DN. Insulin-dependent regulation of insulin receptor concentrations: a direct demonstration in cell culture. *Proc Natl Acad Sci USA* 1974; 7: 84-8.
 163. Flier JS. Insulin receptors and insulin resistance. *Annu Rev Med* 1983; 34: 145-60.

164. Kahn CR. Role of insulin receptors in insulin resistant states. *Metabolism* 1980; 29: 455-66.
165. Fried SK, Russell CD, Grauso NL, Brolin RE. Lipoprotein lipase regulation by insulin and glucocorticoid in subcutaneous and omental adipose tissues of obese women and men. *J Clin Invest* 1993; 92: 2191-8.
166. Ravussin E, Swinburn BA. Pathophysiology of obesity. *Lancet* 1992; 340: 404-8.
167. Wing RR, Matthews KA, Kuller LH, et al. Environmental and familial contributions to insulin levels and change in insulin levels in middle-aged women. *JAMA* 1992; 268: 1890-5.
168. Ravussin E, Valencia ME, Esparza J, Bennett PH, Schulz LO. Effects of a traditional lifestyle on obesity in Pima Indians. *Diabetes Care* 1994; 17: 1067-74.
169. Flatt JP. Dietary fat, carbohydrate balance, and weight maintenance. In: Klimes I, Howard BV, Storlien LH, Šeböková E, eds. *Dietary Lipids and Insulin Action*. New York, NY (USA): New York Acad Sci 1993: 122-40.
170. Swinburn B, Ravussin E. Energy balance or fat balance? *Am J Clin Nutr* 1993; 57(Suppl): 766S-71S.
171. Astrup A, Raben A. Obesity: an inherited metabolic deficiency in the control of macronutrient balance? *Eur J Clin Nutr* 1992; 46: 611-20.
172. Westterterp KR. Food quotient, respiratory quotient, and energy balance. *Am J Clin Nutr* 1993; 57(Suppl): 759S-65S.
173. Flatt JP, Ravussin E, Acheson KJ, Jéquier E. Effects of dietary fat on postprandial substrate oxidation and on carbohydrate and fat balance. *J Clin Invest* 1985; 76: 1019-24.
174. Schutz Y, Flatt JP, Jéquier E. Failure of fat to promote fat oxidation: a factor favoring the development of obesity. *Am J Clin Nutr* 1989; 50: 307-14.
175. Eckel RH. Insulin resistance: an adaptation for weight maintenance. *Lancet* 1992; 340: 1452-3.
176. Astrup A, Buemann B, Western P, Toubro S, Raben A, Christensen NJ. Obesity as an adaptation to a high-fat diet: evidence from a cross-sectional study. *Am J Clin Nutr* 1994; 59: 350-5.
177. Hales CN, Randle PJ. Effects of low-carbohydrate diet and diabetes mellitus on plasma concentrations of glucose, non-esterified fatty acid, and insulin during oral glucose-tolerance tests. *Lancet* 1963; i: 790-4.
178. Fukagawa NK, Anderson JW, Hageman G, Young VR, Minaker KL. High-carbohydrate, high-fiber diets increase peripheral insulin sensitivity in healthy young and old adults. *Am J Clin Nutr* 1990; 52: 524-28.
179. Howard BV, Abbott WGH, Swinburn BA. Evaluation of metabolic effects of substitution of complex carbohydrates for saturated fat in individuals with obesity and NIDDM. *Diabetes Care* 1991; 14: 786-95.
180. Chen M, Bergman RN, Porte D. Insulin resistance and β -cell dysfunction in aging: the importance of dietary carbohydrate. *J Clin Endocrinol Metab* 1988; 67: 951-7.
181. Kolterman OG, Greenfield M, Reaven GM, Saekow M, Olefsky JM. Effect of a high carbohydrate diet on insulin binding to adipocytes and on insulin action *in vivo* in man. *Diabetes* 1979; 28: 731-6.
182. Beck-Nielsen H, Pedersen O, Sørensen NS. Effects of diet on the cellular binding and the insulin sensitivity in young healthy subjects. *Diabetologia* 1978; 15: 289-96.
183. Grey N, Kipnis DM. effect of diet composition on the hyperinsulinemia of obesity. *N Engl J Med* 1979; 285: 827-31.
184. Coulston AM, Liu GC, Reaven RM. Plasma glucose, insulin and lipid responses to high-carbohydrate low-fat diets in normal humans. *Metabolism* 1983; 32: 52-6.
185. Danforth E. Diet and obesity. *Am J Clin Nutr* 1985; 41: 1132-45.
186. Tucker LA, Kano MJ. Dietary fat and body fat: a multivariate study of 205 adult females. *Am J Clin Nutr* 1992; 56: 616-22.
187. Miller WC, Lindeman AK, Wallace J, Niederpruem M. Diet composition, energy intake, and exercise in relation to body fat in men and women. *Am J Clin Nutr* 1990; 52: 426-30.
188. Dreon DM, Frey-Hewitt B, Ellsworth N, Williams PT, Terry RB, Wood PD. Dietary fat: carbohydrate ratio and obesity in middle-aged men. *Am J Clin Nutr* 1988; 47: 995-1000.
189. Romieu I, Willett WC, Stampfer MJ, et al. Energy intake and other determinants of relative weight. *Am J Clin Nutr* 1988; 47: 406-12.
190. Lilienthal Heitman B, Lissner L, Sørensen TIA, Bengtsson C. Dietary fat and weight gain in women genetically predisposed for obesity. *Am J Clin Nutr* 1995; 61: 1213-7.
191. Gazzaniga JM, Burns TL. Relationship between diet composition and body fatness, with adjustment for resting energy expenditure and physical activity, in preadolescent children. *Am J Clin Nutr* 1993; 58: 21-8.
192. Parker DR, Weiss ST, Troisi R, Cassano PA, Vokonas PS, Landsberg L. Relationship of dietary fatty acids and body habitus to serum insulin concentrations: the Normative Aging Study. *Am J Clin Nutr* 1993; 58: 129-36.
193. Mayer EJ, Newman B, Quesenberry CP, Selby JV. Usual dietary fat intake and insulin concentrations in healthy women twins. *Diabetes Care* 1991; 16: 1459-69.
194. Maron DJ, Fair JM, Haskell WL, et al. Saturated fat intake and insulin resistance in men with coronary heart disease. *Circulation* 1991; 84: 2020-7.
195. Lovejoy J, DiGirolamo M. Habitual dietary intake and insulin sensitivity in lean and obese adults. *Am J Clin Nutr* 1992; 55: 1174-9.
196. Feskens EJM, Kromhout D. Habitual dietary intake and glucose tolerance in euglycaemic men: the Zutphen study. *Int J Epidemiol* 1990; 19: 953-59.
197. Marshall JA, Hamman RF, Baxter J. High-fat, low-carbohydrate diet and the etiology of non-insulin-dependent diabetes mellitus: the San Luis Valley Diabetes Study. *Am J Epidemiol* 1991; 134: 590-603.
198. Tsunehara CH, Leonetti DL, Fujimoto WY. Diet of second-generation Japanese-American men with and without non-insulin-dependent diabetes. *Am J Clin Nutr* 1990; 52: 731-8.
199. Field CJ, Ryan EA, Thompson ABR, Clandinin MT. Diet fat composition alters membrane phospholipid composition, insulin binding, and glucose metabolism in adipocytes from control and diabetic animals. *J Biol Chem* 1990; 19: 11143-50.
200. Storlien LH, Kraegen EW, Chisholm DJ, Ford GL, Bruce DG, Pascoe WS. Fish oil prevents insulin resistance induced by high-fat feeding in rats. *Science* 1987; 237: 885-8.
201. Hainault I, Carlotti M, Hajduch E, Guichard C, Lavau M. Fish oil in a high-lard diet prevents obesity, hyperlipemia, and adipocyte insulin resistance in rats. *Ann NY Acad Sci* 1993; 683: 98-101.
202. Pelikanova T, Kohout M, Valek J, Base J, Kazdova L. Insulin secretion and insulin action related to the serum phospholipid fatty acid pattern in healthy men. *Metabolism* 1989; 38: 188-92.
203. Borkman M, Storlien LH, Pan DA, Jenkins AB, Chisholm DJ, Campbell LV. The relation between insulin sensitivity and the fatty-acid composition of skeletal-muscle phospholipids. *N Engl J Med* 1993; 328: 238-44.
204. Ruiz-Gutierrez V, Stifel P, Villar J, Garcia-Donas MA, Acosta D, Carneado J. Cell membrane fatty acid composition type 1 (insulin-dependent) diabetic patients: relationship with

- sodium transport abnormalities and metabolic control. *Diabetologia* 1993; 36: 850-6.
205. Agostoni C, Riva E, Bellu R, Vicenzo SS, Grazia BM, Giovannini M. Relationships between the fatty acid status and insulinemic indexes in obese children. *Prostaglandins Leukot Essent Fatty Acids* 1994; 51: 317-21.
 206. Feskens EJM, Loeber JG, Kromhout D. Diet and physical activity as determinants of hyperinsulinaemia: the Zutphen elderly study. *Am J Epidemiol* 1994; 140: 350-60.
 207. Trevisan M, Krogh V, Freudenheim J, et al. Consumption of olive oil, butter, and vegetable oils and coronary heart disease risk factors. *JAMA* 1990; 263: 688-92.
 208. Popp-Snijders C, Schouten JA, Heine RJ, van der Meer J, van der Veen EA. Dietary supplementation of omega 3 polyunsaturated fatty acids improves insulin sensitivity in non-insulin dependent diabetes. *Diabetes Res* 1987; 4: 141-7.
 209. Schectman G, Kaul S, Kissebah AH. Effect of fish oil concentrate on lipoprotein composition in NIDDM. *Diabetes* 1988; 37: 1567-73.
 210. Pilch PF, Thompson PA, Czech MP. Coordinate modulation of D-glucose transport activity and bilayer fluidity in plasma membranes derived from control and insulin-treated adipocytes. *Proc Natl Acad Sci USA* 1980; 77: 915-8.
 211. Ginsberg BH, Jabour J, Spector AA. Increased membrane fluidity is associated with greater sensitivity to insulin (Abstract). *Diabetes* 1987; 36(Suppl 1): 51A.
 212. Truswell AS. Glycaemic index of foods. *Eur J Clin Nutr* 1992; 46(Suppl 2): S91-101.
 213. Jenkins DA, Wolever TMS, Taylor RH, et al. Glycemic index of foods: a physiological basis for carbohydrate exchange. *Am J Clin Nutr* 1981; 34: 362-6.
 214. Collings P, Williams C, Macdonald I. Effects of cooking on serum glucose and insulin responses to starch. *Br Med J* 1981; 282: 1032-3.
 215. Brand JC, Nicholson PL, Thorburn AW, Truswell AS. Food processing and the glycemic index. *Am J Clin Nutr* 1985; 42: 1192-6.
 216. Crapo PA. Simple versus complex carbohydrate use in the diabetic diet. *Annu Rev Nutr* 1985; 5: 95-114.
 217. Wolever TMS. The glycemic index. In: Bourne GH ed. *Aspects of Some Vitamins, Minerals and Enzymes in Health and Disease. World Rev Nutr Diet (Basel, Karger)* 1990; 62: 120-85.
 218. Jenkins DJA, Wolever TMS, Jenkins AL, Lee R, Wong GS, et al. Glycemic response to wheat products: reduced response to pasta but no effects of fiber. *Diabetes Care* 1983; 6: 155-90.
 219. O'Dea K, Nestel PJ, Antonoff L. Physical factors influencing postprandial glucose and insulin responses to starch. *Am J Clin Nutr* 1980; 33: 760-5.
 220. Jenkins DJA, Wesson V, Wolever TMS, et al. Wholemeal versus wholegrain breads: proportion of whole or cracked grain and the glycaemic response. *Br Med J* 1988; 297: 958-60.
 221. Manolio TA, Savage PJ, Burke GL, et al. Correlates of fasting insulin levels in young adults: the CARDIA study. *J Clin Epidemiol* 1991; 44: 571-8.
 222. Feskens EJ, Kromhout D. Habitual dietary intake and glucose tolerance in euglycemic men: the Zutphen study. *Int J Epidemiol* 1990; 19: 953-9.
 223. Marshall JA, Weiss NS, Hamman RF. The role of dietary fibre in the etiology of non-insulin-dependent diabetes mellitus. *Ann Epidemiol* 1993; 3: 18-26.
 224. Harlan LC, Harlan WR, Landis JR, Goldstein NG. Factors associated with glucose tolerance in US adults. *Am J Epidemiol* 1987; 126: 674-84.
 225. Hill JO, Prentice AM. Sugar and body weight regulation. *Am J Clin Nutr* 1995; 62(Suppl): 264S-74S.
 226. Koivisto VA, Yki-Jarvinen H, DeFronzo RA. Physical training and insulin sensitivity. *Diabetes Metab Rev* 1988; 1: 445-81.
 227. Regensteiner JG, Mayer EJ, Shetterly SM, et al. Relationship between habitual physical activity and insulin levels among nondiabetic men and women. *Diabetes Care* 1991; 14: 1066-74.
 228. Dowse GK, Zimmet PZ, Gareeboo H, et al. Abdominal obesity and physical inactivity are risk factors for NIDDM and impaired glucose tolerance in Indian, Creole, and Chinese Mauritians. *Diabetes Care* 1991; 14: 271-82.
 229. Lindgarde F, Saltin B. Daily physical activity, work capacity and glucose tolerance in lean and obese normo-glycaemic middle-aged men. *Diabetologia* 1981; 20: 137-8.
 230. Wang JT, Ho LT, Tang KT, Wang LM, Chen YDI, Reaven GM. Effect of habitual physical activity on age-related glucose intolerance. *J Am Geriatr Soc* 1989; 37: 203-9.
 231. Manson JE, Rimm E, Stampfer MJ, et al. Physical activity and incidence of non-insulin-dependent diabetes mellitus in women. *Lancet* 1991; 338: 774-8.
 232. Holly JMP, Smith CP, Dunger DB, et al. Relationship between the pubertal fall in sex hormone binding globulin and insulin-like growth factor binding protein-I. A synchronized approach to pubertal development? *Clin Endocrinol* 1989; 31: 277-84.
 233. Belgorosky A, Rivarola MA. Progressive increase in nonsex hormone-binding-globulin-bound testosterone and estradiol from infancy to late puberty in girls. *J Clin Endocrinol Metab* 1988; 67: 234-7.
 234. Bartsch W, Horst HJ, Derwahl KM. Interrelations between sex-hormone-binding globulin and 17 β -estradiol, testosterone, 5(-dihydroxytestosterone, thyroxine, and triiodothyronine in prepubertal and pubertal girls. *J Clin Endocrinol Metab* 1980; 50: 1053-6.
 235. Apter D, Vihko R. Endocrine determinants of fertility: serum androgen concentrations during follow-up of adolescents into the third decade of life. *J Clin Endocrinol Metab* 1984; 71: 970-4.
 236. Apter D, Bolton NJ, Hammond GL, Vihko R. Serum sex hormone binding globulin during puberty in girls and in different types of adolescent menstrual cycles. *Acta Endocrinologica* 1984; 107: 413-9.
 237. Lee IR, Lawder LE, Townend DC, Wetherall JD, Hahnel R. Plasma sex hormone binding globulin concentrations and binding capacity in children before and during puberty. *Acta Endocrinologica* 1985; 109: 276-80.
 238. Holly JMP, Smith CP, Dunger DB, et al. Levels of the small insulin-like growth factor-binding protein are strongly related to those of insulin in prepubertal and pubertal children but only weakly so after puberty. *J Endocrinol* 1989; 121: 383-7.
 239. Argente J, Barrios V, Pozo J, et al. Normative data for insulin-like growth factors (IGFs), IGF-binding proteins, and growth hormone-binding protein in a healthy Spanish pediatric population: age- and sex-related changes. *J Clin Endocrinol Metab* 1993; 77: 1522-8.
 240. Lautala P, Åkerblom HK, Viikari J, et al. Atherosclerosis precursors in Finnish children and adolescents. VII. Serum immunoreactive insulin. *Acta Paediatr Scand* 1985; 318(Suppl): 127-33.
 241. Amiel SA, Sherwin RS, Simonson DC, Lauritano AA, Tamborlane WV. Impaired insulin action in puberty. A contributing factor to poor glycemic control in adolescents with diabetes. *N Engl J Med* 1986; 315: 215-9.
 242. Bloch CA, Clemons P, Sperling MA. Puberty decreases insulin sensitivity. *J Pediatr* 1987; 110: 481-7.

243. Caprio S, Plewe G, Diamond MD, *et al.* Increased insulin secretion in puberty: a compensatory response to reductions in insulin sensitivity. *J Pediatr* 1989; 114: 963-7.
244. Smith CP, Archibald HR, Thomas JM, *et al.* Basal and stimulated insulin levels rise with advancing puberty. *Clin Endocrinol* 1988; 28: 7-14.
245. Laron Z, Aurbach-Klipper Y, Flasterstein B, Litwin A, Dickerman Z, Heding LG. Changes in endogenous insulin secretion during childhood as expressed by plasma and urinary C-peptide. *Clin Endocrinol* 1988; 29: 625-32.
246. Travers SH, Jeffers BW, Bloch CA, Hill JO, Eckel RH. Gender and Tanner stage differences in body composition and insulin sensitivity in early pubertal children. *J Clin Endocrinol Metab* 1995; 80: 172-8.
247. Clemmons DR, van Wyk JJ. Factors controlling blood concentration of somatomedin C. *Clin Endocrinol Metab* 1984; 13: 113-43.
248. Smith CP, Dunger DB, Williams AJK, *et al.* Relationship between insulin, insulin-like growth factor I, and dehydroepiandrosterone sulfate concentrations during childhood, puberty, and adult life. *J Clin Endocrinol Metab* 1989; 68: 932-7.
249. Hesse V, Jahreis G, Schmaabach H, *et al.* Insulin-like growth factor I correlations to changes of the hormonal status in puberty and age. *Exp Clin Endocrinol* 1994; 102: 289-8.
250. Wilson DW, Killen JD, Hammer LD, *et al.* Insulin-like growth factor-I as a reflection of body composition, nutrition, and puberty in sixth and seventh grade girls. *J Clin Endocrinol Metab* 1991; 73: 907-12.
251. Copeland KC, Colletti RB, Devlin JT, McAuliffe TL. The relationship between insulin-like growth factor I, adiposity, and aging. *Metabolism* 1990; 39: 584-7.
252. Hindmarsh P, Di Silvio PJ, Pringle PJ, Kurtz AB, Brook CGD. Changes in serum insulin concentration during puberty and their relationship to growth hormone. *Clin Endocrinol* 1988; 28: 381-8.
253. Arslanian SA, Kalhan SC. Correlations between fatty acid and glucose metabolism: potential explanation of insulin resistance during puberty. *Diabetes* 1994; 43: 908-14.
254. Benassi L, Tridenti G, Orlandi N, Pezzarossa A. Glucose tolerance and insulin release in adolescent female. *J Endocrinol Invest* 1991; 14: 751-6.
255. Rosenfeld RG, Wilson DM, Dollar LA, Bennett A, Hintz RL. Both pituitary growth hormone and recombinant DNA-derived growth hormone cause insulin resistance at a postreceptor site. *J Clin Endocrinol Metab* 1982; 54: 1033-8.
256. Press M, Tamborlane WV, Sherwin RS. Importance of raised growth hormone levels in mediating the metabolic derangements of diabetes. *N Engl J Med* 1984; 310: 810-5.
257. Rizza RA, Mandarino LJ, Gerich JE. Effects of growth hormone on insulin action in man: Mechanisms of insulin resistance, impaired suppression of glucose production, and impaired stimulation of glucose utilization. *Diabetes* 1982; 31: 663-9.
258. Martha PM, Reiter EO. Pubertal growth and growth hormone secretion. *Endocrinol Metab Clin North Am* 1991; 20: 165-82.
259. Bratusch-Marrain PR, Smith D, DeFronzo RA. The effect of growth hormone on glucose metabolism and insulin secretion in man. *J Clin Endocrinol Metab* 1982; 55: 973-82.
260. Zwiauer KF, Pakosta R, Mueller T, Widhalm K. Cardiovascular risk factors in obese children in relation to weight and body fat distribution. *J Am Coll Nutr* 1992; 11(Suppl): 41S-50S.
261. Kikuchi DA, Srinivasan SR, Harsha DW, Webber LS, Sellers TA, Berenson GS. Relation of serum lipoprotein lipids and apolipoproteins to obesity in children: the Bogalusa Heart Study. *Prev Med* 1992; 21: 177-90.
262. Hoffman RP, Stumbo PJ, Janz KF, Nielsen DH. Altered insulin resistance is associated with increased dietary weight loss in obese children. *Horm Res* 1995; 44: 17-22.
263. Wabitsch M, Hauner H, Heinze E, *et al.* Body fat distribution and changes in the atherogenic risk-factor profile in obese adolescent girls during weight reduction. *Am J Clin Nutr* 1994; 60: 54-60.
264. Arslanian S, Suprasongsin C. Insulin sensitivity, lipids, and body composition in childhood: is 'syndrome X' present? *J Clin Endocrinol Metab* 1996; 81: 1058-62.
265. Klesges RC, Klesges LM, Eck LH, Shelton ML. A longitudinal analysis of accelerated weight gain in preschool children. *Pediatrics* 1995; 95: 126-30.
266. Esposito del Puente A, Scalfi L, de Filippo E, *et al.* Familial and environmental influences on body composition and body fat distribution in childhood in southern Italy. *Int J Obes Relat Metab Disord* 1994; 18: 596-601.
267. Pasquali R, Casimirri F, Venturoli S, *et al.* Body fat distribution has weight-independent effects on clinical, hormonal, and metabolic features of women with polycystic ovary syndrome. *Metabolism* 1994; 43: 706-13.
268. Rebuffe-Scrive M, Cullberg G, Lundberg PA, Lindstedt G, Bjorntorp P. Anthropometric variables and metabolism in polycystic ovarian disease. *Horm Metab Res* 1989; 21: 391-7.
269. Bringer J, Lefebvre P, Boulet F, *et al.* Body composition and regional fat distribution in polycystic ovarian syndrome. Relationship to hormonal and metabolic profiles. *Ann N Y Acad Sci* 1993; 687: 115-23.
270. Lanzone A, Fulghese AM, Pappalardo S, *et al.* Growth hormone and somatomedin-C secretion in patients with polycystic ovarian disease. *Gynecol Obstet Invest* 1990; 29: 149-53.
271. Chang RJ, Nakamura RM, Judd HL, Kaplan SO. *J Clin Endocrinol Metab* 1983; 57: 356-9.
272. Rittmaster RS, Deshwal N, Lehman L. The role of adrenal hyperandrogenism, insulin resistance, and obesity in the pathogenesis of polycystic ovarian syndrome. *J Clin Endocrinol Metab* 1993; 76: 1295-1300.
273. Conway GS, Jacobs HS, Holly JMP, Wass JAS. Effects of luteinizing hormone, insulin, insulin-like growth factor I and insulin-like growth factor small binding protein in the polycystic ovary syndrome. *Clin Endocrinol* 1990; 33: 593-603.
274. Tropeano G, Licisano A, Liberale I, *et al.* Insulin, C-peptide, androgens, and beta-endorphin response to oral glucose in patients with polycystic ovary syndrome. *J Clin Endocrinol Metab* 1994; 78: 305-9.
275. Dunaif A, Graf M, Mandeli J, Laumas V, Dobrjanski A. Characterization of groups of hyperandrogenic women with acanthosis nigricans, impaired glucose tolerance, and/or hyperinsulinemia. *J Clin Endocrinol Metab* 1987; 65: 499-507.
276. Dunaif A, Segal KR, Futterweit W, Dobrjansky A. Profound peripheral insulin resistance, independent of obesity, in polycystic ovary syndrome. *Diabetes* 1989; 38: 1165-74.
277. Suikkari AM, Ruutiainen K, Erkkola R, Seppälä M. Low levels of low molecular weight insulin-like growth factor binding protein in patients with polycystic ovarian disease. *Hum Reprod* 1989; 4: 136-9.
278. Franks S, Kiddy D, Sharp P, *et al.* Obesity and polycystic ovary syndrome. *Ann NY Acad Sci* 1991; 626: 201-6.
279. Kiddy DS, Hamilton-Fairley D, Bush A, *et al.* Improvement in endocrine and ovarian function during dietary treatment of obese women with polycystic ovary syndrome. *Clin Endocrinol* 1992; 36: 105-11.

280. Venturoli S, Porcu E, Fabbri R, *et al.* Menstrual irregularities in adolescents: hormonal pattern and ovarian morphology. *Hormone Res* 1986; 24: 269-79.
281. Polson DW, Wadsworth J, Adams J, Franks S. Polycystic ovaries – a common finding in normal women. *Lancet* 1988; i: 870-2.
282. Clayton RN, Ogden V, Hodgkinson J, *et al.* How common are polycystic ovaries in normal women and what is their significance for the fertility of the population? *Clin Endocrinol* 1992; 37: 127-34.
283. Abdel-Gadir A, Khatim MS, Mowafi RS, Alnaser HM, Muharib NS, Shaw RW. Implications of ultrasonically diagnosed polycystic ovaries. I. Correlations with basal hormonal profiles. *Hum Reprod* 1992; 7: 453-7.
284. Nobels F, Dewailly D. Puberty and polycystic ovarian syndrome: the insulin/insulin-like growth factor I hypothesis. *Fertil Steril* 1992; 58: 655-66.
285. Lucky AW, Rosenfield RL, McGuire J, Rudy S, Helke J. Adrenal androgen hyperresponsiveness to adrenocorticotropin in women with acne and/or hirsutism: Adrenal enzyme defects and exaggerated adrenarche. *J Clin Endocrinol Metab* 1986; 62: 840-8.
286. Ehrmann DA, Barnes RB, Rosenfield RL. Polycystic ovary syndrome as a form of functional ovarian hyperandrogenism due to dysregulation of androgen secretion. *Endocr Rev* 1995; 16: 322-53.
287. Pasquali R, Casimirri F. Insulin and androgen relationships with abdominal body fat distribution in women with and without hyperandrogenism. *Horm Res* 1993; 39: 179-87.
288. Barbieri RL, Smith S, Ryan KJ. The role of hyperinsulinemia in the pathogenesis of ovarian hyperandrogenism. *Fertil Steril* 1988; 50: 197-212.
289. Rosenfield RL, Barnes RB, Cara JF, Lucky AW. Dysregulation of cytochrome P450c17(as the cause of polycystic ovarian syndrome. *Fertil Steril* 1990; 53: 785-91.
290. Magoffin DA, Kurtz KM, Erickson GF. Insulin-like growth factor selectively stimulates cholesterol side-chain cleavage expression in ovarian theca-interstitial cells. *Mol Endocrinol* 1990; 4: 489-96.
291. Magoffin DA, Weitsman SR. Differentiation of ovarian theca-interstitial cells in vitro: regulation of 17 alpha-hydroxylase messenger ribonucleic acid expression by luteinizing hormone and insulin-like growth factor-I. *Endocrinology* 1993; 132: 1945-51.
292. Hiney JK, Ojeda SR, Dees WL. Insulin-like growth factor I: a possible metabolic signal involved in the regulation of female puberty. *Neuroendocrinology* 1991; 54: 420-3.
293. Kazer R. The etiology of polycystic ovary syndrome (PCO). *Med Hypotheses* 1992; 30: 151-5.
294. Venturoli S, Porcu E, Fabbri R, *et al.* Postmenarcheal evolution of endocrine pattern and ovarian aspects in adolescents with menstrual irregularities. *Fertil Steril* 1987; 48: 78-85.
295. Apter D, Vihko R. Endocrine determinants of fertility: serum androgen concentrations during follow-up of adolescents into the third decade of life. *J Endocrinol Metab* 1990; 71: 970-4.
296. Simpson JL. Elucidating the genetics of polycystic ovary syndrome. In: Dunaif A, Givens JR, Haseltine FP, Merriam GR, eds. *Polycystic Ovary Syndrome*. Oxford, UK: Blackwell Scientific Publications, 1992: Chapter 6: 59-69.
297. Carey AH, Chan KL, Short F, White D, Williams R, Franks S. Evidence for a single gene effect causing polycystic ovary syndrome and male pattern baldness. *Clin Endocrinol* 1992; 38: 653-8.
298. Carey AH, Waterworth D, Patel K, *et al.* Polycystic ovaries and premature male pattern baldness are associated with one allele of the steroid metabolism gene CYP17. *Hum Mol Genet* 1994; 3: 1873-6.
299. Dunaif A. Hyperandrogenic anovulation (PCOS): a unique disorder of insulin action associated with an increased risk of non-insulin-dependent diabetes mellitus. *Am J Med* 1995; 98: 1A-33S.
300. Prelevic GM, Wurzbarger MI, Balint-Peric I, Ginsburg J. Twenty-four-hour serum growth hormone, insulin, C-peptide and blood glucose profiles and serum insulin-like growth factor-I concentrations in women with polycystic ovaries. *Horm Res* 1992; 37: 125-31.
301. Kazer RR, Unterman TG, Glick RP. An abnormality of the growth hormone/insulin-like growth factor axis in women with polycystic ovary syndrome. *J Clin Endocrinol Metab* 1990; 71: 958-62.
302. Insler V, Shoham Z, Barash A, *et al.* Polycystic ovaries in non-obese and obese patients: possible pathophysiological mechanism based on new interpretation of facts and findings. *Hum Reprod* 1993; 8: 379-84.
303. Insler V, Barash A, Shoham Z, *et al.* Overnight secretion pattern of growth hormone, sex hormone-binding globulin, insulin-like growth factor I, and its binding protein in obese and non-obese women with polycystic ovarian disease. *Isr J Med Sci* 1994; 30: 42-7.
304. Kopelman PG, White N, Pilkington TR, Jeffcoate SL. The effect of weight loss on sex steroid secretion and binding in massively obese women. *Clin Endocrinol Oxf* 1981; 15: 113-6.
305. Bates GW, Whitworth NS. Effect of body weight reduction on plasma androgens in obese, infertile women. *Fertil Steril* 1982; 38: 406-9.
306. Harlass FE, Plymate SR, Fariss BL, Belts RP. Weight loss is associated with correction of gonadotropin and sex steroid abnormalities in the obese anovulatory female. *Fertil Steril* 1984; 42: 649-52.
307. Crave JC, Fimbel S, Lejeune H, Cugnardey N, Dechaud H, Pugeat M. Effects of diet and metformin administration on sex hormone-binding globulin, androgens, and insulin in hirsute and obese women. *J Clin Endocrinol Metab* 1995; 80: 2057-62.
308. Wabitsch M, Hauner H, Heinze E, *et al.* Body fat distribution and steroid hormone concentrations in obese adolescent girls before and after weight reduction. *J Clin Endocrinol Metab* 1995; 80: 3469-75.
309. Cross M, Dexter TM. Growth factors in development, transformation, and tumorigenesis. *Cell* 1991; 64: 271-80.
310. Baserga R, Porcu P, Sell C. Oncogenes, growth factors and control of the cell cycle. *Cancer Surv* 1993; 16: 201-13.
311. King RJB. A discussion of the roles of oestrogen and progesterin in human mammary carcinogenesis. *J Steroid Biochem Molec Biol* 1991; 39: 811-8.
312. Shi YE, Liu YE, Lippman ME, Dickson RB. Progestins and antiprogesterins in mammary tumour growth and metastasis. *Hum Reprod* 1994; 9: 162-73.
313. Henderson BE, Ross R, Bernstein L. Estrogens as a cause of human cancer: the Richard and Hinda Rosenthal Foundation Award lecture. *Cancer Res* 1988; 48: 246-53.
314. Fernig DG, Smith JA, Rudland PS. Relationship of growth factors and differentiation and neoplastic development of the mammary gland. In: Lippman M, Dickson R, eds. *Regulatory Mechanisms in Breast Cancer*. Boston, MA (USA): Kluwer Publishers, 1991: 47-78.
315. Henderson BE, Ross RK, Judd HL, Krailo MD, Pike MC. Do regular ovulatory cycles increase breast cancer risk? *Cancer* 1985; 56: 1206-8.

316. Spicer DV, Pike MC. The prevention of breast cancer through reduced ovarian steroid exposure. *Acta Oncologica* 1992; 31: 167-74.
317. Pike MC, Spicer DV, Dahmouch L, Press MF. Estrogens, progestagens, normal breast cell proliferation, and breast cancer risk. *Epidemiol Rev* 1993; 15: 17-35.
318. Key TJA, Pike MC. The dose-effect relationship between 'unopposed' estrogens and endometrial mitotic rate: its central role in explaining and predicting endometrial cancer risk. *Br J Cancer* 1988; 57: 205-12.
319. Macaulay VM. Insulin-like growth factors and cancer. *Br J Cancer* 1992; 65: 311-20.
320. Yee D, Rosen N, Favoni RE, Cullen KJ. The insulin-like growth factors, their receptors, and their binding proteins in human breast cancer. In: Lippman, Dickson, eds. *Regulatory Mechanisms in Breast Cancer*. Boston, MA (USA): Kluwer Publishers 1991: 93-105.
321. Hilf R. The actions of insulin as a hormonal factor in breast cancer. In: Pike MC, Siiteri PK, Welsch CW, eds. *Hormones and Breast Cancer*. Cold Spring Harbor, NY (USA): Cold Spring Harbor Laboratory, 1981; Banbury Report No. 8: 317-37.
322. Stoll BA. Breast cancer: the obesity connection. *Br J Cancer* 1994; 69: 799-801.
323. Bruning PF, Bonfrère JMG, van Noord PAH, Hart AAM, de Jong-Bakker M, Nooijen WJ. Insulin resistance and breast cancer risk. *Int J Cancer* 1992; 52: 511-6.
324. Adami HO, Rimsten A. Prevalence of hypertension and diabetes in breast cancer: a case-control study in 179 patients and age-matched, non-hospitalized controls. *Clin Oncol* 1978; 4: 243-9.
325. Ragozzino M, Melton LJ, Chu CP, Palumbo PJ. Subsequent cancer risk in the incidence cohort of Rochester, Minnesota, residents with diabetes mellitus. *J Chron Dis* 1982; 35: 13-9.
326. O'Mara BA, Byers T, Schoenfeld E. Diabetes mellitus and cancer risk: a multi-site case-control study. *J Chron Dis* 1985; 38: 435-41.
327. Franceschi S, la Vecchia C, Negri E, Parazzini F, Boyle P. Breast cancer risk and history of selected medical conditions linked with female hormones. *Eur J Cancer* 1990; 26: 781-5.
328. Adami HO, McLaughlin J, Ekblom A, et al. Cancer risk in patients with diabetes mellitus. *Cancer Causes Control* 1991; 2: 307-14.
329. Moseson M, Koenig KL, Shore RE, Pasternack BS. The influence of medical conditions associated with hormones on the risk of breast cancer. *Int J Epidemiol* 1993; 22: 1000-9.
330. La Vecchia C, Negri E, Franceschi S, D'Avanzo B, Boyle P. A case-control study of diabetes mellitus and cancer risk. *Br J Cancer* 1994; 70: 950-3.
331. Andersson B, Marin P, Lissner L, Vermeulen A, Björntorp P. Testosterone concentrations in women and men with NIDDM. *Diabetes Care* 1994; 17: 405-11.
332. Nyholm H, Djursing H, Hagen C, Agner T, Bennett P, Sventrup B. Androgens and estrogens in postmenopausal insulin-treated diabetic women. *J Clin Endocrinol Metab* 1989; 69: 946-9.
333. Djursing H, Hagen C, Nyboe-Andersen A, Sventrup B, Bennet P, Mølsted-Pedersen L. Serum sex hormone concentrations in insulin dependent diabetic women with and without amenorrhea. *Clin Endocrinol* 1985; 23: 147-54.
334. Prelevic GM, Wurzbarger MI, Peric LA. The effect of residual beta-cell activity on menstruation and the reproductive hormone profile of insulin-dependent diabetics. *Arch Gynecol Obstet* 1989; 244: 207-13.
335. Alexander L, Appleton D, Hall R, Ross WM, Wilkinson R. Epidemiology of acromegaly in the Newcastle region. *Clin Endocrinol Oxf* 1980; 12: 71-9.
336. Nabarro JDN. Acromegaly. *Clin Endocrinol Oxf* 1987; 26: 481-512.
337. Bengtsson BÅ, Edén S, Ernest I, Odén A, Sjögren B. Epidemiology and long-term survival in acromegaly. A study of 166 cases diagnosed between 1955 and 1984. *Acta Med Scand* 1988; 223: 327-35.
338. Ron E, Gridley G, Hrubec Z, Page W, Arora S, Fraumeni JF, Jr. Acromegaly and gastrointestinal cancer. *Cancer* 1991; 68: 1673-7.
339. Barzilay J, Heatley GJ, Cushing GW. Benign and malignant tumors in patients with acromegaly. *Arch Intern Med* 1991; 151: 1629-32.
340. Klein I, Parveen G, Gavalier JS, Vanthiel DH. Colonic polyps in patients with acromegaly. *Ann Intern Med* 1982; 97: 27-30.
341. Ituarte EM, Petrini J, Hershman JM. Acromegaly and colon cancer. *Ann Intern Med* 1984; 101: 627-8.
342. Pines A, Rozen P, Ron E, Gilat T. Gastrointestinal tumors in acromegalic patients. *Am J Gastroenterol* 1985; 80: 266-9.
343. Ritter MM, Richter WO, Schwandt P. Acromegaly and colon cancer. *Ann Intern Med* 1987; 106: 636-7.
344. Ziel FH, Peters AL. Acromegaly and gastrointestinal adenocarcinomas. *Ann Intern Med* 1988; 109: 514-5.
345. Brunner JE, Johnson CC, Zafar S, Peterson EL, Brunner JF, Mellinger RC. Colon cancer and polyps in acromegaly: increased risk associated with family history of colon cancer. *Clin Endocrinol Oxf* 1990; 32: 65-71.
346. Ezzat S, Strom C, Melmed S. Colon polyps in acromegaly. *Ann Int Med* 1991; 114: 754-5.
347. Wynder EL, Hebert JR. Homogeneity and nutritional exposure: an impediment in cancer epidemiology? *JNCI* 1987; 79: 605-7.
348. Giovannucci E. Insulin and colon cancer. *Cancer Causes Control* 1995; 6: 164-79.
349. McKeown-Eyssen G. Epidemiology of colorectal cancer revisited: are serum triglycerides and/or plasma glucose associated with risk? *Cancer Epidemiol Biomark Prev* 1994; 3: 687-95.
350. Siiteri PK, MacDonald PC. The role of extraglandular estrogen in human endocrinology. In: Geiger SR, Astwood EB, Geep RO, eds. *Handbook of Physiology*. New York, NY (USA): American Physiological Society, 1973: 615-29.
351. Siiteri PK. Obesity and peripheral estrogen synthesis. In: Frisch RE, ed. *Adipose Tissue and Reproduction*. Basel, Switzerland: Karger 1990; *Progr Reprod Biol Med*: 14: 70-84.
352. Coulam CB, Annegers JF, Kranz JS. Chronic anovulation syndrome and associated neoplasia. *Obstet Gynecol* 1983; 61: 403-7.
353. Gammon MD, Thompson WD. Polycystic ovaries and the risk of breast cancer. *Am J Epidemiol* 1991; 134: 818-24.
354. Ibanez L, Potau N, Virdis R, et al. Postpubertal outcome in girls diagnosed of premature pubarche during childhood: increased frequency of functional ovarian hyperandrogenism. *J Clin Endocrinol Metab* 1993; 76: 1599-603.
355. Lazar L, Kauli R, Bruchis C, Nordenberg J, Galatzer A, Pertzlan A. Early polycystic ovary-like syndrome in girls with central precocious puberty and exaggerated adrenal response. *Eur J Endocrinol* 1995; 133: 403-6.