

Diagnostic Dosages of Protirelin (TRH) Elevate BP by Noncatecholamine Mechanisms

Gary P. Zaloga, MD; Bart Chernow, MD; Russ Zajtcuk, MD; Robert Chin, MD; Thomas G. Rainey, MD; C. Raymond Lake, MD, PhD

• While performing thyroid function tests, we noticed that protirelin (TRH) raised BP, and, therefore, we investigated the effect of diagnostic dosages of protirelin (500 µg) on plasma catecholamine levels and cardiovascular function in eight patients one day before, one day after, and four weeks following heart surgery. Mean arterial pressure (MAP), heart rate (HR), plasma norepinephrine (NE), epinephrine (EPI), dopamine (DA), thyroid hormone (triiodothyronine [T_3], thyroxine), and thyrotropin (TSH) levels were measured before and after the intravenous injection of protirelin. Protirelin increased MAP transiently from 88 ± 2 to 103 ± 3 mm Hg (before surgery), 86 ± 4 to 102 ± 4 mm Hg (one day after surgery), and 86 ± 4 to 104 ± 5 mm Hg (four weeks after surgery). There were no notable changes in HR or plasma NE, EPI, or DA levels. The T_3 and TSH response to protirelin was normal on all three study days. Protirelin raised MAP by an effect on systemic vascular resistance (SVR) rather than an increase in cardiac output. We conclude the following: (1) diagnostic dosages of protirelin transiently elevate MAP and SVR by a noncatecholamine mechanism, (2) clinicians who perform protirelin tests should be aware of protirelin's transient pressor effects.

(Arch Intern Med 1984;144:1149-1152)

Thyrotropin-releasing hormone (TRH) was originally identified as the hypothalamic factor regulating the pituitary gland's secretion of thyrotropin (TSH).¹ The function of TRH was thought to relate solely to thyroid gland homeostasis. Further work, however, demonstrated TRH to be widely distributed throughout the CNS, retina, pancreas, and gastrointestinal tract.²⁻⁷ In addition, its concentration in synaptosomes,^{3,8,9} its analeptic (brain-activating) properties,^{5,7,10} its effect on the electrical activity of single neurons,^{7,11} and its antagonism of opioid effects^{4,5,12,13} led to the concept that TRH is a neurotransmitter or modulator of neurotransmission in addition to its neurohumoral actions.

For editorial comment see p 1138.

Endogenous opioid peptides are implicated in the pathogenesis of shock states¹³⁻¹⁵ and naloxone hydrochloride, an opioid receptor antagonist, improves cardiovascular function and survival in experimental endotoxin,¹⁶⁻¹⁸ hemorrhagic,^{19,20} and spinal²¹⁻²³ shock. The therapeutic effects of naloxone appear to be mediated by the autonomic nervous system. In spinal shock, naloxone acts centrally^{21,24} on opiate receptors to reverse endorphin effects on parasympathetic outflow. Release of dopamine (DA) also plays a role in the therapeutic effects of naloxone in spinal shock.²² Like

naloxone, protirelin (TRH) antagonizes many opioid effects (except antinociception)^{2,4,5,12,13,25} and improves hypotension and survival in experimental endotoxin,²⁶⁻²⁸ hemorrhagic,^{26,29} and spinal³⁰ shock. Naloxone and protirelin differ in that protirelin elevates BP in unstressed normotensive individuals and does not act at the opioid receptor.¹⁴ Thus, protirelin is termed a physiologic opiate antagonist. The mechanism of action of protirelin in shock states remains unknown. A great portion of protirelin's beneficial effects undoubtedly relate to its cardiovascular effects. Some^{31,32} suggest that the cardiovascular effects of protirelin are catecholamine mediated in that plasma norepinephrine (NE) increases following protirelin administration, while others^{7,33,34} think that its effects are independent of adrenergic mechanisms.

Since protirelin or protirelin analogues may have potential therapeutic uses in critically ill patients, we chose to investigate its action before and after surgery in these patients. We specifically were interested in the effect of standard diagnostic dosages (standard dosage of 500 µg used in the evaluation of the hypothalamic-pituitary-thyroid axis) of protirelin on plasma catecholamine levels and cardiovascular function.

PATIENTS AND METHODS

Eight adult patients (age range, 40 to 68 years) (six men, two women) undergoing coronary artery bypass surgery were prospectively studied one day before, one day after, and four weeks following surgery. Patients were studied in the supine position after an overnight fast (except medications). An intravenous (IV) catheter was placed in a vein in the antecubital fossa allowing for blood collection. Arterial pressure was monitored by sphygmomanometer (one day before and four weeks after surgery) or indwelling arterial catheter (one day after surgery). Heart rate (HR) was determined by continuous ECG monitoring. Mean arterial pressure (MAP) was calculated as diastolic pressure plus one third of the pulse pressure (systolic minus diastolic). Following a 20-minute rest period, MAP and HR were measured, and blood samples for plasma catecholamine (NE, epinephrine [EPI], and DA) and thyroid hormone determinations (triiodothyronine [T_3], thyroxine [T_4], TSH) were drawn. This procedure was repeated at 1, 5, 30, and 60 minutes following the IV injection of a diagnostic dose of synthetic protirelin (Thyprinone, 500 µg). In addition, one day after surgery, we measured central venous pressure (CVP), pulmonary capillary wedge pressure, pulmonary artery pressure, and cardiac output (CO) using a thermodilution Swan-Ganz catheter; systemic vascular resistance (SVR, dyne/cm⁻⁵) was calculated from these measurements as $(MAP - CVP)/CO \times 79.9$. It is routine in our hospital for patients after coronary artery bypass surgery to have Swan-Ganz and radial artery catheters placed during surgery. This study was approved by the Protection of Human Subjects and Clinical Investigation Committees of our institutions.

Blood samples for measurement of plasma catecholamine concentrations were obtained as previously described.³⁵ Blood specimens were collected in chilled heparinized test tubes that were immediately placed on wet ice, centrifuged at 4 °C within 30 minutes of collection, and the plasma was separated and stored at -70 °C until assayed. Plasma catecholamine levels were measured by a radioenzymatic technique in which a partially purified enzyme, catechol-O-methyltransferase, catalyzes the transfer of a tritiated methyl group from commercially available tritiated S-adenosylmethionine to the methoxy group of endogenous NE, EPI, and DA, forming tritiated normetanephrine, tritiated metanephrine, and tritiated 3-methoxytyramine, respectively.³⁶

Accepted for publication Sept 20, 1983.

From the Departments of Internal Medicine (Drs Zaloga and Chin) and Critical Care Medicine (Drs Chernow and Rainey), Naval Hospital, Bethesda; the Department of Thoracic Surgery, Walter Reed Army Medical Center, Washington, DC (Dr Zajtcuk); and the Departments of Medicine (Drs Chernow and Rainey), Psychiatry (Dr Lake), Pharmacology (Dr Lake), and Thoracic Surgery (Dr Zajtcuk), Uniformed Services University of the Health Services, Bethesda, Md.

The opinions expressed herein are those of the authors and are not to be construed as reflecting the views of the Navy or Army Departments, of the Naval or Army Services at large, or of the Department of Defense.

Reprint requests to Box 129, Naval Hospital, Bethesda, MD 20814 (Dr Chernow).




Fig 1.—Mean arterial pressure (MAP) and heart rate (HR) responses to injections of protirelin (TRH) (500 µg) before, one day after, and four weeks after coronary artery surgery. After baseline readings (time 0), protirelin was given intravenously and measurements were made at 1, 5, 30, and 60 minutes. (Asterisk indicates $P < .005$ compared with time 0; double asterisk, $P < .02$ compared with time 0.)

Variable	Before Surgery	One Day After Surgery	Four Weeks After Surgery
Serum thyroxine level, µg/dL	7.0 \pm 0.8	4.5 \pm 0.3*	6.4 \pm 0.6
Serum triiodo-thyronine level, ng/dL	111.8 \pm 4.8	51.1 \pm 4.1*	112.0 \pm 9.0
Serum thyrotropin level, µU/mL	2.1 \pm 0.5	1.4 \pm 0.1†	2.1 \pm 0.3
Serum thyrotropin level 30 minutes after intravenous protirelin, µU/mL	8.4 \pm 1.2	7.8 \pm 0.9	9.6 \pm 1.5

* $P < .005$ compared with preoperative value.

† $P < .05$ compared with preoperative value.

These metabolites were separated by thin layer chromatography and quantified by liquid scintillation spectrometry (Packard Tri-carb 460C). The resultant measurements are accurate above catecholamine concentrations of 20 pg/mL.

Thyroid hormone levels were measured by radioimmunoassay. Serum ionized calcium levels were measured in three patients following protirelin injection using an ion selective electrode system (Nova 2).

Standard statistical methods were used to calculate means and SEMs. Data were analyzed by two-tailed paired Student's t tests.

RESULTS

The MAP increased significantly ($P < .005$) at one minute following protirelin injection (Fig 1), while the HR remained unchanged. Similar protirelin-induced increments in MAP (15 to 17 mm Hg) were seen before and after surgery. These increments in MAP occurred despite β -adrenergic blockade for angina pectoris before surgery (propranolol hydrochloride was being taken by six of the eight patients) and aspirin was used to maintain graft

Medications	Before Surgery	One Day After Surgery	Four Weeks After Surgery
Propranolol hydrochloride, mg/24 hr	256 \pm 27 n = 6	0	0
Isosorbide dinitrate, mg/24 hr	130 \pm 26 n = 3	0	0
Transdermal nitroglycerin	1 every day n = 1	0	0
Nifedipine, mg/24 hr	90 n = 1	0	0
Digoxin, mg/24 hr	0.25 n = 1	0	0
Aspirin, mg/24 hr	0	0	1,200 \pm 0 n = 6




Fig 2.—Thyrotropin (TSH) (µU/mL) response to protirelin (TRH) (500 µg) before, one day after, and four weeks after coronary artery surgery. After baseline readings (time 0), protirelin was given intravenously and measurements were made at 30, 60, 90, and 120 minutes.

patency following surgery (given to six of eight patients) (Table 1). All patients were clinically and chemically euthyroid before surgery (Table 2). One day after surgery, patients fulfilled the criteria for the "sick euthyroid syndrome" (low serum T_3 and T_4 levels, normal or low TSH levels, and normal serum TSH response to protirelin^{87,88}) (Table 2, Fig 2). Despite raising MAP, protirelin caused no significant changes in plasma NE, EPI, or DA levels from baseline values (Fig 3). Plasma catecholamine levels were uniformly elevated one day after surgery ($P < .01$) when compared with preoperative levels and returned to normal by four weeks following surgery (Fig 3). Protirelin raised MAP significantly ($P < .005$) by increasing SVR (Table 3). There was no effect of protirelin on serum ionized calcium levels in the three patients in whom this variable was analyzed.

COMMENT

Unlike naloxone, protirelin-induced increases in MAP occurred by an increase in SVR rather than an increase in stroke volume.^{16,23} In contrast to Tuck et al³¹ and Morley et al,³² we found no significant increases in plasma NE, EPI, or DA concentrations following diagnostic dosages of protirelin. It is unlikely that a BP change of the magnitude seen




Fig 3.—Plasma norepinephrine, epinephrine, and dopamine concentrations in response to injection of diagnostic dosages of protirelin (TRH) (500 μ g) before, one day after, and four weeks after coronary artery surgery. After baseline samples were collected (time 0), protirelin was given intravenously and blood samples were collected at 1, 5, 30, and 60 minutes. Plasma norepinephrine, epinephrine, and dopamine were significantly elevated one day after surgery ($P < .01$) compared with before surgery and returned to baseline by four weeks following surgery.

in this study (15 to 17 mm Hg) would be mediated by a change in catecholamine release without detectable changes in plasma catecholamine levels and strongly suggests (but does not prove) that the hemodynamic effects of protirelin are noncatecholamine mediated. If patients in this study had decreasing baseline catecholamine levels, protirelin may have kept the baseline values constant by causing the release of small amounts of catecholamines (rather than allowing them to decrease). This small change would not explain the raised BP but could explain the difference between our results and those of Tuck et al³¹ and Morley et al,³² who had lower baseline NE concentrations (approximately 160 pg/mL) that increased slightly (200 to 275 pg/mL) following protirelin.

Plasma NE has been used as an index of sympathetic nervous system activity.³⁹⁻⁴² Lack of an increase in plasma NE levels argues against sympathetic mediation of protirelin's pressor effects. Beale et al³³ injected protirelin into the cisterna magna of rabbits and observed increases in BP that were not antagonized by chlorpromazine hydrochloride (a central inhibitor of DA and α -adrenergic receptors). Horita and Carino³⁴ also demonstrated a pressor response to intraventricular injection of protirelin in rabbits that was not blocked by phenoxybenzamine hydrochloride (α -adrenergic blocker), guanethidine sulfate (catecholamine-depleting agent), reserpine (catecholamine-depleting agent), or atropine sulfate (antimuscarinic agent).³⁴ We found no effect of propranolol on TRH's pressor response in humans.

Table 3.—Changes in Hemodynamic Variables After Protirelin Injection

Variable*	Time After Injection, min			
	0	1	5	30
MAP, mm Hg	86 \pm 4	103 \pm 4†	89 \pm 5	87 \pm 6
CVP, mm Hg	9.3 \pm 1.1	10.3 \pm 1.2	10.0 \pm 1.3	9.4 \pm 1.1
PCWP, mm Hg	12.4 \pm 0.6	13.0 \pm 0.9	12.5 \pm 1.0	11.6 \pm 1.1
CO, mm Hg	5.0 \pm 0.3	4.8 \pm 0.3	4.9 \pm 0.3	4.8 \pm 0.3
SVR, dyne/s/cm ⁻⁵	1276 \pm 108	1579 \pm 88†	1301 \pm 105	1300 \pm 138

*MAP indicates mean arterial pressure; CVP, central venous pressure; PCWP, pulmonary capillary wedge pressure; CO, cardiac output; and SVR, systemic vascular resistance.

† $P < .005$.

In the study of Horita and Carino,³⁴ the TRH pressor response remained intact in animals with spinal transections below T1 but transections in the cervical cord region abolished the response. MK-771, a protirelin analogue, raises BP in anesthetized cats³⁵ by an effect that is not antagonized by mecamylamine hydrochloride, a ganglionic blocking agent. These data taken together suggest that the pressor response to protirelin is centrally mediated but not expressed through classic autonomic pathways. The pressor response to intraventricular protirelin in endotoxemic rats is abolished by adrenal demedullation.⁴³ However, IV protirelin still produces a pressor response in these animals. Therefore, protirelin appears to increase cardiovascular values via the adrenal medullary system as well as some other effector system.

The pressor response to protirelin was unaffected by the use of aspirin (600 mg orally twice a day in six of eight patients). Since aspirin acetylates and deactivates the cyclo-oxygenase pathway involved in prostaglandin production,^{44,45} these data argue against prostaglandin mediation of the TRH pressor response.

Hypercalcemia may raise BP^{46,47} and thus we investigated the effect of protirelin on ionized calcium levels in three patients. No effect was found.

Protirelin causes release of a variety of hormones including TSH, prolactin, follicle-stimulating hormone, growth hormone (in acromegaly), and glucagon.⁴⁵ Release of these hormones occurs too slowly to explain the pressor effect of protirelin. Goldstein and Pavel⁴⁸ demonstrated release of vasotocin following intracarotid injection of protirelin in cats. Although they do not report hemodynamic variables, CSF vasotocin activity greatly increased within five minutes of protirelin injection. The role of vasotocin in the TRH pressor response is unknown.

Evidence favors a beneficial effect of naloxone hydrochloride¹⁶⁻²³ and protirelin²⁶⁻³⁰ in pharmacologic doses (2 mg/kg/hr) in shock states and suggests that these opioid antagonists favorably increase blood flow to vital organs and improve survival from experimental shock. Naloxone (in pharmacologic dosages) acts at cardiovascular sites within the CNS and/or at peripheral sites to improve myocardial contractility in experimental shock,^{13,14,21,22,24} is associated with DA release,²² and its action depends on adrenal medullary integrity.⁴⁴ Protirelin (a physiologic opiate antagonist)^{13,14} elevates BP in both normotensive and hypotensive individuals. As shown by our data, diagnostic dosages of protirelin increase BP without increasing plasma catecholamine levels or CO. Preliminary evidence⁴⁹ suggests that protirelin and naloxone have additive effects

when used together in the treatment of experimental shock. Although naloxone reverses shock resulting from a variety of causes, it may also potentiate pain processes by blocking opiate receptors. The potential advantage of protirelin over naloxone is that it does not act at opiate receptors and, hence, it does not antagonize antinociception.¹⁴

Protirelin may work via opioid antagonism, release of an unknown humoral substance, or antagonism of some hypotensive factor. Protirelin is effective when given orally at 20 to 30 times the IV dose⁵ and may have potential use in treating shock in field situations. It is not affected by adrenergic or cholinergic blockade^{33,34} and may be useful in the hypotensive patient taking propranolol. Future work into the mechanisms by which protirelin and naloxone work in shock may provide us with both new insights into the pathophysiology of these disease states as well as new therapeutic tools. These agents are not approved for human use in shock; use should be restricted to controlled trials.

We have demonstrated a similar (15- to 20-mm Hg) pressor response in patients with essential hypertension. Clinicians who perform protirelin tests should become aware of the pressor effect described in this report and BP should be measured before and after protirelin injection, especially in patients with hypertension or cardiac disease.

Evaluation of thyroid function in patients undergoing cardiac surgery is the subject of a future report. However, preliminary evidence suggests that TSH and the TSH response to protirelin (Fig 2) remain intact following cardiac surgery and may be of value in ruling out hypothyroidism in these patients.

This work was supported in part by project CIC 82-06-1645 from the Bureau of Medicine and Surgery, Navy Department, Washington, DC.

Kenneth Burman, MD, reviewed the manuscript and Dominic S. D'Angona, Audrey Reid, Duke Owen, and Rose Quirk provided technical support in performing the radioenzymatic assay.

References

- Burgus R, Dunn TE, Desiderio D, et al: Characteristics of bovine hypothalamic hypophysiotropic TSH-releasing factor. *Nature* 1970;336:321-325.
- Jackson I, Reichlin S: Thyrotropin-releasing hormone (TRH): Distribution in hypothalamic and extrahypothalamic brain tissues of mammalian and submammalian chordates. *Endocrinology* 1974;95:854-862.
- Winokur A, Utiger RD: Thyrotropin-releasing hormone: Regional distribution in rat brain. *Science* 1974;185:265-267.
- Morley JE: Extrahypothalamic thyrotropin releasing hormone (TRH): Its distribution and its functions. *Life Sci* 1979;25:1539-1550.
- Jackson IMD: Thyrotropin-releasing hormone. *N Engl J Med* 1982;306:145-155.
- Morley JE: Neuroendocrine control of thyrotropin secretion. *Endocr Rev* 1981;2:396-436.
- Yarbrough GG: On the neuropharmacology of thyrotropin releasing hormone. *Prog Neuro* 1979;12:291-312.
- Barnes A, Ben-Jonathan N, Colston C, et al: Differential subcellular compartmentalization of thyrotropin releasing hormone (TRH) and gonadotropin releasing hormone (LRH) in hypothalamic tissue. *Proc Natl Acad Sci USA* 1975;8:3153-3157.
- Winokur A, Davis R, Utiger RD: Subcellular distribution of thyrotropin releasing hormone (TRH) in rat brain and hypothalamus. *Brain Res* 1977;120:423-434.
- Metcalfe G, Dettmar PW: Is thyrotropin releasing hormone an endogenous ergotropic substance in the brain? *Lancet* 1981;1:586-589.
- Winokur A, Beckman AL: Effects of thyrotropin releasing hormone, norepinephrine and acetylcholine on the activity of neurons in the hypothalamus, septum and cerebral cortex of the rat. *Brain Res* 1978;150:205-209.
- Morley JE, Levine AS: Thyrotropin releasing hormone (TRH) suppresses stress induced eating. *Life Sci* 1980;27:269-274.
- Faden AI: Neuropeptides in shock and CNS injury: Potential clinical implications, in Chernow B, Lake CR (eds): *Pharmacologic Approach to the Critically Ill*. Baltimore, Williams & Wilkins Co, 1983, pp 636-649.
- Zaloga GP, Hostinsky C, Chernow B: Endogenous opioid peptides: Critical care implications. *Heart Lung*, in press.
- Chernow B, Fletcher JR: Catecholamines, opioid peptides and prostaglandins: Implications for critical care medicine, in Shoemaker WC, Thompson WL (eds): *Critical Care: State of the Art*. Fullerton, Calif, Society of Critical Care Medicine, 1983, vol 4, E1-E52.
- Holaday JW, Faden AI: Naloxone reversal of endotoxin hypotension suggests role of endorphins in shock. *Nature* 1978;275:450-451.
- Reynolds DG, Gurll NJ, Vargish T, et al: Blockade of opiate receptors with naloxone improves survival and cardiac performance in canine endotoxic shock. *Circ Res* 1980;7:39-48.
- Faden AI, Holaday JW: Experimental endotoxin shock: The pathophysiological function of endorphins and treatment with opiate antagonists. *J Infect Dis* 1980;142:229-238.
- Faden AI, Holaday JW: Opiate antagonists: A role in the treatment of hypovolemic shock. *Science* 1979;205:317-318.
- Vargish T, Reynolds DG, Gurll NJ, et al: Naloxone reversal of hypovolemic shock in dogs. *Circ Shock* 1980;7:31-38.
- Holaday JW, Faden AI: Naloxone acts at central opiate receptors to reverse hypotension, hyperthermia and hypoventilation in spinal shock. *Brain Res* 1980;189:295-299.
- Faden AI, Jacobs TP, Feuerstein G, et al: Dopamine partially mediates the cardiovascular effects of naloxone after spinal injury. *Brain Res* 1981;213:415-421.
- Faden AI, Jacobs TP, Holaday JW: Opiate antagonist improves neurologic recovery after spinal injury. *Science* 1981;211:493-494.
- Faden AI, Jacobs TP, Holaday JW: Endorphin-parasympathetic interaction in spinal shock. *J Auto Nerv Syst* 1980;2:295-302.
- Holaday JW, Tseng LF, Loh HH, et al: Thyrotropin releasing hormone antagonizes beta endorphin hypothermia and catalepsy. *Life Sci* 1978;22:1537-1544.
- Holaday JW, D'Amato R, Faden AI: Thyrotropin-releasing hormone improves cardiovascular function in experimental endotoxemic and hemorrhagic shock. *Science* 1981;213:216-218.
- Holaday JW, Ruvio BA, Faden AI: Thyrotropin-releasing hormone improves blood pressure and survival in endotoxic shock. *Eur J Pharmacol* 1981;74:101-105.
- Reynolds DG, Gurll NJ, Holaday JW, et al: Thyrotropin-releasing hormone (TRH) in primate endotoxic shock, abstracted. *Physiologist* 1982;25:309.
- Gurll NJ, Reynolds DG, Holaday JW, et al: Improved cardiovascular function and survival using thyrotropin releasing hormone (TRH) in primate hemorrhagic shock, abstracted. *Physiologist* 1982;25:342.
- Faden AI, Jacobs TP, Holaday JW: Thyrotropin-releasing hormone improves neurologic recovery after spinal trauma in cats. *N Engl J Med* 1981;305:1063-1067.
- Tuck M, Morley JE, Mayes D, et al: Thyrotropin releasing hormone (TRH) releases norepinephrine in man, abstracted. *Clin Res* 1979;27:261.
- Morley JE, Tuck ML, Mayes DM, et al: Thyrotropin-releasing hormone increases plasma norepinephrine in man. *Horm Res* 1981;14:18-23.
- Beale JS, White RP, Huang SP: EEG and blood pressure effects of TRH in rabbits. *Neuropharmacology* 1977;16:499-506.
- Horita A, Carino MA: Centrally administered TRH produces a vasopressor response in rabbits. *Proc West Pharmacol Soc* 1977;20:303-304.
- Lake CR, Ziegler MG, Kopin IJ: Use of plasma norepinephrine for evaluation of sympathetic neuronal function in man. *Life Sci* 1976;18:1315-1326.
- Durett LR, Ziegler MG: A sensitive radioenzymatic assay for catechol drugs. *J Neurosci Res* 1980;5:587-591.
- Wartofsky L, Burman K: Alterations in thyroid function in patients with systemic illness: The 'euthyroid sick syndrome'. *Endocr Rev* 1982;3:164-215.
- Kaptein EM: Interpretation and use of thyroid function tests in nonthyroidal disease. *Intern Med Specialist* 1982;3:102-106.
- Goldstein DS: Plasma norepinephrine as an indicator of sympathetic neural activity in clinical cardiology. *Am J Med* 1981;48:1147-1154.
- Burke D, Sundlof G, Wallin BG: Postural effects on muscle nerve synaptic activity in man. *J Physiol* 1977;272:399-414.
- Yamaguchi I, Kopin IJ: Direct stimulation of sympathetic outflow and plasma levels of norepinephrine and epinephrine in pithed rats, in Barches J, Kopin IJ, Usdin E (eds): *Catecholamines: Basic and Clinical Frontiers*. New York, Pergamon Press, 1979, pp 924-926.
- Kopin IJ, Lake CR, Ziegler M: Plasma levels of norepinephrine. *Ann Intern Med* 1978;88:671-680.
- Holaday JW, D'Amato RJ, Ruvio BA, et al: Action of naloxone and TRH on the autonomic regulation of circulation. *Adv Biochem Psychopharmacol* 1982;33:353-361.
- Robertson RP: Prostaglandins, thromboxanes, and eicosanoids: Arachidonic acid metabolites relevant to medicine, in Isselbacher KJ, Adams RD, Braunwald E, et al (eds): *Harrison's Principles of Internal Medicine: Update 1*. New York, McGraw-Hill Book Co, 1981.
- Flower RJ: Drugs which inhibit prostaglandin biosynthesis. *Pharmacol Rev* 1974;26:33-67.
- Agus ZS, Wasserstein A, Goldfarb S: Disorders of calcium and magnesium homeostasis. *Am J Med* 1982;72:473-488.
- Singer FR, Bethune JE, Massry SG: Hypercalcemia and hypocalcemia. *Clin Nephrol* 1977;7:154-162.
- Goldstein R, Pavel S: Vasotocin release into the cerebrospinal fluid of cats induced by luteinizing hormone releasing hormone, thyrotropin releasing hormone and growth hormone release-inhibitory hormone. *J Endocrinol* 1977;75:175-176.
- Holaday JW, Faden AI: Naloxone and thyrotropin-releasing hormone (TRH) have additive effects in reversing endotoxin shock, in Takagi H, et al (eds): *Advances in Endogenous and Exogenous Opioids*. New York, Pergamon Press, in press.