

## Cellular pathways leading to melanoma differentiation: therapeutic implications

P. W. Soballe and M. Herlyn\*

The Wistar Institute of Anatomy and Biology, 3601 Spruce Street, Philadelphia PA 19104 USA. Fax: (+1) 215 898 0980.

**The induction of differentiation, as evidenced by benign growth characteristics, dendritic morphology, pigmentation capability, and a mature antigenic phenotype, is an attractive theoretical basis for therapy in human melanoma. Melanoma differentiation can be experimentally induced by modulating intracellular pathways involving protein kinase C, tyrosine kinases, and protein kinase A, or by modulating nuclear transcription with retinoids, DNA-damaging agents, and chemotherapeutic drugs. Other experimental differentiating agents include the amino acid tyrosine, histamine receptor antagonists, polyamine antagonists, dimethylsulphoxide, caffeic acid ester, and butyrate. The mechanisms involved in the actions of many of these agents are beginning to be understood and the pathways are often intersecting; cross-talk in the form of negative and positive feedback loops is extensive. Uncoupling of pathways is also seen, with some agents leading to simultaneous increases in both differentiated and transformed characteristics. While clinical benefits of this approach have so far been sparse, greater understanding of the cellular pathways of differentiation may open new therapeutic options in melanoma.**

**Key words:** Differentiation, melanocyte, melanoma, transcription factor, transformation, protein kinase A, protein kinase C, tyrosine kinase.

Cancer is histopathologically graded according to the cytological state of differentiation: the less differentiated, the more 'atypical' and the higher the grade; the more a tumour cell resembles a normal cell of its own lineage, the less malignant its properties. The apparent 'de-differentiation' of tumour cells may actually reflect their origin from earlier benign precursors with a superimposed malignant phenotype, and reversion to this benign histotype can be directed in some tumours.<sup>1</sup> Alternatively, therapeutic driving of transformed cells to terminal differentiation, while potentially impairing normal processes such as wound healing, may result in little non-specific killing of normal cells and may thus offer a favourable therapeutic strategy. In human melanocytic tumours, spontaneous complete regression due to terminal differentiation is commonly ob-

served in naevi<sup>2,3</sup> and occasionally in melanomas.<sup>4</sup> Partial spontaneous regression of melanoma is often seen clinically but may actually portend a poor prognosis,<sup>3</sup> suggesting the persistence and selection of non-differentiating, more aggressively proliferating subpopulations. Any therapeutic induction of differentiation will clearly need to be essentially complete and permanent.

Figure 1 depicts current conceptions of melanocytic differentiation. Primary differentiation is an embryologic developmental phenomenon: melanocytes which express the receptor encoded by *c-kit*<sup>5</sup> migrate from the neural crest to the basal epidermis, where the embryonic microenvironment may effect their histotypic commitment as evidenced by a reversion to a benign phenotype in transplanted murine melanoma.<sup>6</sup> There is currently no evidence of a melanocyte stem cell in the skin, although the melanoblast has been defined as a cell which contains premelanosomes but does not express tyrosinase or melanin.<sup>7</sup> 'Premelanocytes'<sup>8</sup> have been identified which are apparently more differentiated and do express the tyrosinase gene. Undifferentiated melanocytes might be found to have characteristics similar to those of the keratinocyte precursor: undifferentiated morphology, slow cycling but responsive to mitogenic stimuli, and localization in a protected, vascularized and innervated anatomical area.<sup>9</sup>

Differentiated melanocytes are antigenically different from their undifferentiated precursors and they are morphologically dendritic, less able to proliferate in response to stimuli, and contain melanosomes and tyrosinase activity. It should be noted, however, that melanin content, unlike tyrosinase, does not correlate with the stage of melanocyte differentiation,<sup>10</sup> nor does pigmentation itself correlate with melanoma differentiation either clinically<sup>11</sup> or *in vitro*.<sup>12</sup>


Houghton and co-workers,<sup>13</sup> noting difficulties with melanin, morphology, and growth characteristics as markers of melanoma differentiation, have proposed the categorization of melanomas based on antigenic phenotypes into stages which correlate with presumed normal melanocyte development. Many systems of antigenic markers have been proposed to distinguish the level of differentiation of

---

Supported in part by NCI grant CA 25874.

PWS is a research fellow in Surgical Oncology of the Health Sciences Education and Training Command of the U.S. Navy, Washington, D.C.

\*To whom correspondence should be addressed



**Figure 1.** Current concept of the stages of melanocytic differentiation. Left column: 'primary differentiation', maturation or developmental differentiation which correlates with anatomic locations. Middle column: 'de-differentiation', or transformation and tumour progression. Non-tumourigenic melanomas include radial growth phase tumours without invasive or metastatic competence. Right column: 're-differentiation' of transformed cells to either the melanocytic parental phenotype, a new (presumably related) phenotype, or to apoptosis.

both benign and malignant melanocytes.<sup>14-16</sup> Antigens present on both melanomas and melanocytes reflect commitment to the primary differentiation programme, while antigens present on melanomas but absent from melanocytes may reflect tumour origin from an earlier precursor, de-differentiation, or re-commitment to a new, non-melanocytic, phenotype ('re-differentiation', Figure 1). As examples of non-melanocyte phenotypes, desmoplastic melanomas<sup>4</sup> and dermal nevocytes may resemble Schwann cells morphologically and antigenically. Melanomas<sup>17</sup> and naevocytes<sup>18</sup> may also be induced to express a fibroblast-like phenotype.

A developmental concept related to differentiation is that of apoptosis. This genetically determined programme leads to proliferative incompetence, which in mammals involves inactivation of the gene *bcl-2*<sup>19</sup> and perhaps *mcl-1*,<sup>20</sup> and a characteristic cellular self-destructive cascade. However it is still not clear whether this process is the same as terminal differentiation.<sup>21,22</sup>

Melanocytic transformation can be defined by cellular phenotypic traits such as those summarized in Table 1.<sup>23</sup> Since the process of transformation results in apparent de-differentiation, these traits are often considered to be markers of the state of melanocytic differentiation. The transformed melanocytic phenotype can be reversed by fusion of melanoma cells with benign fibroblasts<sup>49</sup> or immortalized melanocytes,<sup>50</sup> suggesting that expression of the genetic programme of differentiation is blocked at multiple sites in melanoma and that the blocks are removed by fusion with the differentiated genome.<sup>51</sup> Cell fusions may also restore cellular transcriptional regulation, the loss of which originally leads to transformation.

The concept of a blocked critical junction in the differentiation programme has been developed in the *Drosophila lethal (2) giant larva* mutation which results in absence of a gene product.<sup>52</sup> This mutation has no effect on the organism unless the product is absent at precisely the right stage of larval development and the right anatomic

**Table 1.** Biological characteristics of melanocyte transformation

Marker <sup>a</sup>	Comments
Phorbol independence	Auto-production of PKC-dependent mitogens <sup>22,23</sup>
↓ Growth factor dependence	Autocrine bFGF, IL-8 and MGSA <sup>10</sup>
↑ Plasminogen activator	Autocrine bFGF → activation of bFGF receptor → protease secretion <sup>24</sup> → proliferation <sup>25</sup> and invasion <sup>26</sup>
↑ Glucose transport	High-affinity transporter in cell membranes → facilitated glycolysis, <sup>27</sup> ↑ dependence on glutamine <sup>28</sup>
Culture immortality	SV40-T transfection → immortality, partial transformation; <sup>29,30</sup> ras transfection → immortality, complete transformation <sup>31</sup>
↓ Anchorage dependence	Altered integrin profile: ↑ $\alpha 4$ , <sup>32</sup> ↑ $\beta 3$ , ↓ $\alpha 6$ <sup>33</sup> → src-related kinase, <sup>34</sup> cytoplasmic alkalization <sup>35</sup> → anchorage-like growth signal <sup>36</sup>
↓ Contact inhibition	↑ Muc-18, <sup>37</sup> ↑ GD3 ganglioside, <sup>11</sup> ↑ ICAM-1, <sup>38</sup> → deregulated contact-mediated growth signals
↓ Dendritic morphology	↓ Collagen, ↓ laminin, <sup>39</sup> ↓ fibronectin, <sup>40</sup> altered ganglioside expression, <sup>41,42</sup> → ↓ actin <sup>43</sup>
↑ Surface protein mobility	ABP-280 → submembranous cytoskeletal cross-links <sup>44</sup> Fibronectin receptor mobility → cell motility <sup>45-48</sup> ABP-280 → surface molecule mobility <sup>44</sup>

<sup>a</sup> ↓, decrease; ↑, increase; →, leading to.

location. If the gene is non-functional when needed, its transcriptional machinery is applied elsewhere and cellular transformation occurs.

## Pathways of experimental induction of differentiation in melanoma

Many agents have been shown to induce a more differentiated phenotype in melanoma. (Table 2) Few possess the level of efficacy, safety, and specificity needed clinically, but consideration of their mechanisms of action may focus further in-depth investigations.

### Signal transduction

**Protein kinase C (PKC).** Phorbol esters bind to the regulatory domain of most<sup>53</sup> but not all<sup>54</sup> members of the PKC family of serine-threonine kinases.<sup>55</sup> Long-term exposure of murine melanocytes and melanomas to phorbol esters results in PKC autophosphorylation, proteolysis, and migration to the cell membrane.<sup>56</sup> In human melanocytes chronically exposed to optimal mitogenic levels of phorbol ester, PKC activity is enhanced, whereas it is down-regulated after exposure to higher levels.<sup>57</sup> In human melanoma cells, the effect on PKC of chronic exposure to phorbol esters has not yet been established, and the pathways resulting in the observed differentiated melanoma phenotype following phorbol ester treatment<sup>58,59</sup> remain undefined. However there are possible therapeutic targets in

this system. For example, a major substrate for PKC is the 80 kD MARCKS (myristoylated alanine-rich C-kinase substrate) family of proteins, which has been implicated in a wide range of activities involving cytoskeletal rearrangements.<sup>60</sup> MARCKS proteins may be involved in establishing the differentiated (dendritic) morphology, anchorage and monolayer dependence, and cell-surface protein mobility observed after phorbol ester treatment.

**Protein tyrosine kinases (PTK).** PKC is activated by PTK phosphorylation of phospholipase-C $\gamma$ .<sup>61</sup> PKC in turn transmodulates (inactivates) the PTK cell-surface epidermal growth factor (EGF) receptor.<sup>62,63</sup> Thus an increase in PKC activity, such as that resulting from stimulation by EGF, decreases mitogenic signals from this PTK. Melanoma cells in culture are initially<sup>64</sup> mitogenically stimulated by EGF or the related transforming growth factor- $\alpha$  (TGF- $\alpha$ ), but become less responsive and independent after passage in culture;<sup>65</sup> however, whether these modified responses to EGF are related to PKC activity has not been studied.

Signal modulation by the PTK inhibitor genistein leads to melanoma differentiation.<sup>66</sup> However genistein also decreases intracellular polyamines and nitric oxide production and reverses the action of topoisomerase, which may contribute to the endonuclease-induced internucleosomal damage typical of apoptosis.<sup>67</sup> Thus the mechanism by which genistein affects differentiation remains unclear, and it probably acts at several levels.

Basic fibroblast growth factor (FGF), which plays a central role as a growth factor in the melanocyte system, must be supplied to melanocytes in culture and is pro-

**Table 2.** Melanoma differentiating agents

Agents	Effects and Comments <sup>a</sup>
<b>Signal transduction</b>	
Phorbol esters <sup>56,57</sup>	MARCKS; <sup>58</sup> ERK → AP-1; <sup>87</sup> block p34 <sup>cdc2</sup> & DNA synthesis <sup>85</sup>
Genistein <sup>64</sup>	↓ Tyrosine kinase; ↓ polyamines; ↓ topoisomerase
Antisense bFGF-R <sup>66</sup>	Phorbol esters partially reverse differentiation
Interferon-β <sup>67</sup>	With ↑ PKC → irreversible differentiation
Dibutyl cAMP <sup>73,74</sup>	↑ PKA; <sup>70,71</sup> gene modulation <sup>134</sup>
Theophylline <sup>73</sup>	↓ Phosphodiesterase → ↑ cAMP → ↑ PKA
Caffeine <sup>75</sup>	↓ Phosphodiesterase; ↓ cisplatin activity
MSH <sup>76-79</sup>	↑ Adenyl cyclase → ↑ cAMP in murine melanoma
GTP <sup>83</sup>	↑ Adenyl cyclase; ↑ G proteins
Mycophenolic acid, tizofurin <sup>84</sup>	↓ IMP dehydrogenase → ↑ GTP → ↑ G proteins
<b>Nuclear transcription</b>	
Retinoids <sup>95-97</sup>	↑ PKCα; <sup>94</sup> ↓ AP-1 responsive genes <sup>88</sup>
Beta carotene <sup>103</sup>	→ Retinoids; ↓ cAMP; O <sub>2</sub> radical scavenger <sup>104</sup>
[ <sup>3</sup> H]Thymidine <sup>106</sup>	Direct damage to DNA
BrdU <sup>15,107</sup>	Incorporated into DNA → gene modulation
Novobiocin <sup>108</sup>	↓ Topoisomerase <sup>65</sup>
Anthracyclines <sup>110,111</sup>	Damage microtubules; intercalate into DNA; <sup>112</sup> O <sub>2</sub> radicals <sup>113</sup>
AraC, 5-AzadC, aphidocolin hydroxyurea <sup>109</sup>	↓ DNA synthesis; ↑ apoptosis
<b>Other targets</b>	
Tyrosine <sup>114</sup>	↑ Tyrosinase but not melanin
Phenylacetate <sup>28</sup>	↓ Availability of glutamine; modulates genes
Antihistamines <sup>116</sup>	↑ PKC; ↑ PKA <sup>115</sup>
Antipolyamines <sup>123</sup>	↑ Tyrosinase; <sup>120,121</sup> ↑ apoptosis <sup>122</sup>
DMSO <sup>57,124</sup>	Polar molecule which effects multiple mechanisms
Caffeic acid ester <sup>130</sup>	→ Multiple antigen shifts
Butyrate <sup>124,133</sup>	↑ Histone H1 <sup>131,132</sup>

<sup>a</sup> →, leads to; ↑, increases; ↓, decreases.

duced for autocrine stimulation by melanomas.<sup>12</sup> Antisense inhibition of its PTK receptor leads to melanoma differentiation by undefined mechanisms which can be partly antagonized by phorbol ester.<sup>68</sup>

Interferons (IFN) have shown some limited clinical anti-melanoma potential; although interferons up-regulate expression of histocompatibility antigens, their variable non-immunological anti-proliferative effects *in vitro* are generally unrelated to differentiation.<sup>69</sup> However IFN-β which may be the most effective in terms of anti-melanoma activity,<sup>70</sup> when combined with PKC activation by the non-phorbol drug mezerein, results in irreversible differentiation of cultured melanoma cells.<sup>71</sup> IFN-β triggers a receptor which sends a transcription signal via a non-receptor PTK.<sup>61</sup> Interferon-induced signalling modifications combined with chemotherapeutic agents may thus offer future efficacious strategies.<sup>72</sup>

The ceramide system, which is associated with tumour necrosis factor-α (TNF-α), induces the intracellular PKC inhibitor sphingosin and also phosphorylates the EGF receptor.<sup>73</sup> This pathway leads to differentiation in promyelocytes,<sup>74</sup> and, possibly, also in melanoma cells.

**Cyclic AMP.** Many surface hormone receptors transmit their signals via cAMP. There are both positive and negative receptors and either may effect stimulatory or inhibitory guanine nucleotide regulatory (G) proteins at the cell membrane. G-proteins and the similar GTP-binding *ras* proteins control the activity of adenyl cyclase, which converts 5'AMP to cAMP. Adenyl cyclase and phosphodiesterase, which converts cAMP to 5'AMP, control the intracellular level of cAMP. Binding of two cAMP molecules to each of the two regulatory portions of protein kinase A (PKA) accounts for most of the cAMP signal transduction,<sup>75</sup> and substrate phosphorylation.<sup>76</sup>

The cAMP system interacts with PKC at several levels. PKC inhibits positive and some negative surface receptors; it inactivates the inhibitory G-proteins; it activates the catalytic unit of adenyl cyclase; and it inhibits phosphodiesterase. While most of these interactions tend to increase cAMP levels, the net effect is actually cell-specific and dependent on the expression of individual PKC isoforms.<sup>77</sup>

Many agents that increase cAMP levels lead to differentiation of murine melanoma *in vivo* and *in vitro*. Such

agents include the cAMP analogue dibutyryl cAMP,<sup>78,79</sup> the phosphodiesterase inhibitors theophylline<sup>78</sup> and caffeine,<sup>80</sup> and melanocyte stimulating hormone.<sup>81-84</sup> Elevation of cAMP levels in murine melanoma may lead to both a differentiated phenotype and an increased metastatic potential.<sup>81,82,84</sup> The dual effect might be related to varied levels of cross-talk among transcription factors.<sup>85</sup> The extent to which this simultaneous induction of opposing programmes occurs varies with individual cell lines, and its relevance to human melanoma remains unclear.

**Guanosine nucleotides.** Cyclic guanosine monophosphate (cGMP), the product of nitric oxide-induced guanylyl cyclase, is a differentiating agent for promyelocytes,<sup>86</sup> and the cGMP-dependent kinase shares many substrates with PKA. cGMP may either decrease or increase phosphodiesterase activity, and thus cAMP levels, in a specific cell type.<sup>87</sup> The role of cGMP in human melanoma biology has not yet been defined. Guanosine triphosphate (GTP) treatment results in differentiation of murine melanoma,<sup>88</sup> yet a decrease in GTP produced by inhibition of inosine monophosphate dehydrogenase in the purine synthesis pathway, results in differentiation of human melanoma cells.<sup>89</sup> These apparently conflicting results suggest that the species-specific activity of GTP may depend on either stimulatory or inhibitory G proteins and/or the similar *ras* GTP-binding proteins. Definition of any relationship between GTP levels and *ras* activity in melanoma would be interesting in view of the transforming characteristics of *ras* for human melanocytes<sup>33</sup> and the presence of *ras* mutations in approximately 15% of melanomas.<sup>12</sup>

## Nuclear transcription

**PKC.** At the nuclear level, phorbol esters block phosphorylation of p34<sup>cdc2</sup> in melanoma cells. This is a central regulator of the onset of DNA synthesis,<sup>90</sup> and the block appears to be due to the sustained activation of PKC. The resulting growth arrest is transient and the cells begin to proliferate 20 h after treatment. The ability to restore a regulated cell cycle by this mechanism suggests another potential therapeutic strategy.

Most, but not all, PKC subspecies target I $\kappa$ B, releasing the transcription factor NF- $\kappa$ B.<sup>91</sup> PKC also leads to expression of the generally mitogenic transcription factor AP-1 via phosphorylation of at least one member of the extracellular signal-regulated kinase family (ERKs, also called MAP kinases for mitogen-activated or membrane-associated protein), and phosphorylation of the regulatory portions of *c-jun*.<sup>92</sup> This modulation of transcription factors could account for the activation of proliferation programmes in

melanocytes and differentiation programmes in melanoma. These differences may reflect genomic mutation during transformation or cross-talk between structurally or expression-modified transcription factors, which may occur via overlapping, mutually exclusive DNA binding sites or by direct protein-protein interactions.<sup>93,94</sup> The precise transcriptional mechanisms by which PKC switches from driving a programme of proliferation in melanocytes to differentiation in melanomas is another promising area of investigation.

**cAMP.** PKA phosphorylates cAMP response element binding factors (CREBF), which modulate transcription of genes including *c-fos*, *c-jun*, glucose transporters,<sup>64</sup> and H1<sup>o</sup> histone.<sup>95</sup>

**Retinoids.** Transcription is also modulated by retinoids, which are all-*trans*-retinoic acid and its isomers and synthetic analogues.<sup>96,97</sup> Their actions are attenuated by cytoplasmic binding proteins and are mediated by nuclear receptors which are activated by the formation of dimers with themselves or with the receptors for various hormones.<sup>98</sup> These dimers increase the transcription of PKC $\alpha$ , leading to induction of differentiation in murine<sup>99</sup> and in some human<sup>100-102</sup> melanomas. Retinoids also decrease the transcription of AP-1-responsive genes by cross-coupling of the signal transduction pathways.<sup>93</sup> Thus retinoids increase both PKC and its phosphorylated targets while decreasing the mitogenic transcriptional effect of an important target. The therapeutic implications of this double action are evident, and retinoids have indeed been shown to have clinical potential in many malignancies,<sup>103,104</sup> including melanoma.<sup>105,106</sup> An increase in PKC transcription and simultaneous cross-talk between AP-1 and retinoic acid receptors results in a reversal of some of the transformed phenotypic properties of melanoma cells, except that a differentiated, dendritic morphology is not established.<sup>100,102</sup> This may argue against phosphorylation of MARCKS as a sufficient precursor event to cytoskeletal changes. Other actions of retinoids, for which the significance in melanoma has yet to be determined, include modulation of transcription of the homeobox family of transcriptional regulators and increased transcription of such possibly differentiation-related genes as H1<sup>o</sup> histone,<sup>95</sup> laminin  $\beta$ 1, *c-jun*, tissue plasminogen activator, and various growth factors.<sup>107</sup>

$\beta$ -Carotene, a natural source of vitamin A, is oxidized to retinoids, which may explain its differentiating effect in cultured melanomas.<sup>108</sup> Carotenoids are also potent scavengers of oxygen radicals, and this may allow them to protect the genome from DNA-damaging mutagens.<sup>109</sup> Both carotene and retinol decrease cellular activity

of adenylyl cyclase<sup>108</sup> suggesting a potential role for these compounds in the differentiation pathway related to cAMP.

**DNA.** Many agents that modify DNA have been implicated in melanoma differentiation. Triggering of DNA repair mechanisms or direct arrest of the cell cycle may result in apoptosis,<sup>21</sup> or may create a selective disadvantage, allowing the emergence of differentiating clones which are unaffected by the agent due to low initial levels of DNA synthesis. The actual mechanisms by which these agents induce differentiation are unknown and none have so far achieved therapeutically satisfactory anti-melanoma activity.

Direct injury to DNA can be induced by UV irradiation and by melanin itself,<sup>110</sup> but whether these agents result in differentiation is unknown. Direct damage by radioactive thymidine, however, has produced irreversible differentiation of a human melanoma cell line.<sup>111</sup>

The thymidine analogue 5'-bromodeoxyuridine (BrdU) is incorporated into DNA, where it can induce a gene encoding a specific *trans*-acting DNA-binding factor which then acts in tissue-specific fashion to suppress or enhance other genes.<sup>112</sup> In murine melanoma, BrdU reversibly reduces tumorigenicity and induces a flattened (not dendritic) morphology, increased anchorage dependence, decreased proliferation, and decreased transcription of tyrosinase.<sup>112</sup> Similar results were found in human melanoma along with markedly decreased transcription of *c-myc*.<sup>17</sup> These effects demonstrate induction of an incomplete or modified differentiation programme in melanoma and the ability to separate dendritic morphology and tyrosinase from other elements of melanocytic differentiation.

DNA synthesis can be inhibited by drugs such as novobiocin,<sup>67,113</sup> and cytosine arabinoside (AraC), aphidocolin, hydroxyurea, and 5-aza-2'-deoxycytidine (5-AzaC),<sup>114</sup> all of which induce a differentiated melanoma phenotype as well as apoptosis.

Anthracycline antibiotics such as doxorubicin in murine cells<sup>115</sup> and daunomycin in humans,<sup>116</sup> may induce melanoma differentiation by undetermined mechanisms but have little clinical anti-melanoma activity. They are intercalated into nucleic acids and inhibit DNA repair,<sup>117</sup> damage microtubules,<sup>116</sup> and generate free oxygen radicals.<sup>118</sup>

#### Miscellaneous compounds inducing melanoma differentiation

**Amino acids and derivatives.** High-dose tyrosine induces dendritic morphology, anchorage independence, and in-

creased tyrosinase levels, but not pigmentation, in a human melanoma line.<sup>119</sup>

Tumour cells are more dependent than normal cells on the amino acid glutamine. Phenylacetate conjugates with plasma glutamine and is active against many tumours, including melanoma. Phenylacetate also effects differentiation associated with down-regulation of *c-myc* in leukemic cells,<sup>30</sup> though this has not yet been reported in melanoma.

Histamine, the product of decarboxylation of the amino acid histidine, is the ligand for three specific surface receptors which, by association with different G proteins, activate PKA, PKC, calcium, and ion channels. Histamine also activates serotonin and other related receptors; and the histamine which is retained intracellularly may initiate cellular proliferation.<sup>120</sup> These relationships are so promiscuous that the specific mechanisms responsible for differentiation in cell culture<sup>121</sup> and the clinical efficacy<sup>122</sup> of histamine receptor antagonists in melanoma are difficult to establish.

Polyamines, products of decarboxylation of the amino acid ornithine, have been implicated as either inhibitors or stimulants of growth or differentiation in various cell types.<sup>123</sup> Ornithine decarboxylase is increased by phorbol esters in murine carcinogenesis models,<sup>124</sup> and inhibitors of ornithine decarboxylase, such as  $\alpha$ -difluoromethyl ornithine (DFMO), lead to decreased proliferation and increased tyrosinase and melanin in murine melanomas.<sup>125,126</sup> Murine melanomas also undergo apoptosis in response to high levels of oxidized polyamines when transplanted to an embryonic limb bud.<sup>127</sup> Similarly, polyamine antagonists have been shown to have antitumour effects in human melanoma.<sup>128</sup> Polyamines may therefore have therapeutic potential, although whether or not they are associated with differentiation pathways in the human system remains to be determined.

**Dimethylsulphoxide (DMSO).** The polar molecule DMSO induces phenotypic changes in human melanomas, but, like retinoids, does not induce a dendritic morphology.<sup>59,129</sup> The mechanisms involved are unclear; in other systems, DMSO has been shown to scavenge oxygen radicals, modulate a transcriptional promoter<sup>130</sup> and *c-myc* translation,<sup>131</sup> effect a post-translational decrease of the SV40 T antigen<sup>132</sup> and *c-myb* oncoproteins,<sup>133</sup> and increase polyamine synthesis.<sup>134</sup>

**Caffeic acid phenethyl ester (CAPE).** The naturally occurring product CAPE induces dendritic morphology and growth inhibition in a human melanoma cell line while increasing expression of antigens characterizing both


more- and less-differentiated phenotypes. This agent may thus have a dual therapeutic role as a differentiator and, by inducing surface antigen shifts, an immunomodulator.<sup>135</sup>

**Butyrate.** Sodium butyrate is a differentiating agent for many cell types. In murine melanoma it induces reversible differentiation associated with transcription of histone H1<sup>o</sup>, which stabilized chromatin and is usually associated with irreversible, terminal differentiation.<sup>136,137</sup> Expression of histone H1<sup>o</sup> is promoted by retinoids and by cAMP,<sup>95</sup> but it is unknown whether butyrate exerts its effects via these or other pathways. In human melanoma butyrate results in a spindle (rather than dendritic) morphology; it reduces expression of a melanosomal antigen but does not increase pigmentation,<sup>123</sup> and it induces an antigen which is otherwise found only on benign nevus cells.<sup>138</sup>

## Conclusions

The interactions among the various pathways implicated in melanoma differentiation are summarized in Figure 2. Signalling is modulated by PKC, PTKs and cAMP/PKA. PKC, which phosphorylates the MARCKS substrate,<sup>60</sup> also inactivates the PTK EGF receptor<sup>62,63</sup> and is itself activated by PTK phosphorylation of phospholipase-Cy.<sup>61</sup> The TNF- $\alpha$ /sphingosin pathway inhibits PKC and also phosphorylates the EGF receptor.<sup>73</sup> In the cAMP/PKA pathway, PKC inhibits both positive and negative surface receptors, inhibitory G-proteins, and phosphodiesterase while activating adenyl cyclase.<sup>77</sup> Both inhibitory and stimulatory G proteins depend on GTP, and cGMP may either increase or decrease phosphodiesterase activity.<sup>87</sup>

Nuclear regulatory factors of differentiation are produced



**Figure 2.** Inter-related pathways of melanoma differentiation. Activation of surface membrane receptors results in signal transduction modulated by interactions of PKC, PTKs and cAMP/PKA. All of these plus the nuclear retinoid receptor result in cross-talking nuclear transcriptional regulatory factors. See text for complete descriptions.  $\rightarrow$ , positive/activating pathway;  $\rightarrow \emptyset$ , negative/down-regulating pathway.

by PKC, PTKs, cAMP/PKA, and retinoid receptors. PKC, which blocks the p34<sup>cdc2</sup> DNA synthesis factor,<sup>90</sup> activates NFκB<sup>91</sup> and activates AP-1 via the ERKs pathway.<sup>92</sup> AP-1 is also increased by the cAMP/PKA-dependent CREB<sup>139</sup> and its activity is decreased by retinoids.<sup>93</sup> Retinoids also increase transcription of PKC,<sup>99-102</sup> c-jun, growth-regulatory factors,<sup>107</sup> and the H1<sup>o</sup> histone.<sup>95</sup> Histone H1<sup>o</sup> stabilizes chromatin and is associated with terminal differentiation;<sup>136,137</sup> it is also increased by cAMP/PKA.<sup>95</sup> The complexity of these interrelationships argues against the possibility of any simple linear process that results in universal, irreversible differentiation.

The interactions between these pathways are fluid and there are many examples of uncoupling between an overall differentiating effect and associated characteristics such as decreased proliferation or increased pigmentation. Simultaneous increases in both tumour differentiation and metastatic progression, as seen with cAMP enhancers in murine melanoma,<sup>85</sup> may represent a tool for further investigation as well as a potential hazard for any therapeutic applications.

The expanding knowledge of these mechanisms may well lead to therapeutic approaches which can either drive tumour cells to terminal differentiation or definitively turn back the clock of transformation to a benign melanocytic phenotype.

## References

- Pierce GB, Speers WC. Tumors as caricatures of the process of tissue renewal: prospects for therapy by directing differentiation. *Cancer Res* 1988; **48**: 1996-2004.
- Aso M, Hashimoto K, Eto H, *et al.* Expression of Schwann cell characteristics in pigmented nevus. *Cancer* 1988; **62**: 938-943.
- Clark WH, Elder DE, Guerry D, *et al.* A model predicting survival in stage I melanoma based upon tumor progression. *JNCI* 1989; **81**: 1893-1904.
- Clark WH, Elder DE, Guerry DI, *et al.* A study of tumor progression: the precursor lesions of superficial spreading and nodular melanoma. *Hum Pathol* 1984; **15**: 1147-1165.
- Nishikawa S, Kusakabe M, Yoshinaga K, *et al.* In utero manipulation of coat color formation by a monoclonal anti-c-kit antibody: Two distinct waves of c-kit-dependency during melanocyte development. *EMBO J* 1991; **10**: 2111-2118.
- Gerschenson M, Graves K, Carson SD, *et al.* Regulation of melanoma by the embryonic skin. *Proc Natl Acad Sci USA* 1986; **83**: 7307-7310.
- Hirobe T. Genes involved in regulating the melanocyte and melanoblast-melanocyte populations in the epidermis of newborn mouse skin. *J Exp Zool* 1982; **223**: 257-264.
- Bennett DC, Bridges K, McKay IA. Clonal separation of mature melanocytes from premelanocytes in a diploid human cell strain: spontaneous and induced pigmentation of premelanocytes. *J Cell Sci* 1985; **77**: 167-183.
- Cotsarelis G, Sun T-T, Lavker RM. Label-retaining cells reside in the bulge area of pilosebaceous unit: implications for follicular stem cells, hair cycle, and skin carcinogenesis. *Cell* 1990; **61**: 1329-1337.
- Hearing VJ, Jimenez M. Analysis of mammalian pigmentation at the molecular level. *Pigment Cell Res* 1989; **2**: 75-85.
- Mastrangelo MJ, Baker AR, Katz HR. *Cutaneous Melanoma*. Philadelphia: J.B. Lippincott Co., 1985: 1371-1422.
- Herlyn M. *Molecular and Cellular Biology of Melanoma*. Austin R.G. Landes Co., 1993.
- Lynch SA, Bouchard BN, Vijayasaradhi S, *et al.* Antigens of melanocytes and melanoma. *Cancer Met Rev* 1991; **10**: 141-150.
- Houghton AN, Herlyn M, Ferrone S. *Melanoma Antigens*. Philadelphia: J.B. Lippincott Co., 1992: 130-143.
- Elder DE, Rodeck U, Thurin J, *et al.* Antigenic profile of tumor progression stages in human melanocytic nevi and melanomas. *Cancer Res* 1989; **49**: 5091-5096.
- Carrel S, Dore J-F, Ruiter DJ, *et al.* The EORTC melanoma group exchange program: evaluation of a multicenter monoclonal antibody study. *Int J Cancer* 1991; **48**: 836-847.
- Valyi-Nagy I, Shih IM, Györfi T, *Int J Cancer* 1993; **54**: 159-165.
- Mancianti ML, Herlyn M, Weil D, *et al.* Growth and phenotypic characteristics of human nevus cells in culture. *J Invest Dermatol* 1988; **90**: 134-141.
- Nakayama K-i, Nakayama K, Negishi I, *et al.* Disappearance of the lymphoid system in bcl-2 homozygous mutant chimeric mice. *Science* 1993; **261**: 1584-1588.
- Kozopas KM, Yang T, Buchan HL, *et al.* MCL1, a gene expressed in programmed myeloid cell differentiation, has sequence similarity to bcl2. *Proc Natl Acad Sci USA* 1993; **90**: 3516-3520.
- Gerschenson LE, Rotello RJ. Apoptosis: a different type of cell death. *FASEB J* 1992; **6**: 2450-2452.
- Pezzella F, Turley H, Kuzu I, *et al.* bcl-2 protein in non-small-cell lung carcinoma. *N Engl J Med* 1993; **329**: 690-694.
- Darnell JE, Lodish HF, Baltimore D. *Molecular Cell Biology*. New York: Freeman and Co., 1990: 963-967.
- Brooks G, Birch M, Hart IR. Effects of biologically active tumor-promoting and non-promoting phorbol esters on *in vitro* growth of melanocytic cells. *Pigment Cell Res* 1990; **3**: 98-100.
- Herlyn M, Clark WH, Rodeck U, *et al.* Biology of tumor progression in human melanocytes. *Lab Invest* 1987; **56**: 461-474.
- Sato Y, Rifkin DB. Autocrine activities of basic fibroblast growth factor: regulation of endothelial cell movement, plasminogen activator synthesis and DNA synthesis. *J Cell Biol* 1988; **107**: 1199-1205.
- Kirchheimer JC, Wojta J, Christ G, *et al.* Functional inhibition of endogenously produced urokinase decreases cell proliferation in a human melanoma cell line. *Proc Natl Acad Sci USA* 1989; **86**: 5424-5428.
- Meissauer A, Kramer MD, Hoffman M, *et al.* Urokinase-type and tissue-type plasminogen activators are essential for *in vitro* invasion of human melanoma cells. *Exp Cell Res* 1991; **192**: 435-459.
- Yamamoto T, Yutaka S, Fukumoto H, *et al.* Over-expression of facilitative glucose transporter genes in human cancer. *Biochem Biophys Res Comm* 1990; **170**: 223-230.
- Samid D, Shack S, Sherman LT. Phenylacetate: a novel nontoxic inducer of tumor cell differentiation. *Cancer Res* 1992; **52**: 1988-1992.



31. Melber K, Zhu G, Diamond L. SV40-transfected human melanocyte sensitivity to growth inhibition by the phorbol ester 12- $\alpha$ -tetradecanoyl-phorbol-13-acetate. *Cancer Res* 1989; **49**: 3650-3655.
32. Jambrosic J, Mancianti ML, Ricciardi R, *et al.* Transformation of normal human melanocytes and non-malignant nevus cells by adenovirus 12-SV40 hybrid virus. *Int J Cancer* 1989; **44**: 1117-1123.
33. Albino AP, Sozzi G, Nanus DM, *et al.* Malignant transformation of human melanocytes: induction of a complete melanoma phenotype and genotype. *Oncogene* 1992; **7**: 2315-2321.
34. Mould AP, Wheldon LA, Komoriya A, *et al.* Affinity chromatographic isolation of the melanoma adhesion receptor for the IIICS region of fibronectin and its identification as the integrin  $\alpha_4\beta_1$ . *J Biol Chem* 1990; **265**: 4020-4024.
35. Albelda SM. Biology of disease. Role of integrins and other cell adhesion molecules in tumor progression and metastasis. *Lab Invest* 1993; **68**: 4-17.
36. Guan JL, Trevithick JE, Hynes RO. Fibronectin/integrin interaction induces tyrosine phosphorylation of a 120 kDa protein. *Cell Reg* 1991; **2**: 951-964.
37. Schwartz MA, Rupp EE, Frangioni JV, *et al.* Cytoplasmic pH and anchorage-independent growth induced by v-KI-ras, v-src or polyoma middle T. *Oncogene* 1990; **5**: 55-58.
38. Hynes RO. Integrins: versatility, modulation, and signaling in cell adhesion. *Cell* 1992; **69**: 11-25.
39. Lehmann JM, Riethmuller G, Johnson JP. MUC 18, a marker of tumor progression in human melanoma, shows sequence similarity to the neural cell adhesion molecules of the immunoglobulin superfamily. *Proc Natl Acad Sci USA* 1989; **86**: 9891-9895.
40. Johnson JP, Stade BG, Holzman B, *et al.* De novo expression of intercellular adhesion molecule 1 in melanoma correlates with increased risk of metastasis. *Proc Natl Acad Sci USA* 1989; **86**: 640-644.
41. Schmoeckel C, Stolz W, Sakai LY, *et al.* Structure of basement membranes in malignant melanoma and nevocytic nevi. *J Invest Dermatol* 1989; **92**: 663-668.
42. Rieber M, Gross A, Rieber MS. Relationship of a MR 140 fibronectin receptor and other adhesion-related glyco-proteins to tumor cell-cell interaction. *Cancer Res* 1987; **47**: 5127-5131.
43. Cheresch DA, Pierschbacher MD, Herzig MA, *et al.* Disialogangliosides GD2 and GD3 are involved in the attachment of human melanoma and neuroblastoma cells to extracellular matrix proteins. *J Cell Biol* 1986; **102**: 688-696.
44. Burns GF, Lucas CM, Krissansen GW, *et al.* Synergism between membrane gangliosides and arg-gly-asp-directed glycoprotein receptors in attachment to matrix proteins by melanoma cells. *J Cell Biol* 1988; **107**: 1225-1230.
45. Horowitz A, Duggan K, Buck C, *et al.* Interaction of plasma membrane fibronectin receptor with talin—a trans-membrane linkage. *Nature* 1986; **320**: 531-533.
46. Cunningham CC, Gorlin JB, Kwiatkowski DJ, *et al.* Actin-binding protein requirement for cortical stability and efficient locomotion. *Science* 1992; **255**: 325-327.
47. McCarthy JB, Hagen ST, Furcht LT. Human fibronectin contains distinct adhesion and motility promoting domains for metastatic melanoma cells. *J Cell Biol* 1986; **102**: 179-188.
48. Duband J-L, Nuckolls GH, Ishihara A, *et al.* Fibronectin receptor exhibits high lateral mobility in embryonic locomoting cells but is immobile in focal contacts and fibrillar streaks in stationary cells. *J Cell Biol* 1988; **107**: 1385-1396.
49. Copeman MC, Harris H. The extracellular matrix of hybrids between melanoma cells and normal fibroblasts. *J Cell Sci* 1988; **91**: 281-286.
50. Wakeling WF, Greetham J, Devlin LM, *et al.* Suppression of properties associated with malignancy in murine melanoma-melanocyte hybrid cells. *Br J Cancer* 1992; **65**: 529-537.
51. Harris H. The role of differentiation in the suppression of malignancy. Jean Brachet memorial lecture. *J Cell Sci* 1990; **97**: 5-10.
52. Gateff E, Mechler BM. Tumor-suppressor genes of *Drosophila melanogaster*. *CRC Crit Rev Oncogen* 1989; **1**: 221-245.
53. Ono Y, Fujii T, Igarashi K, *et al.* Phorbol ester binding to protein kinase C requires a cysteine-rich zinc-finger-like sequence. *Proc Natl Acad Sci USA* 1992; **86**: 4868-4871.
54. Nakanishi H, Exton JH. Purification and characterization of the  $\zeta$  isoform of protein kinase C from bovine kidney. *J Biol Chem* 1992; **267**: 16347-16354.
55. Azzi A, Boscoboinik D, Hensey C. The protein kinase C family. *Eur J Biochem* 1992; **208**: 547-557.
56. Brooks G, Wilson R, Dooley TP, *et al.* Protein kinase C downregulation and not transient activation correlates with melanocyte growth. *Cancer Res* 1991; **51**: 3281-3288.
57. Arita Y, O'Driscoll KR, Weinstein IB. Growth of human melanocyte cultures supported by 12- $\alpha$ -tetradecanoylphorbol-13-acetate is mediated through protein kinase C activation. *Cancer Res* 1992; **52**: 4514-4521.
58. Ziegler-Heitbrock HWL, Munker R, Johnson J, *et al.* In vitro differentiation of human melanoma cells analyzed with monoclonal antibodies. *Cancer Res* 1985; **45**: 1344-1350.
59. Huberman E, Heckman C, Langenbach R. Stimulation of differentiated functions in human melanoma by tumor-promoting agents and DMSO. *Cancer Res* 1979; **39**: 2618-2624.
60. Aderem A. The MARCKS brothers: a family of protein kinase C substrates. *Cell* 1992; **71**: 713-716.
61. Pawson T, Gish GD. SH2 and SH3 domains: from structure to function. *Cell* 1992; **71**: 359-362.
62. Countaway JL, McQuilkin P, Girones N, *et al.* Multisite phosphorylation of the epidermal growth factor receptor. *J Biol Chem* 1990; **265**: 3407-3416.
63. Nishizuka Y. The molecular heterogeneity of protein kinase C and its implications for cellular regulation. *Nature* 1988; **334**: 661-665.
64. Singletary SE, Baker FL, Spitzer G, *et al.* Biological effect of epidermal growth factor on the *in vitro* growth of human tumors. *Cancer Res* 1987; **47**: 403-406.
65. Rodeck U, Herlyn M, Menssen HD, *et al.* Metastatic but not primary melanoma cell lines grow *in vitro* independently of exogenous growth factors. *Int J Cancer* 1987; **40**: 687-90.
66. Kiguchi K, Constantinou AI, Huberman E. Genistein-induced differentiation and protein-linked DNA strand breakage in human melanoma. *Cancer Commun* 1990; **2**: 271-277.
67. Alnemri ES, Litwack G. Activation of internucleosomal DNA cleavage in human CEM lymphocytes by glucocorticoid and novobiocin. *J Biol Chem* 1990; **265**: 17323-17333.
68. Becker D, Lee PL, Rodeck U, *et al.* Inhibition of the

- fibroblast growth factor receptor-1 gene in human melanocytes and malignant melanomas leads to inhibition of proliferation and signs indicative of differentiation. *Oncogene* 1992; **7**: 2203–2313.
69. Garbe C, Krasagakis K. Effects of interferons and cytokines on melanoma cells. *J Invest Dermatol* 1993; **100**: 239S–244S.
70. Johns TG, Mackay IR, Callister KA, *et al*. Antiproliferative potencies of interferons on melanoma cell lines and xenografts: higher efficacy of interferon beta. *JNCI* 1992; **84**: 1185–1190.
71. Jiang H, Su Z-Z, Boyd J, *et al*. Gene expression changes induced in human melanoma cells undergoing reversible growth suppression and terminal cell differentiation. *Mol Cell Diff* 1993; **1**: 41–66.
72. Charak BS, Sadowski RM, Mazumder A. Antitumor effect of interferon plus cyclosporine A following chemotherapy for disseminated melanoma. *Cancer Res* 1992; **52**: 6482–6486.
73. Mathias S, Dressler KA, Kolesnick RN. Characterization of a ceramide-activated protein kinase: stimulation by tumor necrosis factor  $\alpha$ . *Proc Natl Acad Sci USA* 1991; **88**: 10009–10013.
74. Obeid LM, Linardic CM, Karolak LA, *et al*. Programmed cell death induced by ceramide. *Science* 1993; **259**: 1769–1771.
75. McKnight GS. Cyclic AMP second messenger systems. *Curr Opin Cell Biol* 1991; **3**: 213–217.
76. Walsh DA, Glass DB, Mitchell RD. Substrate diversity of the cAMP-dependent protein kinase: regulation based upon multiple binding interactions. *Curr Opin Cell Biol* 1992; **4**: 241–251.
77. Houslay MD. 'Crosstalk': a pivotal role for protein kinase C in modulating relationships between signal transduction pathways. *Eur J Biochem* 1991; **195**: 9–27.
78. Kreider JW, Rosenthal M, Lingle N. Cyclic adenosine 3'5'-monophosphate in the control of melanoma replication and differentiation. *JNCI* 1973; **50**: 555–558.
79. Guiffre L, Schreyer M, Mach JP, *et al*. Cyclic AMP induces differentiation *in vitro* of human melanoma cells. *Cancer* 1988; **61**: 1132–1141.
80. Tsuchiya H, Tomita K, Yasutake H, *et al*. Growth inhibition and differentiation of murine melanoma B16-BL6 cells caused by the combination of cisplatin and caffeine. *Jpn J Cancer Res* 1989; **80**: 1246–1251.
81. Hill SE, Rees RC, MacNeil S. A positive association between agonist-induced cyclic AMP production *in vitro* and metastatic potential in murine B16 melanoma and hamster fibrosarcoma. *Clin Exp Metastasis* 1990; **8**: 461–474.
82. Bennett DC, Dexter TJ, Ormerod EJ, *et al*. Increased experimental metastatic capacity of a murine melanoma following induction of differentiation. *Cancer Res* 1986; **46**: 3239–3244.
83. Laskin JD, Mufson RA, Weinstein IB, *et al*. Identification of a distinct phase during melanogenesis that is sensitive to extracellular pH and ionic strength. *J Cell Physiol* 1980; **103**: 467–474.
84. Kameyama K, Vieira WD, Tsukamoto K, *et al*. Differentiation and the tumorigenic and metastatic phenotype of murine melanoma cells. *Int J Cancer* 1990; **45**: 1151–1158.
85. Hart IR. Tumor cell progression and differentiation in metastasis. *Semin Cancer Biol* 1991; **2**: 87–95.
86. Boss GR. cGMP-induced differentiation of the promyelocytic cell line HL-60. *Proc Natl Acad Sci USA* 1989; **86**: 7174–7178.
87. Lincoln TM, Cornwell TL. Intracellular cyclic GMP receptor proteins. *FASEB J* 1993; **7**: 328–338.
88. Giotto GJ, Smith JR, Nicolson GL. Guanosine 5'-triphosphate inhibits growth and stimulates differentiated functions in B16 melanoma cells. *Exp Cell Res* 1978; **112**: 385–393.
89. Kiguchi K, Collart FR, Henning-Chubb C, *et al*. Induction of cell differentiation in human melanoma cells by inhibitors of IMP dehydrogenase expression and activity. *Exp Cell Res* 1990; **178**: 41–50.
90. Coppock DL, Tansey JB, Nathanson L. 12-o-tetradecanoyl phorbol-13-acetate induces transient cell cycle arrest in G<sub>1</sub> and G<sub>2</sub> in metastatic melanoma cells: inhibition of phosphorylation of p34<sup>cdc2</sup>. *Cell Growth Diff* 1992; **3**: 485–494.
91. Baeuerle PA, Baltimore D. I $\kappa$ B: a specific inhibitor of the NF- $\kappa$ B transcription factor. *Science* 1988; **242**: 540–546.
92. Pulverer BJ, Hughes K, Franklin CC, *et al*. Co-purification of mitogen-activated protein kinases with phorbol ester-induced c-jun kinase activity in U937 leukaemic cells. *Oncogene* 1993; **8**: 407–415.
93. Schüle R, Rangarajan P, Yang N, *et al*. Retinoic acid is a negative regulator of AP-1-responsive genes. *Proc Natl Acad Sci USA* 1991; **88**: 6092–6096.
94. Ponta H, Cato AC, Herrlich P. Interference of pathway specific transcription factors. *Biochim Biophys Acta* 1992; **1129**: 255–261.
95. Breur B, Fisher J, Alonso A. Cloning and characterization of the mouse histone H1<sup>o</sup> promoter region. *Gene* 1989; **81**: 307–314.
96. McBurney MW, Costa S, Pratt MAC. Retinoids and cancer: a basis for differentiation therapy. *Cancer Invest* 1993; **1**: 590–598.
97. Gudas L. Retinoids, retinoid-responsive genes, cell differentiation, and cancer. *Cell Growth Diff* 1992; **3**: 655–662.
98. Green S. Nuclear hormone receptors: promiscuous liaisons. *Nature* 1993; **361**: 590–591.
99. Gruber JR, Ohno S, Niles RM. Increased expression of protein kinase C $\alpha$  plays a key role in retinoic acid-induced melanoma differentiation. *J Biol Chem* 1992; **267**: 13356–13360.
100. Hoal E, Wilson L, Dowdle EB. Variable effects of retinoids on two pigmented human melanoma cell lines. *Cancer Res* 1982; **42**: 5191–5195.
101. Lotan R, Lotan D. Stimulation of melanogenesis in a human melanoma cell line by retinoids. *Cancer Res* 1980; **40**: 3345–3350.
102. Meyskens F, Fuller B. Characterization of the effects of different retinoids on the growth and differentiation of human melanoma cell lines and selected subclones. *Cancer Res* 1980; **40**: 2194–2196.
103. Kraemer KH, DiGiovanna JJ, Moshell A, *et al*. Prevention of skin cancer in xeroderma pigmentosum with the use of oral isotretinoin. *N Engl J Med* 1988; **318**: 1663–1667.
104. Smith MA, Parkinson DR, Cheson BD, *et al*. Retinoids in cancer therapy. *J Clin Oncol* 1992; **10**: 839–864.
105. Goodman G. Phase II trial of retinol in patients with advanced cancer. *Cancer Treat Rep* 1986; **70**: 1023–1024.
106. Levine N, Meyskens F. Topical vitamin-A-acid therapy for cutaneous metastatic melanoma. *Lancet* 1980; **2**: 224–226.
107. Gudas LJ. Retinoic acid and teratocarcinoma stem cells. *Semin Dev Biol* 1991; **2**: 171–179.

108. Hazuka MB, Edwards-Prasad J, Newman F, *et al.* Beta carotene induces morphological differentiation and decreases adenylate cyclase activity in melanoma cells in culture. *J Am Coll Nutr* 1990; **9**: 143–149.
109. Sies H, Stahl W, Sundquist AR. Antioxidant functions of vitamins. Vitamins E and C, beta-carotene, and other carotenoids. *Ann NY Acad Sci* 1992; **669**: 7–20.
110. Hubbard-Smith K, Hill HZ, Hill GJ. Melanin both causes and prevents oxidative base damage in DNA: quantification by anti-thymine glycol antibody. *Radiat Res* 1992; **130**: 160–165.
111. Sheridan JW, Simmons RJ. Tritiated thymidine-induced increased DNA content and irreversible differentiation in a human melanoma cell line. *Br J Exp Pathol* 1981; **62**: 289–295.
112. Kidson SH, DeHaan JB. Effect of thymidine analogs on tyrosinase activity and mRNA accumulation in mouse melanoma cells. *Exp Cell Res* 1990; **188**: 36–41.
113. Nordenberg J, Albukrek D, Hadar T, *et al.* Novobiocin-induced anti-proliferative and differentiating effects in melanoma B16. *Br J Cancer* 1992; **65**: 183–188.
114. Steigerwald SD, Pfeifer GD. Variable DNA methylation changes during differentiation of human melanoma cells. *Exp Cell Res* 1988; **178**: 41–50.
115. Mariani M, Supino R. Morphological alterations induced by doxorubicin in B16 melanoma cells. *Cancer Lett* 1990; **51**: 209–212.
116. Molinari A, Calcabrini A, Craeteri P, *et al.* Effects of daunomycin on the microtubular network: a cytochemical study on a human melanoma cell line. *Eur J Cell Biol* 1991; **54**: 291–298.
117. Hande KR. Antitumor antibiotics, epipodophyllins, and vinca alkaloids. *Curr Opin Oncol* 1992; **4**: 1080–1088.
118. Bienvenu P, Caron L, Gasparutto D, *et al.* Assessing and counteracting the prooxidant effects of anticancer drugs. *EXS* 1992; **62**: 257–265.
119. Guerra L, Bover L, Mordoh J. Differentiating effect of L-tyrosine on the human melanoma cell line IIB-MEL. *Exp Cell Res* 1990; **188**: 61–65.
120. Mitsuhashi M, Payan DG. Functional diversity of histamine and histamine receptors. *J Invest Dermatol* 1992; **98**: 8S–11S.
121. McEwan MT, Parsons PG. Regulation of tyrosinase expression and activity in human melanoma cells via histamine receptors. *J Invest Dermatol* 1991; **97**: 868–873.
122. Smith T, Clark JW, Popp MB. Regression of melanoma nodules in patients treated with ranitidine. *Arch Intern Med* 1987; **147**: 1815–1816.
123. Pegg AE. Polyamine metabolism and its importance in neoplastic growth and as a target for chemotherapy. *Cancer Res* 1988; **48**: 759–774.
124. DiGiovanni J. *Modification of Multistage Skin Carcinogenesis in Mice*. Basel: Karger, 1991: 192–229.
125. Sunkhara PS, Chang CC, Prakash NJ, *et al.* Effect of inhibition of polyamine biosynthesis by DL-difluoromethylornithine on the growth and melanogenesis of B16 melanoma *in vitro* and *in vivo*. *Cancer Res* 1985; **45**: 4067–4070.
126. Kapyaho K, Senervista R, Janne J. Effects of inhibitors of polyamine biosynthesis on the growth and melanogenesis of murine melanoma cells. *Cancer Res* 1985; **45**: 1444–1448.
127. Parchment RE, Pierce GB. Polyamine oxidation, programmed cell death and regulation of melanoma in the murine embryonic limb. *Cancer Res* 1989; **49**: 6680–6686.
128. Bernacki RJ, Bergeron RJ, Porter CW. Antitumor activity of N,N'-bis(ethyl)spermine homologues against human MALME-3 melanoma xenografts. *Cancer Res* 1992; **52**: 2424–2430.
129. Takahashi H, Parsons PG. *In vitro* phenotypic alteration of human melanoma induced by differentiating agents: heterogeneous effects on cellular growth and morphology, enzymatic activity, and antigenic expression. *Pigment Cell Res* 1990; **3**: 223–232.
130. Cerosaletti K, Woodward JG, Lord EM, *et al.* Two regions of the H-2 D<sup>d</sup> promoter are responsive to dimethylsulfoxide in line 1 cells by a mechanism distinct from IFN- $\gamma$ . *J Immunol* 1992; **148**: 1212–1221.
131. Spott GD, Hann SR. Enhanced translation and increased turnover of c-myc proteins occur during differentiation of murine erythroleukemia cells. *Mol Cell Biol* 1990; **10**: 3952–3964.
132. Witte MM, Parker RF, Wang H, *et al.* Repression of SV40 T oncoprotein expression by DMSO. *J Cell Physiol* 1992; **151**: 50–55.
133. Boise LH, Gorse KM, Westin EH. Multiple mechanisms of regulation of the human c-myc gene during myelomonocytic differentiation. *Oncogene* 1992; **7**: 1817–1825.
134. Gazit Y, Friend C. Polyamine biosynthesis enzymes in the induction and inhibition of differentiation in Friend erythroleukemia cells. *Cancer Res* 1980; **40**: 1717–1732.
135. Guarini L, Su Z-Z, Zucker S, *et al.* Growth inhibition and modulation of antigenic phenotype in human melanoma and glioblastoma multiforme cells by caffeic acid phenethyl ester. *Cell Mol Biol* 1992; **38**: 513–527.
136. Rousseau D, Khochbin S, Gorka C, *et al.* Induction of H1<sup>o</sup> gene expression in B16 murine melanoma cells. *Eur J Biochem* 1992; **208**: 775–779.
137. Nordenberg J, Wasserman L, Beery E, *et al.* Growth inhibition of murine melanoma by butyric acid and dimethylsulfoxide. *Exp Cell Res* 1986; **162**: 77–85.
138. Nakanishi T, Hashimoto K. The differential reactivity of benign and malignant nevocytic lesions with mouse monoclonal antibody TNKH1. *Cancer* 1987; **59**: 1340–1344.
139. Cornelius P, Marlowe M, Call K, *et al.* Regulation of glucose transport as well as glucose transporter and immediate early gene expression in 3T3-L1 preadipocytes by 8-bromo-cAMP. *J Cell Physiol* 1991; **146**: 298–308.

(Received 25 November 1993; accepted 21 April 1994)