

# The Gonadotropic Function of Insulin\*

LEONID PORETSKY† AND MARCIA F. KALIN

*Division of Endocrinology, Department of Medicine, Beth Israel Medical Center and The Mount Sinai School of Medicine, New York, New York 10003*

## I. Introduction

**D**URING the last decade, it has become apparent that the pituitary gonadotropins, FSH and LH, are not the exclusive regulators of ovarian function. Several other hormones and various growth factors also play a role in ovarian physiology. Among these hormones and growth factors are insulin and insulin-like growth factors (IGFs).

It is conceivable that a deficiency or an excess of insulin or IGFs could significantly alter ovarian function. In fact, insulin-dependent diabetes mellitus (IDDM), a disorder characterized by insulinopenia, is associated with clinical manifestations of ovarian hypofunction: primary amenorrhea, late menarche, anovulation, low pregnancy rate, and early menopause. Furthermore, insulin resistance, the cardinal feature of which is hyperinsulinemia, is associated with clinical manifestations of ovarian hyperstimulation, primarily hyperandrogenism.

It is becoming increasingly clear that insulin plays an important role in ovarian physiology; however, the mechanisms of insulin action on the ovary remain largely hypothetical. We will review clinical observations and experimental studies that shed light on the role of insulin in ovarian function.

## II. Clinical Observations and *in Vivo* Studies

The idea that insulin influences ovarian physiology stems from clinical observations among patients with IDDM and patients with various syndromes of insulin resistance.

### A. IDDM

Early observations among patients with IDDM demonstrated that insulin deficiency was associated with

ovarian hypofunction. In the preinsulin era, amenorrhea was common among young women with diabetes. In a 1925 review of their experience regarding growth, development, and prognosis of children with diabetes, Joslin *et al.* (1) wrote, "Unaided by insulin, no girl in our series developed menstruation after the onset of diabetes." After 2 months to 1 yr of insulin therapy, menstruation commenced in four girls ranging in age from 13 to 17 yr.

As insulin therapy became widely available, this picture changed; menarche developed in most girls with diabetes, although at a somewhat later age than in normal controls. In 1954, Bergqvist (2) reported that among nondiabetic girls ( $n = 114$ ), the average age of menarche was 13.9 yr, whereas among diabetic girls ( $n = 33$ ), the average age of menarche was 15.0 yr; (the difference was statistically significant).

In 1982, Djursing and co-workers (3) reported that menarche still occurred about 1 yr later in diabetic women than it did in nondiabetic women. The average age of menarche among nondiabetic, anovulatory controls ( $n = 45$ ) was 13 yr, whereas among diabetic women, either ovulatory ( $n = 9$ ) or anovulatory ( $n = 22$ ), the average age of menarche was 14 yr (the difference was not statistically significant). The disappearance, from 1954 to 1982, of a significant difference in age of menarche between diabetic women and nondiabetic women could reflect either the later study's smaller number of patients or improved management of diabetes in recent years.

In 1954, Bergqvist (2) examined also the number of pregnancies and the age of onset of menopause among diabetic women. Among 169 diabetic women (ranging in age from 18–45 yr), the number of pregnancies with a duration of 28 weeks or more was 48, instead of a predicted 77. Regarding the age of onset of menopause, the author concluded, "On the whole it seems that the menopause occurs at a slightly lower age in diabetic women than in the comparative group of nondiabetic women" (the difference was not statistically significant) (2).

Sex hormone patterns among women with diabetes have been the subject of surprisingly few studies.

Address requests for reprints to: Leonid Poretsky, M.D., Department of Medicine, Beth Israel Medical Center, First Avenue at 16th Street, New York, New York 10003.

\* Supported in part by a grant from the United Parcel Service.

† Recipient of a First Independent Research Support and Transition (FIRST) Award from the National Institutes of Health (HD-22738).

Walton and co-workers (4), in a study of endocrine abnormalities associated with hemochromatosis, reported that estradiol levels were lower among women with diabetes than among women with hemochromatosis or cirrhosis (122 vs. 146 or 130 pmol/liter). Normal controls were not studied, unfortunately, rendering interpretation of the study's results somewhat difficult. The low estradiol levels among women with diabetes, however, lent support to the idea that ovarian hypofunction was associated with the insulin-deficient state.

Zumoff and co-workers (Zumoff B. Z., L. Miller, L. Poretsky, C. D. Levit, E. H. Miller, U. Heinz, M. Kalin, and R. S. Rosenfeld, personal communication) found that follicular phase progesterone levels were significantly lower among women with IDDM than among normal controls. Progesterone levels correlated negatively with glycosylated hemoglobin, suggesting that some factor involved in glucose metabolism (such as insulin) contributed to the low progesterone levels.

Djursing and co-workers (3) found that median levels of FSH, LH, estradiol, and testosterone were lower among anovulatory diabetic women than among ovulatory diabetic women, ovulatory controls, or anovulatory controls. The authors concluded that the deficiency in ovarian hormone production among the anovulatory diabetic patients was associated with "either a derangement in gonadal-pituitary feedback mechanism or a depression of pituitary function at or above the pituitary level."

To evaluate the relationship between insulin and the pituitary-gonadal axis, Rueda and Urdinola (5) measured insulin levels and LH levels among patients with polycystic ovarian syndrome (PCO); there was no correlation between the two hormones.

Distiller and co-workers (6) examined pituitary responsiveness to LHRH among diabetic women ( $n = 12$ ), diabetic men ( $n = 7$ ), and normal controls. Both the men with diabetes and the women with diabetes had significantly blunted responses of LH and FSH to LHRH. Maximum changes in LH correlated inversely with fasting plasma glucose levels.

Several investigators have examined the effects of insulin deficiency on the ovary in animals. Liu and co-workers (7) demonstrated that ovarian weights and ovary/body weight ratios were significantly lower among rats with Alloxan-induced diabetes than among nondiabetic animals. Response of ovarian weight to FSH also was impaired in diabetic animals. Insulin reversed these abnormalities, producing a dose-related increase in the ovary/body weight ratio, which increased until it normalized. Circulating levels of LH and of FSH were similar among diabetic animals and controls. The authors concluded that "... the decreased ovarian weight in the diabetic immature rats was probably not due to a lack of

pituitary gonadotrophic stimulation, but mainly due to an insulin deficiency." They hypothesized that insulin deficiency in some way decreased responsiveness of the ovaries to gonadotropins.

Bestetti and co-workers (8) evaluated gonadotropic function in female rats with streptozotocin-induced diabetes. Median eminence content of LHRH was low and plasma LH response to LHRH was inadequate, suggesting that gonadotropin deficiency was the cause of ovarian hypofunction. The arcuate nucleus and the median eminence contained numerous degenerate axons, and LH gonadotroph cells were altered (these changes were evident to an even greater degree among diabetic male rats). The authors hypothesized that the abnormalities were caused by insulin deficiency.

Another hypothetical cause of ovarian hypofunction in IDDM is a concomitant deficiency of IGF-1, a substance shown to have gonadotropic activity in *in vitro* studies (9). Maes and co-workers (10) recently showed that rats with experimental IDDM do indeed have low levels of IGF-1. Tan and Baxter (11) reported that the mean serum IGF-1 level was significantly lower among patients with IDDM than among normal controls ( $0.41 \pm 0.17$  U/ml vs.  $0.84 \pm 0.26$  U/ml,  $P < 0.001$ ).

The association between functional ovarian abnormalities and IDDM suggests that insulin plays a role in the regulation of ovarian function. It has been extremely difficult to determine, however, which of the many metabolic derangements of uncontrolled diabetes underlies the abnormalities in ovarian function. Hyperglycemia, insufficient body weight, insulin deficiency, gonadotropin deficiency, IGF-1 deficiency, or any combination of these factors could conceivably affect the ovary. Contributing further confusion to the issue is the fact that some patients with IDDM are hyperinsulinemic, due either to overzealous treatment or to appropriate treatment in response to insulin resistance (12). Thus, patients with IDDM today actually constitute a heterogeneous group in terms of serum insulin levels, which may be low, normal, or high. This heterogeneity may account for the fact that the overall ovarian function in treated patients with IDDM differs only marginally from that of normal controls.

### B. Insulin resistance

Further evidence supporting the idea that insulin affects ovarian function has arisen from clinical observations among patients with various syndromes of insulin resistance and acanthosis nigricans. These patients exhibit features of ovarian hyperstimulation, primarily hyperandrogenism. Patients with syndromes of insulin resistance and acanthosis nigricans are heterogeneous in terms of etiology of insulin resistance and clinical pres-

entation but homogeneous in terms of serum insulin levels: all are hyperinsulinemic. Several different entities can be distinguished.

Patients with the type A syndrome are young, thin women with PCO or hyperthecosis and a genetic deficiency in the number of insulin receptors (13). Patients with the type B syndrome are usually somewhat older than those with the type A syndrome. Type B patients have features of autoimmune disease, and their insulin resistance is caused by insulin-receptor autoantibodies (14–16). Another group comprises several patients whose clinical picture resembles that of the type A syndrome, but their number of insulin receptors is normal (17–23). It is thought that the patients in this latter group have postbinding abnormalities in the insulin action pathway; indeed, in some cases, investigators already have demonstrated abnormalities in the tyrosine-kinase activity of the insulin receptor  $\beta$ -subunit (19, 20, 24). Other unusual syndromes of hyperinsulinemia and hyperandrogenism include lipotrophic diabetes (25, 26), leprechaunism (27, 28), and the Rabson-Mendenhall syndrome (29, 30).

The association between hyperinsulinemia and hyperandrogenism is not limited to unusual syndromes. Acanthosis nigricans, insulin resistance, and hyperandrogenism occur also in approximately 5% of obese women; these women constitute probably the largest subset of patients with the syndrome (31). Their insulin receptor number is down-regulated by the hyperinsulinemia of obesity and normalizes with caloric restriction.

Another common disorder, PCO, also is characterized by varying degrees of hyperandrogenism and insulin resistance (32). Burghen and co-workers (33) found insulin resistance among both obese patients and nonobese patients with PCO. Plasma insulin levels correlated with plasma testosterone levels and androstenedione levels. Dunaif and co-workers (34) similarly demonstrated that insulin resistance, even in the absence of obesity, was common among women with PCO.

It has been of considerable interest that in spite of heterogeneity in clinical picture and in insulin receptor status, various degrees of hyperandrogenism accompany each of the hyperinsulinemic states (25). The source of the excessive androgens initially was unclear. Taylor and co-workers (35) found that among women with the type B syndrome, only those who were premenopausal exhibited hyperandrogenism. The difference between premenopausal women and postmenopausal women pointed to the ovary as the source of excessive androgen formation, a hypothesis confirmed by subsequent studies.

McNatty and co-workers (36) catheterized the ovarian and the adrenal veins in a patient with hyperandrogenism and insulin resistance and demonstrated that the ovary indeed was the source of the patient's excessive

testosterone and androstenedione. Similar findings were observed in a patient with autoantibodies to the insulin receptor (35).

These clinical observations have demonstrated that insulin resistance, regardless of its cause, is closely associated with hyperandrogenism. Two distinct points of view concerning the basis of this association have evolved.

One hypothesis asserts that hyperandrogenism produces insulin resistance (33). This hypothesis has been supported by at least two observations. The first observation, reported by Cole and Kitabchi (37), was that administration of estrogen and progesterone to a patient with PCO caused remission of insulin resistance. The second observation was reported by Shoupe and Lobo (38). They found that administering the androgen antagonist spironolactone to patients with PCO significantly reduced not only testosterone levels, but also fasting insulin levels. These two studies thus showed that reducing the degree of hyperandrogenism reduced also the degree of insulin resistance.

The hypothesis that hyperandrogenism causes insulin resistance does not explain, however, either the cause of hyperandrogenism or the mechanism by which hyperandrogenism could produce insulin resistance. The latter is particularly problematic in cases in which the cause of insulin resistance is apparent (*e.g.* insulin receptor autoantibodies or a genetic deficiency in the number of insulin receptors).

Two recent studies argue against the hypothesis that hyperandrogenism causes insulin resistance. Stuart and co-workers (39) matched patients having acanthosis nigricans with controls according to weight and androgen levels. The hyperandrogenized, obese controls exhibited significantly less hyperinsulinemia than did the acanthotic subjects, suggesting that hyperandrogenism or obesity was not the cause of insulin resistance in the acanthotic group. In the second study, Geffner and co-workers (40) administered a long acting LHRH agonist to three patients with PCO and hyperinsulinemia. Despite suppression of ovarian androgen production, hyperinsulinemia persisted. These two studies thus contradict the hypothesis that hyperandrogenism causes hyperinsulinemia.

The second hypothesis regarding the association between hyperandrogenism and insulin resistance proposes that hyperinsulinemia produces ovarian hyperstimulation, manifested principally as hyperandrogenism (35, 41). A recent study designed by Nestler and co-workers (42) to test this hypothesis failed to find any supporting evidence. These investigators administered insulin via the hyperinsulinemic-euglycemic clamp technique for 12 to 16 hr to four normal women and one woman with hyperandrogenism, insulin resistance, and acanthosis ni-

gricans. Although each woman became hyperinsulinemic (the average serum insulin level was  $1832 \pm 292 \mu\text{U/ml}$ ), none demonstrated a rise in serum testosterone. The meaning of these findings, as noted by the authors, is unclear, because the length of time during which hyperinsulinemia was maintained was extremely brief. Perhaps months or years of hyperinsulinemia are necessary to produce hyperandrogenism.

The hypothesis that hyperinsulinemia produces hyperandrogenism remains attractive, even in view of these findings, because the idea provides a unifying explanation for hyperandrogenism in all of the above mentioned syndromes. The hypothesis raises two questions, however. The first question is: Why would hyperandrogenism be the principal manifestation of ovarian hyperstimulation? There are several proposed mechanisms explaining this phenomenon, and they are discussed below (see *Insulin or IGFs and Ovarian Steroidogenesis*). The second question is: How could the ovary remain sensitive to insulin in insulin-resistant states? To explain this paradox, it was postulated that insulin acts upon the ovary by binding not only to ovarian insulin receptors, but also to ovarian IGF-1 receptors (35). This hypothesis incorporates the knowledge that insulin, in high concentrations, interacts with IGF-1 receptors (the specificity spillover phenomenon) (43) and that IGF-1 stimulates steroidogenesis (9). Thus, one of the first steps in *in vitro* studies examining the effects of insulin on ovarian function was identification of insulin receptors and IGF-1 receptors in the ovary.

### III. *In Vitro* Studies

Insulin initiates its action by binding to an insulin receptor. The insulin-receptor complex is rapidly brought from the plasma membrane into the cell (internalization), and the insulin molecule is degraded. The receptor may then either undergo degradation or be recycled to the cell membrane (44).

It is thought that the insulin signal is transmitted via a recently discovered phenomenon involving phosphorylation of the insulin receptor  $\beta$ -subunit. Kasuga and co-workers (45) reported that insulin-stimulated insulin receptor phosphorylation occurred in a dose-dependent fashion and reached maximum intensity the first minute after insulin binding (45). Found to be a tyrosine-specific protein kinase, the receptor apparently can acutely initiate phosphorylation of various intracellular proteins. It still is unclear as to which postbinding event initiates insulin's metabolic effects; at this time, however, phosphorylation of the insulin receptor  $\beta$ -subunit appears to be the most likely candidate.

Insulin binds also to the IGF-1 receptor, although with an affinity lower than that with which it binds to the

insulin receptor (43, 46). Insulin and IGF-1 are structurally similar: positions 1 to 29 of the IGF-1 molecule are homologous to the insulin  $\beta$ -chain; positions 42 to 62 are homologous to the  $\alpha$ -chain; and the three-dimensional structures of the two peptides are virtually identical (47). Also sharing many features are the peptides' receptors (48). Both insulin receptors and IGF-1 receptors are heterotetramers, consisting of two  $\alpha$ -subunits (mol wt  $\sim 130,000$ ) and two  $\beta$ -subunits (mol wt  $\sim 90,000$ ). The  $\alpha$ -subunits serve as hormone-binding sites, whereas the  $\beta$ -subunits undergo autophosphorylation (of their tyrosine residues) and serve as tyrosine kinases (45, 49). Although insulin and IGF-2 also are structurally similar, their receptors differ, and insulin does not bind to the IGF-2 receptor (44).

#### A. *Insulin receptors and IGF-1 receptors in the ovary*

Because insulin initiates its action by binding to either insulin or IGF receptors, tissues serving as target organs for insulin are expected to contain either one or both receptors.

1. *Insulin receptors and IGF-1 receptors in animal ovarian tissue.* Both insulin receptors and IGF-1 receptors have been identified in animal and in human ovarian cells. Rein and Schomberg (50) identified insulin receptors in porcine granulosa cells. The [ $^{125}\text{I}$ ]insulin binding was saturable and dependent on pH, temperature, and incubation time (optimal conditions were: 2 h, 25 C, pH 7.5). The level of insulin-receptor binding remained constant during the course of follicular development.

Veldhuis and co-workers (51) found IGF-1 receptors (using [ $^{125}\text{I}$ ]somatomedin-C) in swine granulosa cells. Low affinities for insulin and for multiplication-stimulating activity and absent affinity for desoctapeptide insulin demonstrated specificity of the binding sites. Davoren and co-workers (52) found specific high affinity, low capacity IGF-1 binding sites in granulosa cells from immature hypophysectomized estrogen-treated rats.

Insulin receptors and IGF-1 receptors thus have been clearly demonstrated in animal granulosa cells.

2. *Insulin receptors and IGF-1 receptors in human ovarian tissue.* We previously have reported finding specific insulin-binding sites in human ovarian tissue: in stromal tissue from normal women, in stromal tissue from women with PCO, in follicular tissue from normal women, and in purified human granulosa cells (53, 54). Receptors were abundant in all types of ovarian cells and were characterized by the following: high affinity to insulin ( $\text{ED}_{50} \sim 10 \text{ ng/ml}$ ); low affinity to the insulin analog proinsulin ( $\text{ED}_{50} \sim 100 \text{ ng/ml}$ ); and recognition by antiinsulin receptor antibodies (serum B-2), but not by monoclonal anti-IGF-1 receptor antibodies ( $\alpha\text{-IR-3}$ ) (53).

Jarrett and co-workers (55) found insulin-binding sites in cell membrane fractions from normal human ovaries. Maximum binding occurred by 4 h at 15 C and was similar to that of other traditional insulin target tissues. Insulin binding with luteal phase ovarian cell membranes resembled that with follicular phase ovarian cell membranes.

Initial attempts to demonstrate IGF-1 receptors in human ovarian tissue showed only minimal [<sup>125</sup>I]IGF-1 binding in granulosa cells from two women undergoing *in vitro* fertilization procedures (53). Later, however, IGF-1 binding sites were found in ovarian stroma from a patient with hyperthecosis and insulin resistance (Pepper, G. M., L. Poretsky, J. L. Gabrilove, and M. M. Arton, submitted for publication). Subsequent experiments clearly identified high affinity IGF-1 receptors in purified human granulosa cells (Gates, G. S., S. Bayer, M. Seibel, L. Poretsky, J. S. Flier, and A. C. Moses, submitted for publication). Insulin receptors and IGF-1 receptors thus have been clearly demonstrated in the human ovary.

### B. Insulin receptor phosphorylation

The earliest known postbinding event in the insulin action pathway is phosphorylation of the insulin receptor  $\beta$ -subunit (45), a process thought to serve as the signal for insulin action. Among some insulin-resistant patients, insulin receptor binding characteristics are normal, but insulin receptor phosphorylation or tyrosine kinase activity is defective (19–21); these findings suggest that defective insulin receptor phosphorylation or defective tyrosine kinase activity is the cause of insulin resistance in some cases. Insulin receptor phosphorylation or tyrosine kinase activity has been demonstrated in mononuclear cells (56), lymphocytes (56), erythrocytes (57), or fibroblasts (58). Two recent studies have demonstrated insulin receptor phosphorylation in swine ovarian cells or human ovarian cells.

Veldhuis and co-workers (59) showed that insulin stimulated phosphorylation of the insulin receptor  $\beta$ -subunit in cultured swine granulosa cells. Neither somatomedin-C (IGF-1) nor desoctapeptide insulin produced the effect, demonstrating specificity of the reaction. We reported that insulin stimulated phosphorylation of the insulin receptor  $\beta$ -subunit in purified human stromal ovarian membranes (53). These studies thus have demonstrated insulin receptor phosphorylation both in human and in animal ovarian tissues; in regard to phosphorylation, therefore, insulin interaction with its ovarian receptor resembles that with receptors in other target tissues.

### C. Insulin or IGFs and ovarian steroidogenesis

The idea suggested by *in vivo* observations, that insulin affects ovarian hormone production, logically led to *in vitro* studies of the effects of insulin on ovarian cell steroidogenesis. It is known that ovarian granulosa, theca, or medullary tissue from premenopausal women, postmenopausal women, or women with PCO can secrete steroids in culture (60–63). Studies involving short term cultures of animal or human ovarian tissue have examined the effects of insulin or IGFs on ovarian cell steroidogenesis.

1. *Effects of insulin or IGFs on steroidogenesis in animal ovarian tissue.* Insulin-stimulated progesterone secretion has been demonstrated in porcine granulosa cells, in bovine granulosa cells, and in porcine thecal cells. Channing and co-workers (64) reported that insulin produced a 7- to 8-fold increase in progesterone secretion in cultured porcine granulosa cells. Veldhuis and co-workers (65) also studied the effects of insulin on cultured porcine granulosa cells; they found augmentation not only of progesterone secretion but also of progesterone synthesis (65). The peptide's effects were saturable and dose- and time-dependent. The process required synthesis of protein and RNA, augmentation of mitochondrial cytochrome P-450 content, and utilization of cholesterol in the side chain cleavage reaction.

Savion and co-workers (66) reported that insulin increased progesterone secretion by cultured bovine granulosa cells, and Barbieri and co-workers (67) reported that insulin increased progesterone secretion by cultured porcine thecal cells.

Insulin's stimulatory effects on ovarian steroidogenesis are not limited to progesterone production. Davoren and co-workers (52) showed that in cultured rat granulosa cells, insulin increased estrogen production, and Barbieri and co-workers (67) showed that in cultured porcine thecal cells, insulin increased androstenedione production.

These studies thus have demonstrated insulin-stimulated steroidogenesis in cultures of animal ovarian cells.

The regulation of granulosa cell function by IGFs has been the subject of many studies and an excellent review (9). IGFs stimulate steroidogenesis in porcine, bovine, or rat granulosa cells *in vitro* and act synergistically with FSH or LH (68–70).

2. *Effects of insulin or IGFs on steroidogenesis in human ovarian tissue.* Few studies have examined the effects of insulin or IGFs on steroidogenesis in human ovarian tissue. Barbieri and co-workers (71) studied the effects of insulin on incubated ovarian thecal or stromal fragments from a patient with hyperandrogenism, insulin resistance, and acanthosis nigricans. Insulin increased

accumulation of androstenedione, testosterone, progesterone, and estradiol in the thecal cell incubations, and increased accumulation of androstenedione, testosterone, and dihydrotestosterone in the stromal cell incubations. These investigators also incubated stromal fragments from four patients with hyperandrogenism and found that insulin increased accumulation of testosterone and androstenedione (72).

We reported that low concentrations (10 ng/ml) of insulin or IGF-1 stimulated progesterone production by cultured human granulosa cells. Monoclonal antibody to the IGF-1 receptor blocked the steroidogenic effect of IGF-1, but not of insulin, suggesting that each hormone induced progesterone synthesis through its own receptor (73).

These studies have established clearly that insulin or IGFs can stimulate ovarian steroidogenesis in various *in vitro* cell systems. The mechanisms by which insulin could possibly stimulate ovarian steroidogenesis are reviewed below.

**3. Possible mechanisms by which insulin or IGFs stimulate ovarian steroidogenesis.** The mechanisms by which insulin or IGFs stimulate ovarian steroidogenesis remain largely unknown. Possible mechanisms can be broadly classified into nonspecific or specific effects of insulin or IGFs on steroidogenesis. The nonspecific effects comprise the classic actions of insulin on glucose transport, on amino acid uptake, and on DNA synthesis; these effects would improve cell viability and consequently enhance steroidogenesis. The specific effects include a direct effect of insulin on steroidogenic enzymes (*e.g.* aromatase activity), synergism between insulin and LH or FSH, or modulation of LH receptor induction by insulin.

**Nonspecific effects of insulin on ovarian cells.** The first two possible nonspecific effects whereby insulin could enhance steroidogenesis are through enhanced glucose or amino acid metabolism. Allen and co-workers (74) found that insulin stimulated transport of 2-deoxyglucose and  $\alpha$ -amino-isobutyric acid in cultured granulosa cells from unbred heifers. Weber and co-workers (75) showed that IGF-1 (somatomedin-C) or insulin increased glucose oxidation in cultured porcine granulosa cells. These studies thus demonstrated the presence of insulin's classic effects in ovarian cells.

Several studies have assessed the effects of insulin or IGFs on the viability and morphology of cultured ovarian cells.

Channing and co-workers (64) demonstrated that insulin changed the morphology and the viability of cultured porcine granulosa cells. Insulin caused the cells to grow in an epitheloid manner and increased the number of intracellular lipid granules. After 2, 4, or 6 days in

culture, insulin produced a dose-dependent increase in cell number; the effect was attributed to improved plating efficiency and enhanced cell viability.

Savion and co-workers (66) demonstrated that insulin enhanced proliferation of cultured bovine granulosa cells. An insulin concentration of 0.03  $\mu$ g/ml significantly stimulated cell proliferation, whereas a concentration of 1.0  $\mu$ g/ml saturated the effect. Insulin increased cell density 6-fold and shortened the average population doubling time from 41 to 28 h. Somatomedin-C (IGF-1) in concentration as low as 1.0 ng/ml also was mitogenic, and the increase in cell number was dose dependent. Saturation of this effect was not achieved at the highest concentration tested (60 ng/ml). Somatomedin-C appeared to be more potent than insulin in stimulating cell proliferation. Insulin or somatomedin-C acted synergistically with fibroblast growth factor, epidermal growth factor, or high density lipoprotein; any of the latter three factors alone had no stimulatory effect on cell number.

May and Schomberg (76) demonstrated that insulin dramatically influenced monolayer growth and maintenance in porcine granulosa cell cultures; the effect was observed only during 24–146 h of incubation. Insulin also induced cultured swine granulosa cells to adopt an epitheloid appearance.

In summary, these studies showed that insulin or IGF-1 significantly affected the morphology and the viability of cultured porcine or bovine granulosa cells. Under the influence of insulin or of IGF-1, the cells acquired an epitheloid appearance; the number of lipid droplets and vacuoles increased; and cell plating efficiency, cell maintenance, and cell proliferation were stimulated. The mechanisms of these phenomena are the subject of debate (77).

**Specific effects of insulin or IGF-1 on steroidogenesis: Effects of insulin or IGF-1 on aromatase activity.** Insulin possibly stimulates ovarian steroidogenesis by influencing aromatase activity. Garzo and Dorrington (78) demonstrated that in cultured human granulosa cells, insulin alone or with FSH stimulated aromatase activity. Studies by Veldhuis and co-workers (65), in contrast, showed that in swine granulosa cells, insulin suppressed aromatization of testosterone to estradiol.

Adashi and co-workers (79) examined the effects of somatomedin C (IGF-1) on aromatase activity in cultured rat granulosa cells and found that the peptide potentiated FSH-induced aromatase activity. As these experiments varied in cell type studied and dose and type of peptide employed, it is difficult to compare their results or draw conclusions about the direction in which insulin or IGF-1 affects aromatase activity. That insulin or IGF-1 did affect aromatase activity in each of these studies, however, suggests that such an effect is involved in the regulation of ovarian cell steroidogenesis.

*Synergism between insulin or IGF-1 and LH or FSH.* Several studies have demonstrated that insulin and the gonadotropins act synergistically to augment ovarian cell steroidogenesis. Incubation of porcine granulosa cells (64, 65, 80) or rat granulosa cells (52) with insulin and the gonadotropins synergistically augmented progesterone production. Incubation of rat granulosa cells (52) or human granulosa cells (78) with insulin and the gonadotropins synergistically augmented estrogen production. Studies of thecal cells from swine (67) or stromal cells from normally cycling women (71, 72) showed synergistically augmented androstenedione production.

Adashi and co-workers (69) demonstrated synergism between IGF-1 and FSH on induction of progesterone synthesis in cultured rat granulosa cells; progesterone synthesis was stimulated to a level several times higher than the maximal level achieved with FSH alone.

The mechanism for this synergism is largely unknown. Adashi and co-workers (81) showed that IGF-1 enhanced FSH-induced intracellular accumulation of cAMP. This finding is very interesting, as neither insulin nor IGF-1 alone affects intracellular cAMP levels (44), whereas the gonadotropins do (82). Possibly the synergism between the gonadotropins and IGF-1 or insulin (the latter acting through the IGF-1 receptor) is mediated through increased intracellular cAMP production.

It is of interest that IGF-1 acts synergistically not only with the pituitary gonadotropins but also with estradiol [in granulosa cells (83)] and with TSH [in cultured thyroid cells (84)]. These findings suggest the exciting possibility that IGF-1 acts as a universal amplifier of various hormonal stimuli.

*Insulin or IGF-1 modulation of LH-receptor induction.* A third possible specific mechanism whereby insulin or IGF-1 could stimulate ovarian steroidogenesis is through modulation of LH receptor induction. An increased number of LH receptors would enhance steroidogenic cell responsiveness to LH, resulting in stimulation of steroidogenesis. Support for this hypothesis has come from several studies. Davoren and co-workers (52) found that insulin enhanced induction of LH receptors in rat granulosa cells (52). Adashi and co-workers (85) showed that IGF-1 enhanced FSH induction of LH receptors in cultured rat granulosa cells. Mondschein and Schomberg (86, 87) found enhanced induction of LH receptors with other growth factors, suggesting that this mechanism of action may be common for a variety of peptides.

4. *Why does insulin-induced ovarian hyperstimulation manifest primarily as hyperandrogenism?* It is not quite clear why insulin-induced ovarian hyperstimulation manifests primarily as hyperandrogenism. One possible explanation involves the idea that insulin inhibits aromatase and could thus produce a relative preponderance

of androgens. Several studies investigating this hypothesis have yielded differing results (see *Effects of insulin or IGF-1 on aromatase activity*).

A second possible explanation is that insulin affects exclusively the ovary's androgen-producing cells (stromal or thecal cells). Multiple *in vitro* studies with animal granulosa cells and with human granulosa cells have demonstrated that insulin also stimulates progesterone and estrogen production (see *Insulin or IGFs and ovarian steroidogenesis*), suggesting that the peptide's gonadotropic effects are not limited to the ovary's androgen-producing cells.

A third potential explanation is based on the fact that the mass of androgen-producing cells in the ovary exceeds that of estrogen- or progesterone-producing cells. Thus, even if the gonadotropic action of insulin were directed equally at all ovarian cells, hyperandrogenism would be a predominant feature. The merits of this simple explanation have not been tested.

We propose yet another explanation. It is known that LH induces differentiation of ovarian interstitial cells into androgen-producing cells and stimulates androgen production by these cells (88). As insulin acts synergistically with LH *in vitro* (see *Synergism between insulin or IGF-1 and LH or FSH*), perhaps hyperinsulinemia enhances LH-induced differentiation of ovarian interstitial cells and LH-induced androgen production, resulting in androgen excess. The androgen excess would induce follicular atresia, with loss of estrogen-producing cells. Androgen-producing cells would replace the atretic follicles (88) and could perpetuate the cycle by producing more androgens in response to LH and insulin. Hyperinsulinemia thus would produce an excess of androgens, but not of the other ovarian hormones.

#### D. *Insulin and IGFs in ovarian fluid*

The hypothesis that insulin or IGFs affect ovarian steroidogenesis could attain physiological significance only if it were shown that the peptides had access to ovarian steroidogenic cells *in vivo*.

Because some ovarian cells have limited access to the blood supply, it is important to determine whether insulin or IGFs are present in follicular fluid. Diamond and co-workers (89) recently reported that insulin was present in human follicular fluid and that follicular fluid insulin concentrations correlated positively with follicular fluid progesterone levels. Hammond and co-workers (90) showed that *in vitro* porcine granulosa cells produced IGFs and IGF binding-proteins, raising the possibility that the ovary produces IGF-1 *in vivo*. IGF-levels in preovulatory follicular fluid were significantly greater than those in either serum or immature follicles, further supporting this possibility.



These studies provide evidence that insulin and IGFs have access to ovarian steroidogenic cells *in vivo*; the correlation between follicular fluid insulin levels and progesterone levels lends further support to the idea that insulin affects ovarian steroidogenesis *in vivo*.

#### IV. Summary

We have reviewed the role of insulin in ovarian physiology. Clinical observations and experimental data strongly support the hypothesis that insulin possesses gonadotropic activity, when acting alone or with FSH or LH. This idea is further supported by the recent discovery of insulin in follicular fluid.

The idea that insulin has gonadotropic function can explain a variety of clinical observations, which otherwise are difficult to understand. For example, manifestations of ovarian hypofunction (primary amenorrhea, late menarche, anovulation, low pregnancy rate, and early menopause) in IDDM can be understood if it is accepted that insulin is necessary for the ovary to reach its full steroidogenic potential. The idea that insulin affects ovarian steroidogenesis also helps to understand the observation that hyperandrogenism frequently accompanies each of the various insulin-resistant states, regardless of the latter's etiology (e.g. genetic deficiency

in the number of insulin receptors, antiinsulin receptor antibodies, obesity, etc.). The explanation for this association (Fig. 1) is based on the idea that hyperinsulinemia intensifies ovarian steroidogenesis, which manifests clinically as hyperandrogenism. Continuous stimulation of the ovary by insulin over a long period of time possibly produces morphological ovarian changes, such as hyperthecosis or polycystic changes; these changes commonly are observed among women with insulin resistance.

The effects of insulin on ovarian cells are mediated possibly through binding of the peptide to its own receptor or to the IGF-1 receptor (the specificity spillover phenomenon). The latter could be an important mechanism in cases of insulin resistance.

Potential mechanisms underlying the gonadotropic activity of insulin include direct effects on steroidogenic enzymes, modulation of FSH or LH receptor number, synergism with FSH or LH, or nonspecific enhancement of cell viability.

The gonadotropic function of insulin adds yet another note to what has been termed a symphony of insulin action. Further investigation into this new area may yield greater insights not only into normal ovarian physiology, but also into the pathogenesis of such diverse entities as PCO, obesity, diabetes mellitus, and the syndromes of insulin resistance and acanthosis nigricans.

#### Acknowledgments

We wish to thank Dr. Barnett Zumoff of The Mount Sinai School of Medicine and Dr. Jeffrey S. Flier and Dr. Alan C. Moses of Harvard Medical School for their helpful comments.

#### References

1. Joslin EP, Root HF, White P 1985 The growth, development and prognosis of diabetic children. *JAMA* 85:420
2. Bergqvist N 1954 The gonadal function in female diabetics. *Acta Endocrinol [Suppl] (Copenh)* 19:3
3. Djursing H, Nyholm HC, Hagen C, Carstensen L, Pedersen LM 1982 Clinical and hormonal characteristics in women with anovulation and insulin-treated diabetes mellitus. *Am J Obstet Gynecol* 143:876-882.
4. Walton C, Kelly WF, Laing I, Bu'lock DE 1983 Endocrine Abnormalities in idiopathic haemochromatosis. *Q J Med* 205:99
5. Rueda R, Urdinola J, Is insulin response in hyperandrogenism LH mediated? Program of the 41st Annual Meeting of the American Fertility Society, Chicago, IL, 1985, p 49
6. Distiller LA, Sagel J, Morley JE, Joffe BI, Seftel HC 1975 Pituitary responsiveness to luteinizing hormone-releasing hormone in insulin-dependent diabetes mellitus. *Diabetes* 24:378-380
7. Liu FTY, Lin HS, Johnson DC 1972 Serum FSH, LH and the ovarian response to exogenous gonadotropins in alloxan diabetic immature female rats. *Endocrinology* 91:1172
8. Bestetti G, Locatelli V, Tirone F, Rossi GL, Muller EE 1985 One month of streptozotocin-diabetes induces different neuroendocrine and morphological alterations in the hypothalamo-pituitary axis of male and female rats. *Endocrinology* 117:208
9. Adashi EY, Resnick CE, D'Ercole AJ, Svoboda ME, Van Wyk JJ 1985 Insulin-like growth factors as intraovarian regulators of granulosa cell growth and function. *Endocr Rev* 6:400
10. Maes M, Underwood LE, Ketelslegers JM 1986 Low serum soma-




FIG. 1. The proposed pathogenesis of hyperandrogenism in insulin-resistant states is illustrated above. Insulin resistance of any etiology (e.g. genetic abnormalities of the insulin receptor, antiinsulin receptor antibodies, obesity, etc.) produces hyperinsulinemia. The massive quantity of insulin binds to ovarian insulin receptors and ovarian IGF-1 receptors and acts synergistically with FSH and LH to produce ovarian hyperstimulation. The hyperstimulation may produce hyperthecosis or polycystic changes in the ovary. Hyperandrogenism results either through these changes or directly from ovarian hyperstimulation.



- tomedin-C in insulin dependent diabetes: evidence for a postreceptor mechanism. *Endocrinology* 118:377
11. Tan K, Baxter RC 1986 Serum insulin-like growth factor I levels in adult diabetic patients: the effect of age. *J Clin Endocrinol Metab* 63:651
  12. Yki-Jarvinen H, Koivisto VA 1986 Natural course of insulin resistance in type I diabetes. *N Engl J Med* 315:224
  13. Kahn CR, Flier JS, Bar RS, Archer JA, Gorden P, Martin MM, Roth J 1976 The syndromes of insulin resistance and acanthosis nigricans. Insulin-receptor disorders in man. *N Engl J Med* 294:739
  14. Flier JS, Kahn CR, Jarrett DB, Roth J 1976 Characterization of antibodies to the insulin receptor. A cause of insulin-resistant diabetes in man. *J Clin Invest* 58:1442
  15. Flier JS, Kahn CR, Roth J, Bar RS 1975 Antibodies that impair insulin receptor binding in an unusual diabetic syndrome with severe insulin resistance. *Science* 190:63
  16. Flier JS, Kahn R, Roth J 1979 Receptors, antireceptor antibodies and mechanisms of insulin resistance. *N Engl J Med* 300:413
  17. Bar RS, Muggeo M, Roth J, Kahn CR, Havrankova J, Imperato-McGinley J 1978 Insulin resistance, acanthosis nigricans, and normal insulin receptors in a young woman: evidence for a postreceptor defect. *J Clin Endocrinol Metab* 47:620
  18. Flier JS, Young JB, Landsberg L 1980 Familial insulin resistance with acanthosis nigricans, acral hypertrophy, and muscle cramps. *N Engl J Med* 303:970
  19. Grigorescu F, Flier JS, Kahn CR 1984 Defect in insulin receptor phosphorylation in erythrocytes and fibroblasts associated with severe insulin resistance. *J Biol Chem* 259:15003
  20. Grunberger G, Zick Y, Gorden P 1984 Defect in phosphorylation of insulin receptors in cells from an insulin-resistant patient with normal insulin binding. *Science* 223:932
  21. Flier JS, Grigorescu F, Chaiken RL, Moses AC, Minaker KL, Rowe JW, Eastman RC 1985 Pseudoacromegaly: a syndrome apparently caused by hyperinsulinemia and insulin resistance with the receptor kinase defect. *Clin Res* 33:569A (Abstract)
  22. Plehwe WE, Maitland JE, Williams PF, Shearman PR, Turtle JR 1985 Familial hyperinsulinemia complicated by extreme insulin resistance during pregnancy: a probable postreceptor defect. *J Clin Endocrinol Metab* 61:68
  23. Schwenk WF, Rizza RA, Mandarino LJ, Gerich JE, Hayles AB, Hammond MW 1986 Familial insulin resistance and acanthosis nigricans. Presence of a postbinding defect. *Diabetes* 35:33
  24. Grigorescu F, Flier JS, Kahn CR 1986 Characterization of binding and phosphorylation defects of erythrocyte insulin receptors in the type A syndrome of insulin resistance. *Diabetes* 35:127
  25. Taylor SI 1985 Receptor defects in patients with extreme insulin resistance. *Diabetes/Metab Rev* 1:171
  26. Rossini AA, Cahill GF 1979 Lipoatrophic diabetes. In: DeGroot LJ, Cahill GF, Martini L, Nelson DH, Odell WD, Potts JT, Steinberger E, Winegrad AI (eds) *Endocrinology*. Grune & Stratton, New York, 1093-1097
  27. D'Ercole AJ, Underwood LE, Groelke J, Plet A 1979 Leprechaunism: studies of the relationship among hyperinsulinism, insulin resistance, and growth retardation. *J Clin Endocrinol Metab* 48:495
  28. Rechler MM 1982 Leprechaunism and related syndromes with primary insulin resistance: heterogeneity of molecular defects. In: Sheppard J, Anderson EV, Eaton J (eds) *Membranes and Genetic Disease*. Liss, New York, p 245
  29. Rabson SM, Mendenhall EN 1956 Familial hypertrophy of the pineal body, hyperplasia of the adrenal cortex and diabetes mellitus: report of three cases. *Am J Clin Pathol* 26:282
  30. Taylor SI, Underhill LH, Hedo JA, Roth J, Rios MS, Blizzard RM 1983 Decreased insulin binding to cultured cells from a patient with the Rabson-Mendenhall syndrome: dichotomy between studies with cultured lymphocytes and cultured fibroblasts. *J Clin Endocrinol Metab* 56:856
  31. Flier JS, Eastman RC, Minaker KL, Matteson D, Rowe JW 1985 Acanthosis nigricans in obese women with hyperandrogenism: a marker for insulin resistance and distinction from the type A and B syndromes. *Diabetes* 34:101
  32. Chang RJ, Nakamura RM, Judd HL, Kaplan SA 1983 Insulin resistance in nonobese patients with polycystic ovarian disease. *J Clin Endocrinol Metab* 57:356
  33. Burghen GA, Givens JR, Kitabchi AE 1980 Correlation of hyperandrogenism with hyperinsulinism in polycystic ovarian disease. *J Clin Endocrinol Metab* 50:113
  34. Dunaif A, Graf M, Laumas V, Dobrjansky A, Determinants of insulin resistance in hyperandrogenic women. Program and Abstracts of the 67th Annual Meeting of The Endocrine Society, Baltimore, MD, 1985, p 123 (Abstract 622)
  35. Taylor SI, Dons RF, Hernandez E, Roth J, Gorden P 1982 Insulin resistance associated with androgen excess in women with autoantibodies to the insulin receptor. *Ann Intern Med* 97:851
  36. McNatty KP, Smith DM, Makris A, DeGrazia C, Tulchinsky D, Osathanondh R, Schiff I, Ryan KJ 1980 The intraovarian sites of androgen and estrogen formation in women with normal and hyperandrogenic ovaries as judged by *in vitro* experiments. *J Clin Endocrinol Metab* 50:755
  37. Cole C, Kitabchi AE 1978 Remission of insulin resistance with Orthonovum<sup>R</sup> in a patient with polycystic ovarian disease and acanthosis nigricans. *Clin Res* 26:412A (Abstract)
  38. Shoupe D, Lobo RA 1984 The influence of androgens on insulin resistance. *Fertil Steril* 41:385
  39. Stuart CA, Peters EJ, Prince MJ, Richards G, Cavallo A, Meyer WJ 1986 Insulin resistance with acanthosis nigricans: the roles of obesity and androgen excess. *Metabolism* 35:197
  40. Geffner ME, Kaplan SA, Bersch N, Golde DW, Landaw EM, Chang RZ 1986 Persistence of insulin resistance in polycystic ovarian disease after inhibition of ovarian steroid secretion. *Fertil Steril* 45:327
  41. Barbieri RL, Ryan KJ 1983 Hyperandrogenism, insulin resistance, and acanthosis nigricans syndrome: a common endocrinopathy with distinct pathophysiologic features. *Am J Obstet Gynecol* 147:90
  42. Nestler JE, Clore JN, Strauss JF, Blackard WG 1987 The effects of hyperinsulinemia on serum testosterone, progesterone, dehydroepiandrosterone sulfate, and cortisol levels in normal women and in a woman with hyperandrogenism, insulin resistance, and acanthosis nigricans. *J Clin Endocrinol Metab* 64:180
  43. Roth J, Grunfeld C 1985 Mechanism of action of peptide hormones and catecholamines. In: Wilson JD, Foster DW (eds). *Textbook of Endocrinology*. Philadelphia, WB Saunders, p 96
  44. Kahn CR, Crettaz M 1985 Insulin receptors and the molecular mechanism of insulin action. *Diabetes/Metab Rev* 1:5
  45. Kasuga M, Karlsson FA, Kahn CR 1982 Insulin stimulates the phosphorylation of the 95,000-Dalton subunit of its own receptor. *Science* 215:185
  46. Nissley SP, Rechler MM 1984 Somatomedin/insulin-like growth factor tissue receptors. *Clin Endocrinol Metab* 13:43
  47. Zapf J, Schmid CH, Froesch ER 1984 Biological and immunological properties of insulin-like growth factors (IGF) I and II. *Clin Endocrinol Metab* 13:3
  48. Chernauek SD, Jacobs S, Van Wyk JJ 1981 Structural similarities between human receptors for somatomedin C and insulin: analysis by affinity labeling. *Biochemistry* 20:7345
  49. Jacobs S, Kull FC, Earp HS, Svoboda M, Van Wyk JJ, Cuatrecasas P 1983 Somatomedin-C stimulates the phosphorylation of the beta-subunit of its own receptor. *J Biol Chem* 258:9581
  50. Rein MS, Schomberg DW 1982 Characterization of insulin receptors on porcine granulosa cells. *Biol Reprod* 26 [Suppl 1]:113
  51. Velduis JD, Furlanetto RW, Juchter D, Garmuy J, Veldhuis P 1985 Trophic actions of human somatomedin C/insulin-like growth factor I on ovarian cells: *in vitro* studies with swine granulosa cells. *Endocrinology* 116:1235
  52. Davoren JB, Kasson BG, Li CH, Hsueh AJW 1986 Specific insulin-like growth factor (IGF) I- and II-binding sites on rat granulosa cells: relation to IGF action. *Endocrinology* 119:2155
  53. Poretsky L, Grigorescu F, Seibel M, Moses AC, Flier JS 1985 Distribution and characterization of insulin and insulin-like growth factor I receptors in normal human ovary. *J Clin Endocrinol Metab* 61:728
  54. Poretsky L, Smith D, Seibel M, Pazianes A, Moses AC, Flier JS

- 1984 Specific insulin binding sites in human ovary. *J Clin Endocrinol Metab* 59:809
55. Jarrett JC, Ballejo G, Tsibris JCM, Spellacy WN 1985 Insulin binding to human ovaries. *J Clin Endocrinol Metab* 60:460
  56. Grunberger G, Comi RJ, Taylor SI, Gorden P 1984 Tyrosine kinase activity of the insulin receptor of patients with type A extreme insulin resistance: studies with circulating mononuclear cells and cultured lymphocytes. *J Clin Endocrinol Metab* 59:1152
  57. Grigorescu F, White MF, Kahn CR 1983 Insulin binding and insulin-dependent phosphorylation of the insulin receptor solubilized from human erythrocyte. *J Biol Chem* 258:13708
  58. Grigorescu F, King GL, Kahn CR 1984 Demonstration of a genetic defect in the insulin receptor kinase of cultured fibroblasts in type A syndrome of insulin resistance. Program of the 7th Annual International Congress of Endocrinology, Quebec City, Canada, 1984. *Excerpta Medica*, Amsterdam, p 927 (Abstract 1333)
  59. Veldhuis JD, Tamura S, Kolp L, Furlanetto RW, Lerner J 1984 Mechanisms subserving insulin action in the gonad: evidence that insulin induces specific phosphorylation of its immunoprecipitable receptor on ovarian cells. *Biochem Biophys Res Commun* 120:144
  60. McNatty KP, Makris A, DeGrazia C, Osathanondh R, Ryan KJ 1979 The production of progesterone, androgens, and estrogens by granulosa cells, thecal tissue, and stromal tissue from human ovaries *in vitro*. *J Clin Endocrinol Metab* 49:687
  61. Dennefors BL, Janson PO, Knutson F, Hamberger L 1980 Steroid production and responsiveness to gonadotropin in isolated stromal tissue of human postmenopausal ovaries. *Am J Obstet Gynecol* 136:997
  62. Batta SK, Wentz AC, Channing CP 1980 Steroidogenesis by human ovarian cell types in culture: influence of mixing of cell types and effect of added testosterone. *J Clin Endocrinol Metab* 50:274
  63. Wilson EA, Erickson GF, Zarutski P, Finn AE, Tulchinsky D, Ryan KJ 1979 Endocrine studies of normal and polycystic ovarian tissues *in vitro*. *Am J Obstet Gynecol* 134:56
  64. Channing CP, Tsai V, Sachs D 1976 Role of insulin, thyroxine and cortisol in luteinization of porcine granulosa cells grown in chemically defined media. *Biol Reprod* 15:235
  65. Veldhuis JD, Kolp LA, Toaff ME, Strauss JF, Demers LM 1983 Mechanisms subserving the trophic actions of insulin on ovarian cells: *in vitro* studies using swine granulosa cells. *J Clin Invest* 72:1046
  66. Savion N, Liu GM, Laherty R, Gospodarowicz D 1981 Factors controlling proliferation and progesterone production by bovine granulosa cells in serum-free medium. *Endocrinology* 109:409
  67. Barbieri RL, Makris A, Ryan KJ 1983 Effects of insulin on steroidogenesis in cultured porcine ovarian theca. *Fertil Steril* 40:237
  68. Adashi EY, Resnick CE, Svoboda ME, Van Wyk JJ 1984 A novel role for somatomedin-C in the cytodifferentiation of the ovarian granulosa cell. *Endocrinology* 115:1227
  69. Adashi EY, Resnick CE, Svoboda ME, Van Wyk JJ 1985 Somatomedin-C synergizes with follicle-stimulating hormone in the acquisition of progesterone biosynthetic capacity by cultured rat granulosa cells. *Endocrinology* 116:2135
  70. Adashi EY, Resnick CE, Svoboda ME, Van Wyk JJ, Hascall VC, Masaki Y 1986 Independent and synergistic actions of somatomedin-C in the stimulation of proteoglycan biosynthesis by cultured rat granulosa cells. *Endocrinology* 118:456
  71. Barbieri RL, Makris A, Ryan KJ 1984 Insulin stimulates androgen accumulation in incubations of human ovarian stroma and theca. *Obstet Gynecol* 64:73S
  72. Barbieri RL, Makris A, Randall RW, Daniels G, Kistner RW, Ryan KJ 1986 Insulin stimulates androgen accumulation in incubations of ovarian stroma obtained from women with hyperandrogenism. *J Clin Endocrinol Metab* 62:904
  73. Poretsky L, Grigorescu F, Flier JS, Insulin but not IGF-I receptors are widely distributed in normal human ovary. Program of the 67th Annual meeting of The Endocrine Society, Baltimore, MD, p 151 (Abstract 814)
  74. Allen WR, Nilsen-Hamilton M, Hamilton RT 1981 Insulin and growth factors stimulate rapid posttranslational changes in glucose transport in ovarian granulosa cells. *J Cell Physiol* 108:15
  75. Weber SL, LaBarbera AR, Human choriogonadotropin, somatomedin C and insulin increase glucose oxidation by granulosa cells. Program of the 67th Annual Meeting of The Endocrine Society, Baltimore, MD, 1985, p 151 (Abstract 818)
  76. May JV, Schomberg DW 1981 Granulosa cell differentiation *in vitro*: effect of insulin on growth and functional integrity. *Biol Reprod* 25:421
  77. Strauss DS 1981 Effects of insulin on cellular growth and proliferation. *Life Sci* 29:2131
  78. Garzo G, Dorrington JH 1984 Aromatase activity in human granulosa cells during follicular development and the modulation by follicle-stimulating hormone and insulin. *Am J Obstet* 148:657
  79. Adashi EY, Resnick CE, Brodie AMH, Svoboda ME, Van Wyk JJ 1985 Somatomedin-C-mediated potentiation of follicle-stimulating hormone-induced aromatase activity of cultured rat granulosa cells. *Endocrinology* 117:2313
  80. Lino J, Baranao S, Hammond JM 1985 Multihormone regulation of steroidogenesis in cultured porcine granulosa cells: studies in serum-free medium. *Endocrinology* 116:2143
  81. Adashi EY, Resnick CE, Svoboda ME, Van Wyk JJ 1986 Somatomedin-C as an amplifier of follicle-stimulating hormone action: enhanced accumulation of adenosine 3',5'-monophosphate. *Endocrinology* 118:149
  82. Catt KJ, Pierce JG 1986 Gonadotropic hormones of the adenohypophysis. In: Yen SSC, Jaffe RB (eds) *Reproductive Endocrinology*. WB Saunders Co, Philadelphia, pp 75-114
  83. Veldhuis JD, Rodgers RJ, Furlanetto RW, Azimi P, Tuchten D, Garmey J 1986 Synergistic actions of estradiol and the insulin-like growth factor somatomedin-C on swine ovarian (granulosa) cells. *Endocrinology* 119:530
  84. Tramontano D, Cushing GW, Moses AC, Ingbar SH 1986 Insulin-like growth factor-I stimulates the growth of rat thyroid cells in culture and synergizes the stimulation of DNA synthesis induced by TSH and Graves'-IgG. *Endocrinology* 119:940
  85. Adashi EY, Resnick CE, Svoboda ME, Van Wyk JJ 1985 Somatomedin-C enhances induction of luteinizing hormone receptors by follicle-stimulating hormone in cultured rat granulosa cells. *Endocrinology* 116:2369
  86. Mondschein JS, Schomberg DW 1981 Growth factors modulate gonadotropin receptor induction in granulosa cell cultures. *Science* 211:1179
  87. Mondschein JS, Schomberg DW 1981 Platelet-derived growth factor enhances granulosa cell luteinizing hormone receptor induction by follicle-stimulating hormone and serum. *Endocrinology* 109:325
  88. Erickson GF, Magoffin DA, Dyer CA, Hofeditz C 1985 The ovarian androgen producing cells: a review of structure/function relationships. *Endocr Rev* 6:371
  89. Diamond MP, Webster BW, Carr RK, Wentz AC, Osteen KG 1985 Human follicular fluid insulin concentrations. *J Clin Endocrinol Metab* 61:990
  90. Hammond JM, Baranao JLS, Skaleris D, Knight AB, Romanus JA, Rechler MM 1985 Production of insulin-like growth factors by ovarian granulosa cells. *Endocrinology* 117:2553