

# The Presence of Endogenous Morphine Signaling in Animals

George B. Stefano · Patrick Cadet ·  
Richard M. Kream · Wei Zhu

Accepted: 13 March 2008 / Published online: 6 September 2008  
© Springer Science+Business Media, LLC 2008

**Abstract** Recent empirical findings have contributed valuable mechanistic information in support of a regulated *de novo* biosynthetic pathway for chemically authentic morphine and related morphinan alkaloids within animal cells. Importantly, we and others have established that endogenously expressed morphine represents a key regulatory molecule effecting local circuit autocrine/paracrine cellular signaling via a novel  $\mu_3$  opiate receptor coupled to constitutive nitric oxide production and release. The present report provides an integrated review of the biochemical, pharmacological, and molecular demonstration of  $\mu_3$  opiate receptors in historical linkage to the elucidation of mechanisms of endogenous morphine production by animal cells and organ systems. Ongoing research in this exciting area provides a rare window of opportunity to firmly establish essential biochemical linkages between dopamine, a morphine precursor, and animal biosynthetic pathways involved in morphine biosynthesis that have been conserved throughout evolution.

**Keywords** Morphine · Opiate receptors · Nitric oxide ·  $\mu_3$  Opiate receptor

## Demonstration and Characterization of Novel $\mu_3$ Opiate Receptors

The first demonstration by Kosterlitz and coworkers [1] that exogenous morphine can bind to receptors in the

mammalian brain indicated that morphine binds to the same sites as those used by the endogenous opioid peptides (e.g., enkephalins). Since then, demonstration of the multiplicity of receptor types has led to the understanding that, depending on their site of action, opioid peptides as well as opiate alkaloids may bind to more than one opiate receptor subtype [2–5]. Different degrees of selectivity have been recognized for the various ligands by comparing affinity constants as well as relative strength in competitive binding assays. For example, Pasternak and Snyder (see [6]) reported both high and low affinity binding sites for [ $^3$ H]-dihydromorphine ( $^3$ DHM) and [ $^3$ H]-naloxone in the rat brain. The higher affinity type was designated  $\mu_1$  and the lower affinity morphine-selective type was designated  $\mu_2$  [7, 8].

Our laboratory was instrumental in the biochemical identification and characterization of a novel  $\mu$  opiate receptor subtype, designated  $\mu_3$ , located on immunocytes and neural tissues of invertebrates, such as *Mytilus edulis*, as well as human monocytes, granulocytes, vascular endothelial cells and other neuronal and non-neuronal cell types (Table 1) [9, 18, 19]. The  $\mu_3$  receptor differed from previously described neuronal opioid receptor subtypes in that it exhibited non-detectable or exceedingly low affinities for naturally occurring endogenous opioid peptides, peptide analogues,  $\mu_1$  opioid-selective endomorphins 1 and 2, and synthetic opioid alkaloids of the benzomorphan and phenylpiperidine classes [9, 16–19, 29]. In contrast, the  $\mu_3$  opiate receptor displayed high affinity binding for morphine-related morphinan alkaloids such as dihydromorphine and the clinically established antagonists naloxone and naltrexone as well as the active metabolite morphine 6-glucuronide [30]. Finally, it was also established that the  $\mu_3$  receptor is coupled to G protein isoforms, based on guanine nucleotide effects on agonist binding [19].

---

Special issue article in honor of Dr. Ji-Sheng Han.

---

G. B. Stefano (✉) · P. Cadet · R. M. Kream · W. Zhu  
Neuroscience Research Institute, State University of New York –  
College at Old Westbury, P.O. Box 210, Old Westbury, NY  
11568, USA  
e-mail: gstefano@sunynri.org

**Table 1**  $\mu_3$  Receptor binding and expression $\mu_3$  Opiate receptor binding*Tissues or cell lines**Mytilus edulis*: immunocytes [9], ganglia [10–14]

Cat: astrocytes and microglia [15]

Rat: Kupffer cells [15]

Murine: cell lines (J774.2, RAW 264.7, BAC1.2F5) [16], mouse neuroblastoma [17]

Human tissues: vascular [18], white blood cells [9, 19], neural [20]

## Receptor expression

*Mytilus edulis*: ganglia [21, 22]

Human tissues: Sh-Sy5y, [23], heart [24, 25], vascular [23, 24, 26], white blood cells [23, 27], neural [20], stem cells [28]

**Functional Coupling of  $\mu_3$  Opiate Receptors to Constitutive Nitric Oxide Production and Release**

The neurochemical literature has documented a functional association of morphine action with stimulated nitric oxide (NO) production and release. Peripheral morphine analgesia involves NO-stimulated increases in intracellular cGMP [31]. Nitric oxide has been associated with antinociception [32] as well as tolerance and dependence [33]. In addition, the morphine-induced suppression of splenic lymphocyte proliferation has been shown to involve NO [34]. Morphine and NO have been linked in gastrointestinal regulation [35, 36]. Furthermore, morphine, not opioid peptides, stimulate constitutive NO release in macrophages, granulocytes, various types of human and rat

endothelial cells, invertebrate neurons and immunocytes and in rat median eminence fragments, all in a naloxone antagonizable manner [29, 30, 37–44]. These data suggest that the  $\mu_3$  receptor is coupled to constitutive NO release in these cells [39, 45, 46].

Furthermore, morphine's actions in these diverse tissues complements what is known about NO mediating immune and vascular functions, namely that it can down regulate them from an excitatory state or prevent the excitatory state from occurring [30, 42, 44, 47]. Additional information on opiate alkaloid signaling substances can be summarized as follows: Injection of vertebrate animals with morphine results in deficient macrophage function [48] and an alteration of T-cell activity [49]. Morphine also antagonizes interleukin-1 $\alpha$  or tumor necrosis factor- $\alpha$ -induced chemotaxis in human granulocytes and monocytes [50, 51].

**Molecular Cloning, Structural Elucidation, and Functional Cellular Expression of the  $\mu_3$  Opiate Receptor**

Molecular cloning of a  $\mu_3$  receptor encoding cDNA employed a screening probe derived from a conserved region of the human  $\mu_1$  receptor and a human testis cDNA library. A full-length 1,338 base pair cDNA was cloned and subsequently sequenced (Fig. 1 and Table 1) [23]. NCBI Blast analysis indicated that the clone exhibited 100% identity to the  $\mu_1$  opioid receptor subtype in its central conserved region with most of exon 1 spliced out at

**Fig. 1** (a)  $\mu_3$  cDNA sequence (1,338 bp). The *underlined* sequence represents the novel 263 bp segment, and the *bold* letters represents *Homo sapiens* Oprm 3'UTR DNA sequence (nucleotide position 1,625–1,829 of Oprm). (b) The diagram shows the  $\mu_3$  cDNA sequence starting at position 503 of Oprm. The dotted lines represent the novel 263 bp  $\mu_3$  sequence (between nucleotides 1,376 and 1,625 of Oprm), and position 1,625–1,829 is the Oprm 3'UTR DNA sequence that represents the 3'-end of  $\mu_3$  [23]

**A**

5'atacacaagatgaagactgccaccaacatctacattttcaaccttgctctggcagatgccttagccaccagtagccctgccttc  
cagagtgtgaattacctaattggaacatggccatttgaaccatccttgcagatagtgatctccatagattactataacatgttca  
ccagcatattcaccctctgcccatgagtggtgatcgatactgcagcttgccaccctgtcaaggccttagatttccgtactccccg  
aaatgcaaaattatcaatgtctgcaactggatcctctcttcagccattggtcttctgtaatgttcattgctacaacaaaatacaggc  
aaggttccatagattgtacactaacattctctcatccaacctggtagtggaacacctgtgaagatctgtgtttcatcttcgccttca  
ttatgccagtgtcatcattaccgtgtgctatggactgatgatcttgcgcctcaagagtgtccgcatgctctgtgctccaaagaaa  
aggacaggaatcttcgaaggatcaccaggatggtgctggtggtggtggtggtgtgttcacgtctgctggtgactcccatcatttac  
gtcatcattaaagccttggttaacatcccagaaactacgtccagactgtttcttggcacttctgcattgcttaggttacacaaacag  
ctgcctcaaccagctccttattgcatcttggatgaaactcaaacgatgcttcagagagttctgtatcccaacctcttccaacattg  
agcaaaaaactccactcgaattcgtcagaacactagagaccacctccacggccaatacagtgatagaactaatcatcaga  
attattatataattcatagatgttgctgcaataccctcttatttctcaaaagccagcttctgctgtgtgattaaagagagaggggt  
gagtgcccttgcctcactgtggtcatggtgcaagatattcacagaaaattagcatcatagaaaaaaannnnnnnnnnnnnnnnnnnn  
aaaancatgtcggccgctcggcgaacatcgggtcgagcatgcatctaggcggccaattccgcccctctccccccnngcn  
ntttcacaccgaggagtcagtttgtgcaagacaccagcgggaaccaaaccatcgtggtatgtgaatcgaagtcac  
ataaaaggtgaccttctgtctgtaagattttaatttaagcatatatttatgacctcaacaaagacgaacctctttgttaa  
ttaccgtagtaacacataaagttatgctacctctgatcaag-3'

**B**

MOR-1: 1 \_\_\_\_\_ -2162

MOR-3: 1 \_\_\_\_\_ 1338

its 5'-end (Fig. 1). The 3'-end of the  $\mu_3$  receptor cDNA contained the 3'-end of the  $\mu_1$  receptor protein coding sequence with a spliced insert of 263 bases containing a stop codon and terminated by a  $\mu_1$ -specific 202 base untranslated region [23].

RT-PCR and subsequent sequence analysis revealed the presence of this opiate receptor subtype in human vascular endothelial cells, mononuclear cells and polymorphic nuclear cells. To determine if the cDNA clone we isolated was functional and had the biochemical properties expected of the  $\mu_3$  receptor, the cDNA was expressed in a heterologous system (Cos-1 cells). Following exposure to morphine, the transfected Cos-1 cells released NO in a naloxone-reversible manner [23]. Untransfected Cos-1 cells failed to produce any detectable NO upon addition of morphine [23]. The addition of Met-enkephalin, DPDPE, or Leu-enkephalin did not stimulate NO release in the controls or transfected cells [23]. In sum, heterologous cellular expression of the cloned  $\mu_3$  opiate receptor cDNA by Cos-1 cells resulted in morphine-selective stimulation of NO release, consistent with previous biochemical data presented above [23, 52, 53].

### Endogenous Morphine

Prior demonstration of low concentrations of morphine [47, 54–59] and its precursor molecules tetrahydropapaveroline (THP), reticuline, salutaridine, thebaine, and codeine in vertebrate tissues [58, 60–62] provides strong presumptive evidence for the physiological relevance of novel  $\mu_3$  opiate receptors selectively activated by endogenously expressed morphinan alkaloids. In invertebrates, specifically *Mytilus edulis*, the presence of morphine, morphine 6-glucuronide, morphine 3-glucuronide, codeine, THP and reticuline also have been reported [9, 53, 63–65]. Endogenous opiate levels can be induced to change following stimulation [53, 66–70]. Morphine has also been found in human plasma [71, 72], suggesting a hormonal action with immune, vascular and gut tissues as targets [36, 73]. Additionally, immunocytochemical localization of a morphine-like material was reported in neural and immune tissues [74–77] as well as in invertebrate tissues [9]. Taken together, these reports suggest that animals appear to have the ability to synthesize opiate alkaloids (Fig. 2 and Table 2).

Recent studies from our group employing well established ex vivo invertebrate nervous tissue preparations and primary cultures of human white blood cells [78, 86, 100] and those by Zenk and coworkers using human tumor-derived cell lines [84, 101] have markedly facilitated the formulation of an evidence-based model of *de novo* formation of endogenous morphine in animal cells with remarkable similarities to the well-characterized enzymatic

pathway described in *Papaver somniferum* (Fig. 2) [102]. Key observations from these studies indicate that L-tyrosine, its monoamine homolog tyramine, and their respective catechol derivatives, L-DOPA and DA serve as substrates for *de novo* morphine production and that pharmacological characterization of tyramine utilization as a morphine precursor indicates one or more catalytic steps mediated by microsomal CYP 2D6 (Fig. 1) [78, 86]. The significance of tyramine as a biosynthetic intermediate is validated by in vitro enzyme kinetic studies demonstrating dopamine formation via CYP 2D6-catalyzed ring hydroxylation of tyramine [103–106] which in turn lends support to the existence of a potentially important tyrosine hydroxylase-independent pool of cytosolic DA that is available for endogenous morphine expression [78, 86, 100]. These data are complemented by metabolic labeling/isotope enrichment studies employing SH-SY5Y neuroblastoma cells [84, 101], indicating asymmetric isotopic labeling of the benzyl and isoquinoline chemical domains of newly formed morphine that is operationally determined by the type of L-tyrosine-derived precursor molecule that is employed: L-tyrosine and L-DOPA are incorporated in both the benzyl and isoquinoline chemical domains of morphine, whereas dopamine and tyramine are only incorporated into the isoquinoline domain.


Taken together, we have formulated a hypothesis, which states that endogenous morphine, used as either a hormone or neurotransmitter down regulates immune, vascular, neural and gut tissues under normal and following trauma situations [30, 36, 47, 53, 77]. In this regard, as noted in these reports, tolerance is viewed as a mechanism to ensure that morphine's continued presence does not permanently limit a needed tissue excitatory state, since excitation may be required to overcome a traumatic event, which morphine would continue to down regulate. Finally, our laboratory has recently demonstrated the putative role of endogenous morphine in HANS stimulation of invertebrate neural tissues via constitutive NO generation [107].

### Future Directions

Ongoing studies from our group support the existence and biological importance of a distinct morphinergic signaling pathway utilizing endogenously synthesized, chemically authentic, L-morphine and its cognate  $\mu_3$  opiate receptor subtype. Recent data support a novel regulatory role of tonically released NO to maintain various classes of stem cells in metabolically viable, undifferentiated states of biological readiness via activation of a  $\mu_3$ -like receptor found on human stem cells [28].

Sequence analysis indicates striking structural similarities between  $\mu_3$  and  $\mu_3$ -like receptors, most notably the lack

**Fig. 2** Detailed schematic indicating relevant enzymes and chemical intermediates in the morphine biosynthetic pathway



**Table 2** Morphine presence in animal tissues

#### Invertebrates

*Mytilus edulis*: immunocytes [9], ganglia [62, 64, 65, 78, 79], hemolymph [9]

*Modiolus deminensis*: ganglia [63]

*Ascaris suum*: ganglia [80], muscle [81], uterus [81]

*Dracunculus medinensis*: whole body extraction [82]

*Schistosoma mansoni*: whole body extraction [82]

*Homarus americanus*: ganglia [83]

#### Vertebrates

Human: cell line SH-SY-5Y [84], heart [85], white blood cells [86], plasma [71, 76], cerebrospinal fluid [56]

Other mammals (rat, mice, cattle): cell line PC-12 [87, 88], neural [89, 90], adrenal [55, 91–93], brain/neural [54, 55, 57–59, 61, 74, 75, 94–99]

of an extended N-terminal glycosylated domain and transmembrane helical domain expressed from exon 1 of the  $\mu$  opioid receptor gene and unique intracellular C-terminal coupling domains [108]. Future structure–function studies will be designed to elucidate the biochemical basis of the selective molecular recognition profiles of these receptors for morphine and morphinan alkaloids. In conclusion, the demonstration of  $\mu_3$ -like receptors on human stem cells provides compelling evidence in support of both the evolutionary primacy and primordial regulatory role of morphine/NO-coupling in embryogenesis. Taken together with the presence of morphine in human immune, vascular and neuronal tissues demonstrates that the endogenous morphine hypothesis extends beyond analgesic functions. Thus, this discovery opens up a new field of investigation with a very old and familiar signaling molecule.

## References

- Lord JAH, Waterfield AA, Hughes J et al (1977) Endogenous opioid peptides: multiple agonists and receptors. *Nature* 267:495–499
- Clark JA, Liu L, Price M et al (1989) Kappa opiate receptor multiplicity: evidence for two U50, 488-sensitive K1 subtypes and a novel K3 subtype. *J Pharmacol Exp Ther* 251:461–468
- Rothman RB, Bykov V, DeCosta BR et al (1990) Interaction of endogenous opioid peptides and other drugs with four kappa binding sites in guinea pig brain. *Peptides* 11:311–331
- Stefano GB, Melchiorri P, Negri L et al (1992) (D-Ala2)-Deltorphin I binding and pharmacological evidence for a special subtype of delta opioid receptor on human and invertebrate immune cells. *Proc Natl Acad Sci USA* 89:9316–9320
- Traynor JR, Elliott J (1993)  $\delta$ -Opioid receptor subtypes and cross-talk with mu-receptors. *Trends Pharmacol Sci* 14:84–86
- Pasternak GW (1986) Multiple mu opiate receptors: biochemical and pharmacological evidence for multiplicity. *Biochem Pharmacol* 35:361–364
- Wolozin BL, Pasternak GW (1981) Classification of multiple morphine and enkephalin binding sites in the central nervous system. *Proc Natl Acad Sci USA* 78:6181–6185
- Lutz RA, Cruciani RA, Costa T et al (1984) A very high affinity opioid binding site in a rat brain: demonstration by computer modeling. *Biochem Biophys Res Commun* 122:265–269
- Stefano GB, Digenis A, Spector S et al (1993) Opiate-like substances in an invertebrate, an opiate receptor on invertebrate and human immunocytes, and a role in immunosuppression. *Proc Natl Acad Sci USA* 90:11099–11103
- Sonetti D, Ottaviani E, Bianchi F et al (1994) Microglia in invertebrate ganglia. *Proc Natl Acad Sci USA* 91:9180–9184
- Stefano GB (1994) Pharmacological and binding evidence for opioid receptors on vertebrate and invertebrate blood cells. In: Scharrer B et al (eds) *Neuropeptides and immunoregulation*. Springer-Verlag, pp 139–151
- Liu Y, Shenouda D, Bilfinger TV et al (1996) Morphine stimulates nitric oxide release from invertebrate microglia. *Brain Res* 722:125–131
- Stefano GB, Scharrer B (1996) The presence of the  $\mu_3$  opiate receptor in invertebrate neural tissues. *Comp Biochem Physiol* 113C:369–373
- Sonetti D, Ottaviani E, Stefano GB (1997) Opiate signaling regulates microglia activities in the invertebrate nervous system. *Gen Pharmacol* 29:39–47
- Dobrenis K, Makman MH, Stefano GB (1995) Occurrence of the opiate alkaloid-selective  $\mu_3$  receptor in mammalian microglia, astrocytes and kupffer cells. *Brain Res* 686:239–248
- Makman MH, Dvorkin B, Stefano GB (1995) Murine macrophage cell lines contain  $\mu_3$ -opiate receptors. *Eur J Pharmacol* 273:R5–R6
- Cruciani RA, Dvorkin B, Klinger HP et al (1994) Presence in neuroblastoma cells of a  $\mu_3$  receptor with selectivity for opiate alkaloids but without affinity for opioid peptides. *Brain Res* 667:229–237
- Stefano GB, Hartman A, Bilfinger TV et al (1995) Presence of the  $\mu_3$  opiate receptor in endothelial cells: coupling to nitric oxide production and vasodilation. *J Biol Chem* 270:30290–30293
- Makman MH, Bilfinger TV, Stefano GB (1995) Human granulocytes contain an opiate receptor mediating inhibition of cytokine-induced activation and chemotaxis. *J Immunol* 154:1323–1330
- Fricchione G, Zhu W, Cadet P et al (2008) Identification of endogenous morphine and a  $\mu_3$ -like opiate alkaloid receptor in human brain tissue taken from a patient with intractable complex partial epilepsy. *Med Sci Monit* 14:CS45–CS49
- Cadet P, Stefano GB (1999) *Mytilus edulis* pedal ganglia express  $\mu$  opiate receptor transcripts exhibiting high sequence identity with human neuronal  $\mu_1$ . *Mol Brain Res* 74:242–246
- Cadet P, Zhu W, Mantione K et al (2002) Cold stress alters *Mytilus edulis* pedal ganglia expression of  $\mu$  opiate receptor transcripts determined by real-time RT-PCR and morphine levels. *Brain Res Mol Brain Res* 99:26–33
- Cadet P, Mantione KJ, Stefano GB (2003) Molecular identification and functional expression of  $\mu_3$ , a novel alternatively spliced variant of the human mu opiate receptor gene. *J Immunol* 170:5118–5123
- Cadet P, Bilfinger TV, Fimiani C et al (2000) Human vascular and cardiac endothelia express mu opiate receptor transcripts. *Endothelium* 7:185–191
- Cadet P, Weeks BS, Bilfinger TV et al (2001) HIV gp120 and morphine alter  $\mu$  opiate receptor expression in human vascular endothelium. *Int J Mol Med* 8:165–169
- Cadet P, Mantione KJ, Bilfinger TV et al (2004) Differential expression of the human mu opiate receptor from different primary vascular endothelial cells. *Med Sci Monit* 10:BR351–BR355
- Cadet P, Mantione K, Bilfinger TV et al (2001) Real-time RT-PCR measurement of the modulation of mu opiate receptor expression by nitric oxide in human mononuclear cells. *Med Sci Monit* 7:1123–1128
- Cadet P, Mantione KJ, Zhu W et al (2007) A functionally coupled  $\mu_3$ -like opiate receptor/nitric oxide regulatory pathway in human multi-lineage progenitor cells. *J Immunol* 179:5839–5844
- Rialas C, Bilfinger TV, Salzet M et al (1998) Endomorphin 1 and 2 do not interact with the  $\mu_3$  opiate receptor subtype. *Acta Pharmacol Sin* 19:403–407
- Stefano GB, Scharrer B, Smith EM et al (1996) Opioid and opiate immunoregulatory processes. *Crit Rev Immunol* 16:109–144
- Ferreira SH, Duarte ID, Lorenzetti BB (1991) The molecular mechanism of action of peripheral morphine analgesia: stimulation of the cGMP system via nitric oxide release. *Eur J Pharmacol* 201:121
- Przewlocki R, Machelska H, Przewlocka B (1993) Inhibition of nitric oxide synthase enhances morphine antinociception in the rat spinal cord. *Life Sci* 53:PL1–PL5



33. Majeed NH, Przewlocka B, Machelska H et al (1994) Inhibition of nitric oxide synthase attenuates the development of morphine tolerance and dependence in mice. *Neuropharmacology* 33:189–192
34. Fecho K, Maslonek KA, Coussons-Read ME et al (1994) Macrophage-derived nitric oxide is involved in the depressed concanavalin A responsiveness of splenic lymphocytes from rats administered morphine in vivo. *J Immunol* 152:5845–5851
35. Gyires K (1994) The role of endogenous nitric oxide in the gastroprotective action of morphine. *Eur J Pharmacol* 255:33–37
36. Stefano GB, Zhu W, Cadet P et al (2004) Morphine enhances nitric oxide release in the mammalian gastrointestinal tract via the  $\mu 3$  opiate receptor subtype: a hormonal role for endogenous morphine. *J Physiol Pharmacol* 55:279–288
37. Bilfinger TV, Hartman A, Liu Y et al (1997) Cryopreserved veins used for myocardial revascularization: a 5 year experience and a possible mechanism for their increased failure. *Ann Thorac Surg* 63:1063–1069
38. Sawada M, Ichinose M, Stefano GB (1997) Nitric oxide inhibits the dopamine-induced K<sup>+</sup> current via guanylate cyclase in Aplysia neurons. *J Neurosci Res* 50:450–456
39. Stefano GB, Salzet B, Rialas CM et al (1997) Morphine and anandamide stimulated nitric oxide production inhibits presynaptic dopamine release. *Brain Res* 763:63–68
40. Bilfinger TV, Fimiani C, Stefano GB (1998) Morphine's immunoregulatory actions are not shared by fentanyl. *Int J Cardiol* 64:61–66
41. Prevot V, Rialas C, Croix D et al (1998) Morphine and anandamide coupling to nitric oxide stimulated GnRH and CRF release from rat median eminence: neurovascular regulation. *Brain Res* 790:236–244
42. Stefano GB (1998) Autoimmunovascular regulation: morphine and anandamide stimulated nitric oxide release. *J Neuroimmunol* 83:70–76
43. Stefano GB, Salzet M, Bilfinger TV (1998) Long-term exposure of human blood vessels to HIV gp120, morphine and anandamide increases endothelial adhesion of monocytes: uncoupling of Nitric Oxide. *J Cardiovasc Pharmacol* 31:862–868
44. Stefano GB, Salzet M, Magazine HI et al (1998) Antagonist of LPS and IFN- $\gamma$  induction of iNOS in human saphenous vein endothelium by morphine and anandamide by nitric oxide inhibition of adenylate cyclase. *J Cardiovasc Pharmacol* 31:813–820
45. Fimiani C, Mattocks DW, Cavani F et al (1999) Morphine and anandamide stimulate intracellular calcium transients in human arterial endothelial endothelial cells: coupling to nitric oxide release. *Cell Signal* 11:189–193
46. Pryor SC, Zhu W, Cadet P et al (2005) Endogenous morphine: opening new doors for the treatment of pain and addiction. *Expert Opin Biol Ther* 5:893–906
47. Stefano GB, Scharrer B (1994) Endogenous morphine and related opiates, a new class of chemical messengers. *Adv Neuroimmunol* 4:57–68
48. Tubaro E, Borelli G, Croce C et al (1983) Effect of morphine on resistance to infection. *J Infect Dis* 148:656–666
49. Weber RJ, Band LC, DeCosta B et al (1991) Neural control of immune function: opioids, opioid receptors and immunosuppression. *NIDA Res Monogr* 105:96–102
50. Perez-Castrillon JP, Perez-Arellano JP, Garcia-Palomo JD et al (1992) Opioids depress in vitro human monocyte chemotaxis. *Immunopharmacology* 23:57–61
51. Stefano GB, Bilfinger TV (1993) Human neutrophil and macrophage chemokinesis induced by cardiopulmonary bypass: loss of DAME and IL-1 chemotaxis. *J Neuroimmunol* 47:189–198
52. Stefano GB (1999) The  $\mu 3$  opiate receptor subtype. *Pain Forum* 8:206–209
53. Stefano GB, Goumon Y, Casares F et al (2000) Endogenous morphine. *Trends Neurosci* 9:436–442
54. Gintzler AR, Levy A, Spector S (1976) Antibodies as a means of isolating and characterizing biologically active substances: presence of a non-peptide morphine-like compound in the central nervous system. *Proc Natl Acad Sci USA* 73:2132–2136
55. Goldstein A, Barrett RW, James IF et al (1985) Morphine and other opiates from beef brain and adrenal. *Proc Natl Acad Sci USA* 82:5203–5207
56. Cardinale GJ, Donnerer J, Finck AD et al (1987) Morphine and codeine are endogenous components of human cerebrospinal fluid. *Life Sci* 40:301–306
57. Donnerer J, Cardinale G, Coffey J et al (1987) Chemical characterization and regulation of endogenous morphine and codeine in the rat. *J Pharmacol Exp Ther* 242:583–587
58. Donnerer J, Oka K, Brossi A et al (1986) Presence and formation of codeine and morphine in the rat. *Proc Natl Acad Sci USA* 83:4566–4567
59. Oka K, Kantrowitz JD, Spector S (1985) Isolation of morphine from toad skin. *Proc Natl Acad Sci USA* 82:1852–1854
60. Epple A, Nibbio B, Spector S et al (1994) Endogenous codeine: autocrine regulator of catecholamine release from chromaffin cells. *Life Sci* 54:695–702
61. Lee CS, Spector S (1991) Changes of endogenous morphine and codeine contents in the fasting rat. *J Pharmacol Exp Ther* 257:647–650
62. Zhu W, Ma Y, Cadet P et al (2003) Presence of reticuline in rat brain: a pathway for morphine biosynthesis. *Mol Brain Res* 117:83–90
63. Goumon Y, Casares F, Zhu W et al (2001) The presence of morphine in ganglionic tissues of *Modiolus demissus*: a highly sensitive method of quantitation for morphine and its derivatives. *Mol Brain Res* 86:184–188
64. Zhu W, Ma Y, Stefano GB (2002) Presence of isoquinoline alkaloids in molluscan ganglia. *Neuroendocrinol Lett* 23:329–334
65. Zhu W, Baggerman G, Goumon Y et al (2001) Presence of morphine and morphine-6-glucuronide in the marine mollusk *Mytilus edulis* ganglia determined by GC/MS and Q-TOF-MS. Starvation increases opiate alkaloid levels. *Brain Res Mol Brain Res* 88:155–160
66. Stefano GB, Bianchi E, Guarna M et al (2007) Nicotine, alcohol and cocaine coupling to reward processes via endogenous morphine signaling: the dopamine–morphine hypothesis. *Med Sci Monit* 13:RA91–RA102
67. Stefano GB, Kream R (2008) Endogenous opiates, opioids, and immune function: evolutionary brokerage of defensive behaviors. *Semin Cancer Biol* 18:190–198
68. Stefano GB, Kream RM, Mantione KJ et al (2008) Endogenous morphine/nitric oxide-coupled regulation of cellular physiology and gene expression: implications for cancer biology. *Semin Cancer Biol* 18:199–210
69. Esch T, Kim JW, Stefano GB (2006) Neurobiological implications of eating healthy. *Neuro Endocrinol Lett* 27:21–33
70. Stefano GB, Fricchione GL, Esch T (2006) Relaxation: Molecular and physiological significance. *Med Sci Monit* 12:HY21–HY31
71. Brix-Christensen V, Tonnesen E, Sanchez RG et al (1997) Endogenous morphine levels increase following cardiac surgery as part of the antiinflammatory response? *Int J Cardiol* 62:191–197
72. Cadet P, Zhu W, Mantione K et al (2003) Cyclic exercise induces anti-inflammatory signal molecule increases in the plasma of Parkinson's patients. *Int J Mol Med* 12:485–492
73. Stefano GB, Zhu W, Cadet P et al (2002) A hormonal role for endogenous opiate alkaloids: vascular tissues. *Neuroendocrinol Lett* 23:21–26

74. Bianchi E, Alessandrini C, Guarna M et al (1993) Endogenous codeine and morphine are stored in specific brain neurons. *Brain Res* 627:210–215
75. Bianchi E, Guarna M, Tagliamonte A (1994) Immunocytochemical localization of endogenous codeine and morphine. *Adv Neuroimmunol* 4:83–92
76. Liu Y, Bilfinger TV, Stefano GB (1996) A rapid and sensitive quantitation method of endogenous morphine in human plasma. *Life Sci* 60:237–243
77. Esch T, Guarna M, Bianchi E et al (2004) Commonalities in the central nervous system's involvement with complementary medical therapies: limbic morphinergic processes. *Med Sci Monit* 10:MS6–MS17
78. Zhu W, Mantione KJ, Shen L et al (2005) Tyrosine and tyramine increase endogenous ganglionic morphine and dopamine levels in vitro and in vivo: CYP2D6 and tyrosine hydroxylase modulation demonstrates a dopamine coupling. *Med Sci Monit* 11:BR397–BR404
79. Zhu W, Stefano GB (2004) Reticuline exposure to invertebrate ganglia increases endogenous morphine levels. *Neuro Endocrinol Lett* 25:323–330
80. Goumon Y, Casares F, Pryor S et al (2000) *Ascaris suum*, an internal parasite, produces morphine. *J Immunol* 165:339–343
81. Zhu W, Pryor SC, Putnam J et al (2004) Opiate alkaloids and nitric oxide production in the nematode *Ascaris suum*. *J Parasitol* 90:15–22
82. Zhu W, Baggerman G, Secor WE et al (2002) *Dracunculus medinensis* and *Schistosoma mansoni* contain opiate alkaloids. *Ann Trop Med Parasitol* 96:309–316
83. Casares FM, McElroy A, Mantione KJ et al (2005) The American lobster, *Homarus americanus*, contains morphine that is coupled to nitric oxide release in its nervous and immune tissues: evidence for neurotransmitter and hormonal signaling. *Neuroendocrinol Lett* 26:89–97
84. Poeaknapo C, Schmidt J, Brandsch M et al (2004) Endogenous formation of morphine in human cells. *Proc Natl Acad Sci USA* 101:14091–14096
85. Zhu W, Bilfinger TV, Baggerman G et al (2001) Presence of endogenous morphine and morphine 6 glucuronide in human heart tissue. *Int J Mol Med* 7:419–422
86. Zhu W, Cadet P, Baggerman G et al (2005) Human white blood cells synthesize morphine: CYP2D6 modulation. *J Immunol* 175:7357–7362
87. Goumon Y, Weeks BS, Cadet P et al (2000) Identification of morphine in the adrenal medullary chromaffin PC-12 cell line. *Mol Brain Res* 81:177–180
88. Zhu W, Baggerman G, Goumon Y et al (2001) Identification of morphine and morphine-6-glucuronide in the adrenal medullary chromaffin PC-12 cell line by nano electrospray ionization double quadrupole orthogonal acceleration time of flight mass spectrometry. *Eur J Mass Spectrosc* 7:25–28
89. Goumon Y, Bouret S, Casares F et al (2000) Lipopolysaccharide increases endogenous morphine levels in rat brain. *Neurosci Lett* 293:135–138
90. Zhu W, Ma Y, Bell A et al (2004) Presence of morphine in rat amygdala: evidence for the mu3 opiate receptor subtype via nitric oxide release in limbic structures. *Med Sci Monit* 10:BR433–BR439
91. Goumon Y, Stefano GB (2000) Identification of morphine in the rat adrenal gland. *Mol Brain Res* 77:267–269
92. Goumon Y, Strub JM, Stefano GB et al (2005) Characterization of a morphine-like molecule in secretory granules of chromaffin cells. *Med Sci Monit* 11:MS31–MS34
93. Goumon Y, Muller A, Glattard E et al (2006) Identification of morphine-6-glucuronide in chromaffin cell secretory granules. *J Biol Chem* 281:8082–8089
94. Guarna M, Neri C, Petrioli F et al (1998) Potassium-induced release of endogenous morphine from rat brain slices. *J Neurochem* 70:147–152
95. Neri C, Guarna M, Bianchi E et al (2004) Endogenous morphine and codeine in the brain of non-human primate. *Med Sci Monit* 10:MS1–MS5
96. Gintzler AR, Gershon MD, Spector S (1978) A nonpeptide morphine-like compound: immunocytochemical localization in the mouse brain. *Science* 199:447–448
97. Horak P, Haberman F, Spector S (1993) Endogenous morphine and codeine in mice—effect on muramyl dipeptide. *Life Sci* 52:255–260
98. Kodaira H, Spector S (1988) Transformation of thebaine to oripavine, codeine, and morphine by rat liver, kidney, and brain microsomes. *Proc Natl Acad Sci USA* 85:1267–1271
99. Weitz CJ, Lowney LI, Faull KF et al (1986) Morphine and codeine from mammalian brain. *Proc Natl Acad Sci USA* 83:9784–9788
100. Zhu W, Mantione KJ, Shen L et al (2005) In vivo and in vitro L-DOPA exposure increases ganglionic morphine levels. *Med Sci Monit* 11:MS1–MS5
101. Boettcher C, Fellermeier M, Boettcher C et al (2005) How human neuroblastoma cells make morphine. *Proc Natl Acad Sci USA* 102:8495–8500
102. Kream RM, Stefano GB (2006) De novo biosynthesis of morphine in animal cells: An evidence-based model. *Medical Science Monit* 12:RA207–RA219
103. Hiroi T, Imaoka S, Funae Y (1998) Dopamine formation from tyramine by CYP2D6. *Biochem Biophys Res Commun* 249:838–843
104. Miller GP, Hanna IH, Nishimura Y et al (2001) Oxidation of phenethylamine derivatives by cytochrome P450 2D6: the issue of substrate protonation in binding and catalysis. *Biochemistry* 40:14215–14223
105. Guengerich FP, Miller GP, Hanna IH et al (2002) Oxidation of methoxyphenethylamines by cytochrome P450 2D6. Analysis of rate-limiting steps. *J Biol Chem* 277:33711–33719
106. Niwa T, Hiroi T, Tsuzuki D et al (2004) Effect of genetic polymorphism on the metabolism of endogenous neuroactive substances, progesterone and p-tyramine, catalyzed by CYP2D6. *Brain Res Mol Brain Res* 129:117–123
107. Cheng J, Zhang C, Han JS et al (2007) TENS stimulates constitutive nitric oxide release via opiate signaling in invertebrate neural tissues. *Med Sci Monit* 13:BR163–BR167
108. Kream RM, Sheehan M, Cadet P et al (2007) Persistence of evolutionary memory: primordial six-transmembrane helical domain mu opiate receptors selectively linked to endogenous morphine signaling. *Med Sci Monit* 13:SC5–SC6