

The neuroprotective steroid progesterone promotes mitochondrial uncoupling, reduces cytosolic calcium and augments stress resistance in yeast cells

Slaven Stekovic^{1,*}, Christoph Ruckenstein^{1,*,*}, Philipp Royer¹, Christof Winkler-Hermaden¹, Didac Carmona-Gutierrez¹, Kai-Uwe Fröhlich¹, Guido Kroemer³⁻⁸, and Frank Madeo^{1,2,+}

¹Institute of Molecular Biosciences, NAWI Graz, University of Graz, 8010 Graz, Austria.

²BioTechMed Graz, Austria.

³Equipe 11 labellisée par la Ligue contre le Cancer, Centre de Recherche des Cordeliers, Paris, France.

⁴INSERM, U1138, Paris, France.

⁵Université Paris Descartes, Sorbonne Paris Cité, Paris, France.

⁶Cell Biology & Metabolomics Platforms, Gustave Roussy Comprehensive Cancer Center, Villejuif, France.

⁷Pôle de Biologie, Hôpital Européen Georges Pompidou, AP - HP, Paris, France.

⁸Karolinska Institute, Department of Women's and Children's Health, Karolinska University Hospital, 17176 Stockholm, Sweden,

* These authors contributed equally.

+ Corresponding Author:

Frank Madeo, Institute of Molecular Biosciences, University of Graz, Humboldtstrasse 50; 8010 Graz, Austria; E-mail: frank.madeo@uni-graz.at

Christoph Ruckenstein, Institute of Molecular Biosciences, University of Graz, Humboldtstrasse 50; 8010 Graz, Austria; E-mail: ru.ruckenstein@uni-graz.at

ABSTRACT The steroid hormone progesterone is not only a crucial sex hormone, but also serves as a neurosteroid, thus playing an important role in brain function. Epidemiological data suggest that progesterone improves the recovery of patients after traumatic brain injury. Brain injuries are often connected to elevated calcium spikes, reactive oxygen species (ROS) and programmed cell death affecting neurons. Here, we establish a yeast model to study progesterone-mediated cytoprotection. External supply of progesterone protected yeast cells from apoptosis-inducing stress stimuli and resulted in elevated mitochondrial oxygen uptake accompanied by a drop in ROS generation and ATP levels during chronological aging. In addition, cellular Ca^{2+} concentrations were reduced upon progesterone treatment, and this effect occurred independently of known Ca^{2+} transporters and mitochondrial respiration. All effects were also independent of Dap1, the yeast orthologue of the progesterone receptor. Altogether, our observations provide new insights into the cytoprotective effects of progesterone.

doi: 10.15698/mic2017.06.577

Received originally: 10.03.2017;

In revised form: 21.05.2017,

Accepted 22.05.2017,

Published 31.05.2017.

Keywords: TBI; traumatic brain injury, cell protection, cell stress, cell death, neuroprotection, progesterone, mitochondrial uncoupling.

Abbreviations:

DHE – dihydroethidium,
PCD – programmed cell death,
ROS – reactive oxygen species,
TBI – traumatic brain injury.

INTRODUCTION

Progesterone is a sterol-derived hormone that is crucial for female reproductive capacity and plays major regulatory roles in the monthly menstrual cycle and upon conception as well as during pregnancy and embryogenesis. In addition, it also serves as a neurosteroid, thus playing an important role in brain function in both sexes [1]. For instance, progesterone inhibits the neuronal nicotinic acetylcholine receptor and stimulates the synthesis of myelin proteins [1]. Of note, progesterone has been linked

to the gender-specific risk and outcome of brain injuries that is more favorable for females [2]. Interestingly, preclinical data strongly suggest that (high doses of) progesterone may positively affect recovery from traumatic brain injury (TBI) in model organisms [3–7], if administered before or shortly after TBI. Two clinical studies could confirm a neuroprotective effect of progesterone when administered shortly after TBI [8,9], while some more recent clinical data seem to disprove this hypothesis [10–12]. Therefore, it remains an open question




FIGURE 1: Progesterone treatment increases resistance of wildtype yeast to external stressors. ROS accumulation in yeast cells treated with progesterone (10 μ g/ml) or left untreated as shown by the DHE to ethidium turnover rate upon hydrogen peroxide (A) or acetate (B) challenge during logarithmic phase. All data represent mean values ($n = 3 \pm \text{SEM}$). Statistical analysis was conducted using non-paired Student's t-test. * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$; n.s. = non-significant, Prog = progesterone, ctrl = control.

if progesterone affects the recovery and survival after TBI in humans and to which extent it promotes cellular restoration.

In order to investigate the cytoprotective potential of progesterone, we took advantage of *Saccharomyces cerevisiae*, knowing that this organism has repeatedly been shown to be suitable for mechanistic studies of programmed cell death (PCD) [13–19]. Yeast is especially useful as a model to study neuroprotection at the cellular level [20–27]. Here, we describe the positive impact of progesterone on several parameters of cellular physiology. Importantly, our results also suggest a possible receptor-independent mechanism for these effects, since deletion of *DAP1* – a heme-binding protein related to the mammalian membrane progesterone receptor – did not alter susceptibility towards progesterone treatment. Altogether, we reveal that progesterone exerts potent cytoprotective effects in yeast.

RESULTS

Progesterone increases stress tolerance

Traumatic brain injury is connected to elevated PCD and ROS accumulation in the brain tissue [28,29]. Therefore, we tested if progesterone would render yeast cells less susceptible towards different stressors that are connected to an increase in ROS production. Upon addition of progesterone, wild type yeast cultures treated with H₂O₂ or acetate, which are both well-known PCD inducers in yeast [14,30–34], showed reduced ROS accumulation as measured by the ROS-driven conversion of dihydroethidium (DHE) to fluorescent ethidium (Figure 1A and B). Furthermore, under physiological culture conditions, in the absence of PCD inducers, progesterone significantly reduced ROS levels as compared to the

untreated control (Figure 2A). Altogether, progesterone dampens ROS production in yeast, both in normal culture conditions and in the presence of external stress factors.

Progesterone impacts mitochondria by acting as a mild respiration-uncoupler

To further explore the mechanisms underlying progesterone cytoprotection, we next examined the physiology of mitochondria, since these organelles constitute one of the main sources of ROS [35–38]. Interestingly, while O₂ consumption was significantly enhanced during progesterone treatment, ATP levels were reduced (Figure 2B and C). Altogether, this indicates an uncoupling phenotype with diminished oxidative phosphorylation. Accordingly, we observed reduced growth of wild type yeast upon progesterone treatment on a non-fermentable carbon source (glycerol), while no changes were detected on a fermentable carbon source (glucose) (Figure 2D and E). Importantly, this effect was also observed in a mutant strain lacking the heme-binding protein Dap1, which is the sole yeast orthologue of the human progesterone receptor (Figure 2D and E) [39]. Furthermore, we could demonstrate that stress protection by progesterone is respiration-dependent, since progesterone treatment did not confer stress resistance in respiration-deficient ρ^0 cells (Figure 2F). Altogether, it appears that progesterone impacts yeast mitochondrial respiration in a receptor-independent fashion.

Progesterone administration diminishes cytosolic Ca²⁺ concentrations both under physiological as well as under high calcium conditions

Next we investigated progesterone effects on Ca²⁺ homeostasis, knowing that mitochondria are one of the




FIGURE 2: Progesterone impacts energy metabolism and reduces oxygen stress accumulation in wildtype yeast. Wildtype yeast were treated with 10 μg/ml progesterone and assayed for (A) ROS accumulation via DHE to ethidium turnover, (B) oxygen consumption via respirometry, and (C) ATP production. Growth curves of wildtype as well as $\Delta dap1$ strains, with or without progesterone treatment, on glycerol (respiratory carbon source) (D) and glucose (fermentative carbon source) media (E). ROS accumulation in ρ^0 yeast cells +/- progesterone (10 μg/ml) treated or untreated with H₂O₂ or acetate during logarithmic phase (F). All data represent mean values ($n = 3-5 \pm \text{SEM}$). Statistical analysis was conducted using non-paired Student t-test (A-C, F) or using a two-way repeated measurement ANOVA and multiple comparison post-hoc Tukey's test (D, E). * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$; n.s. = non-significant. ROS = reactive oxygen species, rFU = relative fluorescence units, Prog = progesterone, ctrl = control.

organelles

responsible for buffering cytosolic Ca^{2+} under normal conditions [40]. Importantly, TBI, stroke, and even some forms of dementia cause Ca^{2+} accumulation in the cytosol of neurons followed by cell death and neurodegeneration [41]. Thus, we examined the capacity of yeast cells to process Ca^{2+} uptake under the influence of progesterone. Specifically, wild type yeast cell cultures were challenged with 150 mM CaCl_2 and transient concentrations of cytoplasmic Ca^{2+} levels ($[\text{Ca}^{2+}]_{\text{cyt}}$) / responses were monitored. Progesterone caused a significantly reduced Ca^{2+} uptake capacity alongside with a faster reduction of cytoplasmic Ca^{2+} levels (Figure 3A and B). Of note, basal Ca^{2+} levels before and after the Ca^{2+} pulse were already lowered when cells were treated with progesterone (Figure 3B). However, mitochondrial respiration was not involved in this phenotype, since progesterone treatment continued to affect basic cytosolic Ca^{2+} levels in respiration-deficient rho^0 cells (Figure 3C and D).

To further investigate the observed phenotypes, we tested single-gene deletion mutants of all currently known Ca^{2+} channels/transporters in yeast, including the cytoplasmic membrane transporters Cch1 and Mid 1, the organelle transporters Vcx1, Pmr1, Cod1, Yvc1, and Pmc1 as well as Emc7, an ER protein associated to Ca^{2+} homeostasis. Although Ca^{2+} uptake and clearance was influenced by some of these gene deletion, all mutants continued to exhibit significantly reduced Ca^{2+} uptake when treated with progesterone (compare Supplemental Figure 1A-G to H). Thus, the effects observed in wild type cells could not be reversed by single gene deletions in any of these transporters. Similarly, the effects of progesterone treatment on Ca^{2+} homeostasis/uptake were independent of the mammalian membrane progesterone receptor homolog Dap1 (Figure 3E and F). Taken together, progesterone seems to influence Ca^{2+} homeostasis/uptake in a general manner, independently from known Ca^{2+} transporters and respiration capacity.

DISCUSSION

Here, we establish *S. cerevisiae* as a model to investigate cytoprotection by progesterone. We observed that progesterone increased stress tolerance of yeast to the well-known PCD inducers H_2O_2 and acetate [14,30–34] as well as under physiological (control) conditions. Interestingly, progesterone treatment led to a mild uncoupling phenotype with higher O_2 consumption (+50%) but lower ATP levels (-50%), arguing for a mitochondrial uncoupling effect. Indeed, growth on the non-fermentable carbon source glycerol was diminished in the presence of progesterone. Notably, mild uncoupling induced by chemical substances (such as dinitrophenol), caloric restriction or ectopic expression of mammalian uncoupling proteins in yeast - *S. cerevisiae* does not possess any known uncoupling proteins [42] - is known to increase lifespan [43–45]. Similarly, in mammalian aging cells, changes in mitochondrial energy metabolism caused by mitochondrial uncoupling seem to improve cellular fitness [46]. Progesterone treatment of human cells has been

demonstrated to strongly increase the levels of mRNAs coding for uncoupling proteins [47]. Increased O_2 consumption with decreased ^{32}P uptake (as a parameter for ATP production) has been reported for isolated rat mitochondria treated with progesterone [48]. Collectively, our data combined with those reported in the literature highlight the possibility to investigate progesterone-mediated effects in the yeast model. The uncoupling aspect of progesterone, in fact, could represent one of the mechanisms of neuroprotection conferred by this steroid. In fact, the stress tolerance of a respiration-deficient rho^0 strain was not influenced by progesterone treatment.

Progesterone had major effects on Ca^{2+} homeostasis and, in particular, on Ca^{2+} susceptibility/uptake. However, we could not identify any single Ca^{2+} channel in yeast that would influence these effects. However, we cannot exclude that yet unidentified Ca^{2+} channels or a combinations of known Ca^{2+} channels mediate these effects [49]. Another possible mode of action of progesterone on Ca^{2+} homeostasis could reside in its direct interaction with biological membranes. Since the chemical structure of progesterone shows four-ring as well as hydrophobic backbone and polar groups at both ends of the molecule, it could directly interact with cellular and mitochondrial membranes [50] and possibly influence their permeability towards inorganic cations (e.g. Ca^{2+} , H^+). This mode of action could connect our observations of mitochondrial uncoupling and modulation of Ca^{2+} homeostasis. Of note, a progesterone-treated rho^0 strain still showed Ca^{2+} effects but no enhanced stress tolerance, suggesting that altered Ca^{2+} homeostasis may lie upstream of mitochondrial uncoupling. However, these mechanistic hypotheses remain to be empirically tested.

Certainly, the putative relevance of the herein described progesterone effects for TBI pathology remains to be explored. In some mammalian cell types, progesterone leads to a significant increase of intracellular Ca^{2+} [51,52], partly by activating protein kinase C [53] and depleting endocannabinoids by activating α/β hydrolase domain-containing protein 2 (ABHD2) [54]. However, in other cell types, progesterone withdrawal leads to an increased level of cytosolic Ca^{2+} [55]. While progesterone was not able to reduce estrogen-induced Ca^{2+} uptake in the rabbit myometrial smooth muscle cells, it increased the accumulation of Ca^{2+} in mitochondria [55]. This suggests that progesterone withdrawal reduces both myometrial cytosolic Ca^{2+} levels as well as the capacity of these cells to accumulate Ca^{2+} in different cellular compartments. Similar effects were reported for other types of smooth muscles [56,57] and are believed to be caused by regulation of the inward current through L-type Ca^{2+} channels [56,58]. In neurons, the influence on Ca^{2+} signaling and the following inhibition of excitotoxic neuron death seem to be the neuroprotective mechanism induced by acute administration of progesterone after various neuronal injuries [59–61]. Indeed, progesterone might mediate broad neuroprotective effects, not only in the context of TBI but also in other pathologies [62,63].

The role of progesterone in the pathological develop-




FIGURE 3: Cellular Ca^{2+} homeostasis is modulated by progesterone treatment in yeast. Cells were treated with progesterone (10 $\mu\text{g}/\text{ml}$) and challenged with high doses of Ca^{2+} (150 mM). Intake of Ca^{2+} as well as Ca^{2+} -clearance in the cytosol to its basal level were measured in wild type (A,B), a *DAP1*-deletion strain (C,D), as well as in a ρ^0 strain, incapable of mitochondrial respiration (E,F). Data are shown as mean values of at least three replicates including the standard error of the mean. Statistical analysis was conducted using non-paired Student's t-test. * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$; n.s. = non-significant. E_{max} = global maximum of the respective ethanol-treated control; rLU = relative luminiscence units, Prog = progesterone, ctrl = control.

ment of TBI has been well described in recent years. It has been shown that progesterone improves the function of the blood-brain-barrier after TBI [64]. Progesterone also increases the level of circulating endothelial cells, which in turn improves neovascularization and vascular remodeling in the brain [65]. Furthermore, progesterone treatment reduces neuroinflammation and oxidative stress [66] as it improves remyelination and functional recovery [63].

Interestingly, the intracellular effects exerted by progesterone in our model - reduced intracellular Ca^{2+} levels, uncoupled mitochondria and ROS reduction – were not lost when the sole possible yeast orthologue of the human progesterone receptor was removed from the system. This suggests that progesterone mediates its broad cytoprotective effects through other proteins than steroid receptors or perhaps with cellular membrane lipids. We surmise that yeast constitutes an ideal platform for exploring these effects in further detail.

MATERIALS AND METHODS

Growth conditions

S. cerevisiae strains (Table 1) were inoculated to 5×10^5 (for growth curve OD_{600} of 0.05) cells in SC medium containing 0.17% yeast nitrogen base (BD Diagnostics; without ammonium sulfate and amino acids), 0.5% $(\text{NH}_4)_2\text{SO}_4$, 30 mg/L of all amino acids (except 80 mg/L histidine and 200 mg/L leucine), 30 mg/L adenine, and 320 mg/L uracil with 2% glucose (SCD) or alternatively with 3% glycerol (SCGly), w/o treatment with progesterone (10 $\mu\text{g}/\text{ml}$; Sigma Aldrich, Catalogue Nr. P0130). Controls were treated with respective solvent (EtOH). Where

indicated, stress (acetate or H_2O_2) was inflicted as described previously in mid-log phase (~ 6 h of growth, culture density $2-4 \times 10^6$ cells/ml). Due to the inherent reduced respiration-rate of BY4741 strains, TB50a strains were used for respiration-related experiments. *DAP1* deletion was carried out by classical homologous recombination [67,68].

Growth curve

Cells from ONC in SCD media were inoculated to an OD_{600} of 0.05 in SCD media and SCGly media with or without 10 $\mu\text{g}/\text{ml}$ progesterone addition. Untreated cultures were supplemented with 0.1% EtOH for solvent control. To obtain growth curves, 300 μl of respective cultures per well were transferred into Honeycomb® plates, and measured with Bioscreen C MBR system (Oy Growth Curves Ab Ltd.) for a period of 48 hours at 28°C, using continuous shaking and OD_{600} measurements every 30 minutes.

Oxygen consumption measurement

Oxygen consumption was measured using a FireSting oxygen electrode (Pyro-Science) under constant stirring at a temperature of $28.0 \pm 0.2^\circ\text{C}$ in sealed 2 ml bottles. The corresponding cell counts were measured using a CASY Cell Counter, whereas percentage of living cells in the sample were established by flow cytometry with propidium iodide (PI: 100 ng/ml) stained samples. The slope of the oxygen concentration as the function of time in its linear region was calculated and normalized to the number of living cells in the sample.

ROS accumulation (DHE) assay

Oxidation of non-fluorescent di-hydroethidium (DHE) to fluo-

Table 1. Strains used in this study.

Strain	Genotype	Reference
TB50a wildtype	<i>MATa; leu2-3,112 ura3-52 trp1 his3 rme1 HMLa</i>	[69]
TB50a $\Delta dap1$	<i>MATa; leu2-3,112 ura3-52 trp1 his3 rme1 HMLa dap1::kanMX</i>	This study
BY4741 wildtype	<i>MATa his3Δ1 leu2Δ0 met15Δ0 ura3Δ0</i>	Euroscarf
BY4741 $\Delta dap1$	<i>MATa his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 dap1::kanMX</i>	Euroscarf
BY4741 $\Delta cch1$	<i>MATa his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 cch1::kanMX</i>	Euroscarf
BY4741 $\Delta mid1$	<i>MATa his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 mid1::kanMX</i>	Euroscarf
BY4741 $\Delta vcx1$	<i>MATa his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 vcx1::kanMX</i>	Euroscarf
BY4741 $\Delta pmr1$	<i>MATa his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 pmr1::kanMX</i>	Euroscarf
BY4741 $\Delta cod1$	<i>MATa his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 cod1::kanMX</i>	Euroscarf
BY4741 $\Delta yvc1$	<i>MATa his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 yvc1::kanMX</i>	Euroscarf
BY4741 $\Delta pmc1$	<i>MATa his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 pmc1::kanMX</i>	Euroscarf
BY4741 $\Delta emc7$	<i>MATa his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 emc7::kanMX</i>	Euroscarf

rescent ethidium was used to measure ROS accumulation in yeast cells [38]. Approximately 5×10^6 cells from each sample were collected, washed and incubated with DHE solution (2.5 $\mu\text{g/ml}$ in PBS) for 10 min in the dark. After washing samples were re-suspended in PBS buffer and measured using flow cytometry. The relative mean fluorescence measured for the cell population was used for analysis [70].

Boiling ethanol extraction of ATP and ATP measurement

ATP extraction was done with flash-frozen cells by adding 0.5 ml preheated (90°C) BES buffer and incubation at 90°C for 3 minutes. After centrifugation, supernatants were stored at -80°C until the measurement. ATP levels were determined by using the ATP detection kit from Invitrogen in a Luminoskan (Thermo Scientific).

Cytosolic Ca^{2+} measurements

$[\text{Ca}^{2+}]_{\text{cyt}}$ were measured using yeast strains carrying the vector pYX212 encoding the bioluminescent protein aequorin under the control of a TPI promoter. For analysis of the cellular response to high doses of external Ca^{2+} , an equivalent of 6×10^6 cells was harvested, resuspended in 200 μl SCD containing 4 μM coelenterazine and incubated for 1 h in the dark. After washing cells were measured in a Luminoskan for 10 s and then challenged with high dose of Ca^{2+} (pump injection of 150 mM Ca^{2+}). Kinetics were recorded over 120 s. The luminescence signal was normalized to the OD_{600} of each well and reported in relative luminescence units, normalized to the global maximum value of the ethanol treated control of the respective run for better comparability.

ACKNOWLEDGEMENTS

We thank Silvia Dichtinger for technical assistance. FM is grateful to the Austrian Science Fund FWF (Austria) for grants P23490-B20, P29262, P24381, P29203 P27893, I1000 and 'SFB Lipotox' (F3012), as well as to BMWFW and the Karl-Franzens University for grant 'Unkonventionelle Forschung' and grant DKplus Metabolic and Cardiovascular Diseases (W1226). We acknowledge support from NAWI Graz and the BioTechMed-Graz flagship project "EPIAge". GK is supported by the Ligue contre le Cancer Comité de

Charente-Maritime (équipe labellisée); Agence National de la Recherche (ANR) – Projets blancs; ANR under the frame of E-Rare-2, the ERA-Net for Research on Rare Diseases; Association pour la recherche sur le cancer (ARC); Cancéropôle Ile-de-France; Institut National du Cancer (INCa); Inserm (HTE); Institut Universitaire de France; Fondation pour la Recherche Médicale (FRM); the European Commission (ArtForce); the European Research Council (ERC); Fondation Carrefour; the LeDucq Foundation; the LabEx Immuno-Oncology; the RHU Torino Lumière, the SIRIC Stratified Oncology Cell DNA Repair and Tumor Immune Elimination (SOCRATE); the SIRIC Cancer Research and Personalized Medicine (CARPEM); and the Paris Alliance of Cancer Research Institutes (PACRI).

SUPPLEMENTAL MATERIAL

All supplemental data for this article are available online at www.microbialcell.com.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

COPYRIGHT

© 2017 Stekovic *et al.* This is an open-access article released under the terms of the Creative Commons Attribution (CC BY) license, which allows the unrestricted use, distribution, and reproduction in any medium, provided the original author and source are acknowledged.

Please cite this article as: Slaven Stekovic, Christoph Ruckstuhl, Philipp Royer, Christof Winkler-Hermaden, Didac Carmona-Gutierrez, Kai-Uwe Fröhlich, Guido Kroemer, and Frank Madeo (2017). The neuroprotective steroid progesterone promotes mitochondrial uncoupling, reduces cytosolic calcium and augments stress resistance in yeast cells. **Microbial Cell** 4(6): 191–199. doi: 10.15698/mic2017.06.577

REFERENCES

- Baulieu E and Schumacher M (2000). Progesterone as a neuroactive neurosteroid, with special reference to the effect of progesterone on myelination. **Steroids** 65(10–11): 605–612.
- Vagnerova K, Koerner IP, and Hurn PD (2008). Gender and the Injured Brain. **Anesth Analg** 107(1): 201–214.
- Meffre D, Labombarda F, Delespierre B, Chastre A, De Nicola AF, Stein DG, Schumacher M, and Guennoun R (2013). Distribution of membrane progesterone receptor alpha in the male mouse and rat brain and its regulation after traumatic brain injury. **Neuroscience** 231: 111–124.
- Si D, Wang H, Wang Q, Zhang C, Sun J, Wang Z, Zhang Z, and Zhang Y (2013). Progesterone treatment improves cognitive outcome following experimental traumatic brain injury in rats. **Neurosci Lett** 553: 18–23.
- Soltani Z, Khaksari M, Shahrokhi N, Mohammadi G, Mofid B, Vaziri A, and Amiresmaili S (2016). Effect of estrogen and/or progesterone administration on traumatic brain injury-caused brain edema: the changes of aquaporin-4 and interleukin-6. **J Physiol Biochem** 72(1): 33–44.
- Carswell HV, Anderson NH, Clark JS, Graham D, Jeffs B, Dominiczak AF, and Macrae IM (1999). Genetic and gender influences on sensitivity to focal cerebral ischemia in the stroke-prone spontaneously hypertensive rat. **Hypertens Dallas Tex** 1979 33(2): 681–685.
- Alkayed NJ, Murphy SJ, Traystman RJ, Hurn PD, and Miller VM (2000). Neuroprotective effects of female gonadal steroids in reproductively senescent female rats. **Stroke** 31(1): 161–168.
- Xiao G, Wei J, Yan W, Wang W, and Lu Z (2008). Improved outcomes from the administration of progesterone for patients with acute severe traumatic brain injury: a randomized controlled trial. **Crit Care Lond Engl** 12(2): R61.
- Wright DW, Kellermann AL, Hertzberg VS, Clark PL, Frankel M, Goldstein FC, Salomone JP, Dent LL, Harris OA, Ander DS, Lowery DW, Patel MM, Denson DD, Gordon AB, Wald MM, Gupta S, Hoffman SW, and

- Stein DG (2007). ProTECT: A Randomized Clinical Trial of Progesterone for Acute Traumatic Brain Injury. *Ann Emerg Med* 49(4): 391–402.e2.
10. Skolnick BE, Maas AI, Narayan RK, van der Hoop RG, MacAllister T, Ward JD, Nelson NR, Stocchetti N, and SYNAPSE Trial Investigators (2014). A clinical trial of progesterone for severe traumatic brain injury. *N Engl J Med* 371(26): 2467–2476.
11. Lin C, He H, Li Z, Liu Y, Chao H, Ji J, and Liu N (2015). Efficacy of progesterone for moderate to severe traumatic brain injury: a meta-analysis of randomized clinical trials. *Sci Rep* 5: 13442.
12. Zeng Y, Zhang Y, Ma J, and Xu J (2015). Progesterone for Acute Traumatic Brain Injury: A Systematic Review of Randomized Controlled Trials. *PLoS One* 10(10): e0140624.
13. Madeo F, Fröhlich E, and Fröhlich KU (1997). A yeast mutant showing diagnostic markers of early and late apoptosis. *J Cell Biol* 139(3): 729–734.
14. Madeo F, Fröhlich E, Ligr M, Grey M, Sigrist SJ, Wolf DH, and Fröhlich KU (1999). Oxygen stress: a regulator of apoptosis in yeast. *J Cell Biol* 145(4): 757–767.
15. Madeo F, Herker E, Maldener C, Wissing S, Lächelt S, Herlan M, Fehr M, Lauber K, Sigrist SJ, Wesselborg S, and Fröhlich KU (2002). A caspase-related protease regulates apoptosis in yeast. *Mol Cell* 9(4): 911–917.
16. Herker E, Jungwirth H, Lehmann KA, Maldener C, Fröhlich K-U, Wissing S, Büttner S, Fehr M, Sigrist S, and Madeo F (2004). Chronological aging leads to apoptosis in yeast. *J Cell Biol* 164(4): 501–507.
17. Wissing S, Ludovico P, Herker E, Büttner S, Engelhardt SM, Decker T, Link A, Proksch A, Rodrigues F, Corte-Real M, Fröhlich K-U, Manns J, Candé C, Sigrist SJ, Kroemer G, and Madeo F (2004). An AIF orthologue regulates apoptosis in yeast. *J Cell Biol* 166(7): 969–974.
18. Büttner S, Ruli D, Vögtle F-N, Galluzzi L, Moitzi B, Eisenberg T, Kepp O, Habernig L, Carmona-Gutierrez D, Rockenfeller P, Laun P, Breitenbach M, Khoury C, Fröhlich K-U, Rechberger G, Meisinger C, Kroemer G, and Madeo F (2011). A yeast BH3-only protein mediates the mitochondrial pathway of apoptosis. *EMBO J* 30(14): 2779–2792.
19. Galluzzi L, Kepp O, and Kroemer G (2016). Mitochondrial regulation of cell death: a phylogenetically conserved control. *Microb Cell* 3(3): 101–108.
20. Büttner S, Habernig L, Broeskamp F, Ruli D, Vögtle FN, Vlachos M, Macchi F, Küttner V, Carmona-Gutierrez D, Eisenberg T, Ring J, Markaki M, Taskin AA, Benke S, Ruckenstein C, Braun R, Van den Haute C, Bammens T, van der Perren A, Fröhlich K-U, Winderickx J, Kroemer G, Baekelandt V, Tavernarakis N, Kovacs GG, Dengjel J, Meisinger C, Sigrist SJ, and Madeo F (2013). Endonuclease G mediates α -synuclein cytotoxicity during Parkinson's disease. *EMBO J* 32(23): 3041–3054.
21. Büttner S, Broeskamp F, Sommer C, Markaki M, Habernig L, Alavi-Ghavanini A, Carmona-Gutierrez D, Eisenberg T, Michael E, Kroemer G, Tavernarakis N, Sigrist SJ, and Madeo F (2014). Spermidine protects against α -synuclein neurotoxicity. *Cell Cycle Georget Tex* 13(24): 3903–3908.
22. Heinisch JJ and Brandt R (2016). Signaling pathways and posttranslational modifications of tau in Alzheimer's disease: the humanization of yeast cells. *Microb Cell* 3(4): 135–146.
23. Menezes R, Tenreiro S, Macedo D, Santos C, and Outeiro T (2015). From the baker to the bedside: yeast models of Parkinson's disease. *Microb Cell* 2(8): 262–279.
24. Amen T and Kaganovich D (2016). Yeast screening platform identifies FDA-approved drugs that reduce A β oligomerization. *Microb Cell* 3(3): 97–100.
25. Shrestha A and Megeney L (2015). Yeast proteinopathy models: a robust tool for deciphering the basis of neurodegeneration. *Microb Cell* 2(12): 458–465.
26. Bond M, Brown R, Rallis C, Bahler J, and Mole S (2015). A central role for TOR signalling in a yeast model for juvenile CLN3 disease. *Microb Cell* 2(12): 466–480.
27. Carmona-Gutierrez D, Hughes AL, Madeo F, and Ruckenstein C (2016). The crucial impact of lysosomes in aging and longevity. *Ageing Res Rev* 32: 2–12. doi: 10.1016/j.arr.2016.04.009.
28. Raghupathi R (2004). Cell death mechanisms following traumatic brain injury. *Brain Pathol Zurich Switz* 14(2): 215–222.
29. Cheng G, Kong R, Zhang L, and Zhang J (2012). Mitochondria in traumatic brain injury and mitochondrial-targeted multipotential therapeutic strategies: Mitochondria in traumatic brain injury. *Br J Pharmacol* 167(4): 699–719.
30. Ludovico P, Sousa MJ, Silva MT, Leão C, and Côrte-Real M (2001). *Saccharomyces cerevisiae* commits to a programmed cell death process in response to acetic acid. *Microbiol Read Engl* 147(Pt 9): 2409–2415.
31. Rockenfeller P, Ring J, Muschett V, Beranek A, Büttner S, Carmona-Gutierrez D, Eisenberg T, Khoury C, Rechberger G, Kohlwein SD, Kroemer G, and Madeo F (2010). Fatty acids trigger mitochondrion-dependent necrosis. *Cell Cycle Georget Tex* 9(14): 2836–2842.
32. Eisenberg T, Carmona-Gutierrez D, Büttner S, Tavernarakis N, and Madeo F (2010). Necrosis in yeast. *Apoptosis Int J Program Cell Death* 15(3): 257–268.
33. Carmona-Gutiérrez D, Bauer MA, Ring J, Knauer H, Eisenberg T, Büttner S, Ruckenstein C, Reisenbichler A, Magnes C, Rechberger GN, Birner-Gruenberger R, Jungwirth H, Fröhlich K-U, Sinner F, Kroemer G, and Madeo F (2011). The propeptide of yeast cathepsin D inhibits programmed necrosis. *Cell Death Dis* 2: e161.
34. Braun RJ, Sommer C, Carmona-Gutierrez D, Khoury CM, Ring J, Büttner S, and Madeo F (2011). Neurotoxic 43-kDa TAR DNA-binding protein (TDP-43) triggers mitochondrion-dependent programmed cell death in yeast. *J Biol Chem* 286(22): 19958–19972.
35. Holmström KM and Finkel T (2014). Cellular mechanisms and physiological consequences of redox-dependent signalling. *Nat Rev Mol Cell Biol* 15(6): 411–421.
36. Braun RJ, Sommer C, Leibiger C, Gentier RJG, Dumit VI, Paduch K, Eisenberg T, Habernig L, Trausinger G, Magnes C, Pieber T, Sinner F, Dengjel J, van Leeuwen FW, Kroemer G, and Madeo F (2015). Accumulation of Basic Amino Acids at Mitochondria Dictates the Cytotoxicity of Aberrant Ubiquitin. *Cell Rep* 10(9): 1557–1571.
37. Braun R, Sommer C, Leibiger C, Gentier R, Dumit V, Paduch K, Eisenberg T, Habernig L, Trausinger G, Magnes C, Pieber T, Sinner F, Dengjel J, van Leeuwen F, Kroemer G, and Madeo F (2015). Modeling non-hereditary mechanisms of Alzheimer disease during apoptosis in yeast. *Microb Cell* 2(4): 136–138.
38. Ruckenstein C, Büttner S, Carmona-Gutierrez D, Eisenberg T, Kroemer G, Sigrist SJ, Fröhlich K-U, and Madeo F (2009). The Warburg Effect Suppresses Oxidative Stress Induced Apoptosis in a Yeast Model for Cancer. *PLoS ONE* 4(2): e4592.
39. Hand RA, Jia N, Bard M, and Craven RJ (2003). *Saccharomyces cerevisiae* Dap1p, a novel DNA damage response protein related to the mammalian membrane-associated progesterone receptor. *Eukaryot Cell* 2(2): 306–317.
40. Gregor A, Kocylowski M, and Kostorzewska E (1986). Evaluation of the diagnostic usefulness of determining porphobilinogen deaminase activity in the erythrocytes in patients with acute intermittent porphy-

ria and in carriers of the gene of this type of porphyria. **Przegl Lek** 43(11): 703–705.

41. Tubiana N, Mishal Z, le Caer F, Seigneurin JM, Berthoix Y, Martin PM, and Carcassonne Y (1986). Quantification of oestradiol binding at the surface of human lymphocytes by flow cytofluorimetry. **Br J Cancer** 54(3): 501–504.

42. Roussel D, Harding M, Runswick MJ, Walker JE, and Brand MD (2002). Does any yeast mitochondrial carrier have a native uncoupling protein function? **J Bioenerg Biomembr** 34(3): 165–176.

43. Barros MH, Bandy B, Tahara EB, and Kowaltowski AJ (2004). Higher respiratory activity decreases mitochondrial reactive oxygen release and increases life span in *Saccharomyces cerevisiae*. **J Biol Chem** 279(48): 49883–49888.

44. Skulachev VP (1998). Uncoupling: new approaches to an old problem of bioenergetics. **Biochim Biophys Acta BBA - Bioenerg** 1363(2): 100–124.

45. Mookerjee SA, Divakaruni AS, Jastroch M, and Brand MD (2010). Mitochondrial uncoupling and lifespan. **Mech Ageing Dev** 131(7–8): 463–472.

46. Amara CE, Shankland EG, Jubrias SA, Marcinek DJ, Kushmerick MJ, and Conley KE (2007). Mild mitochondrial uncoupling impacts cellular aging in human muscles *in vivo*. **Proc Natl Acad Sci** 104(3): 1057–1062.

47. Rodriguez AM, Monjo M, Roca P, and Palou A (2002). Opposite actions of testosterone and progesterone on UCP1 mRNA expression in cultured brown adipocytes. **Cell Mol Life Sci CMLS** 59(10): 1714–1723.

48. Wade Ruth and Jones Howard W. Jr. (1956). Effect of progesterone on mitochondrial adenosinetriphosphatase. **JBC** 220: 547–551.

49. Liu W (2012). Control of Calcium in Yeast Cells. Introduction to Modeling Biological Cellular Control Systems. Springer Milan, Milano; pp. 95–122.

50. Ren ZW (1992). Radiofrequency ablation of left-sided atrioventricular accessory tract to treat supraventricular tachycardia. **Zhonghua Xin Xue Guan Bing Za Zhi** 20(4): 212–214.

51. Romarowski A, Sánchez-Cárdenas C, Ramírez-Gómez HV, Puga Molina L del C, Treviño CL, Hernández-Cruz A, Darszon A, and Buffone MG (2016). A Specific Transitory Increase in Intracellular Calcium Induced by Progesterone Promotes Acrosomal Exocytosis in Mouse Sperm. **Biol Reprod** 94(3): 63.

52. Li L-F, Xiang C, Zhu Y-B, and Qin K-R (2014). Modeling of progesterone-induced intracellular calcium signaling in human spermatozoa. **J Theor Biol** 351: 58–66.

53. Bonaccorsi L (1998). Progesterone-stimulated intracellular calcium increase in human spermatozoa is protein kinase C-independent. **Mol Hum Reprod** 4(3): 259–268.

54. Miller MR, Mannowetz N, Iavarone AT, Safavi R, Gracheva EO, Smith JF, Hill RZ, Bautista DM, Kirichok Y, and Lishko PV (2016). Unconventional endocannabinoid signaling governs sperm activation via the sex hormone progesterone. **Science** 352(6285): 555–559.

55. Batra S (1986). Effect of estrogen and progesterone treatment on calcium uptake by the myometrium and smooth muscle of the lower urinary tract. **Eur J Pharmacol** 127(1–2): 37–42.

56. Barbagallo M, Dominguez LJ, Licata G, Shan J, Bing L, Karpinski E, Pang PKT, and Resnick LM (2001). Vascular Effects of Progesterone :

Role of Cellular Calcium Regulation. **Hypertens Dallas Tex** 1979 37(1): 142–147.

57. He Y, Gao Q, Han B, Zhu X, Zhu D, Tao J, Chen J, and Xu Z (2016). Progesterone suppressed vasoconstriction in human umbilical vein via reducing calcium entry. **Steroids** 108: 118–125.

58. Wu Z and Shen W (2010). Progesterone inhibits L-type calcium currents in gallbladder smooth muscle cells. **J Gastroenterol Hepatol** 25(12): 1838–1843.

59. Luoma JI, Kelley BG, and Mermelstein PG (2011). Progesterone inhibition of voltage-gated calcium channels is a potential neuroprotective mechanism against excitotoxicity. **Steroids** 76(9): 845–855.

60. Luoma JI, Stern CM, and Mermelstein PG (2012). Progesterone inhibition of neuronal calcium signaling underlies aspects of progesterone-mediated neuroprotection. **J Steroid Biochem Mol Biol** 131(1–2): 30–36.

61. Cai W, Zhu Y, Furuya K, Li Z, Sokabe M, and Chen L (2008). Two different molecular mechanisms underlying progesterone neuroprotection against ischemic brain damage. **Neuropharmacology** 55(2): 127–138.

62. Brotfain E, Gruenbaum SE, Boyko M, Kutz R, Zlotnik A, and Klein M (2016). Neuroprotection by Estrogen and Progesterone in Traumatic Brain Injury and Spinal Cord Injury. **Curr Neuropharmacol** 14(6): 641–653.

63. Wei J and Xiao G (2013). The neuroprotective effects of progesterone on traumatic brain injury: current status and future prospects. **Acta Pharmacol Sin** 34(12): 1485–1490.

64. Pascual JL, Murcy MA, Li S, Gong W, Eisenstadt R, Kumasaka K, Sims C, Smith DH, Browne K, Allen S, and Baren J (2013). Neuroprotective effects of progesterone in traumatic brain injury: blunted *in vivo* neutrophil activation at the blood-brain barrier. **Am J Surg** 206(6): 840–845; discussion 845–846.

65. Li Z, Wang B, Kan Z, Zhang B, Yang Z, Chen J, Wang D, Wei H, Zhang J, and Jiang R (2012). Progesterone increases circulating endothelial progenitor cells and induces neural regeneration after traumatic brain injury in aged rats. **J Neurotrauma** 29(2): 343–353.

66. Webster KM, Wright DK, Sun M, Semple BD, Ozturk E, Stein DG, O'Brien TJ, and Shultz SR (2015). Progesterone treatment reduces neuroinflammation, oxidative stress and brain damage and improves long-term outcomes in a rat model of repeated mild traumatic brain injury. **J Neuroinflammation** 12: 238.

67. Gueldener U, Heinisch J, Koehler GJ, Voss D, and Hegemann JH (2002). A second set of loxP marker cassettes for Cre-mediated multiple gene knockouts in budding yeast. **Nucleic Acids Res** 30(6): e23.

68. Güldener U, Heck S, Fielder T, Beinhauer J, and Hegemann JH (1996). A new efficient gene disruption cassette for repeated use in budding yeast. **Nucleic Acids Res** 24(13): 2519–2524.

69. Loewith R, Jacinto E, Wullschlegel S, Lorbberg A, Crespo JL, Bonenfant D, Oppliger W, Jenoe P, and Hall MN (2002). Two TOR complexes, only one of which is rapamycin sensitive, have distinct roles in cell growth control. **Mol Cell** 10(3): 457–468.

70. Kainz K, Tadic J, Zimmermann A, Pendl T, Carmona-Gutierrez D, Ruckenstein C, Eisenberg T, and Madeo F (2017). Methods to Assess Autophagy and Chronological Aging in Yeast. **Methods in Enzymology** 588: 367–394.