

## Short communication

## Tryptophan ingestion by gestant mothers alters prolactin and luteinizing hormone release in the adult male offspring

Lorenzo Martin <sup>b</sup>, Manuel Rodríguez Díaz <sup>b,\*</sup>, Celsa Santana-Herrera <sup>b</sup>, Antonio Milena <sup>a</sup>,  
Cristo Santana <sup>c</sup>

<sup>a</sup> *Canarian University Hospital, University of La Laguna, La Laguna, 38320 Tenerife, Spain*

<sup>b</sup> *Department of Physiology, School of Medicine, University of La Laguna, La Laguna, 38320 Tenerife, Spain*

<sup>c</sup> *Department of Psychobiology, School of Psychology, Canarian University Hospital, University of La Laguna, La Laguna, 38320 Tenerife, Spain*

Accepted 8 July 1997

---

**Abstract**

The effect of tryptophan administration to pregnant rats on the development of serotonergic systems and serotonin-related hormones in the offspring was studied. The male offspring of rats treated with tryptophan (200 mg/kg/day) during the second half of gestation showed a 4- to 7-fold increase in serum prolactin 40 and 70 days after birth and a 2-fold increase in serum luteinizing hormone 70 days after birth. The forebrain of adult offspring of tryptophan-treated rats showed an increase in serotonin and 5-hydroxyindoleacetic acid levels. Present data suggest that tryptophan regulates serotonergic differentiation during early development. © 1997 Elsevier Science B.V.

**Keywords:** Tryptophan; Serotonin; Prolactin; Luteinizing hormone; Development

---

In addition to their action as neurotransmitters during adulthood, monoamines could act as a differentiation signal during early neurogenesis [1,19,22]. We have previously reported that oral tyrosine administration to pregnant rats induces in the fetus a short-lasting increase in tyrosine and catecholamine brain levels [5] and in the adult offspring a persistent modification of dopamine and 3,4-dihydroxyphenylacetic acid (DOPAC) levels in several brain loci [22], dopamine agonists response [22] and different behavioral patterns [1,21]. Despite tryptophan administration to pregnant rats induces in the fetal brain a similar short-lasting modification of serotonergic neurotransmission [2], its action on the ontogenic development of serotonergic systems has received little attention.

In animals and humans, the serotonergic cell differentiation occurs prenatally and their dendritic growth and terminal synaptogenesis take place during postnatal life [2,8,24,27]. The present study was designed to assess the effect of oral tryptophan administration to pregnant rats on the postnatal development of serotonergic systems in the offspring. The regulation of the neuroendocrine system is one of the main roles of serotonergic cells during adult-

hood. Amongst other hormones, serotonergic neurons regulate prolactin release [4,6,11,13,14,25], and the activity of the hypothalamic-pituitary-gonadal system [23,26]. In order to determine whether prenatal tryptophan action on serotonin systems has persistent functional consequences during postnatal life, the effect of prenatal administration of tryptophan on prolactin and luteinizing hormone (LH) was also examined.

Experiments were carried out on male Sprague–Dawley rats (Letica, Barcelona). Animals were housed at 22°C, two per cage, under normal laboratory conditions on a standard light–dark schedule (12:12 h with 3.00–15.00 h light on) and with free access to food and water.

L-Tryptophan (200 mg/kg; Sigma, St. Louis, MO) was given daily by stomach intubation to young pregnant rats (150–200 g body weight before gestation). The amino acid or its saline vehicle was administered during the light cycle from day 15 to day 21 of gestation. Timing of pregnancies was determined by daily checking vaginal washings for sperm; the day on which sperm was found was regarded as day 0. Only the rats that were born on day 21 of gestation were included in the study. Sixty rats were divided into six groups of 10 rats each. In order to prevent a litter effect each of the animals included in a group descended from a different mother. The rats were sacri-

---

\* Corresponding author.

ficed by decapitation at the onset of the dark period of days 40 or 70 after birth. Blood was collected from the trunk and allowed to clot at room temperature. Serum was separated by centrifugation ( $2000 \times g$ ), divided into aliquots, and stored at  $-40^{\circ}\text{C}$  until assay. Immediately after extraction, the brain was laid on its dorsal surface and the mesencephalon and forebrain dissected from the forebrain structures by the mesencephalic flexure and according to a previously described procedure [2]. The brain pieces were rapidly removed and weighed in conical 5 ml test tubes. Two ml of 0.1 M perchloric acid containing  $4 \times 10^{-5}$  M sodium metabisulphite was pipetted into the tubes to avoid the metabolization or oxidation of monoamines. Thus, 2 min after rat sacrifice the brain monoamines were protected and the solution became stabilized. The mixture was then sonicated at 100 W for about 12 s while on ice, and the homogenate was centrifuged for 15 min at  $15\,000 \times g$ . The supernatant and the pellet were kept in separate tubes. All the biochemical studies were made in the following 2 months after brain dissection and to avoid deterioration of the biological sample, it remained cooled ( $-70^{\circ}\text{C}$ ) until the biochemical determinations were begun. Using this procedure the concentration of the studied compounds is not significantly modified prior to the biochemical determinations [3].

Serum LH and prolactin radioimmunoassays were performed using the materials provided by the NIAMDD and the results were expressed in terms of the respective RP-3 preparation according to previously validated procedures [18].

Brain serotonin (5-HT) and 5-hydroxyindoleacetic acid (5-HIAA) levels were measured by liquid chromatography with electrochemical detection according to previously validated procedures [2,3,22]. An aliquot of the supernatant was injected into the chromatographic column ( $300 \times 3.9$  mm stainless steel column packed with Nova-Pack c18, 4  $\mu\text{m}$  particle size (Waters, Milford, MA, USA). The mobile phase consisted of 0.1 M  $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$ , 0.5 mM EDTA, 1 g/l sodium heptanesulphonate (PIC B7), 6% acetonitrile. The final solution ( $\text{pH} = 4.35$ ) was filtered (0.45  $\mu\text{m}$  Millipore filter) before use. Standards of serotonin and 5-HIAA (Sigma, St. Louis, MO) were dissolved in perchloric acid (PCA) containing  $4 \times 10^{-5}$  M sodium metabisulphite and kept as stock solutions at  $-20^{\circ}\text{C}$ . They were diluted with ice-cold PCA/ $\text{Na}_2\text{O}_5\text{S}_2$  shortly before chromatographic injection. All separations were performed isocratically at a flow-rate of 1.0 ml/min at room temperature. The electrochemical detector used was a Waters Model 460. The detector potential was 0.82 V. Quantifications were performed from standard curves of peak height.




Fig. 1. Top: 5-HIAA and serotonin levels in the forebrain and mesencephalon of 70-day-old rats. Bottom: serum prolactin and LH levels of 40- and 70-day-old rats. The bars represent mean values  $\pm$  S.E.M. \*  $P < 0.05$ ; \*\*  $P < 0.001$ .

The statistical analyses were performed with the Complete Statistical System (CSS) of StatSoft and using a *t*-test. Differences were judged significant when associated with a probability of 5% or less.

As Fig. 1 shows, oral administration of tryptophan to gestant mothers induced, in the male offspring, a marked increase in serum prolactin (days 40 and 70), serum LH (day 70) and forebrain levels of both serotonin and 5-HIAA. No statistical differences were found for serotonin and 5-HIAA, in the forebrain (5-HT  $t_{18} = 1.0$ , 5-HIAA  $t_{18} = 0.87$ ) and mesencephalon (5-HT  $t_{18} = 0.11$ , 5-HIAA  $t_{18} = -0.15$ ) of 40-day-old groups and in the mesencephalon ( $t_{18} = -1.41$ , n.s., serotonin for tryptophan vs. vehicle groups;  $t_{18} = -1.08$ , n.s., 5-HIAA for tryptophan vs. vehicle groups) of 70-day-old groups.

We have previously reported that tryptophan administered to pregnant rats crosses the placental barrier and induces a dose-related increase in tryptophan concentration in the placenta, body and brain of the fetus [2]. Because tryptophan hydroxylase activity is not saturated by its substrate, this increase in tryptophan concentration produces a parallel increase in 5-HT and 5-HIAA levels in the fetal brain [2]. The effect begins 1 h after tryptophan intake and persists for 24 h [2]. The offspring of tryptophan-treated mothers showed 70 days after birth a synthesis (5-HT levels) and metabolism (5-HIAA levels) rise for 5-HT. These data indicate that, in addition to the transitory modification of 5-HT metabolism in the fetal brain, maternal intake of tryptophan during gestation facilitates serotonergic neurotransmission in offspring during adulthood. We found no effects in prepuber rats. Serotonergic neurons differentiate during early ontogenesis and their fibers reach the hypothalamus in the last days of prenatal life [2,8,24,27]. However, the final termination density and the formation of precise termination patterns are not reached until adulthood [10]. Mechanisms involved in the regulation of this delayed synaptogenesis [9] may be at the basis of the prepuber-postpuber differences observed here for the effect of mother intake of tryptophan on serotonergic system development of the offspring.

During adulthood, serotonin is involved in the neural regulation of prolactin release [4,11–13,15,23,25]. The peripheral administration of 5-hydroxytryptophan [12], the intravenous administration of 5-HT [20] or the administration of serotonergic agonists [12] facilitates prolactin release to plasma. The drug blockade of serotonergic neurotransmission decreases the serum prolactin level [7]. Thus, 5-HT facilitates prolactin release from the anterior pituitary gland. Fig. 1 shows a serum prolactin level increase in the offspring of tryptophan mothers. These data suggest that the prenatal administration of tryptophan induces not only a long-lasting activation of serotonergic neurotransmission but also a persistent promotion of the facilitatory activity of 5-HT on prolactin release.

Serotonergic transmission is also involved in the neural regulation of the hypothalamic-pituitary-gonadal system

[18,20,23,26]. The intraventricular injection of high concentrations of 5-HT evokes LH release in male rats [11]. The intraventricular injection of the serotonergic neurotoxin 5,7-dihydroxytryptamine decreases LH release [26]. Thus, and despite serotonergic innervation of preoptic area [18] and hypothalamus [23] probably having a different involvement in the regulation of LH release, these previous data show that an overall facilitation of brain serotonergic neurotransmission during adulthood promotes LH release [14]. Fig. 1 shows that, corresponding to the 5-HT increase, the prenatal administration of tryptophan induces a serum LH concentration increase that can be detected 70 days after birth. Taken together, present data show that prenatal administration of tryptophan induces a long-lasting facilitation of central serotonergic neurotransmission, disrupting the serotonergic regulation of pituitary hormone release.

In conclusion, the present data suggest that excessive ingestion of amino acids during gestation could have important consequences on the brain development of the offspring. In humans, it has been reported that transient tyrosinemia during neonatal life can induce permanent intellectual impairment 8–10 years later [16,17]. Further studies are warranted to evaluate the functional consequences of the neonatal increase in serum tryptophan on human development. According to the hypothesis previously proposed for tyrosine and the catecholaminergic systems [1,5,19,21,22], present data support the idea that serotonin may have a neurotrophic role prior to the evolutionary onset of neurotransmission.

## Acknowledgements

These studies were supported by a grant from the Comisión Interministerial de Ciencia y Tecnología, Plan Nacional I + D, SAL 93-0037, Spain. The generous gift by the NIAMDD National Hormone and Pituitary Program of the materials for LH and prolactin radioimmunoassays is acknowledged.

## References

- [1] R. Arevalo, R. Castro, M.D. Palarea, M. Rodriguez, Tyrosine administration to pregnant rats induces persistent behavioral modifications in the male offspring, *Physiol. Behav.* 39 (1987) 477–481.
- [2] R. Arevalo, D. Afonso, R. Castro, M. Rodriguez, Fetal brain serotonin synthesis and catabolism is under control by mother intake of tryptophan, *Life Sci.* 49 (1991) 53–66.
- [3] D. Afonso, M.A. Castellano, M. Rodriguez, Determination of monoamines and indoles in amniotic fluid by high-performance liquid chromatography-electrochemical detection, *J. Chromatogr.* 528 (1990) 101–109.
- [4] C.L. Chen, J. Meites, Effects of biogenic amines and TRH on release of prolactin and TSH in the rat, *Endocrinology* 96 (1975) 10–18.
- [5] M.V. Garabal, R.M. Arevalo, M.D. Diaz-Palarea, R. Castro, M.

- Rodriguez, Tyrosine availability and brain noradrenaline synthesis in the fetus: control by maternal tyrosine ingestion, *Brain Res.* 457 (1988) 330–337.
- [6] S.E. Gartside, P.J. Cowen, Mediation of ACTH and prolactin response to 5-HTP by 5-HTP2 receptors, *Eur. J. Pharmacol.* 179 (1990) 103–109.
- [7] J. Gid-Ad, F. Zambotti, M.O. Carruba, L. Vicentini, E.E. Muller, Stimulatory role for brain serotonergic system on prolactin secretion in the male rat, *Proc. Soc. Exp. Biol. Med.* 151 (1976) 512–518.
- [8] M. Hyypä, Hypothalamic monoamines in human fetuses, *Neuroendocrinology* 9 (1972) 257–266.
- [9] B.L. Jacobs, E.C. Azmitia, Structure and function of the brain serotonin system, *Physiol. Rev.* 72 (1992) 165–229.
- [10] J.M. Lauder, F.E. Bloom, Ontogeny of monoamine neurons in the locus coeruleus, raphe nuclei and substantia nigra of the rat. II. Synaptogenesis, *J. Comp. Neurol.* 163 (1975) 251–264.
- [11] I.A. Kamberi, R.S. Mical, J.C. Porter, Effect of anterior pituitary perfusion and intraventricular injection of catecholamines on prolactin release, *Endocrinology* 88 (1971) 1012–1020.
- [12] C. Kordon, A. Blake, J. Terkel, C.H. Sawyer, Participation of serotonin-containing neurons in the suckling-induced rise in plasma prolactin levels in lactating rats, *Neuroendocrinology* 13 (1973) 213–223.
- [13] Y. Kato, Y. Nakai, H. Imura, K. Chihara, S.J. Ohgo, Effect of 5-hydroxytryptophan (5-HTP) on plasma prolactin levels in man, *J. Clin. Endocrinol. Metab.* 38 (1974) 695–698.
- [14] S.P. Kalra, P.S. Kalra, Neural regulation of luteinizing hormone secretion in the rat, *Endocr. Rev.* 4 (1983) 311–350.
- [15] K. Lu, J. Meites, Effects of serotonin precursors and melatonin on serum prolactin release in rats, *Endocrinology* 93 (1973) 152–155.
- [16] J.H. Menkes, D.W. Welcher, H.S. Levi, J. Dallas, N.E. Gretskey, Relationship of elevated blood tyrosine to the ultimate intellectual performance of premature infants, *Pediatrics* 49 (1972) 218–224.
- [17] P. Mamunes, P.E. Prince, N.H. Thornton, P.A. Hunt, E.S. Hitchcock, Intellectual deficits after transient tyrosinemia in the term neonate, *Pediatrics* 57 (1976) 675–680.
- [18] M. Mas, M.C. Gonzalez, M. Rodriguez, R. Castro, S. Oaknin, Effects of monoamine neurotoxins injected in different brain areas on gonadotropin and androgen secretion in the male, *Neuroendocrinology* 39 (1984) 156–161.
- [19] M.P. Mattson, Neurotransmitters in the regulation of neural cytoarchitecture, *Brain Res. Rev.* 13 (1988) 179–212.
- [20] N.S. Pilotte, J.C. Porter, Circulating luteinizing hormone and prolactin in systemic plasma of rats treated with 5-hydroxytryptamine, *Endocrinology* 105 (1979) 875–878.
- [21] M. Rodriguez, C. Santana, D. Afonso, Maternal ingestion of tyrosine during rat pregnancy modifies the offspring behavioral laterilization, *Physiol. Behav.* 55 (1994) 607–613.
- [22] C. Santana, L. Martin, M. Rodriguez, Tyrosine ingestion during rat pregnancy alters postnatal development of dopaminergic neurons in the offspring, *Brain Res.* 635 (1994) 96–106.
- [23] L. Van der Kar, S.A. Lorens, A. Vodraska, G. Allers, M. Green, D.E. Van Order, L.S. Van Order, Effect of selective midbrain and diencephalic 5,7-dihydroxytryptamine lesions on serotonin content in individual preoptic hypothalamic nuclei and on serum luteinizing hormone levels, *Neuroendocrinology* 31 (1980) 309–315.
- [24] J.A. Wallace, J.M. Lauder, Development of serotonergic system in the rat embryo: an immunocytochemical study, *Brain Res. Bull.* 10 (1983) 459–479.
- [25] P.D. Wolf, L.J. Lee, Effect of the serotonin precursor, tryptophan, on pituitary hormone secretion, *J. Clin. Endocrinol. Metab.* 45 (1977) 123–126.
- [26] W. Wuttke, J.L. Hancke, K.G. Hohn, H.G. Baumgarten, Effect of intraventricular injection of diencephalic 5,7-dihydroxytryptamine on serum gonadotropins and prolactin, *Ann. NY Acad. Sci.* 305 (1978) 423–436.
- [27] S.H. Zeisel, C. Mauron, C.J. Watkins, R.J. Wurtman, Developmental changes in brain indoles, serum tryptophan and other serum neutral amino acids in the rat, *Dev. Brain Res.* 1 (1981) 551–564.