

Oxidized Lipids in the Diet Are a Source of Oxidized Lipid in Chylomicrons of Human Serum

Ilona Staprāns, Joseph H. Rapp, Xian-Mang Pan, Kelly Y. Kim, Kenneth R. Feingold

Abstract We examined whether oxidized lipids in the diet determine the levels of oxidized lipid in human postprandial serum chylomicrons. After we fed subjects control corn oil containing low quantities of oxidized lipid, the levels of conjugated dienes in the chylomicron fraction were low (9.67 ± 0.92 nmol/ μ mol triglyceride), and no thiobarbituric acid-reactive substances (TBARS) could be detected. However, when subjects were fed a highly oxidized oil, the conjugated diene content in chylomicrons was increased 4.7-fold to 46 ± 5.63 nmol/ μ mol triglyceride, with 0.140 ± 0.03 nmol TBARS/ μ mol triglyceride. When subjects were fed medium-oxidized oil, the degree of oxidation of the chylomicron lipids was moderately increased (21.86 ± 2.03 nmol conjugated

dienes/ μ mol triglyceride). Additionally, we found that chylomicrons isolated after ingestion of oxidized oil were more susceptible to CuSO_4 oxidation than chylomicrons isolated after ingestion of the control oil. The lag time for oxidation decreased from 4.30 ± 0.40 to 3.24 ± 0.51 hours ($P < .05$). These data demonstrate that in humans dietary oxidized lipids are absorbed by the small intestine, incorporated into chylomicrons, and appear in the bloodstream, where they contribute to the total body pool of oxidized lipid. (*Arterioscler Thromb*. 1994;14:1900-1905.)

Key Words • lipid peroxides • oxidized dietary fat • chylomicrons • oxidized serum lipoproteins

Several observations suggest that oxidized lipoproteins are important in the development of atherosclerosis.¹⁻³ Incubation of oxidized low-density lipoprotein (LDL) with macrophages leads to formation of foam cells, whereas incubation with native LDL does not. Oxidized LDL is cytotoxic to endothelial cells and therefore could induce arterial wall injury.⁴ Finally, oxidized LDL stimulates the release of chemotactic proteins, cytokines, and growth factors from endothelial and other arterial wall cells, promoting the inflammatory milieu that characterizes atherosclerotic plaque.⁵⁻⁷

Lipoprotein oxidation is postulated to occur within the microenvironment of the artery wall¹⁻³; however, the source of oxidized lipoproteins in the serum of humans⁸⁻¹⁰ and animals^{11,12} has not been identified. Diet has been suggested as the source of oxidized lipoproteins in the intestinal lymph drainage of rodents.^{13,14} Studies in our laboratory have shown that the levels of oxidized lipids in the serum very-low-density lipoprotein (VLDL) plus LDL fraction of rats correlated with the quantity of oxidized lipids in the diet.¹⁵ Moreover, we observed a direct relation between the levels of oxidized lipid in the diet and the amounts of oxidized lipid in mesenteric lymph chylomicrons,^{14,15} suggesting that in rats oxidized lipids in the diet are absorbed by the small intestine, packaged in chylomicrons, and transported to the bloodstream. Furthermore, our studies have shown that these oxidized chylomicrons are

metabolized similarly to normal chylomicrons, with a substantial portion of oxidized lipids delivered to the liver, where they could potentially be repackaged and secreted in VLDL. Thus, the quantity of oxidized lipids in the diet of rodents has a major effect on the levels of oxidized lipoproteins in the serum. Naruszewicz et al¹⁰ have presented evidence that in humans the ingestion of heat-treated soy oil results in oxidized lipid containing chylomicrons whose uptake and degradation are increased in macrophages.

The hypotheses of the present study were that the amount of oxidized lipid in the diet correlates with the levels of oxidized lipid in human postprandial serum chylomicrons and that the levels of dietary oxidized lipid alter the susceptibility of postprandial chylomicrons to further oxidation.

Methods

Characterization of Study Subjects

All subjects ($n=6$) were male volunteers, 24 to 45 years of age, with serum triglyceride and cholesterol levels within the normal ranges. The average body weight of the subjects was 84.75 ± 8.97 kg; the average age was 33.00 ± 6.68 years. In the fasted state, the serum cholesterol and triglyceride concentrations were 4.90 ± 0.17 and 0.78 ± 0.17 mmol/L, respectively. The HDL cholesterol concentration was 1.18 ± 0.09 mmol/L.

All were nonsmokers, were physically active, and consumed standard American diets. None took excessive amounts of vitamin supplements or any lipid-lowering or hypertension medications.

Experimental Diets

As test meals, one control and two oxidized corn oils were used. The control oil was conventional corn oil that contained vitamin E (0.14 mg/g oil) and low amounts of oxidized lipid (6.5 to 10 nmol conjugated dienes/mg oil). Oxidized oils were α -tocopherol (vitamin E)-depleted corn oil purchased from

Received April 5, 1994; revision accepted September 22, 1994.
From the Department of Veterans Affairs Medical Center and the Departments of Surgery and Medicine, University of California, San Francisco.

Correspondence to Ilona Staprāns, PhD, VA Medical Center (151L), 4150 Clement St, San Francisco, CA 94121.

© 1994 American Heart Association, Inc.

ICN Biochemicals and contained either medium (30 to 50 nmol conjugated dienes/mg oil) or high (80 to 120 nmol of conjugated dienes/mg oil) levels of oxidized lipids, depending on storage and air exposure time before we added vitamin E. Usually, freshly purchased α -tocopherol-depleted oil was used as the medium-oxidized lipid-containing oil, and highly oxidized lipid-containing oil was obtained by air exposure for 6 to 8 weeks. When we examined dietary oils for thiobarbituric acid-reactive substances (TBARS),¹² no significant amount was detected in oils with low-oxidized lipid content. Oil with the medium- and highly oxidized lipid content contained 0.041 and 0.103 nmol TBARS/mg oil, respectively. Direct measurement of peroxides using a color reaction with methylene blue derivative¹⁵ indicated that highly oxidized oil contained 66.0 nmol peroxide/mg oil. We could not detect peroxides in other oils with this method.

Oxidation of dietary oils was further confirmed by demonstrating a reduction of linoleic acid content. Analysis of triglyceride composition showed that the control oil contained 62.45% linoleic acid (18:2) and that oxidized oil containing high levels of peroxides (called highly oxidized oil) contained 57.13% linoleic acid. The oil with medium peroxide concentration (called medium-oxidized oil) contained 59.86% linoleic acid. Thus, as expected, the relative concentration of polyunsaturated linoleic acid decreased in oxidized oils. It should be noted that highly oxidized oil is a relative term compared with the control oil. In absolute terms, the highly oxidized oil used in our experiments is only very mildly oxidized because the oil was not heated and oxidation was achieved by air exposure only. Only approximately 5% of the fatty acids had been degraded in the oxidized oil. To make all ingested oils comparable to the control oil in vitamin E concentration, vitamin E was added to oxidized oils to the same concentration as in the control oil.

Determination of Oxidation

Conjugated dienes were measured with second-derivative UV spectroscopy in a Perkin Elmer 555 Spectrophotometer as described by Corongiu et al.¹⁶ For conjugated diene determination, lipids from the chylomicron fraction were extracted by the method of Dole.¹⁷ Heptane extracts were scanned for absorbance from 260 to 220 nm, and second-derivative spectra were obtained. Absorbance range selected was ± 0.05 . Minima at 233 and 242 nm were identified and quantitated in millimeters by measuring the peak from minimum to the adjacent maximum at the higher wavelength on the printout. This method has been shown to be specific for lipid hydroperoxides. Oxidized linoleic acid was used to generate a standard curve,¹⁸ and the extinction coefficient of $32\,300\text{ L} \cdot \text{mol}^{-1} \cdot \text{cm}^{-1}$ was obtained and used to calculate the results. The results were expressed as nanomoles of conjugated dienes per micromole of triglyceride in the chylomicron fraction. The triglyceride concentration in the heptane extracts was measured as described by Fletcher.¹⁹ All measurements were performed on the same day of lipoprotein isolation. Measurement of conjugated dienes for estimating lipid oxidation was chosen not only because of the specificity but also because of its convenience and sensitivity. As little as 1 nmol of conjugated dienes could be measured. This method allows measurement of peroxide levels in our samples not detectable by direct measurements of lipid peroxides.¹⁵ TBARS were measured by the standard procedure as described by Morel and Chisolm.¹² Additionally, oxidation was confirmed by measuring the linoleic acid degradation in gas-liquid chromatography (GLC).²⁰ Lipid peroxides in the oil with high-oxidized lipid content were determined with a color reaction using methylene blue derivative¹⁵; however, the method was not sufficiently sensitive to detect peroxides in low- or medium-oxidized lipid-containing oils or any of the chylomicron preparations.

Study Protocol

Informed consent was obtained from the study subjects, and the study was approved by the Human Studies Committee of the University of California, San Francisco. Lipid peroxidation in the postprandial serum chylomicron fraction was determined by measuring conjugated dienes 4 hours after administration of the test meal. All six subjects were administered all three oils (the control corn oil, medium-oxidized corn oil, and highly oxidized corn oil). The time interval between administration of each test oil was at least 14 days, and the order of test meal administration was randomly assigned.

After a 12-hour fast (time 0), blood was drawn from each subject for measurement of baseline serum triglycerides and cholesterol. Subjects then were given corn oil (1 g/kg body weight), which was accompanied by 100 g carbohydrate (three to four slices of white bread). The subjects were not permitted to consume any food for the duration of the test period. Water was allowed ad libitum. The subjects tolerated the test meal well; no one had diarrhea or other symptoms of malabsorption. At 4 hours, 50-mL blood samples were obtained and conjugated dienes were measured in the chylomicron fraction ($S_r > 1000$) to determine the oxidized lipid levels. The results were expressed as nanomoles of conjugated dienes per micromole of triglyceride in the chylomicron fraction. For determining the postprandial clearance of conjugated dienes in serum, four subjects were administered the control oil and highly oxidized oil, and blood samples were obtained every 2 hours over an 8-hour period for oxidized lipid measurements. The results were expressed as conjugated dienes (in nanomoles) in the chylomicron fraction per micromole of triglyceride.

Chylomicron Isolation

Chylomicron fraction ($S_r > 1000$) in the postprandial serum was isolated as described previously for lymph.²¹ Briefly, serum samples were overlaid with saline and centrifuged in a swinging bucket rotor (Beckman, SW 40) for 60 minutes at 25 000 rpm at 8°C. The floating chylomicron fraction was removed, layered under saline, recentrifuged, and analyzed immediately for triglycerides, fatty acid composition, and conjugated diene content. When examined by gel electrophoresis,¹⁴ the chylomicron fraction contained a small amount of apolipoprotein B-100, indicating some contamination by large VLDL particles (data not shown).

Oxidation of Chylomicrons by CuSO₄

The susceptibility of chylomicrons to metal oxidation was determined by monitoring the production of conjugated dienes as described by Esterbauer et al.²² Freshly isolated chylomicrons (0.5 mg triglyceride/mL incubation mixture) were incubated with CuSO₄ (final concentration, 0.01 mmol/L) and 0.15 mol/L NaCl containing 0.02 mol/L phosphate buffer, pH 7.2, at 37°C for 8 hours in a total volume of 10 mL. Before incubation, the buffer was purged with oxygen for 30 minutes. Conjugated dienes were measured every 60 minutes in a 1-mL aliquot as described above. Dienes present before addition of copper were subtracted, and the baseline was adjusted to zero. Oxidation rate was expressed as lag time (during which absorption increased minimally) and was determined from an intercept of lines drawn through the linear portions of the lag and propagation phase. During the propagation phase, absorption increased rapidly, indicating the formation of lipid peroxides.²² The lines representing the propagation phase for each subject were generated by a computer (CA CRICKET GRAPH III) using 5-, 6-, 7-, and 8-hour time points (linear portion of the graph). The data were presented as an average lag time for each diet calculated from the results of four subjects.

Analytical Methods

Total cholesterol, HDL cholesterol, and triglycerides in serum were determined by enzymatic assays as described.²¹

Percent Fatty Acid Concentration in Serum Chylomicron Fractions After Ingestion of Oil Containing Low, Medium, or High Quantities of Oxidized Lipid

Fatty Acid	Low	Medium	High
16:0	14.66±0.64	13.74±0.27	17.79±1.79
18:0	2.70±0.45	3.13±0.14	6.05±0.95
18:1	26.38±0.48	28.82±0.44	33.31±4.69
18:2	56.27±0.35*	54.31±0.03	41.45±1.45

Results are mean±SEM and are derived from three preparations.

* $P < .005$ and $P < .001$ when chylomicrons isolated after ingestion of low-peroxide oil are compared with chylomicrons isolated after ingestion of medium- and high-peroxide oil, respectively.

Vitamin E concentration in oils was measured by high-performance liquid chromatography.²³ Fatty acid composition of oils and chylomicrons was determined by GLC (Hewlett Packard 5890).²⁰ Fatty acids were transmethyated with boron trifluoride methanol and were analyzed by GLC equipped with a hydrogen flame ionization detector and an SP-2330 fused silica capillary column (Supelco). The retention time and area of each peak were determined by an HP-3390 integrator. A mixture of fatty acids supplied by Matreya, Inc, was used as standard.

Data are presented as mean±SEM. The mean differences in lipid and oxidation measurements between the control and experimental groups were assessed with Student's *t* test or paired *t* test. Significance was expressed as $P < .05$.

Results

Fatty Acid Composition of the Chylomicron Fraction

The Table shows the fatty acid composition of chylomicrons obtained 4 hours after ingestion of the various diets. The percentage of linoleic acid in the chylomicron fraction after ingestion of the control oil was 56.27 ± 0.35 ; after ingestion of highly oxidized oil, it was $41.45 \pm 1.45\%$ ($P < .001$). Thus, the linoleic acid content in postprandial chylomicrons decreased with decreasing linoleic acid concentration in the dietary oils.

Incorporation of Dietary Oxidized Lipids Into Postprandial Chylomicron Fractions

It has been established that after the first meal of the day the peak triglyceride response is usually 3 to 5 hours after the meal.^{24,25} When subjects were fed corn oil containing increasing amounts of lipid peroxides, TBARS in postprandial serum chylomicrons isolated at 4 hours could be detected only in chylomicrons isolated after ingestion of an oil that contained high amounts of lipid peroxides. These high-peroxide chylomicrons contained 0.140 ± 0.03 nmol TBARS/ μ mol triglyceride. On the other hand, conjugated dienes containing polyunsaturated fatty acid hydroperoxides and hydroxy fatty acids accumulate in much greater amounts (60% to 70% of the degraded polyunsaturated fatty acid) and are the major lipid peroxidation products.^{26,27}

When we measured conjugated dienes, we found that subjects fed corn oil with increasing amounts of lipid peroxides contained significantly increasing amounts of conjugated dienes in their serum chylomicron fraction (Fig 1). When subjects were fed control oil containing low quantities of oxidized lipid, the levels of conjugated dienes in the chylomicron fraction were low (9.67 ± 0.92




Fig 1. Bar graph shows lipid oxidation in the chylomicron fraction measured as conjugated dienes. Six subjects were administered corn oil containing increasing lipid peroxides. Control, medium-oxidized, and highly oxidized oil contained 6.5 to 10, 30 to 50, and 80 to 120 nmol conjugated dienes/mg oil, respectively. Serum samples were obtained at 4 hours, and the chylomicron fraction was isolated. All data are expressed as nanomoles of conjugated dienes per micromole triglyceride and represent mean±SEM. * $P < .005$ when control group fed control oil was compared with the group fed either medium-oxidized oil or highly oxidized oil.

nmol/ μ mol triglyceride). When fed medium-oxidized oil with intermediate quantities of peroxides, the subjects had moderately increased degrees of oxidation of chylomicrons (21.86 ± 2.03 nmol/ μ mol triglyceride). However, when fed the highly oxidized oil, the subjects' conjugated diene content in chylomicrons was increased 4.7-fold to 46.5 ± 5.63 nmol/ μ mol triglyceride. Thus, the content of oxidized lipid in the diet determined the levels of oxidized lipid in postprandial chylomicrons. Moreover, approximately 4% to 5% of the triglyceride is oxidized on a molar basis (46.5 ± 5.63 nmol peroxide/1130 nmol triglyceride), further indicating that the oxidized lipid content in chylomicrons is similar to that of the ingested oil.

Fig 2 shows that there was also a significant increase in conjugated dienes in the circulation over an 8-hour time period after ingestion of the oxidized oil compared with the control oil. The levels of conjugated dienes at 2, 4, and 6 hours were relatively similar, with a decrease seen at 8 hours in the high-peroxide-diet group. Thus, oxidized lipids in the diet result in an increase in oxidized lipids in the circulation for extended postprandial periods, and there is a continuous presence of




Fig 2. Graph shows profiles of postprandial total conjugated dienes in the chylomicron fraction in four subjects who consumed control (6.5 nmol conjugated dienes/mg oil) and highly oxidized oil (108 nmol conjugated dienes/mg oil). Values are expressed as mean±SEM. $P < .01$ for all time points studied.




FIG 3. Graph shows the generation of conjugated dienes during copper oxidation of chylomicrons isolated after four subjects were fed oil containing either low- (6.5 nmol conjugated dienes/mg oil) or high- (111 nmol conjugated dienes/mg oil) lipid peroxides. Chylomicrons (0.5 mg triglyceride) were incubated in phosphate-buffered saline, pH 7.2, containing 0.01 mol/L CuSO_4 at 37°C. Reaction mixture (1 mL) was removed every 60 minutes and extracted, and the conjugated dienes were measured. The values at time 0 were subtracted from all measurements. Values are expressed as mean \pm SEM.

possibly atherogenic oxidized lipids in the blood after a meal containing lipid peroxides.

Oxidation of Chylomicrons by CuSO_4

In all four individuals studied, the lag time before onset of oxidation was shorter for chylomicrons isolated after ingestion of an oil containing high levels of lipid peroxides than for chylomicrons isolated after ingestion of the control oil containing low lipid peroxides (3.24 ± 0.51 versus 4.30 ± 0.40 hours) ($P < .05$, paired t test) (Fig 3). There was no significant increase in conjugated dienes when parallel samples were incubated without CuSO_4 . Thus, oxidized dietary oil results in chylomicrons that not only contain increased lipid peroxides but also are more susceptible to further oxidation. This increased susceptibility to oxidation cannot be explained by a higher linoleic acid content (Table).

Discussion

The present study was undertaken to determine whether in humans oxidized lipids in the diet are also absorbed and incorporated into postprandial serum lipoproteins. When we fed human subjects a test meal containing low amounts of oxidized lipid (6.5 to 10.0 nmol conjugated dienes/mg oil), we found that the serum chylomicrons contained very low amounts of oxidized lipid. In contrast, when the same subjects were fed a test meal containing large quantities of oxidized lipids (80 to 120 nmol conjugated dienes/mg oil), the levels of oxidized lipids in their serum chylomicrons were increased approximately fivefold (Figs 1 and 2). These results, along with the studies of Naruszewicz et al,¹⁰ indicate that in humans oxidized lipids in the diet are absorbed by the small intestine and are transported in chylomicrons to the circulation, where they can contribute to the total body pool of oxidized lipid.

The presence of oxidized lipids in chylomicrons may have adverse consequences. Because of their large size, chylomicrons are not atherogenic, but chylomicron remnants have been implicated repeatedly in the atherogenic

process. Zilversmit²⁸ has proposed that intestinally derived lipoproteins can contribute to atherosclerosis, and experimental animal studies have shown that remnants from triglyceride-rich lipoproteins deposit lipid in the arterial wall.²⁹ Moreover, in humans, familial dysbetalipoproteinemia, which is characterized by an increase in remnant particles of both hepatic and intestinal origin, is associated with premature atherosclerosis.³⁰ It was also shown that patients with coronary artery disease have a delayed clearance of postprandial lipoproteins from their serum.³¹ *In vitro* studies demonstrated that chylomicron remnants are capable of promoting cholesterol esterification and cholesterol ester storage in macrophages³² and smooth muscle cells³³ and that in macrophages the intestinally derived remnants were much more potent than hepatic-derived remnants in inducing foam cell formation.³⁴ Moreover, oxidation of chylomicron remnants has been shown to enhance the delivery of cholesterol to these cells.^{10,35,36}

Additionally, we now demonstrate that partially oxidized chylomicrons are more susceptible to further oxidation. This increased susceptibility could be due to decreases in antioxidant content secondary to the presence of oxidized lipids. Alternatively, the presence of oxidized lipid per se may allow accelerated further oxidation, as suggested by several investigators.³⁷ Of course, it must be recognized that increased susceptibility to copper oxidation may not necessarily correlate with increased potential for oxidation *in vivo*. The mechanism by which chylomicrons and other lipoproteins are oxidized *in vivo* is not clear.

A further adverse consequence of the absorption of oxidized lipids is that the levels of oxidized lipids in endogenous lipoproteins may be increased. In our studies in rats, we observed that the levels of oxidized lipid in VLDL+LDL directly correlated with the quantity of oxidized lipid in the diet.¹⁵ Additionally, studies in humans by Reaven et al³⁸ showed that the fatty acid composition of the diet determines the fatty acid composition of LDL. Whether oxidized lipids in the diet increase the levels of oxidized lipids in the endogenous lipoproteins of humans remains to be determined.

A limited amount of data in the literature suggest that oxidized lipids in the diet may be not only toxic³⁹ but also atherogenic.⁴⁰ Early studies by Kritchevsky et al⁴¹ showed that diets containing polyunsaturated fatty acids heated for 20 minutes at 215°C are more atherogenic in rabbits than diets containing unheated oil. Moreover, similar experiments with olive oil, which contains monounsaturated fatty acids that are not oxidized, did not demonstrate an increase in atherosclerosis with heated oil. These results suggest that oxidized lipids induced by heating of polyunsaturated fatty acids are atherogenic. Unfortunately, these studies were carried out before the potential importance of oxidized lipids was recognized, so the oxidative state of the oils, serum lipoproteins, and lipid in the atherosclerotic lesions were not measured. Some studies also showed that adding fish oils to a cholesterol-rich diet enhances atherosclerosis in rabbits.⁴² Since fish oils are highly susceptible to oxidation, oxidized lipids could account for this observation, but again no measurements to determine the state of oxidation were made. Finally, Blankenhorn et al,⁴³ using dietary recall, reported that development of new coronary artery lesions was increased in patients with a high

consumption of not only saturated but also polyunsaturated fat. It is possible that dietary oxidized lipid contributed to the harmful effects of polyunsaturated fats in this study, but again no data directly addressing the issue of fatty acid oxidation were provided.

The typical American diet contains substantial quantities of polyunsaturated fatty acids that have been subjected to various degrees of processing and heat treatment, particularly deep fat frying, which is known to lead to oxidation. Frankel et al⁴⁴ examined the relative percent of oxidation products in frying oils used in fast food and found that the majority of these foods contain oxidized lipids. For example, french-fried potatoes contain up to 8.2% of oxidized material. Alexander⁴⁵ showed that oils used in many restaurants are kept at 180°C for extended time periods, which results in the appearance of oxidized lipids. Thompson et al⁴⁶ also performed oxidized lipid measurements and showed that oils used for deep-frying in restaurants, institutions, and armed services contained large quantities of oxidized lipids. Finally, Yagi et al⁴⁷ measured peroxides in 30 kinds of foods and found that the peroxide content could be as high as 600 nmoles/g food. Thus, there is abundant evidence that oxidized lipids are frequently present in the typical American diet in both commercial and domestic settings. Furthermore, the quantity of oxidized lipids reported in these studies is higher than the levels of oxidized lipids used in our experiments.

In summary, our studies demonstrate that in humans dietary oxidized lipids are absorbed by the small intestine, incorporated into chylomicrons, and appear in the bloodstream, where they contribute to the total body pool of oxidized lipid. Thus, dietary oxidized lipids could potentially contribute to the development of atherosclerosis.

Acknowledgments

This work was supported by the American Diabetes Association, the Medical Research Service, Veterans Affairs, and the Pacific Vascular Research Foundation. The excellent technical assistance of Agnes Frank and Arthur Moser is appreciated. Dr Marvin D. Siperstein is gratefully acknowledged for helpful discussions.

References

- Steinberg D, Parthasarathy S, Carew T, Khoo JC, Witztum JL. Beyond cholesterol: modification of low density lipoprotein that increases its atherogenicity. *N Engl J Med*. 1989;320:915-924.
- Steinberg D, Witztum JL. Lipoproteins and atherogenesis: current concepts. *JAMA*. 1990;264:3047-3052.
- Witztum JL, Steinberg D. Role of oxidized low density lipoprotein in atherogenesis. *J Clin Invest*. 1991;88:1785-1792.
- Morel DW, DiCorleto PE, Chisolm GM. Endothelial and smooth muscle cells alter low density lipoprotein in vitro by free radical oxidation. *Arteriosclerosis*. 1984;4:357-364.
- Cushing SD, Berliner JA, Valente AJ, Navab M, Parhami R, Gerrity R, Schwartz CJ, Fogelman AM. Minimally modified low density lipoprotein induces monocyte chemotactic protein 1 in human endothelial cells and smooth muscle cells. *Proc Natl Acad Sci U S A*. 1990;87:5134-5138.
- Liao F, Berliner JA, Mehrabian M, Navab M, Demer LL, Lusis AJ, Fogelman AM. Minimally modified low density lipoprotein is biologically active in vivo. *J Clin Invest*. 1991;87:2253-2257.
- Fox PL, Chisolm GM, DiCorleto PE. Lipoprotein-mediated inhibition of endothelial cell production of platelet-derived growth factor-like protein depends on free radical lipid peroxidation. *J Biol Chem*. 1987;262:6046-6054.
- Nishigaki I, Hagihara M, Tsunekawa HT, Maseki M, Yagi K. Lipid peroxide levels of serum lipoprotein fractions of diabetic patients. *Biochem Med Metab Biol*. 1981;25:373-378.
- Sato Y, Hotta N, Sakamoto N, Matsuoka S, Ohishi N, Yagi K. Lipid peroxide level in plasma of diabetic patients. *Biochem Med Metab Biol*. 1979;21:104-107.
- Naruszewicz M, Wozny E, Mirkiewicz G, Nowicka G, Szostak WB. The effect of thermally oxidized soya bean oil on metabolism of chylomicrons: increased uptake and degradation of oxidized chylomicrons in cultured mouse macrophages. *Atherosclerosis*. 1987;66:45-53.
- Higuchi Y. Lipid peroxides and α -tocopherol in rat streptozotocin-induced diabetes mellitus. *Acta Med Okayama*. 1982;3:165-175.
- Morel DW, Chisolm GM. Antioxidant treatment of diabetic rats inhibits lipoprotein oxidation and cytotoxicity. *J Lipid Res*. 1989;30:1827-1834.
- Aw TY, Williams MW, Gray L. Absorption and lymphatic transport of peroxidized lipids by rat small intestine in vivo: role of mucosal GSH. *Am J Physiol*. 1992;262:G99-G106.
- Staprans I, Pan XM, Miller M, Rapp JH. The effect of oxidation on the chylomicron metabolism in rats. *Am J Physiol*. 1993;264:G561-G568.
- Staprans I, Rapp JH, Pan XM, Feingold KR. The effect of oxidized lipid in the diet on serum lipoprotein peroxides in control and diabetic rats. *J Clin Invest*. 1993;92:638-643.
- Corongiu FP, Banni S, Dessi MA. Conjugated dienes detected in tissue lipid extracts by second derivative spectrophotometry. *Free Radic Biol Med*. 1989;7:183-186.
- Dole VP. A relation between non-esterified fatty acids in plasma and the metabolism of glucose. *J Clin Invest*. 1956;35:150-154.
- Boeynaems JM, Brash AR, Oates JA, Hubbard WC. Preparation and assay of monohydroxy-eicosatetraenoic acids. *Anal Biochem*. 1979;104:259-267.
- Fletcher MJ. A colorimetric method for estimating serum triglycerides. *Clin Chim Acta*. 1968;22:393-397.
- Wisneski JA, Gertz EW, Neese RA, Mayr M. Myocardial metabolism of free fatty acids: studies with ¹⁴C labeled substrates in humans. *J Clin Invest*. 1987;79:359-366.
- Staprans I, Rapp JH, Pan XM, Ong DL, Feingold KR. Testosterone regulates metabolism of plasma chylomicrons in rats. *Arteriosclerosis*. 1990;10:591-596.
- Esterbauer H, Striegl G, Puhl H, Rothenegger M. Continuous monitoring of in vitro oxidation of human low density lipoprotein. *Free Radic Res*. 1989;6:67-75.
- Ha YL, Csallany SC. Separation of α -tocopherol and its oxidation products by high performance liquid chromatography. *Lipids*. 1988;23:359-361.
- Ruotolo GH, Zhang V, Bensianov NA, Le NA. Protocol for the study of the metabolism of retinyl esters in plasma lipoproteins during postprandial lipemia. *J Lipid Res*. 1992;33:1541-1549.
- Schneeman BO, Kotite L, Todd KM, Havel RJ. Relationship between the responses of triglyceride-rich lipoproteins in blood plasma containing apolipoprotein B-48 and B-100 to a fat-containing meal in normolipidemic humans. *Proc Natl Acad Sci U S A*. 1993;90:2069-2073.
- Lenz ML, Hughes H, Mitchell JR, Via DP, Guyton JR, Taylor AA, Gotto AM, Smith CV. Lipid hydroperoxy and hydroxy derivatives in copper-catalyzed oxidation of low density lipoprotein. *J Lipid Res*. 1990;31:1043-1050.
- Wang T, Yu W, Powell WS. Formation of monohydroxy derivatives of arachidonic acid, linoleic acid, and oleic acid during oxidation of low density lipoprotein by copper ions and endothelial cells. *J Lipid Res*. 1992;33:525-537.
- Zilversmit DB. Atherosclerosis: a postprandial phenomenon. *Circulation*. 1979;60:473-485.
- Stender S, Zilversmit DB. Comparison of cholesteryl ester transfer from chylomicrons and other plasma lipoproteins to aorta intima-media of cholesterol-fed rabbit. *Arteriosclerosis*. 1982;2:493-499.
- Morganroth J, Levy RI, Fredrickson DS. The biochemical, clinical, and genetic features of type III hyperlipoproteinemia. *Ann Intern Med*. 1975;82:158-174.
- Groot PH, van Stiphout WA, Krauss XH, Jansen H, van Tol A, van Ramshorst E, Chin-On S, Hofman A, Creswell SR, Havekes L. Postprandial lipoprotein metabolism in normolipidemic men with and without coronary artery disease. *Arterioscler Thromb*. 1991;11:653-662.
- Van Lenten BJ, Fogelman AM, Jackson RL, Shapiro S, Haberland ME, Edwards PA. Receptor-mediated uptake of remnant lipoproteins by cholesterol-loaded human monocyte-macrophages. *J Biol Chem*. 1985;260:8783-8788.
- Floren CH, Albers JJ, Bierman EL. Uptake of chylomicron remnants causes cholesterol accumulation in cultured human

- arterial smooth muscle cells. *Biochim Biophys Acta*. 1981;663:336-349.
34. Fainaru M, Mahley RW, Hamilton RL, Innerarity TL. Structural and metabolic heterogeneity of β -very low density lipoproteins from cholesterol-fed dogs and from humans with type III hyperlipoproteinemia. *J Lipid Res*. 1982;23:702-714.
35. Horrigan S, Campbell JL, Campbell GR. Oxidation of β -very low density lipoprotein by endothelial cells enhances its metabolism by smooth muscle cells in culture. *Arterioscler Thromb*. 1991;11:279-289.
36. Haratz D, Stein O, Shwartz R, Berry EM, Stein Y. Preferential metabolism by macrophages of conditioned rabbit hypercholesterolemic remnant lipoproteins. *Biochim Biophys Acta*. 1988;959:127-133.
37. Thomas CE, Jackson RL. Lipid hydroperoxide involvement in copper-dependent and independent oxidation of low density lipoproteins. *J Pharmacol Exp Ther*. 1991;256:1182-1188.
38. Reaven P, Parthasarathy S, Grasse BS, Miller E, Almazan F, Mattson FH, Khoo JC, Steinberg D, Witztum JL. Feasibility of using an oleate-rich diet to reduce the susceptibility of low-density lipoprotein to oxidative modification in humans. *Am J Clin Nutr*. 1991;54:701-706.
39. Andrews JS, Griffith WH, Mead JF, Stein RA. Toxicity of air-oxidized soybean oil. *J Nutr*. 1960;70:199-210.
40. Kubow S. Routes of formation and toxic consequences of lipid oxidation products in foods. *Free Radic Biol Med*. 1992;12:63-81.
41. Kritchevsky D, Moyer AW, Tesar WC, McCandless RFJ, Logan JB, Brown RA, Englert ME. Cholesterol vehicle in experimental atherosclerosis, II: influence of unsaturation. *J Atheroscler Res*. 1967;7:647-651.
42. Theiry J, Seidel D. Fish oil feeding results in an enhancement of cholesterol-induced atherosclerosis in animals. *Atherosclerosis*. 1987;63:53-60.
43. Blankenhorn DH, Johanson RL, Mack WJ, El Zein HA, Vailas RD. The influence of diet on the appearance of new lesions in human coronary arteries. *JAMA*. 1990;263:1646-1652.
44. Frankel EN, Smith LM, Hamblin CL, Creveling RK, Clifford AJ. Occurrence of cyclic fatty acid monomers in frying oils used for fast foods. *J Am Oil Chem Soc*. 1984;61:87-90.
45. Alexander JC. Heated and oxidized fats. *Prog Clin Biol Res*. 1986;222:185-209.
46. Thompson JA, Paulose MM, Reddy BR, et al. A limited survey of fats and oils commercially used for deep fat frying. *Food Technol*. 1967;21:405-407.
47. Yagi K, Kiuchi K, Saito Y, Miike A, Kayahara N, Tanano T, Ohishi N. Use of a new methylene blue derivative for determination of lipid peroxides in foods. *Biochem Int*. 1986;12:367-371.

Arteriosclerosis, Thrombosis, and Vascular Biology



JOURNAL OF THE AMERICAN HEART ASSOCIATION

Oxidized lipids in the diet are a source of oxidized lipid in chylomicrons of human serum.

I Staprāns, J H Rapp, X M Pan, K Y Kim and K R Feingold

Arterioscler Thromb Vasc Biol. 1994;14:1900-1905

doi: 10.1161/01.ATV.14.12.1900

Arteriosclerosis, Thrombosis, and Vascular Biology is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 1994 American Heart Association, Inc. All rights reserved.

Print ISSN: 1079-5642. Online ISSN: 1524-4636

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://atvb.ahajournals.org/content/14/12/1900>

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Arteriosclerosis, Thrombosis, and Vascular Biology* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

Reprints: Information about reprints can be found online at:

<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Arteriosclerosis, Thrombosis, and Vascular Biology* is online at:

<http://atvb.ahajournals.org/subscriptions/>