


Potential antigenic cross-reactivity between SARS-CoV-2 and human tissue with a possible link to an increase in autoimmune diseases

Aristo Vojdani, Datis Kharrazianb



PII: S1521-6616(20)30425-3

DOI: <https://doi.org/10.1016/j.clim.2020.108480>

Reference: YCLIM 108480

To appear in: *Clinical Immunology*

Received date: 21 May 2020

Accepted date: 21 May 2020

Please cite this article as: A. Vojdani and D. Kharrazianb, Potential antigenic cross-reactivity between SARS-CoV-2 and human tissue with a possible link to an increase in autoimmune diseases, *Clinical Immunology* (2020), <https://doi.org/10.1016/j.clim.2020.108480>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Letter to the Editor

Potential Antigenic Cross-Reactivity between SARS-CoV-2 and Human Tissue with a Possible Link to an Increase in Autoimmune Diseases

Aristo Vojdani^{a,b,*} drari@msn.com, Datis Kharrazian^c

^aImmunosciences Lab, Inc. 822 S. Robertson Blvd, Ste. 312, Los Angeles, CA 90035 USA

^bDepartment of Preventive Medicine, Loma Linda University, Loma Linda, CA 92350 USA

^cHarvard Medical School, 25 Shattuck St, Boston, MA 02115, USA

*Corresponding author at: Immunosciences Lab, Inc. 822 S. Robertson Blvd, Ste. 312, Los Angeles, CA 90035 USA.

Since the outbreak of COVID-19 caused by SARS-CoV-2, we tested 5 different blood specimens that were confirmed positive for SARS-CoV-2 IgG and IgM antibodies [1]. The measurements were for anti-nuclear antibody (ANA), anti-extractable nuclear antigen (ENA), anti-double-stranded DNA (dsDNA), actin antibody, mitochondrial antibody, rheumatoid factor (RF), and C1q immune complexes. We were surprised to find out that 3 of the 5 specimens had significant elevations in ANA, ENA, actin and mitochondrial antibodies, but not against dsDNA or RF. This prompted us to investigate patterns of cross-reactivity between SARS-CoV-2 and autoimmune target proteins.

Vaccine-induced autoimmunity from autoimmune cross-reactivity is associated with narcolepsy, Guillain-Barré syndrome, multiple sclerosis, demyelinating neuropathies, systemic lupus erythematosus, and postural orthostatic tachycardia syndrome in susceptible subgroups as reported by Segal and Shoenfeld [2]. Due to the significant red flags for the potential cross-reactive interactions with the current COVID-19 pandemic, we studied the relationships between spike and nuclear proteins of SARS-CoV-2 and autoimmune target proteins.

Commercially available mouse monoclonal antibody made against recombinant SARS coronavirus spike protein and rabbit monoclonal antibody made against SARS coronavirus nucleoprotein were applied at optimal dilution to the SARS-CoV-2 proteins and to 50 different tissue antigens using enzyme-linked immunosorbent assay (ELISA). Recombinant SARS-CoV-2 spike protein S1 and recombinant SARS-CoV-2 nucleocapsid protein were purchased from RayBiotech. ELISA wells were coated with nuclear antigens, dsDNA, F-actin, and mitochondria (M2) antigen purchased from different companies. An additional 45 tissue antigens used in this study have been previously described [9]. Each SARS-CoV-2 antibody was applied to quadruplicate wells. After the completion of all ELISA steps, the developed color was measured at 405 nm.

Looking at the reaction between SARS-CoV-2 spike protein antibody and tissue proteins (Figure 1A), we found that the strongest reactions were with transglutaminase 3 (tTG3), transglutaminase 2 (tTG2), ENA, myelin basic protein (MBP), mitochondria, nuclear antigen (NA), α -myosin, thyroid peroxidase (TPC), collagen, claudin 5+6, and S100B. The reaction of this antibody was not as strong with several other antigens (Figure 1A).

The nucleoprotein antibody showed some overlap in immune cross-reactivity with anti-spike protein antibody. As shown in Figure 1B, nucleoprotein antibody reacted strongly with mitochondria, tTG6, NA, TPC, ENA, TG, actin, and MBP. Similar to spike protein, the nucleoprotein antibody reaction was not as strong with several other antigens as shown in Figure 1A and B.

As the number of SARS-CoV-2 infections increase from day to day, scientists are learning that the damage caused by this virus can extend well beyond the lungs, where infection can lead to pneumonia and the often fatal condition called acute respiratory distress syndrome [3]. The virus can in fact affect the body from head to toe, including the nervous [4], cardiovascular [5], immune [6], and digestive systems [7].

(INSERT FIGURE 1 HERE OR NEAR HERE)

Is it possible that some of the extensive organ, tissue, and cellular damage done by SARS-CoV-2 is due to viral antigenic mimicry with human tissue?

If the answer is yes, then we may face an increase in the rates of autoimmune disease in the future, because any factor that causes chronic inflammation in the body can potentially induce autoimmune disease.

Because SARS-CoV-2 attacks the respiratory system first, in a very interesting letter [8] Kanduc and Shoenfeld suggested that because the SARS-CoV-2 spike glycoprotein and lung surfactant proteins shared 13 out of 24 pentapeptides, the immune response following infection with SARS-CoV-2 may lead to cross-reactions with pulmonary surfactant proteins, followed by SARS-CoV-2-associated lung disease [8]. Based on their findings, they warned against the use of the entire SARS-CoV-2 antigens in the vaccines and cautioned that perhaps the use of only unique peptides would be the most effective way to fight the SARS-CoV-2 infection. Very similar suggestions were made by Razim et al., in designing a vaccine against *Clostridium difficile* [9]. Two sequences, peptide 9 and peptide 10, of *C. difficile* were recognized not only by the sera of patients with *C. difficile* infections but also by the sera of patients with autoimmune disease. Razim et al. concluded that before considering a protein as a vaccine antigen, special care should be taken in analyzing the sequence of tissue cross-reactive epitopes in order to avoid possible future side effects [9].

We agree with Razim et al., and we feel that our own findings that 21 out of 50 tissue antigens had moderate to strong reactions with the SARS-CoV-2 antibodies are a sufficiently strong indication of cross-reaction between SARS-CoV-2 proteins and a variety of tissue antigens beyond

just pulmonary tissue, which could lead to autoimmunity against connective tissue and the cardiovascular, gastrointestinal, and nervous systems.

We live in critical times when the world may be veering towards the very real possibility of requiring immunity certification “passports” earned by prior infection with SARS-CoV-2 or vaccination before being allowed to travel, or perhaps even to work [10].

At the moment, scientists are frantically trying to develop either a definitive cure, neutralizing antibodies, or a vaccine to protect us from contracting the disease in the first place, and they want it right now. We must consider that finding a vaccine for a disease may normally take years. There are reasons for all the cautions involved in developing a vaccine, not the least of which are unwanted side-effects. In light of the information discussed above about the cross-reactivity of the SARS-CoV-2 proteins with human tissues and the possibility of either inducing autoimmunity, exacerbating already unhealthy conditions, or otherwise resulting in unforeseen consequences, it would only be prudent to do more extensive research regarding the autoimmune-inducing capacity of the SARS-CoV-2 antigens. The promotion and implementation of such an aggressive “immune passport” program worldwide in the absence of thorough and meticulous safety studies may exact a monumental cost on humanity in the form of another epidemic, this time a rising tide of increased autoimmune diseases and the years of suffering that come with them.

Declaration of Interests

None

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

- [1] J.D. Whitman, J. Hiatt, C.T. Mowery, B.R. Shy, R. Yu, T.N. Yamamoto, et al. Test performance evaluation of SARS-CoV-2 serological assay. *MedRxiv*. 2020;preprint. doi: 10.1101/2020.04.25.20074856
- [2] Y. Segal, Y. Shoenfeld. Vaccine-induced autoimmunity: the role of molecular mimicry and immune crossreaction. *Cell Mol Immunol*. 2018;15(6):586- 594. doi:10.1038/cmi.2017.151
- [3] P. Zhou, X.-L. Yang, X.-G. Wang, B. Hu, L. Zhang, W. Zhang, et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature*. 2020;579(7798):270-273. doi:10.1038/s41586-020-2012-7.
- [4] A. Nath. Neurologic complications of coronavirus infections. *Neurology*. 2020;94(19):809-810. doi: 10.1212/WNL.0000000000009455
- [5] Y.-Y. Zheng, Y.-T. Ma, J.-Y. Zhang, X. Xie. COVID-19 and the cardiovascular system. *Nat Rev Cardiol*. 2020;17(5):259-260. doi: 10.1038/s41569-020-0360-5.
- [6] Y. Shi, Y. Wang, C. Shao, J. Huang, J. Gan, X. Huang, et al. COVID-19 infection: the perspectives on immune responses. *Cell Death & Differentiation*. 2020;27:1451-1454. doi: 10.1038/s41418-020-0530-3
- [7] L. Lin, X. Jiang, Z. Zhang, S. Huang, Z. Zhang, Z. Fang, et al. Gastrointestinal symptoms of 95 cases with SARS-CoV-2 infection. *Gut*. 2020;69(6):997-1001. doi: 10.1136/gutjnl-2020-321013.
- [8] D. Kanduc, Y. Shoenfeld. On the molecular determinants of the SARS-CoV-2 attack. *Clin Immunol*. 2020; 215:108426. doi: 10.1016/j.clim.2020.108426.
- [9] A. Razim, K. Pacyga, M. Aptekorz, G. Martirosian, A. Szuba, E. Pawlak-Adamska, et al. Epitopes identified in GADPH from *Clostridium difficile* recognized as common antigens

with potential autoimmunizing properties. *Scientific Reports*. 2018;8(1):13946. doi:

10.1038/s41598-018-32193-9.

- [10] M.A. Hall, D.M. Studdert. Privileges and immunity certification during the COVID-19 pandemic. *JAMA*. 2020 May 6. doi: 10.1001/jama.2020.7712.

Aristo Vojdani,^{a, b} Datis Kharrazian^{b, c}

^a*Immunosciences Lab, Inc. 822 S. Robertson Blvd, Ste. 312, Los Angeles, CA 90035 USA*

^b*Department of Preventive Medicine, Loma Linda University, Loma Linda, CA 92350 USA*

^c*Harvard Medical School, 25 Shattuck St, Boston, MA 02115, USA*

E-mail addresses: arari@msn.com (A. Vojdani)

ORCID identifier

Aristo Vojdani 0000-0003-2925-9334

Datis Kharrazian 0000-0001-7985-476X

- Corresponding author at: Immunosciences Lab, Inc. 822 S. Robertson Blvd, Ste. 312, Los Angeles, CA 90035 USA

FIGURE CAPTIONS

Fig. 1. (A) Reaction of anti-SARS-CoV-2 spike protein monoclonal antibody with human tissue antigens. (B) Reaction of anti-SARS-CoV-2 nucleoprotein monoclonal antibody with human tissue antigens.





Figure 1