

The Syrian hamster polymyopathy is a hereditary disease, transmitted by an autosomal recessive gene, involving the heart and the entire musculature. The chronology of the pathologic events in the myocardium and skeletal muscle has been investigated in UM-X7.1 myopathic hamsters aged 0–250 days. A phasic pattern in the progression of the disease process was evident. Microscopic necrotic changes in the heart were visible prior to or at 50 days of age with increasing severity until 100 days of age and subsidence thereafter. More than 50% of the animals died before 250 days of age with signs of cardiac failure. The intensity and extent of myocardial calcific changes together with scar formation were determinant factors in curtailing the survival of animals. Changes in serum creatine kinase (CK) activity followed a phasic pattern similar to the progression of the myopathic disease. Because of the disparity of disease manifestations between the different myopathic hamster lines, it is essential to consider the time course of the heart and skeletal muscle microscopic changes when evaluating the severity of the hamster polymyopathy.

MUSCLE & NERVE 5:20–25 1982

HEREDITARY POLYMYOPATHY AND CARDIOMYOPATHY IN THE SYRIAN HAMSTER. I. PROGRESSION OF HEART AND SKELETAL MUSCLE LESIONS IN THE UM-X7.1 LINE

G. JASMIN, MD, PhD, and L. PROSCHEK, PhD

The hereditary polymyopathy discovered in an inbred strain of Syrian hamsters has generated considerable interest in the field of genetically determined myopathies. This hamster model provides unique possibilities of studying the frequent involvement of the myocardium in human primary muscle disorders.^{5,9,12,33–35,48,52}

The polymyopathy is transmitted by an autosomal recessive gene.¹⁶ This simple mode of inheritance permits the establishment of new myopathic lines by cross-breeding homozygous diseased animals with unrelated healthy hamsters and by recovering the mutant gene in the F2 generation. The resulting lines will have a disease incidence of 100% with no significant difference in the histologic features of muscle lesions, indicating

that the genotype of this hereditary disorder is constant and reliable. However, the time course and severity of the disease appear to vary somewhat between different lines and even between siblings of the same line.¹⁵ Thus, after 28 generations of inbreeding, the BIO 14.6 animals showed a gradual decline in the severity of symptoms with modulation of signs of cardiac failure and a concomitant increase in longevity.^{14,19} This shift is most likely due to a natural selection of the healthiest breeders at weaning.

Survey of the literature shows a great deal of inconsistency in appraisals of the hamster polymyopathy through biochemical or morphologic findings.^{7,10,22,29–32,42,43,45–47,49,50} The objective of the present paper is to describe the essential pathologic features of the heart and skeletal muscles for use as guidelines in assessing the intensity and chronology of the myopathic process. The data have been cumulated over a period of 8 years, ever since the UM-X7.1 hamster line (a subline of BIO 14.6)¹⁷ was established in this laboratory. Emphasis is placed on elucidating the progression of the disease process, which we consider fundamental to the investigation of such subcellular pathologic changes as mitochondrial respiration and calcium load, which is the subject of a separate report.³⁶

From the Département de Pathologie, Faculté de Médecine, Université de Montréal, Montréal, Quebec, Canada

Acknowledgments: This study was supported by a grant from The Medical Research Council of Canada (MA-1827) and The Muscular Dystrophy Association of Canada.

Address reprint requests to Dr. Jasmin at the Département de Pathologie, Faculté de Médecine, Université de Montréal, C.P. 6128, Succursale A, Montréal, P.Q., H3C 3J7, Canada

Received for publication November 24, 1980; revised manuscript accepted for publication March 19, 1981.

0148-639X/0501/0020 \$01.25/0
© 1982 John Wiley & Sons, Inc.




Figure 1. Diagram of skeletal muscle changes and of the relative values of serum creatine kinase activity during the course of the hamster hereditary polymyopathy in the UM-X7.1 line.

MATERIALS AND METHODS

Male and female hamsters of the UM-X7.1 myopathic line were used in this study. Unrelated healthy hamsters (originally from Lakeview Farm, Newfield, NJ) served as controls. They were maintained under controlled housing conditions with free access to Purina laboratory chow and tap water. Groups of 10–20 animals of the same age were sacrificed by exsanguination at intervals of 25 days between 0 and 250 days of age. At autopsy, all animals were skinned for an overall estimation of necrotic foci readily visible by the naked eye. Heart lesions were also recorded. The myocardium and several pieces of skeletal muscle (tongue, diaphragm, obliquus externus abdominis, quadriceps, and tibialis anterior muscles) were fixed either in Lillie's buffered sublimate or in neutral formol for routine histology, using the hematoxylin-phloxinsaffron (HPS), alizarin, and von Kossa calcium stains.¹³ The degree of cardiac hypertrophy was assessed by the overall dimension of heart cavities and the thickness of myocardial wall in tissue sections. Development of heart failure was estimated by the amount of pleural and peritoneal fluids, the liver mass, and subcutaneous edema. The severity of cardiac and skeletal muscle lesions (myolysis, necrosis, and fibrosis) was assessed on an arbitrary scale of 0–3: 0, designating no lesions; 1, occasional isolated fiber degeneration; 2, moderate lesions consisting of small scattered foci of necrosis; and 3, severe involvement with large areas of fiber destruction. The means of these microscopic findings together with the time of occurrence are shown in Figures 1 and 2.

The total creatine kinase (CK) activity in serum




Figure 2. Diagram of heart muscle changes during the course of the hamster hereditary polymyopathy in the UM-X7.1 line.

was determined spectrophotometrically by a kinetic procedure; samples in appropriate dilutions were analyzed at 25°C, using the CK-combination test of Boehringer Mannheim Diagnostica (BMC Diagnostics/Biochemicals Ltd., Montreal, Canada).

RESULTS

Clinical Course of the Disease. The physical appearance and behavior of the myopathic hamster remain normal for a long period despite the presence of degenerative lesions in the skeletal and cardiac muscles. Characteristic gross symptomatology develops progressively with cardiac insufficiency and becomes prominent during heart failure. However, for some undetermined reason, muscular weakness with locomotor impairment is sometimes evident in 150-day-old animals by the end of the necrotic stage. The efficiency of muscle performance is significantly diminished during forced exercise. The development of subcutaneous edema indicates the terminal stage of progressive heart failure; hyperpnea and cyanosis are prominent and the prostrated animals die within 2 weeks.

Skeletal Muscle Changes. In newborn animals until 20 days of age, the initial necrotic changes are confined to the most active respiratory muscles. At autopsy there are scattered white streaks in the muscle. With the progression of the disease, the lesions become more confluent involving bundles of fibers. Light microscopic changes at an early stage consist of hyalinization and fragmentation of isolated myofibers with little or no cellular infiltrate. The necrotizing process extends progressively to




Figure 3. Microscopic degenerative changes in the diaphragm muscle of UM-X7.1 hamsters at late stages of the hamster myopathic disease (hematoxylin-phloxin-saffron stain). (A) Interfibrillar lipomatosis prominent between 200 and 250 days of age. Note in this longitudinal section a thin basophilic fiber with a row of nuclei. Bar = 0.5 mm. (B) Aggregates of calcium apatites sequestered by fibrotic tissue, irregularly found between 125 and 250 days of age. Bar = 0.5 mm.

the whole musculature, reaching its maximum severity in 100-day-old hamsters and receding thereafter (Fig. 1).

The more consistently affected muscles, in order of frequency, are the diaphragm, the intercostal muscles, the tongue, and the limb-girdle groups. Isolated groups of myofibers undergo myolysis in the absence of cellular reaction, but more often the degenerative changes consist of coagulation necrosis with phagocytosis. The phagocytic foci include small regenerative myofibers showing increased basophilia and internal proliferation of nuclei. After 160 days, necrosis and regeneration usually subside; the fibers are dysvolumenic and 80% have central nuclei. Interstitial fibrolipomatosis is prominent in the abdominal and diaphragm muscles (Fig. 3A) but is less prominent in the rest of the musculature. Calcific foci are seen (Fig. 3B) in groups of constantly contracting muscles (diaphragm, tongue) and are otherwise more discreet in the upper or lower limb muscles.

Heart Muscle Changes. Myocardial changes are rarely visible prior to 40 days of age (Fig. 2). Scattered lesions (Fig. 4A) are randomly distributed throughout the wall of both ventricles but invariably gain in severity and extent with progression of the disease. The earliest necrotic changes under the microscope consist of focal myolysis in the absence of cellular infiltration. The cardiocytes disintegrate either by myofibrillar dissolution, leaving sarcolemmal remnants and occasional few nuclei, or by coagulation of the sarcoplasm, which becomes highly granular and calcified. The degenerative changes subside by 150 days of age,


and calcified fragments are sequestered within scar tissue. Interestingly, the myolytic process reappears during later stages of cardiac hypertrophy. In contrast to the skeletal muscle, the pathologic changes in the heart give rise to substantial scarring. The gross signs of circulatory failure consist of liver enlargement and fluid accumulation together with an appreciable increase of heart volume. During the terminal stage, mural thrombi are frequently seen in either cavity of the dilated flabby heart (Fig. 4B). The intensity of subcutaneous edema, ascites, hydrothorax, and hydropericardium does not strictly relate to cardiac enlargement. The best indicators of the severity and duration of heart failure are the size, color, and firmness of the liver.

Serum CK Activity. As shown in Figure 1, the dynamic profile of the relative values of serum CK conform to the intensity of the necrotic process and the phasic pattern of the polymyopathy. A significant rise in enzyme activity, for undetermined reasons, has been noted with occurrence of heart failure.

DISCUSSION

Cumulative data arising from the literature on Syrian hamster polymyopathy have contributed to a better characterization of several pathologic components of this primary, genetically transmitted muscle disease. In the present report, the chronology of pathologic events has been emphasized since the disparity between the disease manifestations of different myopathic hamster strains has been virtually ignored thus far. In our opinion, this may be important when studying the occurrence

Figure 4. (A) Myocardial lesions in a 75-day-old cardiomyopathic hamster (UM-X7.1 line). Several necrotic and myolytic foci are visible in the septum and the left ventricular wall (arrowheads). Bar = 2.5 mm. (B) Typical failing heart in a 250-day-old cardiomyopathic hamster. Note the distension of the heart cavities and a well-organized mural thrombus on the left side. Bar = 2.5 mm.



and progression of the cardiomyopathy and when attempting to interpret microscopic and biochemical findings which are necessarily related to the phasic pattern of the disease. By using the UM-X7.1 myopathic hamster line, which retains all the essential characteristics of the BIO 14.6 line originally described by Bajusz et al.,⁴ we found, for instance, that oxidative phosphorylation of heart mitochondria^{23,36} and the changes in adrenergic nerve activity²⁴ lead to consistent results if related to the chronology of the pathologic events. This similarly applies to our investigation of the plasma membrane defect^{10,44} or the influence of calcium antagonists and β -adrenergic blockers on the course of the cardiomyopathy.^{20,23,26,28}

In spite of the differences between several myopathic lines, all deriving from BIO 1.05,¹⁶ we believe that the genetic defect underlying this polymyopathy is invariable, since all traits subsist under controlled mating conditions, and that variations between different lineages merely represent divergent genetic expressions of the same disease. This phenomenon in turn would suggest that several conditioning factors play an important role in the development and progression of hereditary polymyopathy¹⁻³ and that therapeutic trials to alleviate the symptomatology and to prolong the lifespan of animals would most likely lead to interesting and valuable results. We have already succeeded in interfering with the development and progression of necrotic myocardial changes by administering calcium antagonists.^{19,21,23,28} Thus, it may be postulated that the abnormal calcium influx results from some structural or functional membrane defect. We also investigated different drugs, such as Dibenamine and propranolol, which

act on adrenergic receptors, and certain prostaglandins, which influence adenyl cyclase activity.²⁸ All these agents proved to be effective in reducing the severity of the cardiomyopathy, thus supporting the concept that the disease is related to a basic defect in the plasma membrane.

In our attempts to elucidate the mechanism of heart muscle degeneration in this nonobstructive hereditary cardiomyopathy, we have always been puzzled by the coexistence of myolytic and coagulative necrotic lesions,^{19,27,43} the latter often accompanied by deposits of insoluble calcium apatites. Such a degenerative process is not usually encountered in patients with Duchenne muscular dystrophy whose pathologic myocardial changes essentially comprise variations in fiber size with more or less extensive epimyocardial fibrosis.¹² The present cardiomyolysis is more reminiscent of anoxic lesions in autopsy cases after brain or cardiac surgery^{38,39} or in cases of sudden traumatic death ascribed to "stress cardiomyopathies," presumably due to an excessive liberation of endogenous catecholamines.

We believe that the coagulative muscle changes in hamster cardiomyopathy result from mitochondrial calcium loading which interferes with ATP production and causes calcium precipitation within the mitochondrial matrix. While in light microscopy the pathologic fiber changes are essentially granular, in electron microscopy the collapsed fibers as well as the surrounding histiocytes are filled with calcified mitochondria.²² Myocytolysis, on the other hand, is a more insidious process characterized by myofibrillar dissolution with a concomitant increase of calcium-dependent proteases.^{8,11} The process is possibly due to increased

adrenergic tone, as suggested by Sole et al.⁴⁵ or related to abnormal catecholamine metabolism and turnover.²⁴ Similar myolytic lesions are readily reproduced in experimental animals by administration of isoproterenol,^{6,40} methoxamine, or metaraminol.^{18,25} Whatever the primary causative mechanism, the question is still debated as to whether catecholamines play a primary or a secondary role. It seems, however, that the plasma membrane defect, as evidenced by an increase in cholesterol content, largely accounts for the calcium influx in the cardiocytes and that the modality of cell necrosis probably depends upon the rate of calcium influx.^{37,41,51} In this respect, we found that while calcium antagonists successfully prevent dys-

trophic necrotic changes, these same drugs, when administered over a long period, are less effective in interfering with the myolytic process which parallels the plasma membrane deterioration;²³ the accelerated norepinephrine turnover during the progression of the cardiomyopathy²⁴ as well as the generalized anoxemia due to circulatory failure can only promote the myocytolytic process in the dystrophic heart. These data give insight into the pathologic mechanism of this genetically determined myocardial disease, and we suggest that combined treatment with calcium antagonists and adrenergic blockers may prolong the viability of cardiocytes in which the plasma membrane is primarily defective.

REFERENCES

1. Bajusz E: Hereditary cardiomyopathy: a new disease model. *Am Heart J* 77:686-696, 1969
2. Bajusz E, Baker JR, Nixon CW, Homburger F: Spontaneous, hereditary myocardial degeneration on congestive heart failure in a strain of Syrian hamsters. *Ann NY Acad Sci* 156:105-129, 1969
3. Bajusz E, Homburger F, Baker JR, Bogdonoff P: Dissociation of factors influencing myocardial degeneration and generalized cardiocirculatory failure. *Ann NY Acad Sci* 156:396-420, 1969
4. Bajusz E, Homburger F, Baker JR, Opie LH: The heart muscle in muscular dystrophy with special reference to involvement of the cardiovascular system in the hereditary myopathy of the hamster. *Ann NY Acad Sci* 138:213-231, 1966
5. Berenbaum AA, Horowitz W: Heart involvement in progressive muscular dystrophy. Report of a case with sudden death. *Am Heart J* 51:622-627, 1956
6. Bloom S, Cancilla PA: Myocytolysis and mitochondrial calcification in rat myocardium after low doses of Isoproterenol. *Am J Pathol* 54:373-391, 1969
7. Buchner F, Onishi S, Wada A: *Cardiomyopathy Associated with Systemic Myopathy. Genetic Defect of Actomyosin Influencing Muscular Structure and Function*. Urban & Schwarzenberg, Baltimore, MD, 1978, pp 43-81
8. Dayton WR, Schollmeyer JV, Chan AC, Allen CE: Elevated levels of a calcium-activated muscle protease in rapidly atrophying muscle from vitamin E-deficient rabbits. *Biochim Biophys Acta* 584:216-230, 1979
9. DeWind LT, Jones RJ: Cardiovascular observations in dystrophonia myotonica. *JAMA* 144:299-303, 1950
10. Dhalla NS, Singh A, Lee SL, Anand MB, Bernatsky AM, Jasmin G: Defective membrane systems in dystrophic skeletal muscle of the UM-X7.1 strain of genetically myopathic hamster. *Clin Sci Mol Med* 49:359-368, 1975
11. Ebashi S, Sugita H: The role of calcium in physiological and pathological processes of skeletal muscle. In *Current Topics in Nerve and Muscle Research*. Edited by Aguayo AJ, Karpatis G. Excerpta Medica, Amsterdam, The Netherlands, 1979, pp 73-84
12. Frankel KA, Rosser RJ: The pathology of the heart in progressive muscular dystrophy: epimyocardial fibrosis. *Hum Pathol* 7:375-386, 1976
13. Ganter P, Jollès G: *Histochimie Normale et Pathologique*, Vol 2. Gauthier and Villars, Paris, France, 1970, pp 952-966
14. Gertz W: Cardiomyopathic Syrian hamster: a possible model of human disease. In *Pathology of the Syrian Hamster: Progress in Experimental Tumor Research*, Vol 16. Edited by Homburger F. S Karger, Basel, Switzerland, 1972, pp 242-260
15. Homburger F: Myopathy of hamster dystrophy: history and morphologic aspects. *Ann NY Acad Sci* 317:2-17, 1979
16. Homburger F, Baker JR, Nixon CW, Welgram G: A new hereditary disease of Syrian hamsters: primary generalized polymyopathy and cardiac necrosis. *Arch Intern Med* 110:660-662, 1962
17. Homburger F, Baker JR, Nixon CW, Whitney R: Primary, generalized polymyopathy and cardiac necrosis in an inbred line of Syrian hamsters. *Med Exp (Basel)* 6:339-345, 1962
18. Jasmin G: Cardiac, aortic and pulmonary lesions induced by vasopressors. *Vasc Dis* 2:115-121, 1965
19. Jasmin G, Bajusz E: Polymyopathie et cardiomyopathie héréditaire chez le hamster de Syrie. Inhibition sélective des lésions du myocarde. *Ann Anat Pathol (Paris)* 18:49-66, 1973
20. Jasmin G, Bajusz E: Prevention of myocardial degeneration in hamsters with hereditary cardiomyopathy. In *Pathophysiology and Morphology of Myocardial Cell Alteration: Recent Advances in Studies on Cardiac Structure and Metabolism*, Vol 6. Edited by Fleckenstein A, Rona G. University Park Press, Baltimore, MD, 1975, pp 219-229
21. Jasmin G, Bajusz E, Solymoss B: Selective prevention by Verapamil and other drugs of the hamster hereditary cardiomyopathy. In *Recent Advances in Myology*. Edited by Bradley WG, Gardner-Medwin D, Walton JN. Excerpta Medica, Amsterdam, The Netherlands, 1975, pp 413-417
22. Jasmin G, Eu HY: Cardiomyopathy of hamster dystrophy. *Ann NY Acad Sci* 317:46-58, 1979
23. Jasmin G, Proschek L: Prevention of myocardial degeneration in hamsters with hereditary cardiomyopathy. In *Calcium-Antagonism*. Edited by Fleckenstein A, Roskamm H. Springer-Verlag, Berlin, W. Germany, 1980, pp 144-150
24. Jasmin G, Proschek L: The permissive role of catecholamines in the pathogenesis of the hamster cardiomyopathy. In *Advances in Myocardiology*, Vol 4. Edited by Chazou E, Sask V, Rona G. Plenum Press, New York, NY, in press
25. Jasmin G: Factors influencing the production of cardiomyopathies by methoxamine and metaraminol. *Ann NY Acad Sci* 156:333-343, 1969
26. Jasmin G, Solymoss B: Prevention of hereditary cardiomyopathy in the hamster by Verapamil and other agents. *Proc Soc Exp Biol Med* 149:193-198, 1975
27. Jasmin G, Solymoss B, Eu HY: Effect of thyropara-

- thyroidectomy and of a low-calcium diet on the cardiomyopathy of the Syrian hamster. In *Metabolism of Contraction: Recent Advances in Studies on Cardiac Structure and Metabolism*, Vol 10. Edited by Roy PE, Rona G. University Park Press, Baltimore, MD, 1975, pp 717-729
28. Jasmin G, Solymoss B, Proschek L: Therapeutic trials in hamster dystrophy. *Ann NY Acad Sci* 317:338-348, 1979
29. Lindenmayer GE, Harigaya S, Bajusz E, Schwartz A: Oxidative phosphorylation and calcium transport of mitochondria isolated from cardiomyopathic hamster hearts. *J Mol Cell Cardiol* 1:249-259, 1970
30. Lochner A, Brink AJ, Van der Walt JJ: The significance of biochemical and structural changes in the development of the myocardiopathy of the Syrian hamster. *J Mol Cell Cardiol* 1:47-64, 1970
31. Lossnitzer K, Janke J, Heinz B, Stauch M, Fleckenstein A: Disturbed myocardial calcium metabolism: a possible pathogenic factor in the hereditary cardiomyopathy of Syrian hamster. In *Pathophysiology and Morphology of Myocardial Cell Alteration: Recent Advances in Studies on Cardiac Structure and Metabolism*, Vol 6. Edited by Fleckenstein A, Rona G. University Park Press, Baltimore, MD, 1975, pp 207-217
32. Mendell JR, Higgins R, Sahenk Z, Cosmos E: Relevance of genetic animal models of muscular dystrophy to human muscular dystrophies. *Ann NY Acad Sci* 317:409-430, 1979
33. Perloff JK: Cardiac involvement in hereditary familial neuromyopathic diseases. In *Cardiomyopathy: Cardiovascular Clinics*, Vol 4. Edited by Brest AN. FA Davis Company, Philadelphia, PA, 1972, pp 333-344
34. Perloff JK, De Leon AC, O'Doherty D: The cardiomyopathy of progressive muscular dystrophy. *Circulation* 33:625-648, 1966
35. Perloff JK, Roberts WC, De Leon AC, O'Doherty D: The distinctive electrocardiogram of Duchenne's progressive muscular dystrophy. *Am J Med* 42:179-188, 1967
36. Proschek L, Jasmin G: Hereditary polymyopathy and cardiomyopathy in the Syrian hamster. II. Development of heart necrotic changes in relation to defective mitochondrial function. *Muscle Nerve* 5:26-32, 1982
37. Publicover SJ, Duncan CJ, Smith JL: The use of A23817 to demonstrate the role of intracellular calcium in causing ultrastructural damage in mammalian muscle. *J Neuropathol Exp Neurol* 37:544-557, 1978
38. Reichenbach DD, Benditt EP: Myofibrillar degeneration. A response of myocardial cell to injury. *Arch Pathol* 85:189-199, 1968
39. Reichenbach DD, Benditt EP: Catecholamines and cardiomyopathy: the pathogenesis and potential importance of myofibrillar degeneration. *Hum Pathol* 1:125-150, 1970
40. Rona G, Kahn DS: Experimental studies on the healing of cardiac necrosis. *Ann NY Acad Sci* 156:177-188, 1969
41. Schanne AXF, Kane BA, Young EE, Farber LJ: Calcium dependence of toxic cell death: a final common pathway. *Science* 206:700-702, 1979
42. Schwartz A, Sordahl LA, Crow CA, McCollum WB, Harigaya S, Bajusz E: Several biochemical characteristics of the cardiomyopathic Syrian hamster. In *Myocardiology: Recent Advances in Studies on Cardiac Structure and Metabolism*, Vol 1. Edited by Bajusz E, Rona G. University Park Press, Baltimore, MD, 1972, pp 235-242
43. Singh JN, Dhalla NS, McNamara DB, Bajusz E, Jasmin G: Membrane alteration in failing hearts of cardiomyopathic hamsters. In *Pathophysiology and Morphology of Myocardial Cell Alterations: Recent Advances in Studies on Cardiac Structure and Metabolism*, Vol 6. Edited by Fleckenstein A, Rona G. University Park Press, Baltimore, MD, 1975, pp 259-268
44. Slack BE, Boegman RJ, Downie JW, Jasmin G: Cardiac membrane cholesterol in dystrophic and Verapamil-treated hamsters. *J Mol Cell Cardiol* 12:179-185, 1980
45. Sole MJ, Lo C-M, Laird CW, Sonnenblick EH, Wurtman RJ: Norepinephrine turnover in the heart and spleen of the cardiomyopathic Syrian hamster. *Circ Res* 37:855-862, 1975
46. Strickland KP, Hudson AJ, Thakar JH: Biochemical studies in dystrophic mouse muscle. *Ann NY Acad Sci* 317:187-205, 1979
47. Strobeck JE, Factor SM, Bhan A, Sole M, Liew CC, Fein F, Sonnenblick EH: Hereditary and acquired cardiomyopathies in experimental animals: mechanical, biochemical and structural features. *Ann NY Acad Sci* 317:59-88, 1979
48. Walton JN, Nattrass FJ: On the classification, natural history and treatment of the myopathies. *Brain* 77:169-231, 1954
49. Wrogemann K, Blanchaer MC, Jacobson BE: Oxidative phosphorylation in cardiomyopathic hamsters. *Am J Physiol* 22:1453-1457, 1972
50. Wrogemann K, Nylen EG: Mitochondrial calcium overloading in cardiomyopathic hamsters. *J Mol Cell Cardiol* 10:185-195, 1978
51. Wrogemann K, Pena SDJ: Mitochondrial calcium overload: a general mechanism of cell necrosis in muscle diseases. *Lancet* i:672-674, 1976
52. Zatuchni J, Aegerter EE, Molthan L, Shuman CR: The heart in progressive muscular dystrophy. *Circulation* 3:846-853, 1951