



Mitochondrial signals to nucleus regulate estrogen-induced cell growth

Quentin Felty, Deodutta Roy*

Department of Environmental Health Sciences, University of Alabama at Birmingham, Birmingham, AL 35294-0022, USA

Received 18 August 2003; accepted 18 December 2003

Summary Classical genomic and non-genomic signaling pathways mediated by nuclear and cell membrane estrogen receptors are considered to contribute to estrogen-induced cell proliferation. Here we propose that mitochondrial signals to the nucleus regulate estrogen-induced progression of the cell cycle. The influence of estrogen on mitochondrial oxidative phosphorylation and mitochondrial gene transcription support the idea that mitochondria are significant targets of estrogen. Mitochondria are the major source of reactive oxygen species (ROS) in epithelial cells. Estrogen redox cycling within mitochondria also generates ROS. Antioxidants inhibit estrogen-induced cell growth. A-Raf, Akt, PKC, MEK, ERK, and transcription factors AP-1, NF- κ B, and CREB are targets of both estrogen and ROS. We provide four lines of evidence in support of our hypothesis that estrogen-induced mitochondrial ROS stimulate redox sensor kinase A-Raf, Akt or PKC, which, in turn, activate transcription factors NF- κ B, CREB, or AP-1 via the MEK/ERK pathway. Thus, estrogen-induced mitochondrial ROS leading to the activation of cell cycle genes containing AP-1, NF- κ B, or CREB response elements are involved in the progression of the cell cycle of the estrogen-dependent cells. Our novel concept will contribute to the development of new targets in the prevention and control of estrogen-induced disease including cancer.

© 2004 Elsevier Ltd. All rights reserved.

Introduction

Estrogens elicit different growth responses in various tissues that depend on the cell-type and dose of exposure. For example, the growth of breast, pituitary, endometrial, and testicular Leydig cells is stimulated by an estrogen concentration ranging from 0.1–1.0 pg/ml (0.1–100 pM) to 0.01–100 ng/ml (100 pM to 1 μ M) while a high concentration ranging from 1 to 100 μ g/ml (>1–100 μ M) results in

the inhibition of cell growth [1–3]. Upon hormone binding, the estrogen receptor (ER) α and ER β initiate transcription by directly binding to estrogen response elements (EREs) of genes or indirectly through the phosphorylation of signaling proteins involved in cell cycle progression. However, data obtained from various laboratories show that the receptor binding activity of estrogen does not correlate with its influence on in vitro and in vivo growth of cells. For instance, 17 α -estradiol has a very weak affinity for the estrogen receptor compared to that of diethylstilbestrol (DES), 17 β -estradiol, and 17 α -ethinylestradiol; but it is able to stimulate the growth of Leydig cells and other cells

* Corresponding author. Tel.: +1-205-934-6081; fax: +1-205-975-6341.

E-mail address: royd@uab.edu (D. Roy).

equivalent to that of DES and 17 β -estradiol [1]. Similarly, 4-hydroxyestradiol has a 50% receptor ligand binding affinity when compared to DES or 17 β -estradiol, but is able to produce stronger cell growth than either of these compounds in the uterus of mice [4]. Recently, we have identified that estrogen stimulated the growth of HEK 293 cells in an ER-independent manner given that this cell line does not contain ER (Singh KP, Venkat S, Roy D, unpublished). In contrast to the classical genomic pathways of estrogen action that occur over the course of several hours or days, recent studies have shown evidence of a rapid estrogen signaling pathway mediated by cell surface ERs and non-genomic estradiol-induced signal transduction pathways which contribute to cell proliferation [5]. These findings suggest that in addition to the receptor activity of estrogens, other factor(s) must be involved in the stimulation of cell growth by estrogen. Hence, these findings led to the generation of different perspectives on estrogen-induced signaling and cell proliferation. More recently, mitochondria have been implicated in the control of cell proliferation [6]. For instance, the mitochondrial peripheral benzodiazepine receptor (PBR) has been implicated in the regulation of human breast cancer cell proliferation [7]. Similarly, we have demonstrated that mitochondria can modulate the expression of nuclear cell cycle genes and human breast tumor growth [8,9]. The growth of estrogen-dependent and estrogen-independent cells is inhibited by controlling mitochondrial biogenesis [9]; and estrogen upregulates mitochondrial calcium in MCF-7 cells [8,10]. In this paper, we provide evidence in support of our postulate that signaling between the mitochondria and nucleus is involved in the control of estrogen-dependent cell growth. Estrogen-induced mitochondrial ROS stimulate redox sensor kinases. This in turn activates transcription factors NF- κ B, CREB, or AP-1 which upregulate early genes involved in the progression of the cell cycle of estrogen-dependent cells.

Estrogen and mitochondria

Besides the well-known mitochondrial functions of apoptosis, respiration, and oxidative phosphorylation; mitochondria also regulate inorganic ion homeostasis and the synthesis of heme, lipids, amino acids, and nucleotides. Mitochondria are also an integral component of steroidogenesis. Key enzymes of estrogen biosynthesis (3 β -hydroxysteroid dehydrogenase and aromatase) have been demonstrated in the mitochondria of ovarian tumor epi-

thelial cells [11]. Although estrogen synthesis occurs in the mitochondria, exogenously added estrogen is also transported to this organelle. For instance, in vivo exposure of ovariectomized rats to tritiated estrogen showed with increasing time, the translocation of this hormone from the plasmalemma mainly to the mitochondria (75%) rather than the nuclei in liver, adrenal gland, and spleen tissues [12]. The function of estrogen at the mitochondria is not clear, however, recent studies have identified ER α and ER β within the mitochondria implicating its role in the regulation of mitochondrial genome transcription [13,14]. Partial EREs reported in genes cytochrome oxidase (CO I and CO II) may account for the observed estrogen-induced increase of these mitochondrial transcripts in rat GH4C1 pituitary cells and rat hepatocytes [15–17]. Similarly, long-term estrogen treatment increases mitochondrial transcripts CO II, CO III, and ATP synthase subunit 6 (ATPase 6) in rodents [18]. Other mitochondrial genome regions that also contain EREs include 12S rRNA, 16S rRNA, tRNA-gln, cytochrome oxidase b, unidentified reading frame (URF) 4, URF5, and the D-loop region [15]. In the human mitochondrial genome, we have identified partial or ERE 1/2 sites in the D-loop region, CO II, tRNA-met, 12S rRNA, 7S rRNA, URF1, and URF5 (unpublished Felty Q and Roy D).

Besides transcription estrogen has also been demonstrated to effect mitochondria at the protein level. For instance, estrogen and estrogen receptor agonists have been demonstrated in several studies to inhibit mitochondrial respiratory complex I, II, III, IV, and mitochondrial ATP synthase (F₀F₁-ATPase) [19–22]. Since most of these studies used cytotoxic doses of estrogen (μ M), any physiological responses by mitochondria were probably overlooked. Estrogen is known to act as either an antioxidant or pro-oxidant depending on the concentration [23]. Estrogens redox cycle within mitochondria, which also generate ROS such as hydrogen peroxide, superoxide, and hydroxyl radicals [24–27]. Whether physiological concentrations of estrogen can stimulate mitochondrial ROS is not clear because most studies have been performed with cytotoxic doses. In addition, most experiments were conducted on isolated mitochondria, which neglected the significance of interorganelle cross-talk between the mitochondria and nucleus. Although several studies have reported estrogen specific inhibition of mitochondrial respiratory proteins, it is not clear whether estrogen can modify mitochondrial proteins at the post-translational level. There is a single report that estradiol increased the phosphorylation of a 76 kDa protein in the

mitochondrial fraction of the rat corpus luteum [28]. The presence of protein kinases within the mitochondria along with this last report of estradiol increased phosphorylation, indicates that estrogen may regulate mitochondrial physiology at the post-translational level. In summary, the influence of estrogen on mitochondrial gene transcription and mitochondrial respiratory complexes support the idea that mitochondria are a major target of estrogen within the cell.

Estrogen-induced growth of cells in relation to mitochondria

Four lines of evidence suggest that estrogen-induced mitochondrial ROS is involved in the proliferation of estrogen-dependent cells. A characteristic of rapidly dividing cancer cells is their capacity to produce significant amounts of intracellular reactive oxygen species (ROS), which has been implicated in the promotion of accelerated cell cycle activity in neoplastic cells [30]. Mitochondria have long been suspected to play a role in the development and progression of cancers. In a recent study of renal cell carcinoma, a decreased content of oxidative phosphorylation (OXPHOS) complexes II, III, IV, and V correlated with tumor aggressiveness [31]. In line with their data, Simonnet et al. [31] postulated that a decreased OXPHOS capacity may favor tumor growth and invasiveness. It is of particular interest that estrogen can modulate OXPHOS at the transcriptional level and perhaps the post-translational level since increased free radical formation has been shown to occur in the target organ of cancer in Syrian hamsters treated with estradiol [3].

Growth factor-induced ROS and cell proliferation

The rapid stimulation of intracellular ROS by platelet-derived growth factor (PDGF), epidermal growth factor (EGF), and nerve growth factor (NGF) suggests that this underlying mechanism of cell growth may be shared with other growth factors including estrogen [32]. Exogenous addition of low concentrations of H_2O_2 and/or O_2^- has been demonstrated to stimulate cell growth in a variety of cell types including muscle cells, fibroblasts, amnion cells, prostate cancer cells, and aortic endothelial cells [30]. Tumor necrosis factor alpha (TNF- α) induces gene expression via mitochondrial respiratory chain dependent activation of NF- κ B, AP-1, JNK, and

MAPKK [33]. Although the molecular signaling mechanism that initiates ROS production by mitochondria is not clear, other cell processes besides apoptosis may be coupled to this signaling event. For instance, a recent report of integrin mediated mitochondrial ROS demonstrated that another signaling mechanism besides TNF- α can activate the transcription factor NF- κ B [33]. Similarly, the proliferative response of endothelial cells to hypoxia was demonstrated to be initiated upstream by mitochondrial ROS generation that activates the MEK/ERK pathway [34]. Although other endogenous ROS sources besides mitochondria such as NAD(P)H oxidase exist, mitochondria will be the focus of this paper for the following reasons: (i) mitochondria are the principal source of intracellular ROS in epithelial cells. (ii) the growth of adenocarcinomas occur in tissue of epithelial cell origin.

Mitochondrial ROS and cell proliferation

An increasing body of evidence suggests that a moderate level of mitochondrial ROS is a stimulus for cell proliferation. The ROS molecules hydrogen peroxide and nitric oxide have been demonstrated to stimulate mitochondrial biogenesis, a process that depends on the flow of molecules into and out of the organelle [35,36]. Since mitochondrial proteins are encoded in two separate genomes (mitochondria and nuclear genome), biogenesis is a coordinated effort in which mitochondria transmit signals to the nucleus and vice versa. The question of how mitochondria transmit these signals in the process of cell proliferation has risen from reports of its involvement in cell growth. For example, in the developing rat brain mitochondrial nitric oxide synthase (mtNOS) dependent H_2O_2 production was maximal in newborns and decreased in the adult stage [37]. Cerebral granular cells isolated from newborn rats with high mtNOS activity were reported to exhibit maximal proliferation rates which depended on NO and H_2O_2 levels. In addition, manganese superoxide dismutase (MnSOD) displayed an increased pattern of activity similar to mtNOS [37]. NO has been proposed to inhibit respiratory complex IV in favor of superoxide production [38] and therefore MnSOD may work in unison with mtNOS dismutating superoxide generated by NO-dependent inhibition to H_2O_2 . Since H_2O_2 is a highly diffusible signaling molecule and both MnSOD and NOS are modulated by estrogen [39,40], it is biologically plausible that estrogen-induced redox signaling at the level of mitochondria contributes to cell proliferation in a manner similar to the cerebral granular cells in newborn rats. For

example, an increase in mitochondrial superoxide generation has been demonstrated in cultured rat hepatocytes and HepG2 cells when treated with ethinyl estradiol, estradiol, and estrogen catechol metabolites at a dose range of 0.25–5 μ M [41]. Stilbene estrogen-induced proliferation of Syrian hamster renal proximal tubular cells can be inhibited in vitro when treated with liposomes containing superoxide dismutase or catalase [42]. In human fibroblast cells, exposure to interleukin-1 beta (IL-1 β) or TNF- α causes these cells to release significant levels of superoxide. Co-treatment with an inhibitor of IL-1 β and TNF- α synthesis, pentoxifylline, inhibited stilbene estrogen-induced increase in myeloperoxidase activities, 8-hydroxydeoxyguanosine (8-OHdG) formation, mutations in the testicular genome, and prevented estrogen-induced testicular preneoplastic lesions [3]. Recently, we have shown that estrogen-induced stimulation of macrophage cells and MCF-7 cells in part occurs through reactive oxygen species [43,44]. We have also observed that estrogen-induced growth of MCF-7 cells is inhibited by reactive oxygen scavengers such as *N*-acetylcysteine, ebselen, and catalase (Singh M, Felty Q, Roy D, unpublished). In summary, these findings support the role of ROS as an autocrine growth signal in estrogen-induced cell proliferation.

Estrogen and mitochondrial oxidative stress

Recently, it has been reported that oxidative stress affects mitochondrial proteins of chronically estrogenized Syrian hamster kidney [45]. More specifically, a decrease in thiol/sulfhydryl groups is reported to occur in the mitochondrial fraction at a preneoplastic stage of carcinogenesis. The significance of this finding is that these post-translational modifications of mitochondrial proteins may occur through estrogen redox cycling and/or mitochondrial ROS. In terms of cell signaling, redox reactions involving cysteine thiol groups transduce signals by breaking or forming protein dithiol/disulfide bridges [46]. Since estrogen can induce mitochondrial ROS, we infer that the oxidation of thiols in response to estrogen converts the oxidative stress to a change in protein function involved in cell growth. Oxidative stress modifies mitochondrial matrix protein thiols [47]. Similarly, thiols on protein subunits 51- and 75-kDa of complex I have been recently reported to form mixed disulfides with glutathione (glutathionylation) in response to mitochondrial oxidative stress. This post-translational modification was reversible and correlated with an increase in mitochondrial su-

peroxide production [48]. When MCF-7 human breast tumor cells are co-treated with the mitochondrial inhibitor rotenone (10 nM) and 17- β -estradiol (10 nM) ornithine decarboxylase activity is inhibited by 86% [49]. The inhibition of NADH dehydrogenase activity was suggested to block ROS modulated signal transduction pathways [49]. The dose of rotenone (10 nM) used in this study to produce complex I inhibition is 100-fold lower than micromolar concentrations reported to disrupt microtubule assembly and arrest mammalian cells [50]. Since ornithine decarboxylase activity is a marker for cell growth, it appears that a signal transduction pathway for estrogen-induced cell growth may originate in the mitochondria assuming that rotenone inhibition is specific to complex I. Interestingly, the antitumor arotinoid, mofarotene (Ro 40-8757), down-regulates gene expression of mitochondrial encoded NADH dehydrogenase subunit 1 (MTND1) in breast cancer cell lines MDA-MB-231, ZR-75-I, and MCF-7 [51]. Since MTND1 has been reported to form part of the rotenone-binding site in complex I [52], the absence of MTND1 may remove an important site of estrogen action in mofarotene treated cells and may account for the anti-proliferative effects of this compound. Whether these protein interactions and/or modifications can occur as a result of estrogen exposure remains to be investigated. From these investigations, we infer that estrogen mediated cell growth via mitochondrial generated ROS signaling molecules may exist and merits future exploration to address this novel pathway.

Estrogens and thioredoxin redox balance

Further evidence in support of the role of mitochondria in estrogen-induced cell proliferation comes from the study of the mitochondrial thioredoxin system. In general, the two antioxidant oxidoreductase enzymes thioredoxin (Trx) and thioredoxin reductase (TrxR) that compose the system modulate signal transduction properties of ROS by the reduction of intracellular disulfides. Trx acts as a protein disulfide reductant for ribonucleotide reductase and several transcription factors including p53, NF- κ B, and AP-1 [52]. Once oxidized the active disulfide site is reduced by TrxR re-generating the reductant form of Trx. Enzyme isoforms Trx-2 and TrxR2 are reported to exist in the mitochondria [52]. Recently, a dominant negative form of TrxR2 (TrxR2DN) was expressed in HeLa cells to investigate its biological role [53]. The analysis of TrxR2DN induction by flow cytometry revealed an increase in the G₁ to S phase

transition and an increase in cell growth as measured by the MTT assay. In addition, cell cycle genes involved in the G₁ and S phases of the cell cycle were modulated by TrxR2DN expression. The dominant negative inhibition of TrxR2 was suggested to cause an increase in H₂O₂ accumulation, which activated cell proliferation [54]. Although it is not clear whether estrogen can modulate the H₂O₂ levels of mitochondria, estrogen (<10 nM) is reported to increase the expression of Trx protein in human brain SH-SY5Y cells and may play a role in cell viability [40]. In vitro estrogen (10–100 nM) treatment of primary human endometrial stromal cells show an increase in Trx protein and mRNA which implicate Trx involvement in cell growth and differentiation of estrogen responsive tissue [54]. Further evidence that supports the role of ROS in estrogen responsive tissue comes from an in vivo study that demonstrated sex hormone modulation of redox status in human endometrial cells occurs during the menstrual cycle [55]. Alterations in cellular redox status by increased expression of TrxR2 have been suggested to play a role in the growth of hepatocellular carcinomas [56]. Whether estrogen can specifically increase the expression or activity of Trx2 and/or TrxR2 is not known, but these findings suggest that estrogen may modulate signal transduction of mitochondrial derived ROS via the thioredoxin system. From these studies it appears that mitochondria may be involved in the growth of estrogen-dependent cells.

Relationship between the mitochondrial signals and progression of estrogen-induced cell cycle

The exact mitochondrial signals that control the process of cell proliferation are not clear. However, based on the studies described above it appears that mitochondria may control estrogen-induced proliferation in multiple ways. In this section, we provide evidence in support of mitochondrial signals that may be involved in cell cycle progression of estrogen-dependent cells. Mitochondrial ROS fulfill the prerequisites of a second messenger since they are short-lived (rapidly generated and degraded), produced in response to a stimulus, highly diffusible (H₂O₂), and ubiquitously present in most cell types. Several oxidant sensitive proteins involved in cell signaling have been identified in the mitochondria. For example, src protein tyrosine kinase, which is stimulated by ROS, was recently identified in the intermembrane space of rat brain mitochondria [57]. The close

proximity of src to mitochondrial ROS raises the interesting question of whether this organelle participates in pathways regulated by src family kinases such as cell growth via redox signaling [32]. In addition, other oxidant sensitive kinases are localized at the mitochondria, which include a-raf, Akt, and PKC.

Estrogen-mediated changes in redox sensor proteins and activation of transcription factors

A large class of DNA-binding proteins that include the glucocorticoid receptor (GR), estrogen receptor (ER), and thyroid hormone receptor contain zinc finger domains and are localized in the mitochondria [58]. In addition to hormone receptors, there are several protein kinases such as a-raf and PKC that have zinc finger domains. Zinc finger structures within a protein consist of at least two zinc-coordinated thiolates. Oxidation of a zinc finger converts cysteine thiol groups to disulfide releasing zinc from the protein. A change in protein conformation results from the mobilization of zinc that may affect protein function. Although protein kinases are known to participate in phosphorylation signal cascades, they may also participate in redox signaling networks due to these zinc finger domains. For instance, ROS can activate protein kinase C (PKC) that triggers the release of zinc ions [59]. Another protein kinase, c-Raf, known to participate in the MAPK signal cascade was also demonstrated to be redox activated at the zinc finger domain [60]. Similarly, ROS has been reported to modulate ER α and ER β protein expression in various cell lines. In human breast cancer cells MCF-7 and T-47D, exposure to 2.5 μ M of H₂O₂ increased the protein level of ER β while having a minimal effect on ER α [61]. Endogenous oxidative stress induced by PMA (100 ng/ml) in the macrophage cell line J774A.1 also increased the expression of ER β [61]. Since zinc-finger domains within these proteins were reported to act as oxidant sensors, a mode of cross-communication appears to exist between redox and phosphorylation networks. The mitochondrial localization of protein kinases a-raf, PKC δ , and PKC ϵ is evidence that this subcellular compartment harbors redox sensors [29,62]. Although the role of the protein kinase a-raf in the mitochondria is not clear, a-raf mRNA is highly expressed in normal murine tissues such as the epididymis, ovary, kidney, and urinary bladder [63]. In HeLa cells, epidermal growth factor rapidly (2 min) and transiently activated a-raf, which in turn phosphorylated the MAP kinase activator MEK1

[64]. In addition to a-raf and Akt, other redox sensitive proteins associated with mitochondria, PKC δ and PKC ϵ , have been demonstrated to activate the raf/MEK/ERK pathway or directly activate MAPKs, respectively [62,65]. Interestingly, 17 β -estradiol can stimulate the phosphorylation of a-raf and cell cycle progression in ER positive MCF-7 cells [66]. Rapid effects of estrogen have been demonstrated to mediate the DNA binding activity and phosphorylation of transcription factors. For instance, a 10 nM estradiol treatment of rat adipocytes doubled AP-1 DNA binding in 15 min [67]. In addition, CREB protein was quickly phosphorylated by estradiol (10 nM) treatment within 15 min. Another transcription factor, NF- κ B, is regulated by its interactions with the inhibitory co-factor I κ B that sequesters NF- κ B to the cytoplasm. Akt is known to phosphorylate an upstream kinase, IKK α , which stimulates the degradation of I κ B [68]. Therefore, estrogen-induced mitochondrial ROS may stimulate Akt leading to the degradation of I κ B and activation of the transcription factor NF- κ B [68]. Whether estrogen treatment can activate Akt via mitochondrial derived ROS is not clear, however, phosphorylation and translocation of Akt to the mitochondria was demonstrated when cells are treated with estrogen [69]. Akt can activate raf kinase [68]. Given that estradiol can stimulate mitochondrial ROS generation; ER, a-raf, Akt, and PKC are targets of oxidative stimuli localized at the mitochondria; and the transcription factors AP-1, NF- κ B, and CREB are stimulated by oxidants

[32,70,71]; it is possible that estrogen specific effects at the level of mitochondria can activate these transcription factors. Based on these studies we postulate that estrogen exposure to cells increases mitochondria ROS, which stimulates redox sensor proteins a-raf, Akt, and PKC that sense a change in redox state. Hence, estrogen-induced mitochondrial ROS stimulates a-raf, Akt, or PKC, which in turn activates transcription factors NF- κ B, CREB, or AP-1 via the MEK/ERK pathway (scheme shown in Fig. 1) resulting in the transcription of cell cycle genes containing DNA responsive elements for NF- κ B, CREB, or AP-1 and ultimately estrogen-induced cell proliferation.

ROS mediated reorganization of cytoskeleton

Mitochondrial ROS may also affect cell growth by its actions on actin filaments. Actin is one of the most abundant proteins in eukaryotic cells involved in various processes such as cell cycle control, cell signaling, and cell structure [72]. Mechanical signals associated with cytoskeletal tension and restructuring are a requirement for anchorage-dependent cells to pass through the late G₁ restriction point of the cell cycle [73]. Whether mitochondrial ROS can affect the elasticity of the actin network is not clear, however, hydrogen peroxide was demonstrated to induce actin filament reorganization that increased cell motility




Figure 1 A scheme showing estrogen-induced mitochondrial ROS-mediated stimulation of redox sensor kinases A-Raf, Akt or PKC leading to activation of transcription factors NF- κ B, CREB, or AP-1 via the MEK/ERK pathway, which, in turn, regulate cell cycle genes containing AP-1, NF- κ B, or CREB response elements.

and in vitro angiogenesis [72]. Therefore, we postulate that estrogen-induced mitochondrial ROS may directly regulate progression of cell cycle through mechanical signals associated with cytoskeletal tension.

Conclusion

Although CREB, AP-1, and NF- κ B are responsible for the transcription of a large number of genes in the nucleus, the localization of these transcription factors in the mitochondria raises an interesting question of whether estrogen can modulate mitochondria gene expression independent of the nuclear estrogen receptor [74–76]. Hence, the simultaneous actions of estrogen on both the nuclear genome and mitochondrial genome may be a complex series of events involved in the process of cell growth. We conclude that in addition to the mitochondria being an integral component of steroidogenesis, the influence of estrogen on various components of mitochondrial oxidative phosphorylation and mitochondrial gene transcription supports the idea that mitochondria are a major target of estrogen. Estrogen-induced mitochondrial ROS stimulates redox sensor kinases A-Raf, Akt or PKC leading to activation of transcription factors NF- κ B, CREB, or AP-1 via the MEK/ERK pathway. This, in turn, regulates early cell cycle genes containing AP-1, NF- κ B, or CREB response elements. This concept will contribute to the development of new targets in the prevention and control of estrogen-induced diseases.

References

- [1] DuMond Jr JW, Singh KP, Roy D. Regulation of the growth of mouse Leydig cells by the inactive stereoisomer, 17 α -estradiol: Lack of correlation between the elevated expression of ER α and difference in sensitivity to estradiol isomers. *Oncol Rep* 2001;8:899–902.
- [2] Welshons WV, Thayer KA, Judy BM, Taylor JA, Curran EM, vom Saal FS. Large effects from small exposures. I. Mechanisms for endocrine disrupting chemicals with estrogenic activity. *Env Hlth Perspec* 2003;111:994–1006.
- [3] Roy D, Cai Q. Estrogen, immunoactivation, gene damages and development of breast, endometrial, ovarian, prostate and testicular cancers. *Recent Res Devel Steroid Biochem Mol Biol* 2002;3:1–32.
- [4] Newbold RR, Liehr JG. Induction of uterine adenocarcinoma in CD-1 mice by catechol estrogens. *Cancer Res* 2000;60:235–7.
- [5] Levin ER. Cell localization, physiology, and nongenomic actions of estrogen receptors. *J Appl Physiol* 2001;91:1860–7.
- [6] Rustin P. Mitochondria, from cell death to proliferation. *Nat Genet* 2002;30:352–3.
- [7] Beinlich A, Strohmeier R, Kaufmann M, Kuhl H. Relation of cell proliferation to expression of peripheral benzodiazepine receptors in human breast cancer cell lines. *Biochem Pharmacol* 2000;60:397–402.
- [8] Felty Q, Singh KP, Roy D. Inhibition of human breast tumor cell growth by controlling mitochondrial biogenesis. *The Toxicol* 2003;211.
- [9] Felty Q, Singh KP, Roy D. Modulation of nuclear cell cycle gene expression by arresting mitochondrial function. *Proc Am Assoc Cancer Res* 2003;44:21.
- [10] Parkash J, Felty Q, Roy D. Differential regulation of high and low capacity mitochondrial calcium in breast cancer MCF-7 cells by estrogen. *Proc Am Assoc Cancer Res* 2003;44:715.
- [11] Matsumoto Y. Study on the estrogen production in parenchymatous cells of epithelial ovarian tumor. *Osaka City Med J* 1992;38:27–43.
- [12] Moats II RK, Ramirez VD. Rapid uptake and binding of estradiol-17 β -6- (O-carboxymethyl)oxime:125I-labeled BSA by female rat liver. *Biol Reprod* 1998;58:531–8.
- [13] Monje P, Boland R. Subcellular distribution of native estrogen receptor α and β isoforms in rabbit uterus and ovary. *J Cellular Biochem* 2001;82:467–79.
- [14] Chen J, Yager JD. Mitochondrial localization of estrogen receptor (ER)s α and β . *Proc Am Assoc Cancer Res* 2001;42:882.
- [15] Hatzoglou E, Sekeris CE. The detection of nucleotide sequences with strong similarity to hormone responsive elements in the genome of eubacteria and archaeobacteria and their possible relation to similar sequences present in the mitochondrial genome. *J Theor Biol* 1997;184:339–44.
- [16] Van Itallie CM, Dannies PS. Estrogen induces accumulation of the mitochondrial ribonucleic acid for subunit II cytochrome oxidase in pituitary tumor cells. *Mol Endocrin* 1988;2:332–7.
- [17] Chen J, Gokhale M, Li Y, Trush MA, Yager JD. Enhanced levels of several mitochondrial transcripts and mitochondrial superoxide production during ethinyl estradiol-induced hepatocarcinogenesis and after estrogen treatment of HepG2 cells. *Carcinogenesis* 1998;19:2187–93.
- [18] Thomas RD, Roy D. Base sequence-specific attack of stilbene estrogen metabolite(s) on the mitochondrial DNA: implications in the induction of instability in the mitochondrial genome in the kidney of Syrian hamsters. *Int J Mol Med* 2001;7:389–95.
- [19] De Otamendi ME, Stoppani AO. Action of diethylstilbesterol on the NADH-dehydrogenase region of the respiratory chain. *Arch Biochem Biophys* 1974;165:21–33.
- [20] Garcia MV, Cabezas JA, Perez-Gonzalez MN. Effects of oestradiol, testosterone and medroxyprogesterone on subcellular fraction marker enzyme activities from rat liver and brain. *Comp Biochem Physiol B* 1985;80:347–54.
- [21] Tuquet C, Dupont J, Mesneau A, Roussaux J. Effects of tamoxifen on the electron transport chain of isolated rat liver mitochondria. *Cell Biol Toxicol* 2000;16:207–19.
- [22] Kipp JL, Ramirez VD. Effect of estradiol, diethylstilbestrol, and resveratrol on FoF-ATPase activity from mitochondrial preparations of rat heart, liver, and brain. *Endocrine* 2001;15:165–75.
- [23] Liehr JG, Roy D. Pro-oxidant and antioxidant effects of estrogens. *Meth Mol Biol* 1998;108:432–5.
- [24] Thomas RD, Roy D. Redox cycling of estrogens catalyzed by mitochondrial enzymes. *Carcinogenesis* 1995;16:891–5.
- [25] Roy D, Abul-Hajj Y. Estrogen-nucleic acid adducts: guanine is major site for interaction between 3,4-estrone quinone and Co III gene. *Carcinogenesis* 1997;18:1247–9.

- [26] Roy D, Liehr JG. Estrogens, DNA damages and mutations. *Mut Res* 1999;424:107–15.
- [27] Thomas RD, Roy D. Modifications in mitochondrial DNA by stilbene estrogen and hepatocarcinogenicity. *Oncol Rep* 2001;8:1035–8.
- [28] Steinschneider A, Rao MC, Khan I, McLean MP, Gibori G. Calcium-calmodulin and calcium-phospholipid dependent phosphorylation of membranous fraction proteins related to the tropic regulation by estradiol in the corpus luteum. *Endocrinology* 1991;128:263–72.
- [29] Thomson M. Evidence of undiscovered cell regulatory mechanisms: phosphoproteins and protein kinases in mitochondria. *Cell Mol Life Sci* 2002;59:213–9.
- [30] Sauer H, Wartenberg M, Hescheler J. Reactive oxygen species as intracellular messengers during cell growth and differentiation. *Cell Physiol Biochem* 2001;11:173–86.
- [31] Simonnet H, Alazard N, Pfeiffer K, et al. Low mitochondrial respiratory chain content correlates with tumor aggressiveness in renal cell carcinoma. *Carcinogenesis* 2002;23:759–68.
- [32] Drodge W. Free radicals in the physiological control of cell function. *Physiol Rev* 2002;82:47–95.
- [33] Werner E, Werb Z. Integrins engage mitochondrial function for signal transduction by a mechanism dependent on Rho GTPases. *J Cell Biol* 2003;158:357–68.
- [34] Schafer M, Schafer C, Ewald N, Piper HM, Noll T. Role of redox signaling in the autonomous proliferative response of endothelial cells to hypoxia. *Circ Res* 2003;92:1010–5.
- [35] Lee HC, Yin PH, Chi CW, Wei YH. Increase in mitochondrial mass in human fibroblasts under oxidative stress and during replicative senescence. *J Biomed Sci* 2002;9:517–26.
- [36] Nisoli E, Clementi E, Paolucci C, et al. Mitochondrial biogenesis in mammals: the role of endogenous nitric oxide. *Science* 2003;299:896–9.
- [37] Riobo NA, Melani M, Sanjuan N, et al. The modulation of mitochondrial nitric oxide synthase activity in rat brain development. *J Biol Chem* 2002;277:42447–55.
- [38] Brookes P, Darley-Usmar VM. Hypothesis: the mitochondrial NO (*) signaling pathway, and the transduction of nitrosative to oxidative cell signals: an alternative function for cytochrome c oxidase. *Free Radic Biol Med* 2002;32:370–4.
- [39] Lee SY, Andoh T, Murphy DL, Chiueh CC. 17 beta-estradiol activates ICI 182,780-sensitive estrogen receptors and cyclic GMP-dependent thioredoxin expression for neuroprotection. *FASEB J* 2003;17:947–8.
- [40] Strehlow K, Rotter S, Wassmann S, et al. Modulation of antioxidant enzyme expression and function by estrogen. *Circ Res* 2003;93:170–7.
- [41] Oberley TD, Allen RG, Schultz JL, Lauchner LJ. Antioxidant enzymes and steroid-induced proliferation of kidney tubular cells. *Free Radic Biol Med* 1991;10:79–83.
- [42] Kobiela J, Krajewski J, Kalinska-Blach B, Stefaniak T. Selectivity of oxidative stress targeting in estrogen-induced experimental nephrocarcinogenesis. *Acta Biochim Pol* 2002;49:51–8.
- [43] Sharga A, Felty Q, Roy D. The influence of alcohol on estrogen-induced stimulation of cells in part occurs through reactive oxygen species. *Proc Am Assoc Cancer Res* 2003;44:1200.
- [44] Venkat S, Felty Q, Roy D. Estrogen-induced stimulation of macrophages leading to the generation of reactive oxygen species in the target organ of cancer. *The Toxicologist* 2003:235.
- [45] Pomposiello PJ, Demple B. Redox-operated genetic switches: the SoxR and OxyR transcription factors. *Trends Biotechnol* 2001;19:109–14.
- [46] Lin TK, Hughes G, Muratovska A, et al. Specific modification of mitochondrial protein thiols in response to oxidative stress: a proteomics approach. *J Biol Chem* 2002;277:17048–56.
- [47] Taylor ER, Hurrell F, Shannon RJ, Lin TK, Hirst J, Murphy MP. Reversible glutathionylation of complex I increases mitochondrial superoxide formation. *J Biol Chem* 2003;278:19603–10.
- [48] Rowlands JC, Casida JE. NADH:Ubiquinone oxidoreductase inhibitors block induction of ornithine decarboxylase activity in MCF-7 human breast cancer cells. *Pharm Toxicol* 1998;83:214–9.
- [49] Barrientos A, Moraes CT. Titrating the effects of mitochondrial complex I impairment in the cell physiology. *J Biol Chem* 1999;274:16188–97.
- [50] Uchida T, Inagaki N, Furuichi Y, Eliason JF. Down-regulation of mitochondrial gene expression by the anti-tumor arotinoid mofarotene (Ro 40-8757). *Int J Cancer* 1994;58:891–7.
- [51] Earley FG, Patel SD, Ragan CI, Attardi G. Photolabelling of a mitochondrially encoded subunit of NADH dehydrogenase with [³H] dihydrotorotene. *FEBS Lett* 1987;219:108–13.
- [52] Nordberg J, Arner ES. Reactive oxygen species, antioxidants, and the mammalian thioredoxin system. *Free Radic Biol Med* 2001;31:1287–312.
- [53] Kim MR, Chang HS, Kim BH, et al. Involvements of mitochondrial thioredoxin reductase (TrxR2) in cell proliferation. *Biochem Biophys Res Commun* 2003;304:119–24.
- [54] Maruyama T, Sachi Y, Furuake K, et al. Induction of thioredoxin, a redox-active protein, by ovarian steroid hormones during growth and differentiation of endometrial stromal cells in vitro. *Endocrinology* 1999;140:365–72.
- [55] Serviddio G, Loverro G, Vicino M, et al. Modulation of endometrial redox balance during the menstrual cycle: relation with sex hormones. *J Clin Endocrinol Metab* 2002;87:2843–8.
- [56] Choi JH, Kim TN, Kim S, et al. Overexpression of mitochondrial thioredoxin reductase and peroxiredoxin III in hepatocellular carcinomas. *Anticancer Res* 2002;22:3331–5.
- [57] Salvi M, Brunati AM, Clari G, Toninello A. Interaction of genistein with the mitochondrial electron transport chain results in opening of the membrane transition pore. *Biochem Biophys Acta* 2002;555:187–96.
- [58] Webster KA, Prentice H, Bishopric NH. Oxidation of zinc finger transcription factors: physiological consequences. *Antioxid Redox Signal* 2001;3:535–48.
- [59] Korichneva I, Hoyos B, Chua R, Levi E, Hammerling U. Zinc release from protein kinase C as the common event during activation by lipid second messenger or reactive oxygen. *J Biol Chem* 2002;277:44327–31.
- [60] Hammerling U, Hoyos B, Korichneva I, Chua R, Levi E. The zinc-finger domains of kinases function as redox sensors. *The Oxy Club of California Abstracts*, 2003. p. 13.
- [61] Tamir S, Izrael S, Vaya J. The effect of oxidative stress on ERalpha and ERbeta expression. *J Steroid Biochem Mol Biol* 2002;81:327–32.
- [62] Baines CP, Song CX, Zheng YT, et al. Protein kinase C epsilon interacts with and inhibits the permeability transition pore in cardiac mitochondria. *Circ Res* 2003;92(8):873–80.
- [63] Storm SM, Cleveland JL, Rapp UR. Expression of raf family proto-oncogenes in normal mouse tissues. *Oncogene* 1990;5:345–51.
- [64] Wu X, Noh SJ, Zhou G, Dixon JE, Guan KL. Selective activation of MEK1 but not MEK2 by A-Raf from epidermal

- growth factor-stimulated Hela cells. *J Biol Chem* 1996;271:3265–71.
- [65] Li L, Lorenzo PS, Bogi K, Blumberg PM, Yuspa SH. Protein kinase C δ targets mitochondria, alters mitochondrial membrane potential, and induces apoptosis in normal and neoplastic keratinocytes when overexpressed by an adenoviral vector. *Mol Cell Biol* 1999;19:8547–58.
- [66] Weinstein-Oppenheimer CR, Burrows C, Steelman LS, McCubrey JA. The effects of beta-estradiol on Raf activity, cell cycle progression and growth factor synthesis in the MCF-7 breast cancer cell line. *Cancer Biol Ther* 2002;1:256–62.
- [67] Dos Santos EG, Dieudonne MN, Pecquery R, Le Moal V, Giudicelli Y, Lacasa D. Rapid nongenomic E2 effects on p42/p44 MAPK, activator protein-1, and cAMP response element binding protein in rat white adipocytes. *Endocrinology* 2002;143:930–40.
- [68] Bozinovski S, Jones JE, Vlahos R, Hamilton JA, Anderson GP. Granulocyte/macrophage-colony-stimulating factor (GM-CSF) regulates lung innate immunity to lipopolysaccharide through Akt/Erk activation of NF κ B and AP-1 in vivo. *J Biol Chem* 2002;277:42808–14.
- [69] Sasaki K, Sato M, Umezawa Y. Fluorescent indicators for Akt/protein kinase B and dynamics of Akt activity visualized in living cells. *J Biol Chem* 2003;278:30945–51.
- [70] Chen J, Li Y, Lavigne JA, Trush MA, Yager JD. Increased mitochondrial superoxide production in rat liver mitochondria, rat hepatocytes, and HepG2 cells following ethinyl estradiol treatment. *Toxicol Sci* 1999;51:224–35.
- [71] Bedogni B, Pani G, Colavitti R, et al. Redox regulation of cAMP-responsive element-binding protein and induction of manganese superoxide dismutase in nerve growth factor-dependent cell survival. *J Biol Chem* 2003;278:16510–9.
- [72] Qian Y, Luo J, Leonard SS, et al. Hydrogen peroxide formation and actin filament reorganization by Cdc42 are essential for ethanol-induced in vitro angiogenesis. *J Biol Chem* 2003;278:16189–97.
- [73] Huang S, Ingber DE. A discrete cell cycle checkpoint in late G (1) that is cytoskeleton-dependent and MAP kinase (Erk)-independent. *Exp Cell Res* 2002;275:255–64.
- [74] Cammarota M, Paratcha G, Bevilacqua LR, et al. Cyclic AMP-responsive element binding protein in brain mitochondria. *J Neurochem* 1999;72:2272–7.
- [75] Cogswell PC, Kashatus DF, Keifer JA, et al. NF-kappa B and I kappa B alpha are found in the mitochondria. Evidence for regulation of mitochondrial gene expression by NF-kappa B. *J Biol Chem* 2003;278:2963–8.
- [76] Ogita K, Okuda H, Kitano M, Fujinami Y, Ozaki K, Yoneda Y. Localization of activator protein-1 complex with DNA binding activity in mitochondria of murine brain after in vivo treatment with kainate. *J Neurosci* 2002;22:2561–70.

Available online at www.sciencedirect.com

