

Bone marrow induced osteogenesis in hydroxyapatite and calcium carbonate implants

Jyrki Vuola*, Harry Göransson[†], Tom Böhling[‡] and Sirpa Asko-Seljavaara*

*Department of Plastic Surgery, Helsinki University Central Hospital, Topeliuksenkatu 5, SF-00260; [†]ORTON, Orthopaedic Hospital, Invalid Foundation; [‡]Department of Pathology, University of Helsinki, Helsinki, Finland

In this experimental study, blocks of natural coral (calcium carbonate) and its structurally similar derivate in the form of hydroxyapatite (calcium phosphate) were implanted in rat latissimus dorsi muscle with autogenous bone marrow to compare their bone-forming capability. A block without marrow placed in the opposite latissimus muscle served as a control. The animals were killed at 3, 6 and 12 weeks and, in the hydroxyapatite group, also at 24 weeks. The sections were analysed histologically and histomorphometrically. Bone was found only in implants containing bone marrow. Bone formation was significantly ($p < 0.05$) higher in coral than in hydroxyapatite implants at 3 weeks (10.8% versus 4.8%) and at 12 weeks (13.7% versus 6.3%, bone/total original block area). At 12 weeks all the coral implants had lost their original structure, and the cross-sectional area of the block had diminished to 40% of the original area. © 1996 Elsevier Science Limited.

Keywords: Bone substitute, hydroxyapatite, calcium carbonate, coral, osteogenesis, biodegradation

Received 22 June 1995; accepted 15 November 1995

In structure, the Pacific reef-building coral genus *Porites* resembles osteon-evacuated cortical bone¹. It consists almost completely (99%) of calcium carbonate (aragonite) and can be used as a bone substitute after the remaining 1% of the organic material has been eliminated.

Natural coral (CC) can be converted into structurally similar porous hydroxyapatite (HA). Through a hydrothermal exchange reaction, the CaCO_3 skeleton of CC is changed to HA, $\text{Ca}_{10}(\text{PO}_4)_6(\text{OH})_2$, which is the main mineral of bone². The mean diameter of the pores is ca. $200\text{ }\mu\text{m}$ ($190\text{--}230\text{ }\mu\text{m}$)³. A pore size of over $100\text{ }\mu\text{m}$ was previously found to allow fibrovascular and bone tissue ingrowth, making direct integration with bone possible⁴⁻⁶.

The basic difference between HA and CC is that CC is biodegradable and HA is not at all or only very little. The rate and mechanism of biodegradation of CC is not completely understood but it is assumed that the main factor is carbonic anhydrase, an enzyme that occurs in abundance in osteoclasts⁷. Both CC and HA are known to be osteoconductive, biocompatible and very inert⁸⁻¹⁰. They have been used clinically in dental¹¹⁻¹³, maxillo-facial¹⁴⁻¹⁶, cranio-facial¹⁷⁻²⁰, and orthopaedic surgery²¹⁻²⁵.

Adding bone marrow to porous calcium ceramics

induces membraneous bone formation in extraosseal sites. This has been shown with tricalcium phosphatase blocks implanted subcutaneously in rats²⁶, with HA blocks implanted intramuscularly in rabbits²⁷ and rats^{28,29} and also with CC implanted subcutaneously in rats²⁹.

In modern surgery, prefabricated bone substitutes are needed for the reconstruction of bone defects or as augmentation materials. An implant with existing bone tissue before reconstruction could be used to facilitate integration with the recipient site. This study was designed to compare bone formation inside structurally similar CC and HA blocks in muscle tissue when implanted with autogenous marrow cells. The intramuscular site was chosen to evaluate bone formation in richly vascularized tissue without the direct influence of bone.

MATERIALS AND METHODS

The two implants derive from the same genus of coral, *Porites*. The materials used were Biocoral[®] (calcium carbonate, INOTEB, France), and Interpore 200[®] (hydroxyapatite, INTERPORE INTERNATIONAL, Irvine, California) which are similar in internal structure but differ in chemical constitution. Sixty-seven blocks measuring $3 \times 3 \times 6\text{ mm}$ were implanted.

Correspondence to Dr J. Vuola.

Table 1 Number of implanted blocks

| Weeks | HA | HA contr | CC | CC contr |
|---------|------|----------|----|----------|
| 3 | 6 | 5(6) | 6 | 6 |
| 6 | 5(6) | 5(6) | 6 | 6 |
| 12 | 7 | | 6 | |
| 24 | 6 | | | |
| animals | 25 | | 18 | |

HA, hydroxyapatite; CC, coral. Values in parenthesis indicate original number of implanted blocks. Three blocks were discarded because of technical failures in processing. Figures in last row are number of animals used.

Forty-three male Wistar rats weighing 250–350 g were used for the experiment. The animals were maintained on a standard dry laboratory diet and given water *ad libitum*. They were anaesthetized with an intraperitoneal injection of pentobarbital (50 mg kg⁻¹, Mebumat®, Orion, Finland) and were given procaine penicillin (300,000 IU kg⁻¹, Procapan®, Orion, Finland) intramuscularly before the operation. The numbers of animals and implanted blocks at different follow-up periods are given in Table 1. Autogenous bone marrow was harvested from the left femur through the knee joint. A parapatellar incision was cut and the patella shifted laterally. A hole was drilled into the intercondylar space and the marrow was sucked out with a 20 cm⁻³ syringe and 18-G needle. The bone marrow was mixed with 1 cm⁻³ of saline and the implant was immersed in the solution for 5 min. The left latissimus dorsi muscle was obtained through a dorsolateral incision. It was freed from its distal insertions and the implant was rolled inside the muscle. The muscle was sutured to prevent the implant from dislocating. Right latissimus dorsi muscle with an implant of the same material without bone marrow served as a control.

The animals were killed with an overdose of pentobarbital at 3, 6 and 12 weeks; those in the HA group also at 24 weeks. The muscle around the implant was dissected and the specimen was dehydrated in increasing concentrations of ethyl alcohol–water solution and embedded in methylmethacrylate. Longitudinal 5 µm-thick sections were cut in the centre of the block with microtome (Polycut S, Reichert-Jung, Nussloch, Germany) and stained with the Masson–Goldner method for histological and histomorphometric analysis.

The study followed the principles of the European Convention for the Protection of Vertebrate Animals Used for Experimental and Other Purposes³⁰ and was approved by the ethical committee for animal experiments of Helsinki University Central Hospital.

The specimens were studied by ordinary light microscopy. The number of osteoblasts, osteocytes and giant cells and the amount of fat and bone marrow were evaluated on each sample. Special interest was focussed on the amount of macrophages and lymphocytes/neutrophils as signs of inflammation. The number of each cell type was scored from 0 to 3. The scoring system was only used as an aid to evaluation and the numbers are not presented here. The quality of bone was also assessed.

In quantitative histomorphometric analysis, a Leitz

Diaplan light microscope was linked via video camera to a semiautomatic computerized analysis system (MOP Videoplan, Kontron, Munich, Germany). The entire section area was measured using ×150 magnification.

The variables to be measured were total implant cross-sectional area, bone area, fibrotic tissue area and the void area in pores. Three samples were excluded because processing failed.

Bone formation was compared using the total original cross-sectional area of the implant (before implantation) as a reference, not the true, resorbed area alone.

Values are given as means (±SEM). The Kruskal–Wallis nonparametric test was used for statistical analysis and *p* < 0.05 was considered significant.

RESULTS

Histology

None of the control implants showed bone formation. A giant cell reaction and invasion of macrophages were observed in both materials. In some of the CC implants small foreign-body type granulomas could be detected. No acute inflammation was seen in any of the specimens. The CC implant was successively deformed, whereas the size and form of the HA implants were unchanged. Practically all the pores in both control materials were filled with fibrovascular tissue at 3 weeks.

Bone was formed in all implants with bone marrow. At 3 weeks, abundant osteoblastic activity was seen directly on the surface of the pores. Bone formation had proceeded so far that scattered osteocytes and woven bone could be detected (Figure 1a and b). In both types of implants, numerous giant cells and macrophages were seen, but only in direct contact with the foreign material, never adjacent to new bone formation. Only scattered lymphocytes occurred, and no acute inflammation could be detected. The pores were filled almost completely with fibrovascular and bone tissue. At 6 weeks, bone marrow was also present, and both materials showed less osteoblastic activity than at 3 weeks. In some of the mature CC specimens, lamellar bone could also be seen. In the CC specimens there were fewer giant cells and macrophages than at 3 weeks, but in the HA specimens the number was unchanged. Some fibrovascular stroma, but no fat, could be seen in the pores in the periphery of the blocks of both groups.

At 12 weeks, bone marrow was still present in both types of material. The amount of mature bone had increased and there was less osteoblastic activity. In the CC specimens only a few giant cells or macrophages could be seen. In contrast, giant cells and macrophages were evident in the HA specimens in all areas where bone did not line the implant material. The CC implants were deformed and nearly all the coral material not lined by bone had been phagocytized. In CC blocks the innermost part of the implant resembled an ossicle (Figure 1c and d).

In the HA implants bone marrow was still present at




Figure 1 a, CC at 3 weeks: abundant osteoblastic activity (big arrow). Matrix (white) is partly phagocytized (arrow), $\times 200$. b, HA at 3 weeks: giant cells and macrophages lying directly on matrix (arrow), $\times 260$. c, CC at 12 weeks: implant has lost shape and form, bone marrow occupies central part and is surrounded by lamellar bond (dark) and matrix, $\times 80$. d, Same specimen as in c: osteoblastic activity and osteoid seam is still seen, $\times 400$. e, HA at 24 weeks: form and shape of matrix (grey) is preserved, bone (dark) invades entire implant, $\times 80$. f, Same specimen as in e: fat gradually replaces bone marrow (arrow), $\times 400$.

24 weeks, although fat cells gradually invaded the pores and replaced the marrow. There was some osteoblastic activity, but most of the bone tissue was already quite mature, and contained abundant osteocytes. Some giant cells could still be detected. The size and form of the implant was unchanged (Figure 1e and f). No cartilage was detected in any implant. Bone formation did not show a centripetal pattern; instead bone was scattered all over the sample. New bone always appeared adjacent to the implant matrix.

Histomorphometry

At 3 weeks there was significantly more bone in CC than in HA blocks, 10.8 and 4.8%, respectively, of the original cross-sectional area (Figure 2). The CC implants were reduced to 87% of the original area (Table 2).

At 6 weeks the extent of bone formation had not changed significantly, the mean size of the CC blocks having decreased in the marrow group to 62% and in the control group to 61% of the original. The variation

ranged from 88% to nil. In the control group one block was completely resorbed.

At 12 weeks the bone formation in the HA group was 6.3% and in the CC group 13.7%. The difference between the two materials was significant ($p = 0.032$). The increase was not statistically significant within either group when compared with 3 or 6 week samples.

The size of the CC blocks was reduced to 40% and all the blocks remained. At 24 weeks, 14.4% of bone was found in HA implants. Bone occupied 38% of the porous area (Figure 3).

DISCUSSION

In this study, more bone was produced in CC implants at 3 and 12 weeks than in HA, although at 12 weeks the CC blocks had been reduced to 40% of their original cross-sectional size. The absolute amount of bone did not increase significantly within either group after 3 weeks, but there was a trend of increased bone formation in HA implants at 24 weeks ($p = 0.051$), when the amount of bone was about the same as in CC




Figure 2 Bone ingrowth into CC and HA implants with bone marrow. Bone ingrowth in the implant is given as percentages of mean (\pm SEM) bone area related to original total cross-sectional area of implant. At 3 and 12 weeks the difference between materials is significant.

Table 2 Resorption of coral implants

| Weeks | Mean (range) | |
|-------|--------------|-------------|
| | CC (%) | control (%) |
| 3 | 87 (92–84) | 86 (93–77) |
| 6 | 62 (84–19) | 61 (88–0) |
| 12 | 40 (60–20) | |

Values are mean total cross-sectional areas in percentages related to original area before implantation. Maximum and minimum values are shown in parentheses.

implants at 12 weeks (14.4%/13.7%, Figure 2). As the structure of the two implants is similar, the difference in chemical constitution and in the resorption process of the CC matrix itself may affect bone formation.

Ohgushi *et al.* implanted CC and HA discs with bone marrow into subcutaneous pouches in rats. At 4 weeks, bone occupied 19.1% (HA) and 22.0% (CC) of the porous area²⁹. Our results for intramuscular sites at 3 weeks are 13.4% (HA) and 20.1% (CC). Ohgushi *et al.* found bone only in blocks immersed with bone marrow, not in those without it. This is in accordance

with our findings here and also with previous reports where porous ceramics have been used with bone marrow in extraosseal sites in rats^{26,28}. Van Eeden and Ripamonti recently reported bone formation in porous HA implanted intramuscularly without any osteoinductive factor. Bone was found only in HA blocks but not in granular form. They used baboons in the experiment and postulated that the geometry of the implant is critical. Unfortunately, primates are seldom used in experimental work and further study is needed to explain the bone formation mechanism³¹.

When HA implants without bone marrow have been placed in bone defects, bone ingrowth into HA has been variable. In orthognatic patients, Holmes *et al.* found 18% bone invasion (compared to total measured area) in a follow-up time of 4.7–16.4 (mean 9.3) months in HA implants³². Martin *et al.* placed HA blocks in cortical defects of dog humerus and radius. Bone ingrowth into pores increased from 52% at 16 weeks to 74% at 1 year (amount of bone relative to porous space). In the cancellous site, bone ingrowth was 38% at 4 weeks, then fell to 17% at 1 year³³. It seems that the entire porous space of HA is seldom completely filled with bone¹⁵.

Bone ingrowth into CC is more difficult to measure because of resorption. CC loses its internal porous structure very quickly, in our study after 6 weeks, and after that the bone does not actually invade the pores but replaces the matrix. When implants of CC have been used in posterior vertebral fusions in children or in animal experimental models in bone defects, new bone forms in previous implant sites^{7,24}. Histology has been studied but quantitative data are insufficient.

An ideal bone substitute should resorb gradually after being invaded by bone and allow the new bone to remodel. HA allows bone ingrowth but it does not resorb significantly³. HA blocks and granules used in tibial plateau fractures and bone tumour surgery are well detectable in X-rays after many years^{21,23}. The clinical use of HA is thus limited, because it lacks the ability to remodel the basic quality of living bone.




Figure 3 Tissue ingrowth into implants with bone marrow. Relative amounts of matrix and different tissue components and relative sizes of implants are presented for different follow-up periods. 100% is original total cross-sectional area of implant. Left side: coral implants; right side: hydroxyapatite implants.

In CC, the resorption rate may depend on the animal species used. When CC blocks were implanted in the cortex of the femur and tibia in pig and sheep, resorption at 1 month was 64 and 93%, respectively³⁴; Naaman *et al.* observed complete resorption of CC granules at 24 weeks in connective tissue site in pigs. No inducing material was used and the blocks did not have contact with bone tissue³⁵. In humans, degradation seems to be slower, 50% of the blocks used to fill the 10 mm cranial burr holes having resorbed completely at 1 year. With larger implants the resorption rate was much slower²⁰. In vertebral fusions, small fragments of natural coral blocks were found after 1 year²⁴.

In our experiment the resorption varied widely, one block in the control group being already completely resorbed at 6 weeks, whereas the largest implant in the same group still retained 88% of its original size. The resorption pattern was noteworthy. At 12 weeks bone marrow occupied the central part of the implant and was surrounded by lamellar bone. The blocks were deformed and consisted of only 23% of matrix (originally 58%). Active bone formation continued and the implant resembled an ossicle (Figure 1c). It has previously been postulated that the main factor in the resorption process is carbonic anhydrase enzyme, abundant in osteoclasts⁸. The control implants did not show any bone formation but the average resorption rate was very similar in both coral groups. Giant cells were also found in the control specimens but whether these cells were osteoclasts or not could not be established.

When porous hydroxyapatite or natural coral are used as bone substitutes, quick and effective bone formation is advantageous, in particular for calcium carbonate. A bone substitute that resorbs before bone has filled the pores is useless. This favours the use of some inducing factor. At the moment autogenous bone marrow is the most readily available inducing material, and more use could be made of it in clinical applications to induce bone formation inside porous CC or HA implants.

ACKNOWLEDGEMENTS

Thus study was supported by Paulo Foundation (J. Vuola) and by Finska Läkarsällskapet (T. Böhling).

REFERENCES

- Holmes RE. Bone regeneration within a coralline hydroxyapatite implant. *Plast Reconstr Surg* 1979; **63**: 626–633.
- Roy DM, Linnehan SK. Hydroxyapatite formed from coral skeletal carbonate by hydrothermal exchange. *Nature* 1974; **247**: 220–222.
- White E, Shors EC. Biomaterial aspects of Interpore-200 porous hydroxyapatite. *Dent Clin North Am* 1986; **30**: 49–67.
- Hulbert SF, Young FA, Mathews RS, Klawitter JJ, Talbert CD, Stelling FH. Potential of ceramic materials as permanently implantable skeletal prostheses. *J Biomed Mater Res* 1970; **4**: 433–456.
- Chiroff RT, White EW, Weber KN, Roy DM. Tissue ingrowth of Replamineform implants. *J Biomed Mater Res* 1975; **9**: 29–45.
- Daculsi G, Passuti N. Effect of the macroporosity for osseous substitution of calcium phosphate ceramics. *Biomaterials* 1990; **11**: 86–87.
- Guillemin G, Patat JL, Fournie J, Chetail M. The use of coral as a bone graft substitute. *J Biomed Mater Res* 1987; **21**: 557–567.
- Guillemin G, Fournie J, Patat JL, Chetail M. [Fate of a fragment of madreporic coral skeleton implanted in the diaphysis of long bones in dogs]. *C R Seances Acad Sci III* 1981; **293**: 371–376.
- Jarcho M. Calcium phosphate ceramics as hard tissue prosthetics. *Clin Orthop* 1981; **157**: 259–278.
- Doherty MJ, Schlag G, Schwartz N, Mollan RA, Nolan PC, Wilson DJ. Biocompatibility of xenogeneic bone, commercially available coral, a bioceramic and tissue sealant for human osteoblasts. *Biomaterials* 1994; **15**: 601–608.
- Meffert RM, Thomas JR, Hamilton KM, Brownstein CN. Hydroxylapatite as an alloplastic graft in the treatment of human periodontal osseous defects. *J Periodontol* 1985; **56**: 63–73.
- Kenney EB, Lekovic V, Carranza FJ, Dimitrijevic B, Han T, Takei H. A comparative clinical study of solid and granular porous hydroxylapatite implants in human periodontal osseous defects. *J Biomed Mater Res* 1988; **22**: 1233–1243.
- Issahakian S, Ouhayoun JP. [Clinical and histological evaluation of a new filling material: natural coral]. *J Parodontol* 1989; **8**: 251–259.
- Salzer KE, Hall CD. Porous hydroxyapatite as an onlay bone-graft substitute for maxillofacial surgery. *Plast Reconstr Surg* 1989; **84**: 236–244.
- Rosen HM, McFarland MM. The biological behavior of hydroxyapatite implanted into the maxillofacial skeleton. *Plast Reconstr Surg* 1990; **85**: 718–723.
- Levet, Y, Guero S, Guillemin G, Jost G. [Use of coral as replacement for bone grafts in facial surgery. 4 years' follow-up]. *Ann Chir Plast Esthet* 1988; **33**: 278–282.
- Byrd HS, Hobar PC, Shewmake K. Augmentation of the craniofacial skeleton with porous hydroxyapatite granules. *Plast Reconstr Surg* 1993; **91**: 15–22.
- Honig JF, Merten HA. Subperiosteal versus epiperiosteal forehead augmentation with hydroxylapatite for aesthetic facial contouring: experimental animal investigation and clinical application. *Aesthetic Plast Surg* 1993; **17**: 93–98.
- Waite PD, Morawetz RB, Zeiger HE, Pincock JL. Reconstruction of cranial defects with porous hydroxylapatite blocks [see comments]. *Neurosurgery* 1989; **25**: 214–217.
- Roux FX, Brasnu D, Loty B, George B, Guillemin G. Madreporic coral: a new bone graft substitute for cranial surgery. *J Neurosurg* 1988; **69**: 510–513.
- Bucholz RW, Carlton A, Holmes R. Interporous hydroxyapatite as a bone graft substitute in tibial plateau fractures. *Clin Orthop* 1989; **240**: 53–62.
- Kamegaya M, Shinohara Y, Shinada Y, Moriya H, Koizumi W, Tsuchiya K. The use of a hydroxyapatite block for innominate osteotomy. *J Bone Joint Surg Br* 1994; **76**: 123–126.
- Uchida A, Araki N, Shinto Y, Yoshikawa H, Kurisaki E, Ono K. The use of calcium hydroxyapatite ceramic in bone tumour surgery. *J Bone Joint Surg Br* 1990; **72**: 298–302.
- Pouliquen JC, Noat M, Verneret C, Guillemin G, Patat JL. [Coral substituted for bone grafting in posterior vertebral arthrodesis in children. Initial results]. *Rev Clin Orthop* 1989; **75**: 360–369.

- 25 Patel A, Honnart F, Guillemin G, Patat JL. [Use of madreporaria coral skeletal fragments in orthopedic and reconstructive surgery: experimental studies and human clinical application (author's transl.)]. *Chirurgie* 1980; **106**: 199–205.
- 26 McDavid PT, Boone MW, Kafrawy AH, Mitchell DF. Effect of autogenous marrow and calcitonin on reactions to a ceramic. *J Dent Res* 1979; **58**: 1478–1483.
- 27 Nade S, Armstrong L, McCartney E, Baggaley B. Osteogenesis after bone and bone marrow transplantation. The ability of ceramic materials to sustain osteogenesis from transplanted bone marrow cells: preliminary studies. *Clin Orthop* 1983; **181**: 255–263.
- 28 Okumura M, Ohgushi H, Tamai S. Bonding osteogenesis in coralline hydroxyapatite combined with bone marrow cells. *Biomaterials* 1991; **12**: 411–416.
- 29 Ohgushi H, Okumura M, Yoshikawa T et al. Bone formation process in porous calcium carbonate and hydroxyapatite. *J Biomed Mater Res* 1992; **26**: 885–895.
- 30 European Convention for the Protection of Vertebrate Animals Used for Experimental and Other Scientific Purposes. Council of Europe, Strasbourg. *European Treaty Series* 1987; **No. 123**.
- 31 van Eeden SP, Ripamonti U. Bone differentiation in porous hydroxyapatite in baboons is regulated by the geometry of the substratum: implications for reconstructive craniofacial surgery. *Plast Reconstr Surg* 1994; **93**: 959–965.
- 32 Holmes RE, Wardrop RW, Wolford LM. Hydroxylapatite as a bone graft substitute in orthognathic surgery: histologic and histometric findings. *J Oral Maxillofac Surg* 1988; **46**: 661–671.
- 33 Martin RB, Chapman MW, Sharkey NA, Zissimos SL, Bay B, Shors EC. Bone ingrowth and mechanical properties of coralline hydroxyapatite 1 yr after implantation. *Biomaterials* 1993; **14**: 341–348.
- 34 Guillemin G, Meunier A, Dallant P, Christel P, Pouliquen JC, Sedel L. Comparison of coral resorption and bone apposition with two natural corals of different porosities. *J Biomed Mater Res* 1989; **23**: 765–779.
- 35 Naaman BAN, Patat JL, Guillemin G, Issahakian S, Forest N, Ouhayoun JP. Evaluation of the osteogenic potential of biomaterials implanted in the palatal connective tissue of miniature pigs using undecalcified sections. *Biomaterials* 1994; **15**: 201–207.