

Alzheimer's Disease Is Associated With a Selective Increase in $\alpha 7$ Nicotinic Acetylcholine Receptor Immunoreactivity in Astrocytes

THANASAK TEAKTONG,¹ ALISON GRAHAM,¹ JENNIFER COURT,¹
ROBERT PERRY,² EVELYN JAROS,² MARY JOHNSON,¹ ROS HALL,¹
AND ELAINE PERRY^{1*}

¹MRC Building, Centre Development in Clinical Brain Aging, Newcastle General Hospital,
Newcastle Upon Tyne, U.K.

²Department of Neuropathology, Newcastle General Hospital, Newcastle Upon Tyne, U.K.

KEY WORDS cholinergic receptors; dementia; limbic system; parahippocampal gyrus; neuroglia

ABSTRACT Alzheimer's disease (AD) and dementia with Lewy bodies (DLB) are common forms of dementia in the elderly associated with cholinergic dysfunction, including reductions in nicotinic acetylcholine receptors (nAChRs). In AD, astrocytes are implicated in the formation of senile plaques, one of the core pathological features. Using immunohistochemistry, we have investigated astrocytic expression of the two major nicotinic receptor α subunits in the human hippocampus and entorhinal cortex. $\alpha 7$, but not $\alpha 4$, subunit immunoreactivity was associated with astrocytes. An increase in the proportion of astrocytes expressing $\alpha 7$ immunoreactivity was observed in AD compared with age-matched controls. A similar increase was not evident in DLB. Elevated $\alpha 7$ nAChRs on astrocytes in AD may contribute to alterations in calcium homeostasis and nitric oxide production, which in turn could affect β -amyloid-mediated inflammatory processes in AD. *GLIA* 41:207–211, 2003. © 2003 Wiley-Liss, Inc.

Alzheimer's disease (AD) and dementia with Lewy bodies (DLB) (McKeith et al., 1996; Winblad et al., 2001) together account for more than half of the dementia in the elderly. Both disorders are affected by pathology associated with impaired cholinergic function, including deficits in brain nicotinic acetylcholine receptors (nAChRs) in cortical and other brain areas (Court et al., 2001). nAChRs are ligand-gated cation channels that are expressed not only on neurons, but also on astrocytes where they participate in calcium signaling (Sharma and Vijayaraghavan, 2001). Astrocytes are likely to be involved in the pathology of AD; they surround senile plaques (Kato et al., 1998), appear to promote amyloid plaque maturation (Terai et al., 2001; Wegiel et al., 2001), and can induce neurotoxicity cascades involving many inflammatory mediators (Sutton et al., 1999; Combs et al., 2000). DLB is characterized by α -synuclein-positive Lewy bodies and neurites in the cerebral cortex and brain stem and with cortical

senile plaques to a lesser degree than in AD (Ince et al., 1991). The present study explores the possible involvement of nicotinic receptors in astrocytosis in AD and makes comparison with DLB. The expression of the two major nAChR α subunits ($\alpha 4$ and $\alpha 7$) (Gotti et al., 1997) was investigated in the hippocampus and entorhinal cortex using immunohistochemistry.

Autopsy samples of temporal cortex were obtained from six cases of AD (age, 83.7 ± 2.7 years), six cases of DLB (age, 76.5 ± 3.1 years), and four age-matched controls (age, 82.5 ± 9.5 years). Mean postmortem de-

Grant sponsor: Janssen Pharmaceutical Research Foundation.

*Correspondence to: Elaine Perry, MRC Building, Centre Development in Clinical Brain Aging, Newcastle General Hospital, Westgate Road, Newcastle Upon Tyne, NE4 6BE, U.K. E-mail: e.k.perry@ncl.ac.uk

Received 2 May 2002; Accepted 15 July 2002

DOI 10.1002/glia.10132




Fig. 1. $\alpha 7$ -immunoreactive astrocytes in CA1 (A–C), CA4 (D–F), and entorhinal cortex layer 3 (G–I) of control (A, D, and G), Alzheimer (B, E, and H), and DLB (C, F, and I) cases. $\alpha 7$ -immunoreactive astrocytes were found in all groups but were significantly increased in AD cases compared with controls. Arrows indicate $\alpha 7$ -immunoreactive astrocytes. Scale bars, 20 μ m. [Color figure can be viewed in the online issue, which is available at www.interscience.wiley.com].

lays of control, AD, and DLB cases were 46.0 ± 18.9 , 58.8 ± 24.5 , and 41.2 ± 33.4 h, respectively. We have previously shown that this range of postmortem delay does not affect nAChR immunohistochemistry (Graham et al., 2002). None of the controls smoked tobacco or had any history of a neurological or psychiatric disease or significant neuropathological abnormality. AD and DLB were diagnosed according to neuropathological examination and clinical diagnostic criteria (Perry et al., 1990; McKeith et al., 1996). Ten μ m thick formalin-fixed paraffin-embedded sections were immunostained with monoclonal antibodies against the $\alpha 4$ (mAb 299, 1:8,000; Cambridge Bioscience, Cambridge, U.K.) and $\alpha 7$ nAChR subunits (mAb 306, 1:4,000; Cambridge Bioscience) and with polyclonal antiglial fibrillary acidic protein (GFAP; 1:4,000; Dako, Cambridge, U.K.) as a marker of astrocytes, using the Vectastain Elite kit method. The specificity of the monoclonal antibodies has been established elsewhere; for mAb 299, Whiting and Lindstrom (1988), Peng et al.

(1994), and Schroder et al. (2001); for mAb 306, Schoepfer et al. (1990), McLane et al. (1992), Burghaus et al. (2000). For double labeling of $\alpha 7$ and GFAP, antibodies were applied sequentially using the anti- $\alpha 7$ followed by anti-GFAP. Controls from which primary antibodies were omitted showed no immunoreactivity. Controls in which the anti-GFAP was replaced by rabbit IgG confirmed no cross-reactivity between the two secondary antisera. The percentage of $\alpha 7$ -immunoreactive astrocytes was assessed in double-stained sections using a light microscope at $20\times$ magnification in CA1, CA4, and entorhinal cortex. The entorhinal cortex was subdivided into three layers: layer 1 corresponded to the superficial acellular layer, layer 2 the layer of islands of small pyramidal cells, and layer 3 the layers of medium and large pyramidal cells extending below layer 2 to the white matter. Group differences were evaluated using the Mann-Whitney test.

$\alpha 7$ immunoreactivity was present on astrocytes in all three groups (Figs. 1 and 2A, C, and E). In controls, the




Fig. 2. Immunohistochemistry of $\alpha 7$ nAChR and GFAP double labeling (A, C, and E) and $\alpha 4$ nAChR (B, D, and F) in CA4 of control (A and B), Alzheimer (C and D), and DLB (E and F) cases. Red and blue/gray stainings show $\alpha 7$ and GFAP immunoreactivity, respectively. Both $\alpha 7$ and GFAP immunoreactivity were increased in Alzheimer's disease but not DLB compared with controls. Arrows indicate double-immunolabeled astrocytes in control, AD, and DLB. No $\alpha 4$ immunoreactive astrocytes were found in all three groups. Scale bars, 20 μ m.

percentage of $\alpha 7$ -labeled astrocytes varied between 0% and 54% depending on area, being low in CA1 and layer 3 of the entorhinal cortex and highest in CA4 and entorhinal cortex layer 1 (which contained the pial

surface; Table 1), possibly reflecting regional differences in astrocytic function or activation. In DLB, a similar pattern was observed in controls. However, in AD, the percentage of $\alpha 7$ -immunoreactive astrocytes

TABLE 1. Percentage of $\alpha 7$ and GFAP Double-Labeled Compared With the Total Number of GFAP-Labeled Astrocytes

Brain area	Control	AD	DLB
Pyramidal cell layer			
CA1	12.5 \pm 25	61.1 \pm 9.6 ^a	11.1 \pm 19.2
CA4	37.5 \pm 8.4	80.6 \pm 17.3 ^b	25.0 \pm 43.3
Entorhinal cortex			
Layer 1	54.2 \pm 8.4	100.0 \pm 0.0 ^b	77.8 \pm 38.5
Layer 2	0.0 \pm 0.0	83.3 \pm 28.9 ^b	16.7 \pm 28.9
Layer 3	20.8 \pm 25.0	61.1 \pm 9.6 ^a	22.2 \pm 19.2

^a $P < 0.05$.

^b $P < 0.01$, compared to control.

was markedly greater in the hippocampus and entorhinal cortex compared with control and DLB cases. In contrast to the large numbers of $\alpha 7$ -reactive astrocytes found in the hippocampus and entorhinal cortex in AD, no $\alpha 4$ -immunoreactive astrocytes were observed in these areas in AD, controls, or DLB (Fig. 2B, D, and F). As expected, astrocytes assessed by GFAP immunoreactivity were more numerous in most subfields of the hippocampal formation of AD compared with both DLB and control groups; however, in CA4 (Fig. 2A, C, and E), astrocyte density was similar between groups.

Wevers et al. (1999) also noted $\alpha 7$ expression on cells with astrocytic profiles in the frontal cortex in AD, some associated with amyloid plaques, but not in age-matched controls. That $\alpha 7$ expression on astrocytes is greater in both controls and AD cases in hippocampus and entorhinal cortex than the neocortex is likely to reflect the greater age-related pathology of the archicortex. The upregulation of mRNA for $\alpha 7$ observed by Hellstrom-Lindahl et al. (1999) in hippocampus but not in the temporal cortex or cerebellum in AD is consistent with such a mechanism. The present findings contrast with the net reduction in $\alpha 7$ protein expression and α -bungarotoxin binding previously observed in some studies in the hippocampus in AD (Court et al., 2001). Nevertheless, there was no relationship between increased $\alpha 7$ astrocytic immunoreactivity in entorhinal cortex and the severity of β -amyloidosis as indicated by the numbers of neocortical plaques stained histologically (Perry et al., 1990) in either AD or DLB ($r = 0.098$, $P = 0.854$ in AD; $r = -0.099$, $P = 0.852$ in DLB).

The increased expression of $\alpha 7$ on astrocytes in AD could be in response to the reduction in acetylcholine release. However, if this were the case, a similar increase in astrocytic $\alpha 7$ expression would be expected to occur in DLB in which cholinergic innervation is equally reduced (Tiraboschi et al., 2000). Beta-amyloid induces many proinflammatory cytokines such as interleukin- 1β (IL- 1β) and tumor necrosis factor α (TNF α) (Sutton et al., 1999; Szczepanik et al., 2001), which can induce nitric oxide production and release from astrocytes (Casamenti et al., 1999; Pazmany et al., 1999). Astrocytic expression of the calcium-dependent NOS enzymes is reported to be increased in both archi- and neocortex in AD (de la Monte and Bloch, 1997; Simic et al., 2000). However, if β -amyloid accumulation is the sole mechanism whereby astrocytic $\alpha 7$ expression is upregulated in AD, it is surprising that

this does not also occur in DLB. It is possible that the difference observed between AD and DLB at least in part reflects distinct pathological processes in the two disorders.

Stimulation of $\alpha 7$ nAChRs on astrocytes has been shown to increase intracellular calcium released from intracellular stores (Sharma and Vijayaraghavan, 2001), hence elevated expression of $\alpha 7$ nAChR on astrocytes may lead to abnormally high levels of intracellular calcium, which might contribute to a number of inflammatory cascades.

Further studies are required to explore the potential mechanisms of astrocytic $\alpha 7$ nAChR upregulation in AD and the contribution that this might make to AD pathology.

ACKNOWLEDGMENTS

The authors thank Mrs. Jean Dawes for demographic data of patients and Professor Mondhon Sanguanserm Sri (Naresuan University, Thailand) for his support.

REFERENCES

- Burghaus L, Schutz U, Krempel U, de Vos RA, Jansen Steur EN, Wevers A, Lindstrom J, Schroder H. 2000. Quantitative assessment of nicotinic acetylcholine receptor proteins in the cerebral cortex of Alzheimer patients. *Mol Brain Res* 76:385–388.
- Casamenti F, Prosperi C, Scali C, Giovannelli L, Colivicchi MA, Fausone-Pellegrini MS, Peepeu G. 1999. Interleukin- 1β activates forebrain glial cells and increases nitric oxide production and cortical glutamate and GABA release in vivo: implications for Alzheimer's disease. *Neuroscience* 91:831–842.
- Combs CK, Johnson DE, Karlo JC, Cannady SB, Landreth GE. 2000. Inflammatory mechanisms in Alzheimer's disease: inhibition of β -amyloid-stimulated proinflammatory responses and neurotoxicity by PPAR γ agonists. *J Neurosci* 20:558–567.
- Court J, Martin-Ruiz C, Piggott M, Spurdin D, Griffiths M, Perry E. 2001. Nicotinic receptor abnormalities in Alzheimer's disease. *Biol Psychiatry* 49:175–184.
- de la Monte SM, Bloch KD. 1997. Aberrant expression of the constitutive endothelial nitric oxide synthase gene in Alzheimer disease. *Mol Chem Neuropathol* 30:139–159.
- Gotti C, Fornasari D, Clementi F. 1997. Human neuronal nicotinic receptors. *Prog Neurobiol* 53:199–237.
- Graham A, Court J, Jaros E, Perry R, Volsen S, Ince P, Kuryatov A, Lindstrom J, Perry E. 2002. Immunohistochemical localization of nicotinic acetylcholine receptor subunits in human cerebellum. *Neuroscience* 113:493–507.
- Hellstrom-Lindahl E, Mousavi M, Zhang X, Ravid R, Nordberg A. 1999. Regional distribution of nicotinic receptor subunit mRNAs in human brain: comparison between Alzheimer and normal brain. *Mol Brain Res* 66:94–103.
- Ince P, Irving D, MacArthur F, Perry RH. 1991. Quantitative neuropathological study of Alzheimer-type pathology in the hippocampus: comparison of senile dementia of Alzheimer type, senile dementia of Lewy body type, Parkinson's disease and non-demented elderly control patients. *J Neurol Sci* 106:142–152.
- Kato S, Gondo T, Hoshii Y, Takahashi M, Yamada M, Ishihara T. 1998. Confocal observation of senile plaques in Alzheimer's disease: senile plaque morphology and relationship between senile plaques and astrocytes. *Pathol Int* 48:332–340.
- McKeith IG, Galasko D, Kosaka K, Perry EK, Dickson DW, Hansen LA, Salmon DP, Lowe J, Mirra SS, Byrne EJ, Lennox G, Quinn NP, Edwardson JA, Ince PG, Bergeron C, Burns A, Miller BL, Lovestone S, Collerton D, Jansen EN, Ballard C, de Vos RA, Wilcock GK, Jellinger KA, Perry RH. 1996. Consensus guidelines for the clinical and pathologic diagnosis of dementia with Lewy bodies (DLB):

- report of the consortium on DLB international workshop. *Neurology* 47:1113–1124.
- McLane KE, Wu X, Lindstrom JM, Conti-Tronconi BM. 1992. Epitope mapping of polyclonal and monoclonal antibodies against two α -bungarotoxin-binding α subunits from neuronal nicotinic receptors. *J Neuroimmunol* 38:115–128.
- Pazmany T, Mechtler L, Tomasi TB, Kosa JP, Turoczy A, Urbanyi Z. 1999. Differential regulation of major histocompatibility complex class II expression and nitric oxide release by β -amyloid in rat astrocyte and microglia. *Brain Res* 835:213–223.
- Peng X, Gerzanich V, Anand R, Whiting PJ, Lindstrom J. 1994. Nicotine-induced increase in neuronal nicotinic receptors results from a decrease in the rate of receptor turnover. *Mol Pharmacol* 46:523–530.
- Perry RH, Irving D, Blessed G, Fairbairn A, Perry EK. 1990. Senile dementia of Lewy body type: a clinically and neuropathologically distinct form of Lewy body dementia in the elderly. *J Neurol Sci* 95:119–139.
- Schoepfer R, Conroy WG, Whiting P, Gore M, Lindstrom J. 1990. Brain α -bungarotoxin binding protein cDNAs and MAbs reveal subtypes of this branch of the ligand-gated ion channel gene superfamily. *Neuron* 5:35–48.
- Schroder H, Schutz U, Burghaus L, Lindstrom J, Kuryatov A, Monteggia L, deVos RA, van Noort G, Wevers A, Nowacki S, Happich E, Moser N, Arneric SP, Maelicke A. 2001. Expression of the $\alpha 4$ isoform of the nicotinic acetylcholine receptor in the fetal human cerebral cortex. *Dev Brain Res* 132:33–45.
- Sharma G, Vijayaraghavan S. 2001. Nicotinic cholinergic signaling in hippocampal astrocytes involves calcium-induced calcium release from intracellular stores. *Proc Natl Acad Sci USA* 98:4148–4153.
- Simic G, Lucassen PJ, Krsnik Z, Kruslin B, Kostovic I, Winblad B, Bogdanovic N. 2000. nNOS expression in reactive astrocytes correlates with increased cell death related DNA damage in the hippocampus and entorhinal cortex in Alzheimer's disease. *Exp Neurol* 165:12–26.
- Sutton ET, Thomas T, Bryant MW, Landon CS, Newton CA, Rhodin JA. 1999. Amyloid-beta peptide induced inflammatory reaction is mediated by the cytokines tumor necrosis factor and interleukin-1. *J Submicrosc Cytol Pathol* 31:313–323.
- Szczepanik AM, Rampe D, Ringheim GE. 2001. Amyloid- β peptide fragments p3 and p4 induce pro-inflammatory cytokine and chemokine production in vitro and in vivo. *J Neurochem* 77:304–317.
- Terai K, Iwai A, Kawabata S, Sasamata M, Miyata K, Yamaguchi T. 2001. Apolipoprotein E deposition and astrogliosis are associated with maturation of β -amyloid plaques in β APPswe transgenic mouse: Implications for the pathogenesis of Alzheimer's disease. *Brain Res* 900:48–56.
- Tiraboschi P, Hansen LA, Alford M, Sabbagh MN, Schoos B, Masliah E, Thal LJ, Corey-Bloom J. 2000. Cholinergic dysfunction in diseases with Lewy bodies. *Neurology* 54:407–411.
- Wegiel J, Wang K-C, Imaki H, Rubenstein R, Wronska A, Osuchowski M, Lipinski WJ, Walker LC, LeVine H. 2001. The role of microglial cells and astrocytes in fibrillar plaque evolution in transgenic APP(SW) mice. *Neurobiol Aging* 22:49–61.
- Wevers A, Monteggia L, Nowacki S, Bloch W, Schutz U, Lindstrom J, Pereira EFR, Eisenberg H, Giacobini E, de Vos RAI, Jansen Steur ENH, Maelicke A, Albuquerque EX, Schroder H. 1999. Expression of nicotinic acetylcholine receptor subunits in the cerebral cortex in Alzheimer's disease: histotopographical correlation with amyloid plaques and hyperphosphorylated-tau protein. *Eur J Neurosci* 11:2551–2565.
- Winblad B, Brodaty H, Gauthier S, Morris JC, Orgogozo J, Rockwood K, Schneider L, Takeda M, Tariot P, Wilkinson D. 2001. Pharmacotherapy of Alzheimer's disease: is there a need to redefine treatment success? *Int J Geriatr Psychiatry* 16:653–666.
- Whiting PJ, Lindstrom JM. 1988. Characterization of bovine and human neuronal nicotinic acetylcholine receptors using monoclonal antibodies. *J Neurosci* 8:3395–3404.