

Nitric Oxide: Target for Therapeutic Strategies in Alzheimer's Disease

A.P. Fernandez^{1,†}, A. Pozo-Rodrigalvarez, J. Serrano^{1,†} and R. Martinez-Murillo*

Department of Molecular, Cellular, and Developmental Neurobiology, Instituto Cajal, Spanish Council for Scientific Research (CSIC), 28002 Madrid, Spain

Abstract: Alzheimer's disease (AD) constitutes a progressive neurodegenerative disorder and the main cause of dementia. Numerous studies have focused on the pathogenic mechanism of AD to cure or prevent this devastating disease. But, despite recent advances, our understanding on the pathophysiology of this genetically complex and heterogeneous disorder is rather limited and treatment of the disease consists of medications to control the symptoms. Acetylcholinesterase inhibitors and N-methyl-D-aspartate (NMDA) receptor antagonists are the only available treatments recommended to manage the cognitive deficits caused by the disease. Therefore, the production of new drugs that may be able to cure the underlying cause of this chronic disease, not just the symptoms, is a matter of clinical interest. There is data implicating nitric oxide (NO) in the progression of the disease. The three isoforms of the NO-synthesizing enzyme (NOS) operate as central mediators of amyloid beta-peptide (A β) action, giving rise to elevated levels of NO that contributes to the maintenance, self-perpetuation and progression of the disease. Reducing A β production and the cholinergic deficit is a goal in the treatment of AD. In addition, a possible way to delay the progression of the illness must include a rationale design of enzyme inhibitors, subtype selective, targeting NOS isoforms implicated in damage to brain cells in AD.

We are now presenting an overview regarding approved drugs for AD treatment and substances that although are not in use for the treatment of AD, including NOS inhibitors, may represent useful tools to unravel the pathophysiologic enigma of AD.

Keywords: Alzheimer's disease, Nitric oxide, Nitric oxide synthase inhibitors, Acetylcholinesterase, Amyloid beta-peptide, Neuroinflammation, Neurodegeneration, Immunotherapy.

INTRODUCTION

Alzheimer's Disease: General Topics

Alzheimer's disease (AD), first described by Alois Alzheimer in 1907 [1], constitutes the most common progressive neurodegenerative disorder and the main cause of dementia [2]. This chronic disorder syndrome evolves with loss of neurons and their synapses and with cognitive deficits: memory, intellectual and emotional dysfunctions [3]. The disease appears usually associated with characteristic risks factors including old age, family history, apolipoprotein E epsilon 4 genotype, angiopathy, etc. [4]. Environmental factors and/or epigenetic phenomena may also contribute to AD pathology and phenotypic expression of dementia [5, 6].

Numerous studies have focused on the pathogenic mechanism of AD but, despite recent advances, our understanding on the pathophysiology of this genetically complex and heterogeneous disorder is rather limited and the currently available treatments approved by the US Food and Drug Administration and the European Medicines Agency do not halt progression or cure the illness. Histopathologically, the brains of AD patients share specific characteristics, namely amyloid plaques and neurofibrillary tangles (NFT), which are insoluble deposits made of an abnormally folded, fibrillar form of the amyloid beta-peptide (A β), as shown in Figs. (1-2), and hyperphosphorylated Tau, respectively. A β peptides are composed of 39–43 amino acids derived from the proteolytic processing of the amyloid precursor protein (APP) by β and γ secretases that give rise two structural variants, A β 1-40 and A β 1-42 [7]. Characteristically, the ratio of these structural variants differed by 10-fold between brains from nondemented controls and those with sporadic AD [8]. High ratio of the structural variants A β 1-40 and A β 1-42 is considered a biomarker of AD




Fig. (1). A β expression and distribution in the cerebral temporal cortex from postmortem brain of CDR3 Alzheimer's disease patient showing immunoreactive plaques in all cortical layers. Scale bar: 100 μ m.

*Address correspondence to this author at the Department of Cellular, Molecular, and Developmental Neurobiology, Instituto Cajal, Spanish Council for Scientific Research (CSIC), Avda. Doctor Arce 37, 28002-Madrid, Spain; Tel: 34 + 915854714; Fax: 34 + 915854754; E-mail: r.martinez@cajal.csic.es

[†]Both authors contributed equally to this study and should be considered first authors.

[9]. Although it is not clear whether abnormal processing of the APP is an initial cause or rather a late event in the pathophysiology of AD [10], intraneuronal NFT together with accumulation of A β and neurotoxicity constitutes the major neuropathological hallmark of AD [11, 12]. Overwhelming evidence indicates that the progressive abnormal deposits of insoluble aggregates of A β within the neuropil and in the cerebrovascular walls (amyloid angiopathy) is the basis of the formation of senile plaques (amyloids) containing massive quantities of A β structural variants [8, 13] that causes neuronal injury and loss, inflammation and characteristic activation of microglia and astrocytic cells that occurs in AD [14-17], as shown in Figs. (3-5).

A growing body of evidence has suggested a critical role for APP metabolism and oxidative stress in the pathogenesis and advancement of the disease [18-22]. Together with aberrant A β -deposits in the neuropil originated in an anomalous proteolytic processing of APP [23], a genetic basis is essential in influencing to onset and/or modifying the progression of the disease [24, 25]. Early-onset dominant/familial AD (FAD) and the late-onset/sporadic AD (LAD) share identical pathology, thus suggesting common pathogenic pathways for both forms of AD. Mutations of three genes that influence the accumulation of the A β have been recognized tightly linked to FAD [10], which include the APP and the catalytic components of the γ secretase complex, presenilins 1 (PS1) and 2 (PS2) that are responsible for the intramembranous proteolysis of APP. The e4 allele of apolipoprotein E (ApoE), contains a polymorphism which increases the risk to develop sporadic AD [26]. In addition, genomic factors induced by environmental factors and epigenetic phenomena, might be responsible for AD pathogenesis leading to premature neuronal death [5, 6]. Extracellular insoluble aggregates of long A β 1-42, known to induce neurotoxicity [27], are exhibited by both FAD and LAD patients together with intracellular NFT [28] triggered by A β through the action of protein kinases [29-31]. APP was the first gene associated with FAD and the recent description of an extra copy of the APP gene in FAD [32] provided further support that increased A β production can primarily drive the disease. Abnormal APP and presenilins linked to AD lead invariably to the production of insoluble amyloidogenic form A β 1-42 peptide in a pathological manner [33-36], which strongly supports the amyloid hypothesis of AD [37]. A β also mediates synaptic dysfunction leading to cognitive deficits [38-40], which correlates very well with synapse loss [41, 42]. As a result of anomalous deposits accumulation of A β , neuronal damage and dysfunction occur through the activation of microglia and astrocytic cells [43], synaptic failure [44] and associated amyloid angiopathy [45, 46]. The current A β hypothesis suggests that the soluble oligomers can disrupt synaptic connections between neurons and, simultaneously, may aggregate in amyloid plaques which can trigger a local inflammatory response [47] essential in the progression of the disease [48]. It has been reported an increased expression of the pro-apoptosis gene products p53 and Bax associated with A β deposits in senile plaques [49-51]. We have also found elevated expression of a mutated form of p53 in older AD mice, suggesting an alternative mechanism for the predisposition of AD brains to develop brain neoplasms [52]. An established mechanism of cell loss in both neurons and glial cells in AD involves Bax and p53-mediated apoptosis [53, 54]. Both p53 and Bax inhibit the expression of Bcl-2, a cell survival gene product, thereby permitting Bax homodimers to enter the nucleus, promote cytochrome c release into the cytoplasm, and activate caspases. Caspases cause proteolytic damage, cytoskeletal collapse, and nuclease release followed by DNA fragmentation. Therefore, A β depletes the neurons of one of its anti-apoptotic mechanisms downregulating Bcl-2 [55]. It has been also proved that p53 upregulation in AD induces tau-phosphorylation [56]. It remains to be determined as to whether A β -deposits directly induces apoptotic cell death or triggers an alternative pathway, *via* nitric oxide (NO) systems, that then leads to degeneration [57, 58].




Fig. (2). Confocal microscopy microphotograph of a histological section through the frontal cerebral cortex of a postmortem brain of an Alzheimer's disease patient, CDR3 case, showing double immunostaining for A β (red colour), and for nNOS (green colour). Notice that nNOS immunopositive neurons are in close relation with senile plaques. Scale bar: 20 μ m.

As stated above, the generation of A β and other fragments requires sequential intramembrane cleavages of the APP by β - and γ -secretases. The γ -secretase complex operates as a cleaving multiprotein complex that cleaves APP and a large number of other type I integral transmembrane proteins including Notch-1, ErbB4, neuregulin, E- and N-cadherines, CD44, Low-density lipoprotein (LDL)-receptor related protein (LRP), nectin-1 and growth hormone receptor [59]. The Notch receptor undergoes a series of proteolytic events, in a manner similar to APP, ultimately releasing the intracellular domain that mediated the expression of genes controlling many types of cellular differentiation [60]. The first step in the production of A β peptides from APP is initiated by the β -site APP-cleaving enzyme 1 (BACE-1) [47] that catalyzes the rate-limiting reaction in the generation of A β [61]. β -Secretase cleavage of APP occurs predominantly in endosomes, and endocytosis of APP and β -secretase is essential for β cleavage and A β production [62-64]. Genetic dysregulation of APP near the β -secretase cleavage site make APP a better substrate for this enzyme and lead to increased production of all variants of A β . However, mutations near γ -secretase cleavage site lead to increase in the more aggregation-prone A β 1-42 relative to A β 1-40. Mutations in the A β itself change the biophysical properties of the peptide to render it more likely to aggregate [65]. The intramembrane γ -secretase complex includes presenilins together with Nicastrin, Aph-1 and presenilin enhancer 2 [66-74]. Alterations in the presenilin genes appear to be the major cause of FAD [75]. Presenilins are involved not only in the processing of APP but also in a number of pathways for cell death and survival modulating various cell signal transduction pathways as well [74, 76, 77], including the intracellular calcium signalling [78-80]. In addition, presenilins regulate phospholipase C and consequently Protein kinase C activity [81]. Therefore, binding of presenilins to APP also plays a central role inducing intracellular signalling [82]. In this regard, genetic mutation of presenilins have been associated with calcium signaling abnormalities that might play an early proximal and perhaps central role in AD pathogenesis [83, 84].

It has been also proposed that PS1 has a role for NO synthase (NOS) activation in neurons and confers oxidative stress-resistance on neurons in a calcium/NO-dependent manner [67]. Thus, oxida-

tive stress plays a critical role in the pathogenesis of AD related with APP proteolytic dysfunction.

Reducing A β production is a goal in the treatment of AD. Both β - and γ -secretase are thus propitious therapeutic targets [85, 86] but may be detrimental for AD therapy. It has been shown that nonselective γ -secretase inhibitors exert deleterious effects on embryogenesis in zebrafish and on lymphoid and gastrointestinal tissues in mammals [87, 88], and also may impair neurogenesis. Numerous neurogenic players, such as Notch-1, are cleaved by PS1/ γ secretase. In this context, PS1 familial AD mutation leads to defective associative learning and impairs adult neurogenesis in PS1M146V knockin mice [89]. In view of the multiple functions of γ -secretase, β -secretase might then be considered as the preferred therapeutic target [88]. A β plaques induce BACE-1 in neighbouring neurons at early stages of disease, suggesting that BACE-1 induction triggered by the amyloid pathway may drive a positive-feedback loop in AD [90-92]. In this regard, morphine that down-regulates the expression of BACE-1 *via* NO could be a protective instrument for AD [93]. In addition, compounds like pioglitazone and ibuprofen that decrease BACE-1 mRNA and protein levels in APPV717I transgenic mice [94] might be of potential use. Therefore, identification and characterization of secretases is a focus on interest in AD basic research and therapeutic use in clinic.

A number of neurotransmitter specific systems are implicated in AD, including the glutamatergic, cholinergic and nitrergic systems [95-104]. Typically, the cholinergic deficit is a trait of the AD. During the evolution of the disease, there is a conspicuous decrease in the cholinergic innervation of the brain, particularly in the hippocampus and neocortex, where the number of cholinergic elements and cholinergic activity dramatically decrease [105]. Based on the latter, drugs have been developed to repair the cholinergic deficit with the aim to obtain symptomatic benefits based on improving cholinergic function by increasing acetylcholine levels. In this field of research, many efforts are being performed to develop acetylcholinesterase (AChE) inhibitors (AChEIs) with clinical applications [106]. However, a recent meta-analysis on the efficacy of AChEIs indicates that these treatments can result in statistically significant but clinically marginal improvement [107]. Therefore, a "causative therapy" should be promoted. The discovery of new drugs to modify the pathological steps leading to AD by acting on the evolution of the disease, represents a paradigm shift away from symptomatic treatment. By interfering with the pathogenic steps of AD, these products collectively termed as "disease-modifying" agents [108] might be able to interact with different therapeutic targets, thus blocking the progression and evolution of the disease, even pre-clinical. An update on disease-modifying drugs has been recently published [108]. Multifunctional disease-modifying agents that simultaneously inhibit the human AChE and the β A aggregation and that also display neuroprotective properties against the mitochondrial oxidative stress are now being tested [109]. In this regard, this review provides information concerning AChEIs targeting NO metabolism.

The Nitrergic System in Alzheimer's Disease

The short-lived free radical messenger NO functions as an atypical diffusible neurotransmitter in an autocrine/paracrine manner in different tissues [110, 111]. NO is involved in a multitude of inter- and intra-cellular signalling pathways participating in considerable physiological and pathophysiological scenarios such as blood pressure control, neurotransmission, learning and memory, and as a defensive cytotoxin [99]. This highly diffusible gas, that crosses the biological membranes without any difficulty, plays a central role in neurodegenerative diseases, including Parkinson's and Alzheimer's diseases [112]. NO is synthesized by a specific enzyme, NOS (NOS; EC 1.14.13.39), present in the mammalian brain in three different isoforms, two constitutive enzymes (i.e., neuronal, type-I, nNOS, and endothelial, type III, eNOS) and one inducible enzyme

(type II, iNOS) which is the catalyst of high-output pathway of NO production [113]. From the discovery by Vodovotz in 1996 that iNOS is localized in AD lesions, other studies confirmed this observation identifying iNOS protein in neurons, astrocytes, and microglia that are related with the anomalous A β deposits [12, 17, 94, 114-117], but at far lower incidence, extent, and intensity in brains from age-matched controls. Whereas nNOS and eNOS are expressed constitutively in normal brain, a widespread expression of iNOS in the central nervous system is considered pathologic. The oxidative stress caused by NO in the brain has been proposed as a pathogenic mechanism in AD [118, 119], being suggested that NO plays a central role in the cascade of events leading from enhanced intracellular A β accumulation to enhanced vulnerability to apoptosis and cell death [120]. NO can be scavenged in a rapid reaction with superoxide ($O_2^{\cdot-}$) to generate peroxynitrite (ONOO $^-$) [121], that may act as a potent source for oxidative stress in AD [12, 122, 123] and the production of nitrotyrosine. At the cellular level, the mitochondrion appears to be of importance when considering the deleterious effects of both A β and induced NO [124]. Generation of highly reactive nitrogen species secondary to high concentrations of NO causes irreversible inhibition of mitochondrial respiration and damage to various mitochondrial components *via* oxidizing reactions leading to cell death. Nitrotyrosine, that derives from peroxynitrite-mediated nitration of protein tyrosine residues [125, 126], is considered as an indicator of cell damage and inflammation secondary to overproduction of NO [122]. Nitration may impair protein function, thus leading to neuron degeneration [12, 123]. A progressive elevation of NO markers and protein nitration in reactive astrocytes and microglia, as shown in Figs (2-5), was reported in Tg2576 transgenic mice [123] and in clinically evaluated cases of AD in human [12, 127, 128]. This is in agreement with previous reports that A β can stimulate NO production from astrocytes [129], and that activated astrocytes are able to produce NO levels that are detrimental to neurons [130]. *In vitro* studies in rat glioma C6 cells support this conclusion showing that A β treatment leads to increased production of NO through iNOS activation, yielding cytotoxicity in these cells [131]. In this context, a primary possibility to explain how A β accumulation leads to neurotoxicity is that A β may triggers oxidative and/or nitrosative injury in both neurons and glial cells.




Fig. (3). Inflammation in AD. Microphotograph showing immunostaining for glial fibrillary acid protein (GFAP) in the cerebral cortex of Tg2576 transgenic mice. An amyloid plaque is surrounded by reactive astrocytes showing immunoreactions product. Scale bar: 25 μ m.




Fig. (4). Microphotograph showing immunostaining for nitrotyrosine in the cerebral cortex of Tg2576 transgenic mice. Nitrotyrosine immunoreactivity appears in stellate shaped cells, resembling reactive astrocytes, surrounding a senile plaque. Scale bar: 10 μ m.

There is evidence implicating NO as a mediator of glutamate-mediated neuronal death (glutamatergic excitotoxicity) *via* N-methyl-D-aspartate (NMDA) receptors. Influx of calcium followed by activation of nNOS may conduct to the generation of peroxynitrite, which has been implicated in the mechanism of neuron loss in neurodegenerative diseases. All the three isoforms of the NO-synthesizing enzyme are bizarrely expressed in AD models, giving rise to elevated levels of NO. However, it is controversial as to whether aberrant expression of NOS isoforms constitutes a primary event in the pathogenesis cascade of AD, or merely reflects a secondary effect shown at more advanced stages of the disease process [132]. In both AD patients and transgenic mice models of the disease, eNOS-expressing astrocytes exceeded those expressing iNOS in number [17]. Double immunolabeling studies revealed that in astrocytes iNOS and eNOS are co-localized with nitrotyrosine [122]. In some cases of AD, genetics defects in PS1 on microglial cells resulted in enhanced NO through superinduction of iNOS and eNOS [17, 133, 134].

Astrocytes with elevated levels of iNOS or eNOS were persistently seen in direct association with clusters of A β -deposits in AD patients and transgenic mice. The expression of both iNOS and eNOS is increased in activated astrocytes under experimental conditions in association with elevated expression of APP or bulk of A β -deposits in APP23 transgenic mice [17]. Whereas it is clear from the data in the literature that NO systems are involved in the pathophysiology of AD [98, 135] acting iNOS and eNOS as central mediators of A β action [49, 124, 136, 137], it has been also reported that changes in the activity of nNOS by the action of A β is responsible for the increased NO production in neurons [98]. Thus, reducing oxidative stress should be of potential value for limiting the incidence and progression of AD [138].

Taken together, studies on NO system in AD suggest that NO contributes to the maintenance, self-perpetuation and progression of the neurodegenerative process, and also that the altered expression of NOS isoforms might be secondary to the amyloid pathology. In view of the wide range of isoform-specific NOS inhibitors, the determination of the most responsible isoform of NOS for the formation of peroxynitrite in AD could be of therapeutic importance in the treatment of AD. It therefore appears important to gain a better understanding of the pathogenic mechanisms involving the NO systems, since they may represent an interesting therapeutic avenue for patients.




Fig. (5). Microphotograph taken from the cerebral cortex of Tg2576 transgenic mice showing immunostaining for iNOS. Notice that the immunoreaction product is distributed outside the core of the senile plaque labelling numerous round shaped small cells resembling microglia. Scale bar: 10 μ m.

NITRIC OXIDE AND ITS RELATIONSHIP WITH CONSIDERED ALZHEIMER'S DISEASE TREATMENTS

Cholinergic Therapy and Nitric Oxide

Amongst AChEIs, four products: tacrine, donepezil, rivastigmine, and galantamine are approved for clinical treatment, and the latter three are widely recommended for clinical use [139]. These products together with N-methyl-D-aspartate receptor antagonist (memantine) represent the available treatments for Alzheimer's disease [140]. Other cholinergic drugs such as muscarinic agonists have been explored, and although they are not approved, there is robust preclinical evidence for a beneficial, perhaps disease-modifying effect [141].

Acetylcholine (ACh) plays a role regulating the information flow and processing during learning and memory. There is certainly a link between cholinergic neurotransmission and cyclic guanosine monophosphate (cGMP) synthesis. The NO-signaling cascade has a role in learning and memory processes as well, which is also regulated by cGMP. Therefore, NO chimeras, for example GT 1061 and hybrid nitrates, may be a potential therapy for AD [142]. The nNOS isoform is ubiquitous in the brain and involved in glutamate neurotoxicity, learning and memory processes.

As stated above, AChEIs are the main recommended drugs for AD treatment. However, sometimes their efficacy has been questioned. Neurogenic vasodilation mediated by alpha 7-neuronal ACh nicotinic receptor has been described to be NO mediated and was blocked by AChEIs. Statins can prevent this blockage at the receptor level [143]. Statins have been found alter APP metabolism by lowering cholesterol levels (FDA Phase: Phase II/IIa/IIb).

Donepezil, galanthamine, and tacrine are AChEIs used widely for the treatment of AD. All these compounds have neuroprotective properties which have been related with glutamate neurotoxicity. Donepezil and galanthamine protect cortical neurons steps before NO formation and tacrine after [144]. In addition, Dimebon (3,6-dimethyl-9-(2-methyl-pyridyl-5)-ethyl-1,2,3,4-tetrahydro- γ -carboline dihydrochloride), shows activity as an inhibitor of cholinesterase and NMDA receptors. Through mitochondrial-mediated inhibition of apoptosis restrain neuronal death (FDA Phase III).

The cholinergic deficit that occurs in AD affects the cerebral vasoregulation. In fact, in AD there is a vasoregulative deficit that could be related with a lower eNOS expression. It has been shown

that the use of AChEIs, such as Donepezil, improves cerebrovascular regulation in Alzheimer's patients [106].

More recently, a new promising dimeric AChEI: bis(7)-Cognitin [145] has been reported. At neuroprotective concentrations, it reverses the over-activation of nNOS caused by glutamate without interfering with the basal activity of NOS. Through inhibition of AChE, NMDA receptor, nNOS, and APP/ β A cascade concurrently, it possesses remarkable neuroprotective activities. The synergism between these targets might serve as one of the most effective therapeutic strategies to arrest/modify AD pathology.

Lipoic acid function has been related as a mediator in acetylcholine production by activating of choline acetyltransferase. In fact, lipoic acid has been shown to ameliorate AD symptoms in patients with mild dementia and also in animal models [146].

Anti-Inflammatory Therapy and Nitric Oxide

Neuroinflammation is an essential component of many neurodegenerative diseases, including AD. This has been supported largely by observational studies that anti-inflammatory approaches may be protective against the development of AD [147, 148]. In the pathogenesis of neuroinflammation, free radicals are playing a relevant role, particularly NO produced through the activation of microglia and astrocytic cells. Thus, the increased understanding of the effects of NO in AD will lead to the development of novel anti-inflammatory therapeutic strategies.

Anomalous deposition of A β peptide has been widely described responsible for AD pathogenic process [13]. As a result of such anomalous deposits accumulation, neuronal damage and dysfunction occur through the activation of microglia and astrocytic cells [44]. Detrimental neuroinflammatory factors are characteristically expressed by activated glia in such a process, including pro-inflammatory cytokines and chemokines, oxidative stress-related enzymes, acute phase proteins, and components of the immunologic and complement cascades [40, 92]. Amyloid plaques are responsible for a local inflammatory response [48] essential in the progression of the disease [149, 150]. Particularly, astrocytes and microglial cells in close apposition to the extracellular accumulation of A β account for the synthesis and pathological release of variety of cytokines, chemokines and reactive oxygen species which may all lead to degeneration of neurotransmitter specific neuronal populations. It is discussed whether overproduction of cytokines (IL-1, IL-6, TGF- β , and TNF- α) secondary to cerebral inflammatory processes, may be involved in the development and progression of the disease [151]. A β deposits influence NO system leading to degeneration [123]. Because of the pathogenic inflammatory response that takes place in AD, it is possible that therapeutic intervention in such an inflammatory process hold great promise for the amelioration of AD symptoms. The compounds listed down here are described to exert a pharmacological therapeutic in AD through affecting inflammatory mechanism in which NO appears to be concerned.

The nonsteroidal anti-inflammatory drugs (NSAIDs) may lessen the inflammatory response acting on A β peptide deposition and therefore protecting against AD. NSAIDs antiinflammatory action results by inhibiting cyclooxygenase-1 and cyclooxygenase-2 and by activating the peroxisome proliferator γ nuclear transcription factor (PPAR α). These drugs have been proposed for the treatment of AD [155]. Cyclooxygenase-2 upregulation, an enzyme that plays a crucial role in mediating inflammatory response, seems to be associated with A β plaque formation in AD. This well described pro-inflammatory mediator acts close together with iNOS [152]. NSAIDs also possess inhibitory effects in the generation of NO radicals [153]. Ibuprofen, the most commonly used NSAIDs, is a cyclooxygenase-2 (COX-2) inhibitor that decreases the production of NO. This compound protects neurons against glutamate toxicity, decreases the

production of proinflammatory cytokines and suppresses neuritic plaques pathology and inflammation in AD [154]. This compound is clinical trial phase III. Flurbiprofen is and additional NSAIDs that influences A β peptide deposition and metabolism. Flurbiprofen racemate, its *R*-enantiomer and its NO-releasing derivatives, HCT 1026 and NCX 2216, are effective on AD amyloid pathology [155] (FDA Phase: discontinued).

Resveratrol (3,5,4'-trihydroxy-trans-stilbene), a naturally occurring phytoalexin found in grapes and wine, has a neuroprotective action against A β induced toxicity in C6 glioma cells by inhibiting iNOS and COX-2 expression [130]. (FDA Phase III).

Retinoic acid has an inhibitory effect on TNF- α and iNOS expression *in vitro* preparations of activated rat microglia by A β peptide [156].

Lipoic acid acts downregulating iNOS activity and pro-inflammatory proteins [146]. Thanks to its neuroprotective action, lipoic acid has been proposed as an alternative for AD treatment.

Catalpol, an iridoid glycoside isolated from the root of Rehmannia glutinosa, protects neuronal cells by decreasing the A β 1-42 and inhibiting the to some extent glial activation [151].

Gemfibrozil, a member of the fibrate class of hypolipidaemic drugs, inhibits LPS-induced expression of iNOS as well as pro-inflammatory cytokines in mouse BV-2 microglial cells and primary microglia [157].

The pharmacological blockage with the iNOS inhibitor aminoguanidine reduced the cognitive deficit evoked by intracerebroventricular injection of A β 1-40 in mice. Similar results were obtained in iNOS knock-out mice [89].

Vitamin E (alpha-tocopherol) inhibits iNOS expression and therefore NO production in astrocytes stimulated with A β [158]. Therefore, prevents brain cell damage by destroying toxic free radicals. (FDA Phase III).

Poly(ADP-ribose) polymerase-1 (PARP-1) has been shown to play a key role in NF- κ B driven expression of inflammatory mediators by glia. Inhibitors of PARP-1 enzymatic activity reduce the expression of inflammatory mediators such as iNOS and APP among others, and the neurotoxic potential of activated glia *in vitro* [159].

Induced cholinergic degeneration produces increased expression of inflammatory mediators, among others iNOS. The nonsteroidal anti-inflammatory drug, Nimesulide, attenuate iNOS activity [160].

Phospholipases: a Target for Therapeutic Intervention

Phospholipases hydrolyse fatty acids of membrane phospholipids. Activation of phospholipase A₂ (PLA₂) mediates arachidonic acid (AA) release from phospholipids sn-2 position. NO affects the intracellular level of AA through alteration of PLA₂ activity. Excess of activation of PLA₂ produces damage of cell membrane integrity. PLA₂ acts by altering membrane permeability and calcium homeostasis increasing the levels of intracellular Ca²⁺, that leads an excitotoxic phenomena involved in a lot of pathologies [161]. The interaction of A β with cell surface receptors results in the stimulation of phospholipases [162]. Particularly, a marked stimulation of PLA₂ activity has been shown to occur in AD [163, 164]. The nNOS isoform is one of the target enzymes of the PLA₂ activation and responsible for excitotoxicity-induced neurodegenerative diseases, including AD [165]. Therefore, phospholipases may represent a therapeutic target for AD. However, using the selective inhibitors of PLA₂ activity, indomethacin and nor-dihydroguaiaretic acid, the A β peptide continues inducing cell death *in vitro* as well as with the NOS inhibitors NG-monomethyl-L-arginine and N omega-nitro-L-arginine [166]. DP-55, a lipid compound which is cleaved by PLA₂, has been described as a compound of potential therapeutic action for AD. This compound is efficacious in re-

ducing levels of A β in transgenic AD mouse (Tg2576) brains as well as in reducing neurodegeneration, and inflammation in an *in vitro* AD model [167]. It has been described that Resveratrol prevents NO dependent AA action [168].

Phosphodiesterases (PDE) Inhibitors

The cyclic nucleotide phosphodiesterase (PDE) comprise a group of enzymes that degrade phosphodiester bond in the second messenger molecules cyclic adenosine monophosphate (cAMP) and cGMP. They regulate the localization, duration, and amplitude of cyclic nucleotide signaling within subcellular domains. Phosphodiesterases (PDEs) are therefore important regulators of signal transduction mediated by these second messenger molecules. The PDE superfamily of enzymes is classified into 11 families in mammals, namely PDE1-PDE11. PDEs have different substrate specificities. Some are cAMP selective hydrolases (PDE4, 7 and 8), others are cGMP selective (PDE5, 6 and 9). Others can hydrolyse both cAMP and cGMP (PDE1, 2, 3, 10 and 11). The potential for selective PDE inhibitors to be used as therapeutic agents was predicted as early as 1977 by Weiss and Hait [169, 170].

NO, memory enhancement, and PDE are correlated *via* cGMP signaling. Activation of NMDA glutamate receptors causes increased intracellular calcium that binds to calmodulin activating nNOS and the production of NO. NO in turn activates soluble guanylyl cyclase followed by an increased formation of cGMP. Increased formation of cGMP leads to protein synthesis, synaptogenesis and memory enhancement [171].

Phosphodiesterase-5 (PDE5) belongs to an important family of proteins that regulates the intracellular level of cGMP. Recent studies have shown that PDE-5 inhibitors can counteract memory deficits in an AD model [172]. Sildenafil, a PDE5 inhibitor, increases the level of cGMP leading to beneficial effects in some targeting organs [173]. Sildenafil is the first oral drug approved by the United States Food and Drug Administration for the therapeutic treatment of erectile dysfunction. Besides, it may offer a novel strategy in the therapeutic treatment of a lot of pathologies: age-related memory impairment, pain, pulmonary hypertension, multiple sclerosis and progression, prevention and, eventually, AD [174]. This drug attenuates memory impairment induced by NOS inhibition, among others [175]. Propentofylline, another PDE5 inhibitor, prevents the neuronal damage NO-related in dementing processes [176]. However, in a clinical trial, this compound has shown a very limited capacity for improve cognition and global function in AD [177]. (FDA Phase: Discontinued). Phosphodiesterase-4 (PDE4) belongs to an important family of proteins that regulates the intracellular level of cAMP. Targeting PDE4 with selective inhibitors may offer a novel therapy aimed at slowing progression and, eventually, therapy of AD. In this sense, Rolipram, a specific PDE4 inhibitor improves memory in young and aged mice [178] as well as in double-transgenic mice APP/PS1 mice [179].

Caffeine, an adenosine receptor antagonist, inhibits phosphodiesterases action thus being a potential candidate for the treatment of AD. Caffeine reduces A β deposition and improves cognitive behavior in a mouse model of AD as well as in neuronal cultures from such a model [180].

Besides caffeine, other selective adenosine A (2A) antagonists prevent the toxic effects of A β peptide. For example, the compound SCH58261 avoid A β induced cognitive deficits following intracerebroventricular administration of this peptide in mouse [181]. Also, the compound ZM 241385 protects rat cerebellar neurons from the toxic effects of A β peptide [182].

FUTURE DIRECTIONS IN ALZHEIMER'S DISEASE TREATMENT: DRUGS TO RESTRAIN NITRIC OXIDE SYNTHESIS

Regarding the involvement of NO in the progression of the disease, research is directed towards the production of new drugs

that may be able to treat the underlying cause of this chronic disease, not just the symptoms. We are describing the progress being made in the scientific community and pharmaceutical industry on the development of any breakthrough product of potential use.

NOS Inhibitors

Because of the harmful effects of excess NO, selective restraint of NO biosynthesis through a controlled inhibition of the NOS isoforms, particularly iNOS [183], might provide an important therapeutic opportunity. Many efforts have been focused to the design of high affinity drugs, most based on strategies carried out to modify the enzyme's substrate. The inhibitory action of a putative drug on iNOS and nNOS activities may be beneficial in various forms of shock and inflammation and may protect against neuroinjury, respectively [184]. But since NO plays certain central physiological roles, a complete inhibition of its production may be detrimental. Therefore, to design drugs that target a specific NOS isoenzyme in a therapeutic dosage is critical to avoid side effects. These drugs should reach a proper homeostasis of NO metabolism in the tissue where endogenous production of NO is excessively increased. The use of non-specific NOS inhibitors, such as L-nitroarginine methyl ester (L-NAME), impairs long-term memory consolidation [185]. It is also noteworthy the role of eNOS in maintaining vascular tone and spatial memory function [186]. On the other hand, complete removal of iNOS, which is primarily associated with the immune response, from an APP transgenic mouse (APPSwDI/NOS2^{-/-}) results in development of a much greater spectrum of AD-like pathology and behavioral impairments [187]. Studies on the Tg2576 APP mouse with an iNOS knock-out background also support the evidence suggesting that NO generated by iNOS under conditions of long-term injury or disease reduces functional loss and mitigates pathological changes in brain [98].

In summary, a possible way to delay the progression of AD must include a rationale design of enzyme inhibitors, subtype selective, targeting NOS isoforms implicated in damage to brain cells [188]. The catalytic activity of NOS can be tone down by a pharmacological attack on different enzyme sites, in example, the arginine, heme, and tetrahydrobiopterin (BH₄) sites.

We are now presenting an overview of enzyme inhibitors targeting NOS isoforms. These products, although are not in use for the treatment of AD, may represent useful tools for investigating the biological functions of NO in AD.

From a chemical point of view, these chemicals are divided into two principal groups: aminoacid-based inhibitors and non-aminoacid-based inhibitors.

1. Aminoacid-Based Inhibitors

This type of derivatives have been grouped, based on their chemical structures, into three categories of compounds [188, 189]: L-arginine analogues, conformationally restricted arginine analogues and dipeptides.

1.1. L-Arginine Analogues

Specific modifications of the enzyme's substrate became a strategy outlook to produce NOS inhibitors that have become a putative targeted therapy strategy in AD for the pharmaceutical industry. The first NOS inhibitors derived from L-arginine were mono- or di-substituted guanidino analogues, most notable L-N^G monomethyl arginine (L-NMMA) and L-N^G nitroarginine (L-NOARG) and its methyl ester (L-NAME). These products showed a potent inhibitory action but a poor selectivity among the NOS isoforms.

Arginine analogues encompass L-NAME [190] and its derivatives; N-methyl-L-arginine (L-NMA) [191], N^δ-(1-iminoethyl)-L-ornithine (L-NIO), N-nitro-L-arginine (L-NNA) [190], N-alkyl-L-arginines, N-(1-imino-3-butenyl)-L-ornithine (L-VNIO), L-thiocitrulline [192], L-homothiocitrulline [193], N-(1-iminoethyl)-

L-lysine (L-NIL) [194], and S-alkyl-L-thiocitrullines [195]. The development of N^ω-2-nitroaryl amino acid derivatives and heterocyclic analogues of L-citrulline [196] were obtained by means of modifications of the guanidine moiety of L-arginine. N^ω-2-nitroaryl amino acid showed a good inhibitory potency against nNOS, and less active against eNOS and iNOS, maintaining the same order of the potency and selectivity of the prototypic NOS inhibitors L-NMA and L-NNA.

1.2. Conformationally Restricted Arginine Analogues of Endogenous NOS Substrate L-Arginine

Guanidine, N^G-methylguanidine and amidine L-arginine analogues have been synthesized. With some exceptions, these structurally related analogues were, generally, weak inhibitors with moderate selectivity, if compared to the parent molecules. Conformational restriction appears to prevent the molecules from assuming the appropriate discriminatory binding orientations needed for high selectivity of the NOS isoenzymes [197]. Thus, restricted conformations do not seem to be a proper approach for the increase of inhibitors selectivity.

1.3. Dipeptides

This group includes L-NNA-containing dipeptides. Because of the selectivity of L-NNA for nNOS vs iNOS, in order to improve selectivity in favour of nNOS, [197] designed L-NNA (Arg^{NO2})- and phenylalanine (PHE)-containing dipeptides and dipeptide esters.

A library of 152 dipeptides amides containing nitroarginine and amino acids different from PHE were synthesized and screened for activity [198]. Excellent inhibitory potency and selectivity for nNOS over eNOS and iNOS is achieved with the dipeptide amides containing a basic amine side chain. Among these compounds, the most potent nNOS inhibitor is L-Arg^{NO2}-L-Dbu-NH₂, which also exhibits the highest selectivity over eNOS (>1500-fold) with a 192-fold selectivity over iNOS. These compounds do not exhibit time-dependent inhibition. This compound can be considered the most selective nNOS inhibitor known. Structure Activity Relationship (SAR) studies demonstrated the importance of the α-amino group and the NH moiety of the peptide bond for binding at the active site.

Several L-arginine analogs are known as potent inhibitors of NOS. Since dipeptide containing are potent and isozyme-selective NOS inhibitors, following the dipeptide approach, it has been described additional dipeptides containing arginine-analogues [199]. S-methyl-L-isothiocitrullinyl-L-phenylalanine showed 66-fold selectivity for iNOS over nNOS, while S-methyl-L-isothiocitrullinyl-L-leucine and N^G-nitro-L-argininyl-L-phenylalanine showed 20- and 14-fold selectivity, respectively. It is suggested that each NOS isozyme has a cavity of different size near the C-terminal of the L-arginine binding site, and that the selectivity of inhibitors is due to the differences in the size of the cavity.

Recently, a new cyclic dipeptide product, cyclo(dehydrohistidyl-1-tryptophyl) (CDHT), was found to inhibit NO in lypopolysaccharide (LPS)-treated BV-2 cells by preventing iNOS dimerization [200]. Characteristically, this product has no effect on iNOS expression or enzyme activity. In contrast, CDHT did not inhibit eNOS activity. These results reveal that CDHT could be a useful therapeutic agent for inflammation-mediated diseases.

2. Non-Aminoacid-Based Inhibitors

In order to improve the selectivity and therapeutic profile of L-arginine derivatives, non-amino acid-based inhibitors of NOS have been studied; these include a continually growing list of compounds that, from a chemical point of view, may be divided into two principal groups: amidinic and heterocyclic compounds.

2.1. Amidinic Compounds

2.1.1. Guanidines

Despite the fact that simple guanidines are only weak inhibitors [201], guanidine moiety can mimic L-Arg for recognition at the binding site so that it can be considered a potential pharmacophore for NOS inhibition. Aminoguanidine is a selective inhibitor of mouse inducible NOS and numerous *in vivo* studies have demonstrated that this inhibition could be useful in the treatment of disease states characterized by the pathological overproduction of NO from iNOS [202].

2.1.2. Isothioureas

Inhibition by non-amino acid isothioureas has provided compounds which are potent and selective inhibitors of human iNOS vs eNOS [203]. The inhibitory effect of these compounds is caused not only by their competition with the substrate for the L-arginine-binding site and/or oxidizing center of the enzyme (heme) but also by interaction with peptide motifs of the enzyme that influence its dimerization, affinity for cofactors, and interaction with associated proteins. The clinical use of these isothiourea-based compounds is limited owing to their poor cellular penetration and high toxicity.

2.1.3. Amidines

There are a great number of derivatives of amidines, but the most active compounds are: 2-iminopiperidine, 2-iminohomopiperidine and (1*S*,5*S*,6*R*,7*R*)-7-Chloro-3-imino-5-methyl-2-azabicyclo [4.1.0]heptane hydrochloride (ONO-1714). These are more potent inhibitors of human iNOS than L-NMA with little preference for iNOS vs eNOS when compared with L-NMA.

ONO-1714 constitutes a novel cyclic amidine analogue that inhibits human iNOS with a *K_i* of 1.88 nM and rodent iNOS with similar potency *in vitro*. ONO-1714 was found to be 10-fold selective for human iNOS over human eNOS. When the inhibitory activity of ONO-1714 was compared for iNOS, it was found to be 451-fold and >20,000-fold more potent than L-NMMA and aminoguanidine, respectively [204].

2.2. Heterocyclic Compounds

This group of non-amino acid-based NOS inhibitors includes a range of heterocyclic compounds among which the most represented are: indazoles, imidazoles and analogues of BH₄.

2.2.1. Indazoles

Despite the absence of *in vitro* selectivity in enzymatic or functional assays [205]; 7-nitroindazole named 7-NI, may be considered as the first selective inhibitor of nNOS *in vivo*; in fact, several studies have demonstrated its protective effects in pain, experimental stroke [206] and in mouse model of Parkinson's disease [207] with minimal systemic pressure effects.

2.2.2. Imidazoles

Since it has been demonstrated that NOS contains a very similar heme site to that of cytochrome-P₄₅₀ [208], imidazole, 1-phenyl, 2-phenyl and 4-phenylimidazole and antimycotic drugs (myconazole, ketoconazole and clotrimazole) were tested as inhibitors [209]. In fact, it is well known that imidazoles inhibit the activity of various heme-containing proteins by binding to the heme group [210].

2.2.3. Analogues of Tetrahydrobiopterin

Owing to the structural relation with the enzymatic cofactor BH₄, pteridine based compounds have been described as NOS inhibitors. This approach appears promising because of the much lower affinity and selectivity of BH₄ binding to other pteridine dependent enzymes [211]. Among a series of tested compounds, the 4-amino analogue of BH₄, 5,6,7,8-tetrahydro-6-(D-theo-1,2-dihydroxypropyl)pterin, was a potent inhibitor of the recombinant rat brain

NOS both *in vitro* and *in vivo* and, although to a lesser extent, also of the other isoform.

Other pteridine-based compounds, named anti-pterines, structurally related to 5,6,7,8-tetrahydro-6-(D-theo-1,2-dihydroxypropyl) pterin, were described as NOS inhibitors. Some of these compounds specifically interacted with the BH₄ binding site of NOS without interference with any other known cofactors or substrate binding sites [212].

Most recently, a novel antipterin NOS-inhibitor, 4-amino-(6R,S)-5,6,7,8-tetrahydro-L-biopterin (VAS203) has been described [23] to inhibit NOS working over the BH₄ cofactor binding site. Characteristically, much lower concentrations of VAS203 are required to inhibit iNOS as compared to the constitutive NOS isoforms.




Fig. (6). Represents a low-power magnification micrograph taken from a histological section through the temporal cortex of a postmortem brain of AD patient, CDR3 case, showing the distribution of anti-human IgG immunolabeling. Notice immunoreactivity in neurons, senile plaques and surrounding glial cells. Scale bar: 100 μ m.

IMMUNOTHERAPY

Although little is known on the etiology and pathogenesis of AD, several reports indicate the involvement of immunological mechanisms playing an important role in the disease process. In the temporal cortex from postmortem brain of CDR3 AD patients, we have found cortical neurons (both interneurons and pyramidal cells), senile plaques and its surrounding glial cells exhibiting autoantibodies (human IgGs), as shown in Fig. (6) (unpublished results), which might represent an autoimmune aggression. In addition, a prominent innate immune response occurs in the brain in association with A β deposition and plaque formation. This response includes the activation of complement, secretion of proinflammatory cytokines and NO, and the expression of the chemokines [213]. These observations support a close relationship between both the immune system and the pathophysiology of AD. There are data that support the capacity of anti-A β antibodies to reduce A β aggregation and neurotoxicity. Based on the latter, it has been suggested that immunotherapy with a vaccine that avoid toxic side effects might generate specific and effective immune response against A β , the causative agent of synaptic loss and cognitive decline, providing a powerful tool for the treatment of AD [214]. This is particularly consistent given findings that microglia-mediated release of NO by toxic A β contributes to AD neurotoxicity.

A β stimulates inflammation and the production of neurotoxic reactive oxygen species by inducing the pathological over-expression of iNOS [215, 218]. Therefore, patients might benefit with immunotherapy which enhances clearing of this neurotoxic peptide

from the brain [215-217]. Since only a small percentage of patients with AD show symptomatic benefits from current drug therapy and for only a relatively short time, immunotherapy is among the leading therapeutic directions for the disease. The use of intravenous Immunoglobulin (Gammagard, IVIg) is in FDA Phase III.

CONCLUSIONS

Presently, there is no cure for AD and no proven treatment has been developed to slow or halt its progression. Strictly, five drugs have been approved by the US Food and Drug Administration and the European Medicines Agency that improve the mental function of people with AD, these include: tacrine (Cognex), rivastigmine (Exelon), galantamine (Razadyne), donepezil (Aricept), and memantine (Namenda). All except memantine are approved for the treatment of mild to moderate AD. Memantine is approved for the treatment of moderate to severe AD only. All of these drugs, except memantine, act by increasing brain levels of acetylcholine, a neurotransmitter that is abnormally low in patients with AD. Memantine works differently interacting with the NMDA receptor.

The production of new specific drugs that cure the underlying cause of this chronic disease, not just the symptoms, is a matter of clinical interest. The three isoforms of the NOS operate as central mediators of A β action, giving rise to elevated levels of NO that contributes to the maintenance, self-perpetuation and progression of the disease. Indeed, NO links mitochondrial dysfunction to AD. At the cellular level, the mitochondrion appears to be of importance when considering the deleterious effects of both A β and induced NO. Generation of highly reactive nitrogen species secondary to high concentrations of NO causes irreversible inhibition of mitochondrial respiration and damage to various mitochondrial components *via* oxidizing reactions leading to cell death. Attacks on the mitochondrial protein dynamin-related protein 1 (Drp1) by NO, which causes a chemical reaction called S-nitrosylation, mediates neurodegeneration associated with AD [219]. Prior to this study, the mechanism by which A β caused synaptic damage to neurons in AD was unknown. These findings suggest that preventing S-nitrosylation of Drp1 may reduce or even prevent neurodegeneration in Alzheimer's patients. Therefore, a possible way to delay the progression of AD must include a rationale design of enzyme inhibitors, subtype selective, targeting NOS isoforms implicated in damage to brain cells in AD.

ACKNOWLEDGEMENTS

We gratefully acknowledge Ms Beatriz Lázaro López for her valuable technical help. This work was supported by Spanish Ministry of Science and Innovation Grant SAF2007-60010, and Instituto de Salud Carlos III Grant RD06/0026/1001.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ABBREVIATIONS

A β	=	Amyloid beta-peptide
AA	=	Arachidonic Acid
ACh	=	Acetylcholine
AChE	=	Acetylcholinesterase
AChEIs	=	Acetylcholinesterase Inhibitors
AD	=	Alzheimer Disease
ApoE	=	Apolipoprotein E
APP	=	Amyloid precursor protein
BACE-1	=	β -site APP-cleaving enzyme 1
BH ₄	=	Tetrahydrobiopterin
cAMP	=	cyclic adenosine MonoPhosphate
CDHT	=	Cyclo (dehydrohistidyl-1-tryptophyl

CDR3	=	Clinical dementia rating 3 (severe dementia)
cGMP	=	cyclic Guanosine MonoPhosphate
COX-2	=	Cyclooxygenase-2
eNOS	=	endothelial Nitric Oxide Synthase
FAD	=	Familial Alzheimer Disease
GFAP	=	Glial fibrillary acid protein
iNOS	=	inducible Nitric Oxide Synthase
LAD	=	Late-onset/sporadic Alzheimer Disease
LDL	=	Low-Density Lipoprotein
L-NAME	=	L-nitroarginine methyl ester
L-NIL	=	N-(1-iminoethyl)-L-lysine
L-NIO	=	N ⁸ -(1-iminoethyl)-L-ornithine
L-NMA	=	N-methyl-L-arginine
L-NMMA	=	L-N ^G monomethyl arginine
L-NNA	=	N-nitro-L-arginine
L-NOARG	=	L-N ^G nitroarginine
LPS	=	Lipopolysaccharide
LRP	=	LDL-receptor related protein
L-VNIO	=	N-(1-imino-3-3-butenyl)-L-ornithine
NFT	=	Neurofibrillary tangles
NMDA	=	N-methyl-D-aspartate
nNOS	=	neuronal Nitric Oxide Synthase
NO	=	Nitric Oxide
NOS	=	Nitric Oxide Synthase
NSAIDs	=	Nonsteroidal anti-inflammatory drugs
ONO-1714	=	(1S,5S,6R,7R)-7-Chloro-3-imino-5-methyl-2-azabicyclo[4.1.0] heptane hydrochloride
ONOO ⁻	=	Peroxynitrite
PARP-1	=	Poly(ADP-ribose) polymerase-1
PDE	=	Phosphodiesterase
PDE-4	=	Phosphodiesterase-4
PDE-5	=	Phosphodiesterase-5
PHE	=	Phenylalanine
PKC	=	Protein Kinase C
PLA ₂	=	Phospholipase A ₂
PS1	=	Presenilins 1
PS2	=	Presenilins 2
SAR	=	Structure Activity Relationship
TNF- α	=	Tumor Necrosis Factor-alpha
VAS203	=	4-amino-(6R,S)-5,6,7,8 tetrahydro-L-biopterin

REFERENCES

- [1] Alzheimer A, Stelzmann RA, Schnitzlein HN, Murtagh FR. An English translation of Alzheimer's 1907 paper, "Über eine eigenartige Erkrankung der Hirnrinde". Clin Anat 1995; 8(6): 429-31.
- [2] Morris JC, Ernesto C, Schafer K, *et al.* Clinical dementia rating training and reliability in multicenter studies: the Alzheimer's Disease Cooperative Study experience. Neurology 1997; 48(6): 1508-10.
- [3] Selkoe DJ. Cell biology of the amyloid beta-protein precursor and the mechanism of Alzheimer's disease. Annu Rev Cell Biol 1994; 10: 373-403.
- [4] Henderson S. Epidemiology of dementia. Ann Med Interne (Paris) 1998; 149(4): 181-6.
- [5] Cacabelos R. Influence of pharmacogenetic factors on Alzheimer's disease therapeutics. Neurodegener Dis 2008; 5(3-4): 176-8.
- [6] Zawia NH, Lahiri DK, Cardozo-Pelaez F. Epigenetics, oxidative stress, and Alzheimer disease. Free Radic Biol Med 2009; 46(9): 1241-9.
- [7] Kang J, Lemaire HG, Unterbeck A, *et al.* The precursor of Alzheimer's disease amyloid A4 protein resembles a cell-surface receptor. Nature 1987; 325(6106): 733-6.
- [8] Naslund J, Schierhorn A, Hellman U, *et al.* Relative abundance of Alzheimer A beta amyloid peptide variants in Alzheimer disease and normal aging. Proc Natl Acad Sci USA 1994; 91(18): 8378-82.
- [9] Sun X, Chiu CC, Liebson E, *et al.* Depression and plasma amyloid beta peptides in the elderly with and without the apolipoprotein E4 allele. Alzheimer Dis Assoc Disord 2009; 23(3): 238-44.
- [10] Tanzi RE, Bertram L. Twenty years of the Alzheimer's disease amyloid hypothesis: a genetic perspective. Cell 2005; 120(4): 545-55.
- [11] Lukiw WJ. Emerging amyloid beta (Ab) peptide modulators for the treatment of Alzheimer's disease (AD). Expert Opin Emerg Drugs 2008; 13(2): 255-71.
- [12] Fernandez-Vizarra P, Fernandez AP, Castro-Blanco S, *et al.* Expression of nitric oxide system in clinically evaluated cases of Alzheimer's disease. Neurobiol Dis 2004; 15(2): 287-305.
- [13] Hoozemans JJ, Rozemuller JM, van Haastert ES, Veerhuis R, Eikelenboom P. Cyclooxygenase-1 and -2 in the different stages of Alzheimer's disease pathology. Curr Pharm Des 2008; 14(14): 1419-27.
- [14] Pasinetti GM. From epidemiology to therapeutic trials with anti-inflammatory drugs in Alzheimer's disease: the role of NSAIDs and cyclooxygenase in beta-amyloidosis and clinical dementia. J Alzheimers Dis 2002; 4(5): 435-45.
- [15] Velez-Pardo C, Arellano JI, Cardona-Gomez P, Jimenez Del RM, Lopera F, De FJ. CA1 hippocampal neuronal loss in familial Alzheimer's disease presenilin-1 E280A mutation is related to epilepsy. Epilepsia 2004; 45(7): 751-6.
- [16] Hu J, Akama KT, Krafft GA, Chromy BA, Van Eldik LJ. Amyloid-beta peptide activates cultured astrocytes: morphological alterations, cytokine induction and nitric oxide release. Brain Res 1998; 785(2): 195-206.
- [17] Luth HJ, Holzer M, Gartner U, Staufenbiel M, Arendt T. Expression of endothelial and inducible NOS-isoforms is increased in Alzheimer's disease, in APP23 transgenic mice and after experimental brain lesion in rat: evidence for an induction by amyloid pathology. Brain Res 2001; 913(1): 57-67.
- [18] Muller T, Meyer HE, Egensperger R, Marcus K. The amyloid precursor protein intracellular domain (AICD) as modulator of gene expression, apoptosis, and cytoskeletal dynamics-Relevance for Alzheimer's disease. Prog Neurobiol 2008; 85(4): 393-406.
- [19] Butterfield DA, Perluigi M, Sultana R. Oxidative stress in Alzheimer's disease brain: new insights from redox proteomics. Eur J Pharmacol 2006; 545(1): 39-50.
- [20] Butterfield DA, Reed TT, Perluigi M, *et al.* Elevated levels of 3-nitrotyrosine in brain from subjects with amnesic mild cognitive impairment: implications for the role of nitration in the progression of Alzheimer's disease. Brain Res 2007; 1148: 243-8.
- [21] Sultana R, Perluigi M, Butterfield DA. Protein oxidation and lipid peroxidation in brain of subjects with Alzheimer's disease: insights into mechanism of neurodegeneration from redox proteomics. Antioxid Redox Signal 2006; 8(11-12): 2021-37.
- [22] Hardy JA, Higgins GA. Alzheimer's disease: the amyloid cascade hypothesis. Science 1992; 256(5054): 184-5.
- [23] Gandy S. The role of cerebral amyloid beta accumulation in common forms of Alzheimer disease. J Clin Invest 2005; 115(5): 1121-9.
- [24] Oddo S, Caccamo A, Cheng D, Laferla FM. Genetically altering abeta distribution from the brain to the vasculature ameliorates tau pathology. Brain Pathol 2008.
- [25] Bertram L, Tanzi RE. Alzheimer's disease: one disorder, too many genes? Hum Mol Genet 2004; 13 Spec No 1: R135-41.
- [26] Nakajima M, Shirasawa T. Presenilin-1-deficient neurons are nitric oxide-dependently killed by hydrogen peroxide *in vitro*. Neuroscience 2004; 125(3): 563-8.
- [27] Yankner BA, Duffy LK, Kirschner DA. Neurotrophic and neurotoxic effects of amyloid beta protein: reversal by tachykinin neuropeptides. Science 1990; 250(4978): 279-82.
- [28] Wolfe MS. Shutting down Alzheimer's. Sci Am 2006; 294(5): 72-9.
- [29] Perez M, Moran MA, Ferrer I, Avila J, Gomez-Ramos P. Phosphorylated tau in neuritic plaques of APP(sw)/Tau (v1w) transgenic

- mice and Alzheimer disease. *Acta Neuropathol* 2008; 116(4): 409-18.
- [30] Avila J. Tau kinases and phosphatases. *J Cell Mol Med* 2008; 12(1): 258-9.
- [31] Hernandez F, Avila J. Tauopathies. *Cell Mol Life Sci* 2007; 64(17): 2219-33.
- [32] Rovelet-Lecrux A, Hannequin D, Raux G, *et al.* APP locus duplication causes autosomal dominant early-onset Alzheimer disease with cerebral amyloid angiopathy. *Nat Genet* 2006; 38(1): 24-6.
- [33] Levy-Lahad E, Wasco W, Poorkaj P, *et al.* Candidate gene for the chromosome 1 familial Alzheimer's disease locus. *Science* 1995; 269(5226): 973-7.
- [34] De SB, Saftig P, Craessaerts K, *et al.* Deficiency of presenilin-1 inhibits the normal cleavage of amyloid precursor protein. *Nature* 1998; 391(6665): 387-90.
- [35] Duff K, Eckman C, Zehr C, *et al.* Increased amyloid-beta42(43) in brains of mice expressing mutant presenilin 1. *Nature* 1996; 383(6602): 710-3.
- [36] Borchelt DR, Thinakaran G, Eckman CB, *et al.* Familial Alzheimer's disease-linked presenilin 1 variants elevate Abeta1-42/1-40 ratio *in vitro* and *in vivo*. *Neuron* 1996; 17(5): 1005-13.
- [37] Hardy J, Selkoe DJ. The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science* 2002; 297(5580): 353-6.
- [38] Townsend M, Shankar GM, Mehta T, Walsh DM, Selkoe DJ. Effects of secreted oligomers of amyloid beta-protein on hippocampal synaptic plasticity: a potent role for trimers. *J Physiol* 2006; 572(Pt 2): 477-92.
- [39] Walsh DM, Townsend M, Podlisny MB, *et al.* Certain inhibitors of synthetic amyloid beta-peptide (Abeta) fibrillogenesis block oligomerization of natural Abeta and thereby rescue long-term potentiation. *J Neurosci* 2005; 25(10): 2455-62.
- [40] Shankar GM, Li S, Mehta TH, *et al.* Amyloid-beta protein dimers isolated directly from Alzheimer's brains impair synaptic plasticity and memory. *Nat Med* 2008; 14(8): 837-42.
- [41] Tanzi RE. The synaptic Abeta hypothesis of Alzheimer disease. *Nat Neurosci* 2005; 8(8): 977-9.
- [42] Knafo S, Ono-Nanclares L, Gonzalez-Soriano J, *et al.* Widespread changes in dendritic spines in a model of Alzheimer's disease. *Cereb Cortex* 2008.
- [43] Malchiodi-Albedi F, Domenici MR, Paradisi S, Bernardo A, Jimenez-Cat MA, Minghetti L. Astrocytes contribute to neuronal impairment in beta A toxicity increasing apoptosis in rat hippocampal neurons. *Glia* 2001; 34(1): 68-72.
- [44] Snyder EM, Nong Y, Almeida CG, *et al.* Regulation of NMDA receptor trafficking by amyloid-beta. *Nat Neurosci* 2005; 8(8): 1051-8.
- [45] Deane R, Zlokovic BV. Role of the blood-brain barrier in the pathogenesis of Alzheimer's disease. *Curr Alzheimer Res* 2007; 4(2): 191-7.
- [46] Thomas T, Thomas G, McLendon C, Sutton T, Mullan M. beta-Amyloid-mediated vasoactivity and vascular endothelial damage. *Nature* 1996; 380(6570): 168-71.
- [47] Haass C, Selkoe DJ. Soluble protein oligomers in neurodegeneration: lessons from the Alzheimer's amyloid beta-peptide. *Nat Rev Mol Cell Biol* 2007; 8(2): 101-12.
- [48] Rosenberg RN. The molecular and genetic basis of AD: the end of the beginning: the 2000 Wartenberg lecture. *Neurology* 2000; 54(11): 2045-54.
- [49] de la Monte SM, Lu BX, Sohn YK, *et al.* Aberrant expression of nitric oxide synthase III in Alzheimer's disease: relevance to cerebral vasculopathy and neurodegeneration. *Neurobiol Aging* 2000; 21(2): 309-19.
- [50] Lanni C, Racchi M, Mazzini G, *et al.* Conformationally altered p53: a novel Alzheimer's disease marker? *Mol Psychiatry* 2008; 13(6): 641-7.
- [51] Uberti D, Lanni C, Racchi M, Govoni S, Memo M. Conformationally altered p53: a putative peripheral marker for Alzheimer's disease. *Neurodegener Dis* 2008; 5(3-4): 209-11.
- [52] Serrano J, Fernandez AP, Martinez-Murillo R, Martinez A. High sensitivity to carcinogens in the brain of a mouse model of Alzheimer's disease. *Oncogene* 2010; 29(15): 2165-71.
- [53] Su JH, Deng G, Cotman CW. Bax protein expression is increased in Alzheimer's brain: correlations with DNA damage, Bcl-2 expression, and brain pathology. *J Neuropathol Exp Neurol* 1997; 56(1): 86-93.
- [54] de la Monte SM, Sohn YK, Wands JR. Correlates of p53- and Fas (CD95)-mediated apoptosis in Alzheimer's disease. *J Neurol Sci* 1997; 152(1): 73-83.
- [55] Paradis E, Douillard H, Koutroumanis M, Goodyer C, LeBlanc A. Amyloid beta peptide of Alzheimer's disease downregulates Bcl-2 and upregulates bax expression in human neurons. *J Neurosci* 1996; 16(23): 7533-9.
- [56] Hooper C, Meimaridou E, Tavassoli M, Melino G, Lovestone S, Killick R. p53 is upregulated in Alzheimer's disease and induces tau phosphorylation in HEK293a cells. *Neurosci Lett* 2007; 418(1): 34-7.
- [57] Messmer UK, Brune B. Attenuation of p53 expression and Bax down-regulation during phorbol ester mediated inhibition of apoptosis. *Br J Pharmacol* 1997; 121(4): 625-34.
- [58] Ranjan P, Shrivastava P, Singh SM, Sodhi A, Heintz NH. Baculovirus P35 inhibits NO-induced apoptosis in activated macrophages by inhibiting cytochrome c release. *J Cell Sci* 2004; 117(Pt 14): 3031-9.
- [59] Lammich S, Okochi M, Takeda M, *et al.* Presenilin-dependent intramembrane proteolysis of CD44 leads to the liberation of its intracellular domain and the secretion of an Abeta-like peptide. *J Biol Chem* 2002; 277(47): 44754-9.
- [60] Lathia JD, Mattson MP, Cheng A. Notch: from neural development to neurological disorders. *J Neurochem* 2008; 107(6): 1471-81.
- [61] Vassar R, Bennett BD, Babu-Khan S, *et al.* Beta-secretase cleavage of Alzheimer's amyloid precursor protein by the transmembrane aspartic protease BACE. *Science* 1999; 286(5440): 735-41.
- [62] Small SA, Gandy S. Sorting through the cell biology of Alzheimer's disease: intracellular pathways to pathogenesis. *Neuron* 2006; 52(1): 15-31.
- [63] Rajendran L, Honsho M, Zahn TR, *et al.* Alzheimer's disease beta-amyloid peptides are released in association with exosomes. *Proc Natl Acad Sci USA* 2006; 103(30): 11172-7.
- [64] He X, Cooley K, Chung CH, Dashti N, Tang J. Apolipoprotein receptor 2 and X11 alpha/beta mediate apolipoprotein E-induced endocytosis of amyloid-beta precursor protein and beta-secretase, leading to amyloid-beta production. *J Neurosci* 2007; 27(15): 4052-60.
- [65] Roberson ED, Mucke L. 100 years and counting: prospects for defeating Alzheimer's disease. *Science* 2006; 314(5800): 781-4.
- [66] Haass C, Steiner H. Alzheimer disease gamma-secretase: a complex story of GxGD-type presenilin proteases. *Trends Cell Biol* 2002; 12(12): 556-62.
- [67] Wolfe MS. When loss is gain: reduced presenilin proteolytic function leads to increased Abeta42/Abeta40. Talking Point on the role of presenilin mutations in Alzheimer disease. *EMBO Rep* 2007; 8(2): 136-40.
- [68] Li YM, Xu M, Lai MT, *et al.* Photoactivated gamma-secretase inhibitors directed to the active site covalently label presenilin 1. *Nature* 2000; 405(6787): 689-94.
- [69] Esler WP, Kimberly WT, Ostaszewski BL, *et al.* Transition-state analogue inhibitors of gamma-secretase bind directly to presenilin-1. *Nat Cell Biol* 2000; 2(7): 428-34.
- [70] Xia W, Ray WJ, Ostaszewski BL, *et al.* Presenilin complexes with the C-terminal fragments of amyloid precursor protein at the sites of amyloid beta-protein generation. *Proc Natl Acad Sci USA* 2000; 97(16): 9299-304.
- [71] Shen J, Bronson RT, Chen DF, Xia W, Selkoe DJ, Tonegawa S. Skeletal and CNS defects in Presenilin-1-deficient mice. *Cell* 1997; 89(4): 629-39.
- [72] Baki L, Shioi J, Wen P, *et al.* PS1 activates PI3K thus inhibiting GSK-3 activity and tau overphosphorylation: effects of FAD mutations. *EMBO J* 2004; 23(13): 2586-96.
- [73] Burgering BM, Coffey PJ. Protein kinase B (c-Akt) in phosphatidylinositol-3-OH kinase signal transduction. *Nature* 1995; 376(6541): 599-602.
- [74] Dehvari N, Cedazo-Minguez A, Isacson O, *et al.* Presenilin dependence of phospholipase C and protein kinase C signaling. *J Neurochem* 2007; 102(3): 848-57.
- [75] Rocchi A, Pellegrini S, Siciliano G, Murri L. Causative and susceptibility genes for Alzheimer's disease: a review. *Brain Res Bull* 2003; 61(1): 1-24.
- [76] Cowburn RF, Popescu BO, Ankarcrona M, Dehvari N, Cedazo-Minguez A. Presenilin-mediated signal transduction. *Physiol Behav* 2007; 92(1-2): 93-7.

- [77] Cheung KH, Shineman D, Muller M, *et al.* Mechanism of Ca²⁺ disruption in Alzheimer's disease by presenilin regulation of InsP3 receptor channel gating. *Neuron* 2008; 58(6): 871-83.
- [78] Ebinu JO, Yankner BA. A RIP tide in neuronal signal transduction. *Neuron* 2002; 34(4): 499-502.
- [79] Marambaud P, Shioi J, Serban G, *et al.* A presenilin-1/gamma-secretase cleavage releases the E-cadherin intracellular domain and regulates disassembly of adherens junctions. *EMBO J* 2002; 21(8): 1948-56.
- [80] Tandon A, Fraser P. The presenilins. *Genome Biol* 2002; 3(11): reviews3014.
- [81] Dewji NN, Singer SJ. Specific intercellular binding of the beta-amyloid precursor protein to the presenilins induces intercellular signaling: its significance for Alzheimer's disease. *Proc Natl Acad Sci USA* 1998; 95(25): 15055-60.
- [82] Laferla FM. Calcium dyshomeostasis and intracellular signalling in Alzheimer's disease. *Nat Rev Neurosci* 2002; 3(11): 862-72.
- [83] St George-Hyslop PH. Molecular genetics of Alzheimer's disease. *Biol Psychiatry* 2000; 47(3): 183-199.
- [84] Rogaev EI, Sherrington R, Rogaeva EA, *et al.* Familial Alzheimer's disease in kindreds with missense mutations in a gene on chromosome 1 related to the Alzheimer's disease type 3 gene. *Nature* 1995; 376(6543): 775-8.
- [85] Geling A, Steiner H, Willem M, Bally-Cuif L, Haass C. A gamma-secretase inhibitor blocks Notch signaling *in vivo* and causes a severe neurogenic phenotype in zebrafish. *EMBO Rep* 2002; 3(7): 688-94.
- [86] Wong GT, Manfra D, Poulet FM, *et al.* Chronic treatment with the gamma-secretase inhibitor LY-411,575 inhibits beta-amyloid peptide production and alters lymphopoiesis and intestinal cell differentiation. *J Biol Chem* 2004; 279(13): 12876-82.
- [87] Vassar R. Beta-secretase (BACE) as a drug target for Alzheimer's disease. *Adv Drug Deliv Rev* 2002; 54(12): 1589-602.
- [88] Zhao J, Fu Y, Yasvoina M, *et al.* Beta-site amyloid precursor protein cleaving enzyme 1 levels become elevated in neurons around amyloid plaques: implications for Alzheimer's disease pathogenesis. *J Neurosci* 2007; 27(14): 3639-49.
- [89] Wang Q, Rowan MJ, Anwyl R. Beta-amyloid-mediated inhibition of NMDA receptor-dependent long-term potentiation induction involves activation of microglia and stimulation of inducible nitric oxide synthase and superoxide. *J Neurosci* 2004; 24(27): 6049-56.
- [90] Parameswaran K, Dhanasekaran M, Suppiramaniam V. Amyloid beta peptides and glutamatergic synaptic dysregulation. *Exp Neurol* 2008; 210(1): 7-13.
- [91] Parameswaran K, Sims C, Kanju P, *et al.* Amyloid beta-peptide Abeta(1-42) but not Abeta(1-40) attenuates synaptic AMPA receptor function. *Synapse* 2007; 61(6): 367-74.
- [92] Gasparini L, Dityatev A. Beta-amyloid and glutamate receptors. *Exp Neurol* 2008; 212(1): 1-4.
- [93] Pak T, Cadet P, Mantione KJ, Stefano GB. Morphine *via* nitric oxide modulates beta-amyloid metabolism: a novel protective mechanism for Alzheimer's disease. *Med Sci Monit* 2005; 11(10): BR357-66.
- [94] Heneka MT, Wiesinger H, Dumitrescu-Ozimek L, Riederer P, Feinstein DL, Klockgether T. Neuronal and glial coexpression of argininosuccinate synthetase and inducible nitric oxide synthase in Alzheimer disease. *J Neuropathol Exp Neurol* 2001; 60(9): 906-16.
- [95] Francis PT, Palmer AM, Snape M, Wilcock GK. The cholinergic hypothesis of Alzheimer's disease: a review of progress. *J Neurol Neurosurg Psychiatry* 1999; 66(2): 137-47.
- [96] Rupasingh R, Borrie M, Smith M, Wells JL, Bartha R. Reduced hippocampal glutamate in Alzheimer disease. *Neurobiol Aging* 2009.
- [97] Schaeffer EL, Gattaz WF. Cholinergic and glutamatergic alterations beginning at the early stages of Alzheimer disease: participation of the phospholipase A2 enzyme. *Psychopharmacology (Berl)* 2008; 198(1): 1-27.
- [98] Nathan C, Calingasan N, Nezezon J, *et al.* Protection from Alzheimer's-like disease in the mouse by genetic ablation of inducible nitric oxide synthase. *J Exp Med* 2005; 202(9): 1163-9.
- [99] Moncada S. Nitric oxide. *J Hypertens Suppl* 1994; 12(10): S35-9.
- [100] Schmidt HH, Walter U. NO at work. *Cell* 1994; 78(6): 919-25.
- [101] Moncada S, Palmer RM, Higgs EA. Nitric oxide: physiology, pathophysiology, and pharmacology. *Pharmacol Rev* 1991; 43(2): 109-42.
- [102] Law A, Gauthier S, Quirion R. Say NO to Alzheimer's disease: the putative links between nitric oxide and dementia of the Alzheimer's type. *Brain Res Brain Res Rev* 2001; 35(1): 73-96.
- [103] Togo T, Katsuse O, Iseki E. Nitric oxide pathways in Alzheimer's disease and other neurodegenerative dementias. *Neurol Res* 2004; 26(5): 563-6.
- [104] Lipton SA. Failures and successes of NMDA receptor antagonists: molecular basis for the use of open-channel blockers like memantine in the treatment of acute and chronic neurologic insults. *NeuroRx* 2004; 1(1): 101-10.
- [105] Hellstrom-Lindahl E. Modulation of beta-amyloid precursor protein processing and tau phosphorylation by acetylcholine receptors. *Eur J Pharmacol* 2000; 393(1-3): 255-63.
- [106] Rosengarten B, Paulsen S, Molnar S, Kaschel R, Gallhofer B, Kaps M. Acetylcholine esterase inhibitor donepezil improves dynamic cerebrovascular regulation in Alzheimer patients. *J Neurol* 2006; 253(1): 58-64.
- [107] Raina P, Santaguida P, Ismaila A, *et al.* Effectiveness of cholinesterase inhibitors and memantine for treating dementia: evidence review for a clinical practice guideline. *Ann Intern Med* 2008; 148(5): 379-97.
- [108] Galimberti D, Scarpini E. Treatment of Alzheimer's disease: symptomatic and disease-modifying approaches. *Curr Aging Sci* 2010; 3(1): 46-56.
- [109] Fernandez-Bachiller MI, Perez C, Campillo NE, *et al.* Tacrine-melatonin hybrids as multifunctional agents for Alzheimer's disease, with cholinergic, antioxidant, and neuroprotective properties. *ChemMedChem* 2009; 4(5): 828-41.
- [110] Chabrier PE, Merle-Pallardy C, Auguet M. Nitric oxide synthases: targets for therapeutic strategies in neurological diseases. *Cell Mol Life Sci* 1999; 55(8-9): 1029-35.
- [111] Law A, Gauthier S, Quirion R. Neuroprotective and neurorescuing effects of isoform-specific nitric oxide synthase inhibitors, nitric oxide scavenger, and antioxidant against beta-amyloid toxicity. *Br J Pharmacol* 2001; 133(7): 1114-24.
- [112] Xie QW, Cho HJ, Calaycay J, *et al.* Cloning and characterization of inducible nitric oxide synthase from mouse macrophages. *Science* 1992; 256(5054): 225-8.
- [113] Vodovotz Y, Lucia MS, Flanders KC, *et al.* Inducible nitric oxide synthase in tangle-bearing neurons of patients with Alzheimer's disease. *J Exp Med* 1996; 184(4): 1425-33.
- [114] Chiarini A, Dal P I, Marconi M, Chakravarthy B, Whitfield JF, Armato U. Calcium-sensing receptor (CaSR) in human brain's pathophysiology: roles in late-onset Alzheimer's disease (LOAD). *Curr Pharm Biotechnol* 2009; 10(3): 317-26.
- [115] Ali AK, Banks WA, Kumar VB, *et al.* Nitric oxide activity and isoenzyme expression in the senescence-accelerated mouse p8 model of Alzheimer's disease: effects of anti-amyloid antibody and antisense treatments. *J Gerontol A Biol Sci Med Sci* 2009; 64(10): 1025-30.
- [116] Lee SC, Zhao ML, Hirano A, Dickson DW. Inducible nitric oxide synthase immunoreactivity in the Alzheimer disease hippocampus: association with Hirano bodies, neurofibrillary tangles, and senile plaques. *J Neuropathol Exp Neurol* 1999; 58(11): 1163-9.
- [117] Wong A, Luth HJ, Uther-Conrad W, *et al.* Advanced glycation endproducts co-localize with inducible nitric oxide synthase in Alzheimer's disease. *Brain Res* 2001; 920(1-2): 32-40.
- [118] Shibata N, Kobayashi M. [The role for oxidative stress in neurodegenerative diseases]. *Brain Nerve* 2008; 60(2): 157-70.
- [119] Wang B, Tan S, Yang Z, *et al.* Association between Alzheimer's disease and the NOS3 gene Glu298Asp polymorphism in Chinese. *J Mol Neurosci* 2008; 34(2): 173-6.
- [120] Murphy MP, Packer MA, Scarlett JL, Martin SW. Peroxynitrite: a biologically significant oxidant. *Gen Pharmacol* 1998; 31(2): 179-86.
- [121] Koppal T, Drake J, Yatin S, *et al.* Peroxynitrite-induced alterations in synaptosomal membrane proteins: insight into oxidative stress in Alzheimer's disease. *J Neurochem* 1999; 72(1): 310-7.
- [122] Luth HJ, Munch G, Arendt T. Aberrant expression of NOS isoforms in Alzheimer's disease is structurally related to nitrotyrosine formation. *Brain Res* 2002; 953(1-2): 135-43.
- [123] Rodrigo J, Fernandez-Vizarra P, Castro-Blanco S, *et al.* Nitric oxide in the cerebral cortex of amyloid-precursor protein (SW) Tg2576 transgenic mice. *Neuroscience* 2004; 128(1): 73-89.

- [124] Law A, Gauthier S, Quirion R. Say NO to Alzheimer's disease: the putative links between nitric oxide and dementia of the Alzheimer's type. *Brain Res Brain Res Rev* 2001; 35(1): 73-96.
- [125] Flirski M, Sobow T. Biochemical markers and risk factors of Alzheimer's disease. *Curr Alzheimer Res* 2005; 2(1): 47-64.
- [126] Mesenge C, Charriaud-Marlangue C, Verrecchia C, Allix M, Boulu RR, Plotkine M. Reduction of tyrosine nitration after N(omega)-nitro-L-arginine-methylester treatment of mice with traumatic brain injury. *Eur J Pharmacol* 1998; 353(1): 53-7.
- [127] Murphy GM, Jr., Yang L, Cordell B. Macrophage colony-stimulating factor augments beta-amyloid-induced interleukin-1, interleukin-6, and nitric oxide production by microglial cells. *J Biol Chem* 1998; 273(33): 20967-71.
- [128] Serrano J, Fernandez AP, Martinez-Murillo R, MartInez A. High sensitivity to carcinogens in the brain of a mouse model of Alzheimer's disease. *Oncogene* 2009; In press.
- [129] Meda L, Cassatella MA, Szendrei GI, et al. Activation of microglial cells by beta-amyloid protein and interferon-gamma. *Nature* 1995; 374(6523): 647-50.
- [130] Kim YA, Lim SY, Rhee SH, Park KY, Kim CH, Choi BT et al. Resveratrol inhibits inducible nitric oxide synthase and cyclooxygenase-2 expression in beta-amyloid-treated C6 glioma cells. *Int J Mol Med* 2006; 17(6): 1069-75.
- [131] Apelt J, Bigl M, Wunderlich P, Schliebs R. Aging-related increase in oxidative stress correlates with developmental pattern of beta-secretase activity and beta-amyloid plaque formation in transgenic Tg2576 mice with Alzheimer-like pathology. *Int J Dev Neurosci* 2004; 22(7): 475-84.
- [132] Lee J, Chan SL, Mattson MP. Adverse effect of a presenilin-1 mutation in microglia results in enhanced nitric oxide and inflammatory cytokine responses to immune challenge in the brain. *Neuromolecular Med* 2002; 2(1): 29-45.
- [133] de la Monte SM, Wands JR. Molecular indices of oxidative stress and mitochondrial dysfunction occur early and often progress with severity of Alzheimer's disease. *J Alzheimers Dis* 2006; 9(2): 167-81.
- [134] Duncan AJ, Heales SJ. Nitric oxide and neurological disorders. *Mol Aspects Med* 2005; 26(1-2): 67-96.
- [135] Medeiros R, Prediger RD, Passos GF, et al. Connecting TNF-alpha signaling pathways to iNOS expression in a mouse model of Alzheimer's disease: relevance for the behavioral and synaptic deficits induced by amyloid beta protein. *J Neurosci* 2007; 27(20): 5394-404.
- [136] Stepanichev MY, Onufriev MV, Yakovlev AA, et al. Amyloid-beta (25-35) increases activity of neuronal NO-synthase in rat brain. *Neurochem Int* 2008; 52(6): 1114-24.
- [137] de la Monte SM, Chiche J, von dem BA, et al. Nitric oxide synthase-3 overexpression causes apoptosis and impairs neuronal mitochondrial function: relevance to Alzheimer's-type neurodegeneration. *Lab Invest* 2003; 83(2): 287-98.
- [138] Lahiri DK, Chen D, Ge YW, et al. Does nitric oxide synthase contribute to the pathogenesis of Alzheimer's disease?: effects of beta-amyloid deposition on NOS in transgenic mouse brain with AD pathology. *Ann N Y Acad Sci* 2003; 1010: 639-42.
- [139] Kadoszkiewicz H, Zimmermann T, Beck-Bornholdt HP, van den BH. Cholinesterase inhibitors for patients with Alzheimer's disease: systematic review of randomised clinical trials. *BMJ* 2005; 331(7512): 321-7.
- [140] Walker LC, Ibegbu CC, Todd CW, et al. Emerging prospects for the disease-modifying treatment of Alzheimer's disease. *Biochem Pharmacol* 2005; 69(7): 1001-8.
- [141] Sabbagh MN, Farlow MR, Relkin N, Beach TG. Do cholinergic therapies have disease-modifying effects in Alzheimer's disease? *Alzheimers Dement* 2006; 2(2): 118-25.
- [142] Thatcher GR, Bennett BM, Reynolds JN. Nitric oxide mimetic molecules as therapeutic agents in Alzheimer's disease. *Curr Alzheimer Res* 2005; 2(2): 171-82.
- [143] Das UN. Folic acid and polyunsaturated fatty acids improve cognitive function and prevent depression, dementia, and Alzheimer's disease--but how and why? *Prostaglandins Leukot Essent Fatty Acids* 2008; 78(1): 11-19.
- [144] Takatori Y. [Mechanisms of neuroprotective effects of therapeutic acetylcholinesterase inhibitors used in treatment of Alzheimer's disease]. *Yakugaku Zasshi* 2006; 126(8): 607-16.
- [145] Li W, Mak M, Jiang H, et al. Novel anti-Alzheimer's dimer Bis(7)-cognitin: cellular and molecular mechanisms of neuroprotection through multiple targets. *Neurotherapeutics* 2009; 6(1): 187-201.
- [146] Maczurek A, Hager K, Kenkies M, et al. Lipoic acid as an anti-inflammatory and neuroprotective treatment for Alzheimer's disease. *Adv Drug Deliv Rev* 2008; 60(13-14): 1463-70.
- [147] in t, V, Ruitenbergh A, Hofman A, Launer LJ, et al. Nonsteroidal antiinflammatory drugs and the risk of Alzheimer's disease. *N Engl J Med* 2001; 345(21): 1515-1521.
- [148] Zandi PP, Anthony JC, Hayden KM, Mehta K, Mayer L, Breitner JC. Reduced incidence of AD with NSAID but not H2 receptor antagonists: the Cache County Study. *Neurology* 2002; 59(6): 880-6.
- [149] Griffin WS, Nicoll JA, Grimaldi LM, Sheng JG, Mrak RE. The pervasiveness of interleukin-1 in alzheimer pathogenesis: a role for specific polymorphisms in disease risk. *Exp Gerontol* 2000; 35(4): 481-7.
- [150] Wyss-Coray T, Masliah E, Mallory M, et al. Amyloidogenic role of cytokine TGF-beta1 in transgenic mice and in Alzheimer's disease. *Nature* 1997; 389(6651): 603-6.
- [151] Jiang B, Du J, Liu JH, Bao YM, An LJ. Catalpol attenuates the neurotoxicity induced by beta-amyloid(1-42) in cortical neuron-glia cultures. *Brain Res* 2008; 1188: 139-47.
- [152] Chung HY, Cesari M, Anton S, et al. Molecular inflammation: underpinnings of aging and age-related diseases. *Ageing Res Rev* 2009; 8(1): 18-30.
- [153] Asanuma M, Miyazaki I, Tsuji T, Ogawa N. New aspects of neuroprotective effects of nonsteroidal anti-inflammatory drugs. *Nihon Shinkei Seishin Yakurigaku Zasshi* 2003; 23(3): 111-9.
- [154] Dokmeci D. Ibuprofen and Alzheimer's disease. *Folia Med (Plovdiv)* 2004; 46(2): 5-10.
- [155] Gasparini L, Ongini E, Wilcock D, Morgan D. Activity of flurbiprofen and chemically related anti-inflammatory drugs in models of Alzheimer's disease. *Brain Res Brain Res Rev* 2005; 48(2): 400-8.
- [156] Dheen ST, Jun Y, Yan Z, Tay SS, Ling EA. Retinoic acid inhibits expression of TNF-alpha and iNOS in activated rat microglia. *Glia* 2005; 50(1): 21-31.
- [157] Jana M, Jana A, Liu X, Ghosh S, Pahan K. Involvement of phosphatidylinositol 3-kinase-mediated up-regulation of I kappa B alpha in anti-inflammatory effect of gemfibrozil in microglia. *J Immunol* 2007; 179(6): 4142-52.
- [158] Ayasolla K, Khan M, Singh AK, Singh I. Inflammatory mediator and beta-amyloid (25-35)-induced ceramide generation and iNOS expression are inhibited by vitamin E. *Free Radic Biol Med* 2004; 37(3): 325-38.
- [159] Chiarugi A, Moskowitz MA. Poly(ADP-ribose) polymerase-1 activity promotes NF-kappaB-driven transcription and microglial activation: implication for neurodegenerative disorders. *J Neurochem* 2003; 85(2): 306-17.
- [160] Scali C, Prosperi C, Vannucchi MG, Pepeu G, Casamenti F. Brain inflammatory reaction in an animal model of neuronal degeneration and its modulation by an anti-inflammatory drug: implication in Alzheimer's disease. *Eur J Neurosci* 2000; 12(6): 1900-12.
- [161] Farooqui AA, Yang HC, Horrocks L. Involvement of phospholipase A2 in neurodegeneration. *Neurochem Int* 1997; 30(6): 517-22.
- [162] Hicks JB, Lai Y, Sheng W, et al. Amyloid-beta peptide induces temporal membrane biphasic changes in astrocytes through cytosolic phospholipase A2. *Biochim Biophys Acta* 2008; 1778(11): 2512-9.
- [163] Stephenson DT, Lemere CA, Selkoe DJ, Clemens JA. Cytosolic phospholipase A2 (cPLA2) immunoreactivity is elevated in Alzheimer's disease brain. *Neurobiol Dis* 1996; 3(1): 51-63.
- [164] Ross BM, Moszczynska A, Erlich J, Kish SJ. Phospholipid-metabolizing enzymes in Alzheimer's disease: increased lysophospholipid acyltransferase activity and decreased phospholipase A2 activity. *J Neurochem* 1998; 70(2): 786-93.
- [165] Zhou L, Zhu DY. Neuronal nitric oxide synthase: structure, subcellular localization, regulation, and clinical implications. *Nitric Oxide* 2009; 20(4): 223-30.
- [166] Lockhart BP, Benicourt C, Junien JL, Privat A. Inhibitors of free radical formation fail to attenuate direct beta-amyloid25-35 peptide-mediated neurotoxicity in rat hippocampal cultures. *J Neurosci Res* 1994; 39(4): 494-505.
- [167] Dvir E, Elman A, Simmons D, et al. DP-155, a lecithin derivative of indomethacin, is a novel nonsteroidal antiinflammatory drug for

- analgesia and Alzheimer's disease therapy. *CNS Drug Rev* 2007; 13(2): 260-77.
- [168] Chalimoniuk M, Glowacka J, Zabiela A, Eckert A, Strosznajder JB. Nitric oxide alters arachidonic acid turnover in brain cortex synaptoneurosomes. *Neurochem Int* 2006; 48(1): 1-8.
- [169] Devan BD, Bowker JL, Duffy KB, *et al.* Phosphodiesterase inhibition by sildenafil citrate attenuates a maze learning impairment in rats induced by nitric oxide synthase inhibition. *Psychopharmacology (Berl)* 2006; 183(4): 439-45.
- [170] Corbin JD, Beasley A, Blount MA, Francis SH. Vardenafil: structural basis for higher potency over sildenafil in inhibiting cGMP-specific phosphodiesterase-5 (PDE5). *Neurochem Int* 2004; 45(6): 859-63.
- [171] Mullershausen F, Russwurm M, Koesling D, Friebe A. *In vivo* reconstitution of the negative feedback in nitric oxide/cGMP signaling: role of phosphodiesterase type 5 phosphorylation. *Mol Biol Cell* 2004; 15(9): 4023-30.
- [172] Puzzo D, Staniszewski A, Deng SX, *et al.* Phosphodiesterase 5 inhibition improves synaptic function, memory, and amyloid-beta load in an Alzheimer's disease mouse model. *J Neurosci* 2009; 29(25): 8075-86.
- [173] Uthayathas S, Karuppagounder SS, Thrash BM, Parameswaran K, Suppiramaniam V, Dhanasekaran M. Versatile effects of sildenafil: recent pharmacological applications. *Pharmacol Rep* 2007; 59(2): 150-63.
- [174] Domek-Lopacinska K, Strosznajder JB. Cyclic GMP metabolism and its role in brain physiology. *J Physiol Pharmacol* 2005; 56 Suppl 2: 15-34.
- [175] Kleppisch T, Wolfgruber W, Feil S, *et al.* Hippocampal cGMP-dependent protein kinase I supports an age- and protein synthesis-dependent component of long-term potentiation but is not essential for spatial reference and contextual memory. *J Neurosci* 2003; 23(14): 6005-12.
- [176] Schubert P, Morino T, Miyazaki H, *et al.* Cascading glia reactions: a common pathomechanism and its differentiated control by cyclic nucleotide signaling. *Ann N Y Acad Sci* 2000; 903: 24-33.
- [177] Frampton M, Harvey RJ, Kirchner V. Propentofylline for dementia. *Cochrane Database Syst Rev* 2003; (2): CD002853.
- [178] Barad M, Bourtochouladze R, Winder DG, Golan H, Kandel E. Rolipram, a type IV-specific phosphodiesterase inhibitor, facilitates the establishment of long-lasting long-term potentiation and improves memory. *Proc Natl Acad Sci USA* 1998; 95(25): 15020-5.
- [179] Gong B, Vitolo OV, Trinchese F, Liu S, Shelanski M, Arancio O. Persistent improvement in synaptic and cognitive functions in an Alzheimer mouse model after rolipram treatment. *J Clin Invest* 2004; 114(11): 1624-34.
- [180] Arendash GW, Schleif W, Rezai-Zadeh K, *et al.* Caffeine protects Alzheimer's mice against cognitive impairment and reduces brain beta-amyloid production. *Neuroscience* 2006; 142(4): 941-52.
- [181] Dall'Igna OP, Fett P, Gomes MW, Souza DO, Cunha RA, Lara DR. Caffeine and adenosine A(2a) receptor antagonists prevent beta-amyloid (25-35)-induced cognitive deficits in mice. *Exp Neurol* 2007; 203(1): 241-5.
- [182] Poucher SM, Keddie JR, Singh P, *et al.* The *in vitro* pharmacology of ZM 241385, a potent, non-xanthine A2a selective adenosine receptor antagonist. *Br J Pharmacol* 1995; 115(6): 1096-102.
- [183] Baratti CM, Boccia MM, Blake MG. Pharmacological effects and behavioral interventions on memory consolidation and reconsolidation. *Braz J Med Biol Res* 2009; 42(2): 148-54.
- [184] Southan GJ, Szabo C. Selective pharmacological inhibition of distinct nitric oxide synthase isoforms. *Biochem Pharmacol* 1996; 51(4): 383-94.
- [185] de la Torre JC, Aliev G. Inhibition of vascular nitric oxide after rat chronic brain hypoperfusion: spatial memory and immunocytochemical changes. *J Cereb Blood Flow Metab* 2005; 25(6): 663-72.
- [186] Wilcock DM, Lewis MR, Van Nostrand WE, *et al.* Progression of amyloid pathology to Alzheimer's disease pathology in an amyloid precursor protein transgenic mouse model by removal of nitric oxide synthase 2. *J Neurosci* 2008; 28(7): 1537-45.
- [187] Colton CA, Vitek MP, Wink DA, *et al.* NO synthase 2 (NOS2) deletion promotes multiple pathologies in a mouse model of Alzheimer's disease. *Proc Natl Acad Sci USA* 2006; 103(34): 12867-72.
- [188] Salerno L, Sorrenti V, Di GC, Romeo G, Siracusa MA. Progress in the development of selective nitric oxide synthase (NOS) inhibitors. *Curr Pharm Des* 2002; 8(3): 177-200.
- [189] Moore PK, al-Swayeh OA, Chong NW, Evans RA, Gibson A. L-NG-nitro arginine (L-NOARG), a novel, L-arginine-reversible inhibitor of endothelium-dependent vasodilatation *in vitro*. *Br J Pharmacol* 1990; 99(2): 408-12.
- [190] Babu BR, Griffith OW. N5-(1-Imino-3-butenyl)-L-ornithine. A neuronal isoform selective mechanism-based inactivator of nitric oxide synthase. *J Biol Chem* 1998; 273(15): 8882-9.
- [191] Furfine ES, Harmon MF, Paith JE, Garvey EP. Selective inhibition of constitutive nitric oxide synthase by L-NG-nitroarginine. *Biochemistry* 1993; 32(33): 8512-7.
- [192] Narayanan K, Griffith OW. Synthesis of L-thiocitrulline, L-homothiocitrulline, and S-methyl-L-thiocitrulline: a new class of potent nitric oxide synthase inhibitors. *J Med Chem* 1994; 37(7): 885-7.
- [193] Narayanan K, Spack L, McMillan K, *et al.* S-alkyl-L-thiocitrullines. Potent stereoselective inhibitors of nitric oxide synthase with strong pressor activity *in vivo*. *J Biol Chem* 1995; 270(19): 11103-10.
- [194] Furfine ES, Harmon MF, Paith JE, *et al.* Potent and selective inhibition of human nitric oxide synthases. Selective inhibition of neuronal nitric oxide synthase by S-methyl-L-thiocitrulline and S-ethyl-L-thiocitrulline. *J Biol Chem* 1994; 269(43): 26677-83.
- [195] Ulhaq S, Chinje EC, Naylor MA, Jaffar M, Stratford IJ, Threadgill MD. Heterocyclic analogues of L-citrulline as inhibitors of the isoforms of nitric oxide synthase (NOS) and identification of N(delta)-(4,5-dihydrothiazol-2-yl)ornithine as a potent inhibitor. *Bioorg Med Chem* 1999; 7(9): 1787-96.
- [196] Eustache J, Grob A, Lam C, Sellier O, Schulz G. Conformationally constrained NO synthase inhibitors: rigid analogs of L-N-iminoethylornithine. *Bioorg Med Chem Lett* 1998; 8(21): 2961-6.
- [197] Huang H, Martasek P, Roman LJ, Masters BS, Silverman RB. N(omega)-Nitroarginine-containing dipeptide amides. Potent and highly selective inhibitors of neuronal nitric oxide synthase. *J Med Chem* 1999; 42(16): 3147-53.
- [198] Kobayashi N, Higuchi T, Urano Y, Kikuchi K, Hirobe M, Nagano T. Dipeptides containing L-arginine analogs: new isozyme-selective inhibitors of nitric oxide synthase. *Biol Pharm Bull* 1999; 22(9): 936-40.
- [199] Sohn MJ, Hur GM, Byun HS, Kim WG. Cyclo(dehydrohistidyl-L-tryptophyl) inhibits nitric oxide production by preventing the dimerization of inducible nitric oxide synthase. *Biochem Pharmacol* 2008; 75(4): 923-30.
- [200] Cross AH, Misko TP, Lin RF, Hickey WF, Trotter JL, Tilton RG. Aminoguanidine, an inhibitor of inducible nitric oxide synthase, ameliorates experimental autoimmune encephalomyelitis in SJL mice. *J Clin Invest* 1994; 93(6): 2684-90.
- [201] Garvey EP, Oplinger JA, Tanoury GJ, *et al.* Potent and selective inhibition of human nitric oxide synthases. Inhibition by non-amino acid isothioureas. *J Biol Chem* 1994; 269(43): 26669-76.
- [202] Moore WM, Webber RK, Fok KF, *et al.* 2-Iminopiperidine and other 2-iminoazaheterocycles as potent inhibitors of human nitric oxide synthase isoforms. *J Med Chem* 1996; 39(3): 669-72.
- [203] Naka M, Nanbu T, Kobayashi K, *et al.* A potent inhibitor of inducible nitric oxide synthase, ONO-1714, a cyclic amide derivative. *Biochem Biophys Res Commun* 2000; 270(2): 663-7.
- [204] Babbidge RC, Bland-Ward PA, Hart SL, Moore PK. Inhibition of rat cerebellar nitric oxide synthase by 7-nitro indazole and related substituted indazoles. *Br J Pharmacol* 1993; 110(1): 225-8.
- [205] Yoshida T, Limmroth V, Irikura K, Moskowitz MA. The NOS inhibitor, 7-nitroindazole, decreases focal infarct volume but not the response to topical acetylcholine in pial vessels. *J Cereb Blood Flow Metab* 1994; 14(6): 924-9.
- [206] Schulz JB, Beal MF. Neuroprotective effects of free radical scavengers and energy repletion in animal models of neurodegenerative disease. *Ann N Y Acad Sci* 1995; 765: 100-10.
- [207] White KA, Marletta MA. Nitric oxide synthase is a cytochrome P-450 type hemoprotein. *Biochemistry* 1992; 31(29): 6627-31.
- [208] Wolff DJ, Datto GA, Samatovicz RA, Tempick RA. Calmodulin-dependent nitric-oxide synthase. Mechanism of inhibition by imidazole and phenylimidazoles. *J Biol Chem* 1993; 268(13): 9425-9.
- [209] Rogerson TD, Wilkinson CF, Hetarski K. Steric factors in the inhibitory interaction of imidazoles with microsomal enzymes. *Biochem Pharmacol* 1977; 26(11): 1039-42.
- [210] Mayer B, Werner ER. In search of a function for tetrahydrobiop-terin in the biosynthesis of nitric oxide. *Naunyn Schmiedeberg Arch Pharmacol* 1995; 351(5): 453-63.

- [211] Bommel HM, Reif A, Frohlich LG, *et al.* Anti-pterins as tools to characterize the function of tetrahydrobiopterin in NO synthase. *J Biol Chem* 1998; 273(50): 33142-9.
- [212] Terpolilli NA, Zweckberger K, Trabold R, *et al.* The novel nitric oxide synthase inhibitor 4-amino-tetrahydro-L-biopterine prevents brain edema formation and intracranial hypertension following traumatic brain injury in mice. *J Neurotrauma* 2009; 26(11): 1963-75.
- [213] Monsonogo A, Weiner HL. Immunotherapeutic approaches to Alzheimer's disease. *Science* 2003; 302(5646): 834-8.
- [214] Gelinas DS, DaSilva K, Fenili D, St George-Hyslop P, McLaurin J. Immunotherapy for Alzheimer's disease. *Proc Natl Acad Sci USA* 2004; 101 Suppl 2: 14657-62.
- [215] Weiner HL, Frenkel D. Immunology and immunotherapy of Alzheimer's disease. *Nat Rev Immunol* 2006; 6(5): 404-16.
- [216] Kellner A, Matschke J, Bernreuther C, Moch H, Ferrer I, Glatzel M. Autoantibodies against beta-amyloid are common in Alzheimer's disease and help control plaque burden. *Ann Neurol* 2009; 65(1): 24-31.
- [217] Klyubin I, Walsh DM, Lemere CA, *et al.* Amyloid beta protein immunotherapy neutralizes Abeta oligomers that disrupt synaptic plasticity *in vivo*. *Nat Med* 2005; 11(5): 556-61.
- [218] Weldon DT, Rogers SD, Ghilardi JR, *et al.* Fibrillar beta-amyloid induces microglial phagocytosis, expression of inducible nitric oxide synthase, and loss of a select population of neurons in the rat CNS *in vivo*. *J Neurosci* 1998; 18(6): 2161-73.
- [219] Cho DH, Nakamura T, Fang J, *et al.* S-nitrosylation of Drp1 mediates beta-amyloid-related mitochondrial fission and neuronal injury. *Science* 2009; 324(5923): 102-5.

Received: July 9, 2010

Accepted: July 21, 2010