



ARTICLE

Unsaturated fatty acids: Nutritionally essential, or toxic?

In 1929 George and Mildred Burr published a paper claiming that unsaturated fats, and specifically linoleic acid, were essential to prevent a particular disease involving dandruff, dermatitis, slowed growth, sterility, and fatal kidney degeneration.

In 1929, most of the B vitamins and essential trace minerals were unknown to nutritionists. The symptoms the Burrs saw are easily produced by deficiencies of the vitamins and minerals that they didn't know about.

What really happens to animals when the "essential fatty acids" are lacking, in an otherwise adequate diet?

Their metabolic rate is very high.

Their nutritional needs are increased.

They are very resistant to many of the common causes of sickness and death.

They are resistant to the biochemical and cellular changes seen in aging, dementia, autoimmunity, and the main types of inflammation.

The amount of polyunsaturated fatty acids often said to be essential (Holman, 1981) is approximately the amount required to significantly increase the incidence of cancer, and very careful food selection is needed for a diet that provides a lower amount.

When I was studying the age pigment, lipofuscin, and its formation from polyunsaturated fatty acids, I saw the 1927 study in which a fat free diet practically eliminated the development of spontaneous cancers in rats (Bernstein and Elias). I have always wondered whether George and Mildred Burr were aware of that study in 1929, when they published their claim that polyunsaturated fats are nutritionally essential. The German study was abstracted in Biological Abstracts, and the Burrs later cited several studies from German journals, and dismissively mentioned two U.S. studies* that claimed animals could live on fat-free diets, so their neglect of such an important claim is hard to understand. (*Their bibliography cited, without further comment, Osborne and Mendel, 1920, and Drummond and Coward, 1921.)

Since 1927, others have demonstrated that the polyunsaturated fats are essential for the development of cancer (and some other degenerative diseases), but the Burrs' failed to even mention the issue at any time during their careers. How could they, studying fat-free diets, have missed an important contemporary publication, if I, 40 years later, saw it? There were very few publications on dietary fats in those years, so it was hardly possible to miss it.

When researchers at the Clayton Foundation Biochemical Institute at the University of Texas demonstrated that "Burr's disease" was actually a vitamin B6 deficiency, rather than a fatty acid deficiency, the issue was settled. Later studies failed to confirm the existence of the Burr disease caused by a deficiency of fatty acids, though many similar conditions were produced by a variety of other dietary defects. In 1938, a group in Burr's own laboratory (Brown, et al.) failed to produce dermatitis in a man during a six month experiment. Neither of the other major features of the Burr disease, male sterility and kidney degeneration, has been subsequently confirmed. The claim that polyunsaturated fatty acid deficiency caused sterility of male animals ("A new and uniform cause of sterility is shown") was quickly dropped, probably because an **excess** of polyunsaturated fats was discovered to be an important cause of testicular degeneration and sterility.

HOME
ARTICLES
ABOUT RAY PEAT
ART GALLERY
LINKS

Loading

www.RayPeat.com
©2006-16 Ray Peat
All Rights Reserved

One of the features of the Burrs' rats on the fat-free diet was that they ate more calories and drank much more water than the rats that received polyunsaturated fatty acids in their diet. They believed that the animals were unable to synthesize fat without linoleic acid, although in another context they cited a study in which the fat of rats on a fat-free diet was similar in composition to lard: "McAmis, Anderson, and Mendel [37] fed rats a high sucrose, fat-free diet and rendered the fat of the entire animal. This fat had an iodine number of 64 to 71, a fairly normal value for lard."

The "wasteful" food consumption, and the leanness of animals that weren't fed polyunsaturated fats became fairly common knowledge by the late 1940s, but no one repeated the Burrs' claim that the absence of those fatty acids led quickly to the animals' death. Meanwhile, "crazy chick disease" caused by feeding an excess of polyunsaturated fats, and a little later, "yellow fat disease," caused by too much fish fat, were being recognized by farmers. In the 1950s, the seed oil industry created the anti-cholesterol diet culture, and a few decades later, without any new "Burr-like" publications, the omega minus 3 oils, especially fish oils, were coming to be represented as the overlooked essential fatty acids, which were capable of preventing the toxic effects of the original "essential" linoleic acid.

Although the 1929 Burr paper is still often cited as proof of the essentiality of PUFA, Burr's younger colleague (at the University of Minnesota Hormel Institute), Ralph Holman, has cited an infant (1970), and a 78 year old woman (in 1969), who developed dermatitis while receiving fat-free intravenous feedings. Dermatitis, with dandruff, similar to Burr's disease, has been produced by various nutritional deficiencies besides vitamin B6, including a trace mineral deficiency and a biotin deficiency, so there is no valid reason to associate dermatitis with a fat deficiency. The cases of "EFA deficiency" produced by intravenous feedings that have been widely cited were probably the result of a deficiency of zinc or other trace mineral, since so-called "Total Parenteral Nutrition" was in use for many years before the trace minerals were added to the "total" formula. In 1975, I learned that our local hospital was putting all premature babies on what they called total intravenous feeding, without trace minerals, for weeks, or months. There is still more emphasis on polyunsaturated fat in intravenous feeding than on the essential trace nutrients.

Holman and the Hormel Institute have been extremely influential in promoting the doctrine of the essentiality of PUFA, including fish oils and the omega -3 oils, but their best evidence, the Burr experiment, doesn't make their case. Far worse than that is the effect it has had in distracting attention from the profoundly toxic effects of the so-called essential fatty acids. Long after he should have known better, Holman was arguing that butter was a nutritionally inferior fat.

When the Burrs were doing their study, Raymond Pearl was one of the most famous biologists in the country, and his "rate of living" theory of aging was very widely known. According to that theory, an organism has an intrinsic potential to produce a certain total amount of energy during its lifetime, and if it metabolizes at a higher than normal rate, its life span will be proportionately shorter than normal.

There is general agreement that animals on a fat free diet have a very high metabolic rate, but the people who believe the "rate of living" theory will be inclined to see the increased rate of metabolism as something harmful in itself. It is clear that this is what the Burrs thought. They didn't attempt to provide a diet that provided increased amounts of all vitamins and minerals, in proportion to the increased metabolic rate.

Pearl did an experiment, sprouting cantaloupe seeds in a dish with water. The sprouts that grew rapidly died sooner than those that grew more slowly. They died as soon as the nutrients stored in the endosperm had been consumed. Naturally, when nutrients are depleted, growth and metabolism must stop. If food and air and water are rationed, then slow metabolizers are going to live longer. But when nutritional needs are met, the organisms with the highest metabolic rate generally are healthier and live longer. In a study of nurses, those who habitually consumed the most calories lived longer than those who consumed the least. Even while Pearl was promoting his theory, other famous biologists, for example John Northrup in Jacques Loeb's lab at the

Rockefeller Institute, were making observations that contradicted the rate of living theory. For example, around 1916, Northrup observed that fruit flies that metabolized at the highest rate lived the longest. Northrup was doing biology, Pearl was doing propaganda, following Weismannism.

The idea of extending life span by slowing metabolism and growth was a logical implication of the "rate of living" theory of aging, and it's an idea that is still popular. Many people have supposed that eating less would slow metabolism. Caloric restriction does extend the life span of many species, but it generally preserves the high metabolic rate of youth, so that at a given age the calorie-restricted animal has a higher rate of oxygen consumption per gram of body weight than the unrestricted eaters.

Roy Walford, a gerontologist who wrote about extending the human life span to 120 years by caloric restriction, spent 30 years limiting his diet to about 1600 calories, with little animal protein, almost no saturated fat--fish once or twice per week, poultry or beef about once, and a fat free milkshake for breakfast--and after about 15 years, began developing a degenerative brain disease, ALS, one of the nerve diseases involving lipid peroxidation and excitotoxicity. When he died from the disease, he had lived a year longer than the normal life expectancy.

V. Stefannson, one of the early polar explorers, spent a winter living entirely on caribou meat, and felt that it had prevented the scurvy that had killed so many of the other explorers, who had counted on fruit and vegetables to prevent it. But he believed that meat was a metabolic stimulant that made people age prematurely, as Pearl's rate of living theory predicted. Stefannson said that Eskimo women were getting old in their twenties, and that at the age of 60 they looked as old as Europeans did at 80. He was a well informed anthropologist, and his observations were probably accurate. The Eskimos he observed ate large amounts of fish, and other unsaturated fats, and sometimes ate highly decayed fish. An accelerated rate of aging would be expected from such a diet, because of the toxic lipid peroxides.

Calorie-restricted animals (on a diet of normal composition) have a lower degree of fat unsaturation in their mitochondria as they age, preserving the relatively more saturated fats of youth.

Birds' mitochondrial fats are much less polyunsaturated than those of mammals, and birds' metabolic rates are much higher, and they live much longer than mammals of a similar size.

With aging, the highly peroxidizable fatty acids, arachidonic and docosahexaenoic acid, increase greatly in a variety of tissues, and lipid peroxidation increases with aging. Peroxidation slows mitochondrial respiration, lowering the metabolic rate. Caloric restriction slows the accumulation of the highly unsaturated fatty acids in mitochondria, and reduces peroxidation.

Over the years, it has become evident that the polyunsaturated fats are not very compatible with a high rate of metabolism, though they are necessary for organisms that live at low temperatures and metabolize slowly, such as fish and vegetables. The saturated fats solidify at low temperature; beef fat is very stiff at refrigerator temperature, and in a fat fish, such stiffness would be lethal.

Even some hibernating rodents can stay alive with their body tissues close to the freezing point, and their stored fats have to be unsaturated. When their diet doesn't allow them to store enough polyunsaturated fat, they fail to go into hibernation. This is probably a clue to some of the general biological effects of the PUFA.

A series of studies about 20 years ago showed that the functions of the thyroid hormone are all inhibited by unsaturated fats, with the inhibition increasing in proportion to the number of unsaturations (double bonds) in the fat molecule.

When the tissues are saturated with those antithyroid fats, metabolism slows, especially when any stress, such as cold or hunger, increases the concentration of free fatty acids in the blood stream. Stress and hypothyroidism increase the formation of serotonin, which is an important factor in producing the torpor of hibernation, and lowering the body temperature. The polyunsaturated fatty acids themselves directly contribute to the formation of serotonin, for example by increasing the

ability of tryptophan to enter the brain. In a certain cold climate, the PUFA are essential for hibernation, but under other conditions, the rodent would be able to continue gathering food and eating, instead of hibernating.

The direct effects of the PUFA on the endocrine and nervous systems, as illustrated by the hibernating squirrel, interact with their effects on intercellular communication (including the formation of prostaglandins and related substances), and the effects of their oxidative breakdown products, such as acrolein. But the people who claim that they are absolutely, rather than conditionally, essential, base their argument on the idea that they are needed for the formation of prostaglandins and cell membranes. The fact that cells can replicate in fat free conditions shows that the argument from membranes is unfounded. The argument from prostaglandins is more complex, but has no firmer foundation.

When a dose of PUFA is administered to a lizard, which isn't a hibernator, the lizard's body temperature is lowered by several degrees. There are probably many ways in which the PUFA produce that effect, besides increasing serotonin and decreasing thyroid. The PUFA are increased by estrogen, and they increase estrogen, and have some directly estrogen-like effects. Estrogen itself tends to lower body temperature and shift metabolism away from oxidative energy production. Aging, like estrogen, increases the body's content of the PUFA: Linoleic, linolenic, dihomo-gamma-linolenic, docosahexaenoic and docosapentaenoic acids are increased by age, and the longer chain acids increase more rapidly in women than in men (Bolton-Smith, et al., 1997). (Women are apparently relatively protected by progesterone, which inhibits lipolysis and prostaglandin formation, and protects the brain, thymus, and other tissues from lipid peroxidation and other effects of the PUFA.)

Aging involves a decreasing metabolic rate, an increased tendency toward inflammation, and a decreased ability to synthesize proteins. Inflammation contributes to the decreasing ability to use oxygen, and the slowed renewal of proteins combined with lower ability to produce energy impair the organism's ability to control peroxidative damage and inflammation.

The fragments of deteriorating PUFA combine with proteins and other cell materials, producing immunogenic substances. The so-called "advanced glycation end products," that have been blamed on glucose excess, are mostly derived from the peroxidation of the "essential fatty acids." The name, "glycation," indicates the addition of sugar groups to proteins, such as occurs in diabetes and old age, but when tested in a controlled experiment, **lipid peroxidation of polyunsaturated fatty acids produces the protein damage about 23 times faster than the simple sugars do** (Fu, et al., 1996).

Several autoimmune disease models in animals (involving the eye, kidney, and pancreas) have been prevented by a deficiency of the EFA (Schreiner, et al., 1989, Bazan, et al., 1990, Benhamou, et al., 1995).

Besides causing a general slowing of metabolism, aging and toxic PUFA have specific actions on the detoxifying system. The enzymes that help to detoxify PUFA and estrogen and serotonin are inhibited by both PUFA and estrogen. All systems, including blood vessels and the intestine, are made leaky by estrogen and the PUFA and their products. A reduced ability to regulate the excitatory amino acids, resulting from PUFA toxins, tends to produce excitotoxicity, damaging nerves (Ou, et al., 2002).

Although the interplay of the various types of nerve is very complex, a variety of experiments suggest that the PUFA are acting directly on serotonergic nerves, rather than just increasing the conversion of tryptophan to serotonin.

For example, a deficiency of the so-called essential fatty acids, EFA, makes animals more sensitive to some anesthetics, and more resistant to others. It makes them resistant to the anesthetics that act by promoting the actions of serotonin, but it prolongs the effects of those that don't act through serotonin, and these are the anesthetics such as xenon and nitrous oxide, that apparently act by stabilizing the structure of water, as described by Linus Pauling. Progesterone and the saturated fats seem to act partly through the stabilizing of cell water, and estrogen and the PUFA have opposing effects, creating cellular excitation while

interfering with the stable cellular water structure.

Serotonin interferes with slow wave sleep, and promotes cortisol, both of which can be harmful to brain cells. (Hypothyroidism is one of the causes of a decrease in slow wave sleep.) Babies whose mothers' serum contained more DHA were more wakeful on their second day of life, than the babies of low-DHA mothers. The amide of oleic acid is a sleep promoter, with apparent antiserotonin activity (Yang, et al., 2003), and since oleic acid tends to be displaced by diets high in PUFA, this suggests another way in which the highly unsaturated fatty acids could promote serotonin's effects.

People who don't have a normal amount of slow wave sleep are likely to have slow reaction times when they are awake, and quickness of reactions is a good indicator of general intelligence.

Manufacturers of baby formulas are claiming that the highly unsaturated fatty acids accelerate brain development, but they neglect to mention studies that show either no effect, or retardation of development. In some of the tests that are used to measure infant development, a generalized state of arousal or anxiety could be interpreted as "more mature."

In one experiment, animals that received less than 0.32% of their calories as EFA grew slightly less than rats on a standard diet, but their brains were as large as those of normal rats (Bruckner, et al., 1984). That is, their brain to body ratio was a little larger than normal, which is a typical feature of individuals with a higher metabolic rate. That result is very different from the claims of the baby food industry, that the brain is the organ most easily damaged by a PUFA deficiency.

One of the standard signs of toxicity is the enlargement of the spleen and liver, and that effect is produced by larger amounts of the EFA. The weight of the thymus is reduced by PUFA in the diet (Guimarães, et al., 1990). Thymus cells tend to be easily killed by a combination of stress and EFA, and bone marrow cells, though less sensitive than thymic cells, are damaged by lipid peroxidation of the PUFA. The effects of PUFA on the thymus were compared to those of radiation by Soviet researchers. Immunodeficiency, produced largely by damage to thymic cells, increases when larger amounts of PUFA are eaten for a prolonged time.

The growth and metastasis of a variety of tumors are inhibited by saturated fatty acids, and increased by fish oil--as much as 10 times in number of metastases, 1000 times in size (Griffini, et al., 1998).

Mothers whose breast milk contains more long-chain n-3 fatty acids are more likely to have allergic children (Stoney, et al., 2004). (And children whose mothers are allergic have higher levels of DHA and EPA in their tissues.) These associations aren't mentioned by the manufacturers who speak of those fats as essential.

When animals have been "deprived" of the EFA during gestation and nursing, and then given a standard diet, they develop larger bones, with a thicker cortex and more trabecular bone, both of which would suggest a lower level of stress. Many types of inflammation and stress are significantly reduced in "EFA deficient" animals. Inflammation caused by the injection of carrageenan is decreased, partly because of the absence of prostaglandins in these animals. The absence of the EFA protects against colitis and nephritis (). The kidneys are more effective in several ways in the deficient animals.

Shock, caused by the injection of endotoxin, which is 100% lethal to normally fed animals, is only 24% lethal to the deficient animals.

Poisons are much less harmful to deficient animals, for example, a cobra venom factor causes less tissue damage to their lungs.

Concussive trauma and burns cause much less damage to deficient animals.

The endothelial lining of blood vessels is protected by saturated fats and oleic acid, damaged by polyunsaturated, and their barrier function is improved by the absence of PUFA.

Alzheimer's disease, retinal degeneration, cataracts, and liver cirrhosis all involve reactions of oxidized PUFA with proteins. Saturated fats help to heal alcoholic liver cirrhosis.

The lesions of atherosclerosis and cataracts contain some of the same oxidized lipids as the age pigment itself. When large deposits of age pigment become visible, it's probably because the general reduction of metabolism and protein synthesis has interfered with the normal processes for removing debris. The age pigment contributes to degeneration by wasting energy and oxygen, weakening the antioxidation, antiglycation, and other defensive systems.

The EFA amplify nearly all kinds of injury and stress, and the results of many recent publications make it look as though serotonin interacts harmfully with the EFA in most of these situations. The specific balance of polyunsaturated fatty acids, and their various breakdown products, from carbon monoxide, glyoxal, and acrolein, to the larger aldehydes and radicals, and the stress-induced substances such as serotonin, histamine, estrogens, can produce an immense variety of biological problems.

When the various claims of an EFA "deficiency disease" or syndrome or symptom are examined, their inconsistency over the years makes skepticism seem increasingly justified. The Burrs' publications were typical of others, in failing to describe and account for the evidence that contradicted their claims. Claiming that certain fatty acids are essential, a scientific approach would require showing what was wrong with the experiments that showed that they were not essential, and especially, those that showed that they were positively harmful.

In this culture that repeatedly makes such claims of essentiality, the growing number of reports of biological superiority of "deficient" animals suggests that nutritional research may be near the point at which it can resume the line of study begun by Northrup, Osborne, Mendel, Drummond, Bernstein, Elias, and others, that was interrupted for 60 years by industrial interests that promoted antiscientific opinions.

For example, in 1914 F.P. Rous showed that limiting food intake reduced the incidence of cancer, and then in 1915 and 1917, Osborne and Mendel showed that food restriction extended the fertility and longevity of female rats. The association between estrogen and cancer had become known during this time, and vitamin E, which was originally known as the fertility vitamin, was soon recognized to have antiestrogenic properties, as well as to prevent the deadly effects of excessive polyunsaturated fats in the diet. My endocrinology professor, A.S. Soderwall, who had found that excess estrogen prevented (or interrupted) pregnancy, demonstrated that increased vitamin E extended fertility in aging female rodents.

By the time I began my research, it seemed clear that it had been the reduction of PUFA in the diet which, like the addition of vitamin E, had prevented sterility in the calorie restriction experiments, and that those treatments had limited the effects of estrogen in the aging organisms.

Estrogen, by activating phospholipase A2, acts to amplify the toxic effects of PUFA in the tissues, and these effects increase with age, and with decreased amounts of thyroid and progesterone.

Antioxidants can slightly retard the cumulative degenerative effects of the fats interacting with estrogen, serotonin, and other mediators of inflammation, but real elimination of the degenerative diseases will require an exploration of the effects of the entire series of lipid signalling substances derived from the saturated and omega minus 9 fatty acids.

REFERENCES

J. Nutr. 2000 Nov;130(11):2855-65. **Low arachidonic acid rather than alpha-tocopherol is responsible for the delayed postnatal development in offspring of rats fed fish oil instead of olive oil during pregnancy and lactation.** Amusquivar E, Ruperez FJ, Barbas C, Herrera E.

Dev. Psychobiol. 8(1), 59-66, 1985. **"Effects of a commercial soy lecithin preparation on development of sensorimotor behavior and brain biochemicals in the rat,"** Bell, J. M. and P. K. Lundberg.

Zeitschr. Krebsforsch. 28(1), 1-14, 1927 **"Lipoids and carcinoma growth,"** Bernstein, S. and Elias, H.

J. Biol. Chem. 82:345-367.(1929) **A new deficiency disease produced by the rigid exclusion of fat from the diet.** Burr, G. & Burr, M.

J. Biol. Chem. 86:587-621. (1930) **On the nature and role of the fatty acids essential in nutrition.** Burr, G. & Burr, M.

J Natl Cancer Inst. 1984 Jul;73(1):185-91. **Dietary lipid effects on the growth, membrane composition, and prolactin-binding capacity of rat mammary tumors.** Cave WT Jr, Jurkowski JJ. "Our results indicated that 1) when the polyunsaturated lipid component (corn oil) of the diet exceeded 3%, it was the quantitative level of total lipid, rather than the level of polyunsaturated lipid alone, that best correlated with the observed reduction in tumor latent period; 2) when the polyunsaturated lipid content of the diet fell below 3%, there was a decrease in tumor incidence and an increase in the mean latent period...."

Lipids. 1997 Sep;32(9):979-88. **Modulation of adjuvant-induced arthritis by dietary arachidonic acid in essential fatty acid-deficient rats.** Chinn KS, Welsch DJ, Salsgiver WJ, Mehta A, Raz A, Obukowicz MG. "Controlled feeding of linoleic acid (LA) or arachidonic acid (AA) to essential fatty acid-deficient (EFAD) rats was used to define the relationship between dietary AA and the inflammatory response evoked during adjuvant-induced arthritis." "Compared to rats fed the standard laboratory chow diet (Control), edema in the primary hind footpads was decreased by 87% in EFAD, 71% in EFAD + 1x AA, 45% in EFAD + 10x AA, and 30% in EFAD + 0.5x LA. The decrease in edema in the footpads of EFAD rats was nearly identical to the decrease in edema in the footpads of Control rats dosed with indomethacin. Hind footpad edema correlated with the final AA plasma level and eicosanoid levels extracted from hind footpad tissue, but not with neutrophil infiltration."

J Nutr. 1996 Jun;126(6):1534-40. **Dietary (n-9) eicosatrienoic acid from a cultured fungus inhibits leukotriene B4 synthesis in rats and the effect is modified by dietary linoleic acid.** Cleland LG, Gibson RA, Neumann MA, Hamazaki T, Akimoto K, James MJ.

J Nutr. 1981 Nov;111(11):2039-43. **Effects of dietary fatty acids on delayed-type hypersensitivity in mice.** Dewille JW, Fraker PJ, Romsos DR. Effects of an essential fatty acid deficient (EFAD) [0% corn oil (CO)] diet and a diet high in polyunsaturated fatty acids [PUFA (50% CO)] on one aspect of in vivo T cell function [delayed-type hypersensitivity (DTH)] were assessed. After a 70-day feeding trial, **DTH was reduced by 30% in mice fed the EFAD diet**, but the response of mice fed the high PUFA diet equaled that of control mice fed a diet containing 13% CO. **The time required for the EFAD diet to reduce DTH was 42 days. Although consumption of the EFAD diet reduced DTH**, this reduction was rapidly reversed, within 7 days, by switching the EFAD mice to the control diet. These results indicate that :1) consumption of the EFAD diet reduces one aspect of in vivo T cell function (DTH), but the effect can be reversed by refeeding the control diet; and 2) a high PUFA diet does not adversely affect DTH.

Biochem. J., 1917, xi, 325. **A comparative study of tumour and normal tissue growth.** Drummond, JC

J. Physiol. 1920, liv, p. xxx, Proc. Physiol. Soc. **Nutrition on diets practically devoid of fats.** Drummond, JC.

J. Gerontol. 30(6), 647-54, 1975, **"Rat brain fatty acid composition: effect of dietary fat and age,"** Eddy, D. E., and D. Harman.

J Biol Chem 1996 Apr 26;271(17):9982-6. **The advanced glycation end product, Nepsilon-(carboxymethyl)lysine, is a product of both lipid peroxidation and glycoxidation reactions.** Fu MX, Requena JR, Jenkins AJ, Lyons TJ, Baynes JW, Thorpe SR.

Pediatrics 31:171-192, 1963. Hansen AE, Wiese HF, Boelsche AN, Haggard ME, Adam DJD, Davis H.

Am J Clin Nutr 49:290-300, 1982. Holman RT, Johnson SB, Hatch TF.

Cancer Res. 45(5), 1997-2001, 1985. **"Requirement of essential fatty acids for mammary tumor,"** Ip, C., et al.

Food and Nutrition Board. 1989. **Recommended Dietary Allowances** 10th ed. Washington, DC: National Academy of Sciences. p 48.

Am J Clin Nutr. 2004 Nov;80(5):1167-74. **Docosahexaenoic acid concentrations are higher in women than in men because of estrogenic effects.** Giltay EJ, Gooren LJ, Toorians AW, Katan MB, Zock PL. "The proportion of DHA was 15 +/- 4% (x +/- SEM; P < 0.0005) higher in the women than in the men. **Among the women, those taking oral contraceptives had 10 +/- 4% (P = 0.08) higher DHA concentrations than did those not taking oral contraceptives. Administration of oral ethinyl estradiol, but not transdermal 17beta-estradiol, increased DHA by 42 +/- 8% (P < 0.0005), whereas the antiandrogen cyproterone acetate did not affect DHA.** Parenteral testosterone decreased DHA by 22 +/- 4% (P < 0.0005) in female-to-male transsexual subjects. Anastrozole decreased estradiol concentrations significantly and DHA concentrations nonsignificantly (9 +/- 6%; P = 0.09). **CONCLUSION: Estrogens cause higher DHA concentrations in women than in men, probably by upregulating synthesis of DHA from vegetable precursors.**"

J. American Geriatrics Soc. 24(1) 292-8, 1976, **"Free radical theory of aging: effect of dietary fat on central nervous system function,"** Harman, D., et al.

Adv Exp Med Biol 266:3-15, 1989, **"Lipofuscin and ceroid formation: the cellular recycling system,"** Harman, D.

J Theor Biol. 2005 May 21;234(2):277-88. Epub 2005 Jan 24. **On the importance of fatty acid composition of membranes for aging.** Hulbert AJ.

Biochim Biophys Acta. 2005 Jan 5;1686 3:248-54. **Perinatal essential fatty acid deficiency influences body weight and bone parameters in adult male rats.** Korotkova M, Ohlsson C, Gabrielsson B, Hanson LA, Strandvik B. Department of Pediatrics, Goteborg University, The Queen Silvia Children's Hospital, SE 41685 Goteborg, Sweden. Marina.Korotkova@cmm.ki.se Fetal and postnatal nutrition have long-term effects on the risk for development of diseases late in life in humans and animals. The aim of the present study was to investigate the effect of dietary deficiency of essential fatty acids (EFA) in the perinatal period on later body weight and bone mass. **During late gestation and throughout lactation**, rats were fed a control or an EFA-deficient (EFAD) diet. At 3 weeks of age the offspring were weaned onto an ordinary chow and followed until adult age. The mean body weight of adult rats receiving the EFAD diet during the perinatal period was significantly increased from 12 weeks of age compared to the controls ($P < 0.05$). Analysis by peripheral quantitative computerized tomography (pQCT) at 44 weeks of age showed that the trabecular volumetric bone mineral density (BMD) of the femur was significantly decreased ($P < 0.05$) **but the cortical bone mineral content, cortical area, and cortical thickness were increased** ($P < 0.05$) in the EFAD group of rats. The length of the femur was not affected. **In conclusion, neonatal EFA deficiency was in adult rats associated with increased body weight and significant changes in both cortical and trabecular bone. The results indicate that regulatory mechanisms related to bone mass seemed to be programmed by EFA in the perinatal period.** The nature of this modulation needs to be identified.

Biochem Biophys Res Commun. 1987 Jun 30;145(3):1185-91. **Anti-lipoperoxidation action of food restriction.** Laganier S, Yu BP. "Chronic food restriction inhibited the age-related increase of malondialdehyde production and lipid hydroperoxides in liver mitochondrial and microsomal membranes of ad libitum fed Fischer 344 rats. The anti-lipoperoxidation action of food restriction could not be attributable to the changes in membrane lipid content nor vitamin E status. **Restricting calories modified membrane fatty acid composition by increasing linoleic acid and decreasing docosapentaenoic acid** content in both membranes."

Lipids 22(3), 133-6, 1987.. **"Effects of parenteral nutrition with high doses of linoleate on the developing human liver and brain,"** Martinez, M., and A. Ballabriga.

J Nutrit 10:63(1935). **"The effect of retarded growth upon length of the life span and upon the ultimate body size."** McCay, CM., Crowell, MF., and Maynard, LA.

McCollum EV. 1957. **A History of Nutrition.** Boston: Houghton Mifflin. p 374.

J Clin Invest 73:272-276,1984. Neuringer M, Connor WE, Van Petten C, Barstad L.

J. Biol. Chem., 1912, xii, 81. **Feeding experiments with fat-free food mixtures.** Osborne, TB., Mendel, LB.

Science 45:294-295. (1917). **"The effect of retardation of growth upon the breeding period and duration of life in rats."** Osborne, TB., Mendel, LB., and Fey, EL.

Gerontology 1993;39(1):7-18. **Modulation of membrane phospholipid fatty acid composition by age and food restriction.** Laganier S, Yu BP. H.M. "Phospholipids from liver mitochondrial and microsomal membrane preparations were analyzed to further assess the effects of age and lifelong calorie restriction on membrane lipid composition." **"The data revealed characteristic patterns of age-related changes in ad libitum (AL) fed rats: membrane levels of long-chain polyunsaturated fatty acids, 22:4 and 22:5, increased progressively, while membrane linoleic acid (18:2) decreased steadily with age. Levels of 18:2 fell by approximately 40%, and 22:5 content almost doubled making the peroxidizability index increase with age."** **"We concluded that the membrane-stabilizing action of long-term calorie restriction relates to the selective modification of membrane long-chain polyunsaturated fatty acids during aging."**

Free Radic Biol Med 1999 Feb;26(3-4):260-5. **Modulation of cardiac mitochondrial membrane fluidity by age and calorie intake.** Lee J, Yu BP, Herlihy JT. "The fatty acid composition of the mitochondrial membranes of the two **ad lib fed groups differed: the long-chain polyunsaturated 22:4 fatty acid was higher in the older group, although linoleic acid (18:2) was lower. DR eliminated the differences.**" "Considered together, these results suggest that DR maintains the integrity of the cardiac mitochondrial membrane fluidity by **minimizing membrane damage through modulation of membrane fatty acid profile.**"

Lipids 2001 Jun;36(6):589-93. **Effect of dietary restriction on age-related increase of liver susceptibility to peroxidation in rats.** Leon TI, Lim BO, Yu BP, Lim Y, Jeon EJ, Park DK.

Neurobiol Aging. 1982 Fall;3(3):173-8. **Lipid peroxides in brain during aging and vitamin E deficiency: possible relations to changes in neurotransmitter indices.** Noda Y, McGeer PL, McGeer EG. "Lipid peroxide levels, were found to be **significantly higher in brains of 18 month old as compared to 4 month old rats, with particularly large increases** occurring in the olfactory bulb, globus pallidus, cerebral cortex and caudate-putamen (CP). Eighteen month old rats fed a **vitamin E deficient diet for 9 months before sacrifice had lipid peroxide levels significantly higher than age-matched controls in the cerebral cortex, hippocampus and**

hypothalamus." "Age-related decreases were seen in choline acetyltransferase, acetylcholinesterase and 3H-QNB binding in some but not all brain regions, while GABA transaminase and MAO showed age-related increases." "As compared with controls, vitamin E deficient rats showed decreases of 38% in cortical 3H-DHA binding, of 33% in 3H-QNB binding in the CP and of 23% and 12% in choline acetyltransferase in the CP and cerebellum, respectively."

J Immunol, 1989 Nov 15, 143:10, 3192-9. **The antiinflammatory effects of essential fatty acid deficiency in experimental glomerulonephritis. The modulation of macrophage migration and eicosanoid metabolism.** Schreiner GF, Rovin B, Lefkowitz JB.

Br J Nutr. 2003 Aug;90(2):385-94. **Dietary docosahexaenoic acid-induced production of tissue lipid peroxides is not suppressed by higher intake of ascorbic acid in genetically scorbutic Osteogenic Disorder Shionogi/Shi-od/od rats.** Sekine S, Kubo K, Tadokoro T, Maekawa A, Saito M.

Cancer Res. 2004 Sep 15;64(18):6482-8. **Estrogen enhances angiogenesis through a pathway involving platelet-activating factor-mediated nuclear factor-kappaB activation.** Seo KH, Lee HS, Jung B, Ko HM, Choi JH, Park SJ, Choi IH, Lee HK, Im SY.

Clin Exp Allergy. 2004 Feb;34(2):194-200. **Maternal breast milk long-chain n-3 fatty acids are associated with increased risk of atopy in breastfed infants.** Stoney RM, Woods RK, Hosking CS, Hill DJ, Abramson MJ, Thien FC.

Proc Natl Acad Sci U S A. 1988 Aug;85(16):6137-41. **Essential fatty acid deficiency prevents multiple low-dose streptozotocin-induced diabetes in CD-1 mice.** Wright JR Jr, Lefkowitz JB, Schreiner G, Lacy PE.

Mech Ageing Dev. 2005 May 10; [Epub ahead of print] **Membrane alteration as a basis of aging and the protective effects of calorie restriction.** Yu BP.

Acta Paediatr 1996 Jun;85(6):679-83. **Phospholipid fatty acids in cord blood: family history and development of allergy.** Yu G, Kjellman NI, Bjorksten B.

Fiziol Cheloveka. 2005 Mar-Apr;31(2):108-15. **[Age-related changes in lipid peroxidation in various regions of the central nervous system]** [Article in Russian] Volchegorskii IA, Shemiakov SE, Telesheva IB, Malinovskaia NV, Turygin VV.

Surg Today. 2003;33(8):600-5. **Beneficial effects of n-9 eicosatrienoic acid on experimental bowel lesions.** Yoshida H, Soh H, Sando K, Wasa M, Takagi Y, Okada A. **PURPOSE:** Dietary fortification of n-9 polyunsaturated fatty acids (PUFA) or 5,8,11-eicosatrienoic acid (ETrA) as well as n-3 PUFA might contribute to the suppression of leukotriene B4 (LTB4) synthesis and thereby reduce inflammatory bowel lesions. As a result, the effect of an ETrA-enriched diet on experimental bowel lesions was examined in this study. **METHODS:** In Expt. 1, rats were freely fed either an ETrA-enriched or a standard diet. After 7 days of feeding, acute bowel lesions were induced by the subcutaneous injection of 10 mg/kg indomethacin. In Expt. 2, chronic bowel lesions were made by performing subcutaneous injections of 7.5 mg/kg indomethacin twice. After the first injection, the rats were freely fed either an ETrA-enriched or a standard diet for 7 days. **RESULTS:** In both experiments, the rats fed an ETrA-enriched diet showed increased levels of ETrA in the plasma and intestinal mucosa, and a decreased inflammation score. However, there was no significant decrease in plasma and intestinal mucosal LTB4 in the ETrA-enriched diet-fed rats. **CONCLUSION:** These results suggest that the dietary supplementation of ETrA may have both prophylactic and therapeutic effects on experimentally produced bowel lesions. Further investigations are necessary to clarify the effects of ETrA on bowel lesions and its mechanisms.

© Ray Peat Ph.D. 2007. All Rights Reserved. www.RayPeat.com

^^^ **Top** ^^^