

Ray Peat's Newsletter

If others had not been foolish, we should be so. William Blake

Copyright 2011

Raymond Peat P.O. Box 5764 Eugene OR 97405

May 2011

Not for republication without written permission.

Endotoxin, stress, depression: Serotonin, starches, fatty acids, and antidotes

Increasingly, intestinal serotonin is being recognized as a factor in osteoporosis, migraine, irritable bowel syndrome, appendicitis, and cancer. From ancient times until about 100 years ago, the bowel had a very important place in medical practice, with purgatives and enemas being recommended for headaches, persistent coughs, seizures, and many chronic diseases. Pavlov studied the digestive system as a way to understand the mind, because it's where we interact most closely with the environment.

Up to Pavlov's time, traditional medicine thought in terms of achieving balance in the organism, using metaphors such as the four humors (phlegm, blood, yellow bile and black bile) and the four personality types---phlegmatic, sanguine, choleric, and melancholic.

In western Europe and the US during the 19th century, divisions in medical doctrine began to appear between traditional, "empirical" practitioners and advocates of a new "rational," reductionist allopathic doctrine who were coming to believe that disease had its origin in cells and their defects of form or function. These "scientific physicians" began a major propaganda campaign to ridicule blood-letting and purging as examples of the unscientific practices of their competitors. The idea of internal toxins itself was ridiculed. A gastroenterology professor (Walter C. Alvarez) used his newspaper column to argue against the old ideas, for example that toxins absorbed from the intestine caused headaches. He described putting wads of cotton into the rectums of his medical students, and a considerable number of them developed headaches while the cotton was in

place. That, he asserted, disproved the idea that absorbed toxins were the cause of headaches. It didn't really disprove anything, but it contributed to the culture that was discouraging the use of laxatives. Russian researchers, who were still working in Pavlov's holistic tradition, found that inflating a balloon in an animal's intestine didn't have any observable effect in a normal animal, but when the animal's blood sugar was lowered, the stretching of the intestine would produce a wide range of symptoms, from asthma to seizures. It is likely that some of Alvarez's students had low blood glucose, and were more susceptible than others to the abnormal stretching of the rectum.

Inflammation of the bowel used to be recorded frequently as the cause of death of older people, but now it's seldom mentioned in autopsy reports as the cause of death, if it's mentioned at all.

Hans Selye had noticed that intestinal bleeding was a standard feature of being sick, and it became a defining feature of his concept of stress. Old people sometimes get ulcers in their colon or rectum, adjacent to lumps of hardened fecal material. Selye devised animal experiments, in which he showed that an interference with blood circulation that normally didn't produce serious harm, would cause similar (stercoral) ulcers, and also paralysis of the hind legs, if he injected serotonin into the animal. If the anti-serotonin drug, cyproheptadine, was given to the animal before the serotonin, they were protected. The ulcers required the combination of restricted blood circulation, excess serotonin, and the presence of feces in the lower intestine. Without the serotonin, the other two factors failed to produce the ulcers (1966). In a previous experiment, he had demonstrated that the combination of impaired circulation with serotonin would cause severe muscular dystrophy (1965).

Selye's research group showed how closely the digestive system and the nervous system interact, but they didn't go very far in exploring the effects of diet on the nervous system.

Just friction, or scratching or stretching the intestine is enough to cause it to release serotonin into the bloodstream. Serotonin increases the permeability of the intestine and blood vessels, and so is likely to be a major cause of the absorption of endotoxin (and other harmful material) during intestinal irritation or stress. The biological meaning of serotonin might be very different without endotoxin, but that hasn't been investigated.

Bacterial endotoxin increases serotonin release from the intestine, and increases its synthesis in the brain (Nolan, et al., 2000) and liver (Endo, 1983). It also stimulates its release from platelets, and reduces the lungs' ability to destroy it. The formation of serotonin in the intestine is also stimulated by the lactate, propionate and butyrate that are formed by bacteria fermenting fiber and starch, but these bacteria also produce endotoxin. The inflammation-producing effects of lactate, serotonin, and endotoxin are overlapping, additive, and sometimes synergistic, along with histamine, nitric oxide, bradykinin, and the cytokines.

There have been some studies showing that bacterial fermentation in the intestine can cause many symptoms, including behavioral changes.

For example, a diet of soy protein increases aggression in monkeys (Simon, et al., 2004) and chickens (McKeegan, et al., 2001).

When various fermentable carbohydrates were fed to rats, they became anxious and aggressive, and these changes in behavior corresponded to the fermentation of these materials by bacteria in the lower intestine, with the production of lactic acid (Hanstock, et al., 2003, 2004).

Dietary fiber has been promoted for a long time, and for many reasons---John Harvey Kellogg argued that it prevented dirty thoughts, the cholesterol people argued that it would lower cholesterol, Denis Burkitt, in the 1970s, suggested that it could prevent colon cancer. When Burkitt visited the US and saw that oat bran was being recommended, he commented that the main fiber

in the African diet he had studied was potato. By the 1980s, studies in animals were showing that some kinds of fiber, including **oat bran, pectin, corn bran, metamucil, undegraded carrageenan, agar, psyllium, guar gum, and alfalfa**, were carcinogenic, increasing the incidence of colon cancer, and possibly ovarian and breast cancer (Toth, 1984, Jacobs, 1986, Goodman, et al., 2002).

Many advocates of increasing dietary fiber argue that butyric acid is a protective factor produced by bacterial decomposition of starch and fiber, and it does (at least in vitro) sometimes cause cancer cells to differentiate into normal cells. However, it accelerates the growth of intestinal cells, and when that is combined with toxins it might act as a tumor promoter. The lactic acid produced by bacteria and by the inflamed intestine itself is now recognized as an important factor in the development of cancer.

The upper part of the small intestine is sterile in healthy people. In the last 40 years, there has been increasing interest in the "contaminated small-bowel syndrome," or the "small intestine bacterial overgrowth syndrome." When peristalsis is reduced, for example by hypothyroidism, along with reduced secretion of digestive fluids, bacteria are able to thrive in the upper part of the intestine. Sugars are very quickly absorbed in the upper intestine, so starches and fibers normally provide most of the nourishment for bowel bacteria.

Besides avoiding gums and other "soluble fiber," and starches that aren't quickly digested, there are several things that can protect against the toxic effects of bacterial degradation products. Beans are famous for their gas-producing polysaccharides, but all grains and vegetables and firm under-ripe fruits contain starches that resist our digestive enzymes, but that can provide food for bacteria. Undercooked starch, and starches combined with enzyme inhibitors (including polyunsaturated fats), tend to be fermented by bacteria. New technology in the orange juice industry has converted normal pulp to soluble fiber that supports bacterial fermentation.

A high ratio of calcium to phosphorus reduces the breakdown of fibers, and this probably contributes to some of the protective effects of a

large amount of calcium in the diet, for example reduction in the incidence of bowel and ovarian cancer.

A few plant fibers have a high proportion of cellulose, that resists bacterial attack, combined with antibiotics that suppress bacterial growth, but in most grains and vegetables, and many fruits, cellulose is combined with fibers that support bacterial growth. Fats, especially saturated fatty acids, have antiseptic properties, and when they are combined with starches and fibers they help to suppress bacterial growth. Their effects can be seen in various ways, including protection against the anxiety caused by fermentation (Prasad and Prasad, 1996).

Thyroid hormone increases digestive activity, including stomach acid and peristalsis, and both thyroid and progesterone increase the ability of the intestine to absorb sugars quickly; their deficiency can permit bacteria to live on sugars as well as starches.

Pavlov's and Anokhin's understanding of the organism in terms of its integration of perceptions of the actual situation, including its needs, and its projection of the possibility of satisfying those needs, gives us a way to understand some of the ways that our digestive system interacts with our physical health and our moods and emotions. Anokhin, in analyzing Pavlov's Reflex of Purpose and Exploratory Reflex, described the way that the cortex of the brain creates a model of the world, including information about the present state of the organism. Anokhin called this model of the situation the Acceptor of Action, because it was the framework for evaluating behavior. A properly functioning organism is always projecting or imagining future possible situations, and projecting or imagining its actions in those situations, evaluating the possibilities.

Any particular external situation can create stress or not, depending on what it means to the individual. The perception of a possibly good outcome mobilizes the creative adaptive abilities, in a functional system that includes an external goal and relevant physiological processes; the failure to see possibility in the situation--anticipating failure rather than success--demobilizes the systems of adaptation, and the

excitation caused by unsatisfied needs produces generalized catabolic processes.

As the organism prepares to enact these imagined possible futures, it mobilizes the appropriate resources, activating some systems and inactivating other systems, constructing, consuming and redistributing its own substance, while assimilating substances from the environment. The anabolic hormones, such as thyroid, and the tissue-catabolic hormones such as cortisol, act both locally and systemically, according to various regulatory processes, mainly nervous processes. A functional system is created for achieving the goal. As the organism achieves its purposes or adapts, the catabolic hormones will subside.

Sensory nerves in the intestine contribute to the organism's assessment of the situation--the Acceptor of Action--and motor nerves adjust the state of the intestine and other organs. The interplay between the state of the intestine and the assessed state of the stressful situation determines whether the individual will withdraw (often with incontinence of the bowel and bladder) or persist and struggle to adapt.

Defensive aggression is probably a response intermediate between fearful giving up and confident achievement. When a rat is restrained, held down on its back, it quickly develops ulcers, but if it has a stick to bite, it is very resistant to the formation of the ulcers. The ability to do something with a defensive meaning prevents the excessive production of serotonin and its consequences, such as increased production of cortisol and other stress hormones, and disturbance of circulation and energy production. Endotoxin and prostaglandins activate these same systems, and progesterone and aspirin are among the protective factors that can oppose those effects.

When physical and mental activity are intensified to meet an adaptive crisis, the adrenergic sympathetic nervous system optimizes muscular energy production, but inhibits the digestive system. In the state of anxious-helpless withdrawal, the parasympathetic and serotonergic systems demobilize the active neuro-skeletomuscular systems, and activate the bladder, intestine and stomach, tending to empty the

contents by causing diarrhea, vomiting, and urination. Serotonin reduces energy production in muscles; this is probably related to the muscular dystrophy it produced in Selye's experiments, and the myopathy produced by tryptophan (Castot, et al., 1991). Antiserotonin drugs are used for controlling diarrhea and vomiting, and some of them are protective against ulcers (Ramesh, et al., 2009).

Chronically elevated cortisol is commonly seen in depressed people, but giving a supplement of cortisol is effective in relieving depression. Both cortisol and the pituitary corticotrophic hormone that stimulates its production, ACTH, have some antidepressant effects, and they inhibit the hypothalamic corticotropin release hormone, CRH. CRH is more directly associated with depression than cortisol is, and it by itself activates many inflammatory processes, including the release of histamine, cytokines, and nitric oxide. CRH is promoted in the hypothalamus (and in many other tissues) by inflammation, endotoxin, serotonin, interleukins, and prostaglandins, but also by the perception of unavoidable difficulties.

Besides cortisol, progesterone and androgens are internal factors that decrease the activity of CRH. Estrogen and hypothyroidism increase its activity.

The convergence, interaction, and summation of inflammatory molecules with the perceived meaning of a situation is a subject carefully avoided by those who think of the organism as a sum of its molecules and cells. They accept that transmitter molecules and receptor molecules regulate behavior, and influence perception, but they can't explain how understanding and interpretation can make the profound differences in feeling, behavior, and biochemistry that they in fact do make.

The model of nerves as on-off switches, governed by the state of their membrane, has made the problem of integrating meaning with chemistry seem impossibly complex. In recent years it has been recognized that nerves can produce or respond to more than one kind of signal, but few people have understood nerve cells in a way that could allow individual cells to

handle complex and subtle signals, and to make rational judgments based on such signals. P.K. Anokhin, G. Albrecht-Buehler and a few others have observed the importance of recognizing cellular complexity, and the ability of cells to integrate complex information. The cell physiology of Gilbert Ling and Albert Szent-Gyorgyi demonstrated the basis on which these complex integrating processes can be understood.

Nerves and muscle cells should be considered together, because they respond to many things in similar ways. The membrane people don't like to think that nerves have any contractile properties, but in fact they do twitch slightly when stimulated, showing that it is the entire cytoplasm that responds to information, not the hypothetical plasma membrane. When they are overstimulated, they swell, as muscles do when they are fatigued. When a muscle is stretched while it's trying to contract (as in running downhill; this is called "eccentric contraction") it becomes inflamed, and the structural damage can be cumulative. By exercising muscles with "concentric contractions," allowing them to shorten against resistance, the cellular damage can be repaired.

Stretching a nerve causes some metabolic changes similar to the muscle in eccentric contraction, but nerves are normally protected against injuries of that sort. However, very intense stimulation of a nerve fiber can disrupt its internal structure, causing "beading," an intermittent clumping of the nerve filaments and microtubules along the axon, as if the internal structure is contracting in segments, while the overall axon doesn't contract. With intense stress, the beaded fiber can separate into individual pieces, like droplets of cytoplasm, instead of the elongated cylinder. Long before stress causes it to disintegrate in that manner, it is likely that it causes it to lose its functional coherence.

The excessive stimulation of a cell increases its internal alkalinity, causing it to take up more water. The mediators of inflammation, such as CRH, serotonin, and endotoxin cause cell swelling and increased alkalinity. CRH and endotoxin can increase the susceptibility to seizures, but they can also block the ability of cells to respond to normal stimulation.

Serotonin, as an alkaloid, is bound and inactivated by acidic cells. Carbon dioxide, produced by respiration, normally keeps cells slightly acidic. Hyperventilation, or the production of less carbon dioxide and more lactic acid, causes serotonin to be released from platelets and other cells where it has been inactive, allowing it to enter other cells where it has its characteristic effects, increasing lactate, decreasing ATP (Assouline-Cohen, et al. 1998, Koren-Schwartz, et al., 1994), increasing pH, and stimulating contraction or mitosis.

One of the very neglected mechanisms of serotonin's action is its ability to activate carbonic anhydrase, a class of enzymes that convert carbon dioxide to bicarbonate, and tend to increase the alkalinity of cells. Inhibitors of carbonic anhydrase are currently being investigated for their ability to stop the growth of cancer cells and to permit their recovery of normal functions. One of the carbonic anhydrase inhibitors, acetazolamide, has been used for many years to treat epilepsy, glaucoma, migraine, sleep apnea, mountain sickness--and even psychosis (Inoue, et al., 1984).

Serotonin's effects appear to be the result of its action on the microtubules or other elements of the cell skeleton (Bianchi, et al., 2003, Grass, et al., 2004), which are the main integrating elements of the cell. Chronic disturbance of the microtubules and neurofilaments, with inflammation and impaired energy production, is characteristic of Alzheimer's disease.

If the internal and external causes of stress converge, additively, on the cell's internal communication and integration system, then the basic resistance of the organism to stress can be increased by any of the factors which oppose the signals of stress.

Carbon dioxide, progesterone, and thyroid act on many of the factors that interfere with our ability to handle stress constructively. A diet that reduces fermentation and endotoxin, with an abundance of calcium--fruit, milk, and cheese, for example--can help to shift the balance away from lactic acid, estrogen, and serotonin, toward carbon dioxide, progesterone, and thyroid.

REFERENCES

- Aust N Z J Med. 1984 Dec;14(6):888-95. **Serotonin antagonists.** Anthony M. "The realisation that serotonin plays a role not only in the carcinoid syndrome but also in migraine, nociception, dumping syndrome, vascular disease and hypertension, has led to an enormous amount of activity in search of serotonin antagonists."
- Mol Genet Metab. 1998 Mar;63(3):235-8. **Activation of membrane skeleton-bound phosphofructokinase in erythrocytes induced by serotonin.** Assouline-Cohen M, Ben-Porat H, Beitner R. "We show here that serotonin, both in vivo and in vitro, induced a marked activation of phosphofructokinase, the rate-limiting enzyme in glycolysis, in the membrane-skeleton fraction from erythrocytes. Concomitantly, the hormone induced a striking increase in lactate content, reflecting stimulation of glycolysis." "These results suggest a defense mechanism in the erythrocytes against the damaging effects of serotonin, whose concentration in plasma increases in many diseases and is implicated as playing an important role in circulation disturbances."
- Synapse. 2003 Sep 1;49(3):188-94. **Cytoskeletal changes in the hippocampus following restraint stress: role of serotonin and microtubules.** Bianchi M, Heidbreder C, Crespi F.
- Nutr Cancer 1998;31(1):24-30. **Dietary lignin, and insoluble fiber, enhance uterine cancer but did not influence mammary cancer induced by N-methyl-N-nitrosourea in rats.** Birt DF, Markin RS, Blackwood D, Harvell DM, Shull JD, Pennington KL "Mammary carcinogenesis was not altered by dietary lignin. However, uterine endometrial adenocarcinoma was observed only in the MNU-lignin diet group (4 carcinomas/40 effective rats) ($p < 0.05$)."
- Science. 1941 Nov 14;94(2446):462. **Diminution in ability of the liver to inactivate estrone in vitamin D complex deficiency.** Biskind MS, Biskind GR.
- Am J Clin Pathol. 1946 Dec;16(12):737-45. **The nutritional aspects of certain endocrine disturbances.** Biskind GR, Biskind MS.
- J R Soc Health 2000 Dec;120(4):242-7. **Dietary fibre, physicochemical properties and their relationship to health.** Blackwood AD, Salter J, Dettmar PW, Chaplin MF. "Dietary carbohydrates that escape digestion and absorption in the small intestine include **non-digestible oligosaccharides (carbohydrates with a degree of polymerisation between**

three and ten), resistant starch and non-starch polysaccharides."

Lancet 2000 Oct 14;356(9238):1300-6. **Calcium and fibre supplementation in prevention of colorectal adenoma recurrence: a randomised intervention trial.** European Cancer Prevention Organisation Study Group. Bonithon-Kopp C, Kronborg O, Giacosa A, Rath U, Faivre J.

Braz J Med Biol Res. 1982 Oct;15(4-5):281-3. **Effect of serotonergic drugs on the aggressiveness induced by delta 9-tetrahydrocannabinol in rem-sleep-deprived rats.** Carlini EA, Lindsey CJ. 1. delta 9-Tetrahydrocannabinol (THC) induced aggressive behavior in rats previously deprived of REM sleep. This aggressiveness was significantly potentiated by tryptophan and fluoxetine, drugs which increase brain serotonin availability. 2. Conversely, drugs which decrease serotonergic function such as D,L-p-chlorophenylalanine, cinanserin and cyproheptadine strongly blocked the aggressive behavior. 3. On the basis of previous data indicating an involvement of dopaminergic mechanisms in this type of aggressiveness and the present results showing a role for serotonin, it is concluded that REM deprivation-THC aggression is under the control of at least these two neurotransmitters.

Therapie 1991 Sep-Oct;46(5):355-65. **["Eosinophilia-myalgia" syndrome due to L-tryptophan containing products. Cooperative evaluation of French Regional Centers of Pharmacovigilance. Analysis of 24 cases].** Castot A, Bidault I, Bournerias I, Carlier P, Efthymiou ML

Nutr Cancer 1984;6(2):77-85. **Enhancement of 1,2-dimethylhydrazine-induced large bowel tumorigenesis in Balb/c mice by corn, soybean, and wheat brans.** Clapp NK, Henke MA, London JF, Shock TL "This study was designed to determine the effects of four well-characterized dietary brans on large bowel tumorigenesis induced in mice with 1,2-dimethylhydrazine (DMH)." "In DMH-treated mice, tumors were found almost exclusively in the distal colon. Tumor incidences were as follows: controls, 11%; soybean group, 44%; soft winter wheat group, 48%; hard spring wheat group, 58%; and corn group, 72%. Tumors per tumor-bearing mouse ranged from 1.4 to 1.6, except in the corn group, which had 2.1."

Mol Genet Metab. 2007 Dec;92(4):364-70. **Serotonin stimulates mouse skeletal muscle 6-phosphofructo-1-kinase through tyrosine-phosphorylation of the enzyme altering its**

intracellular localization. Coelho WS, Costa KC, Sola-Penna M.

Neurosci Biobehav Rev. 2005;29(4-5):761-70. **Endotoxin- or pro-inflammatory cytokine-induced sickness behavior as an animal model of depression: focus on anhedonia.** De La Garza R 2nd.

Brain Behav Immun. 2010 May;24(4):558-63. Epub 2010 Jan 4. **Inflammation and social experience: an inflammatory challenge induces feelings of social disconnection in addition to depressed mood.** Eisenberger NI, Inagaki TK, Mashal NM, Irwin MR.

Biol Psychiatry. 2010 Oct 15;68(8):748-54. **Inflammation-induced anhedonia: endotoxin reduces ventral striatum responses to reward.** Eisenberger NI, Berkman ET, Inagaki TK, Rameson LT, Mashal NM, Irwin MR.

Proc R Soc Med. 1967 Jun;60(6):553-5. **A psychological setting of somatic disease: the 'giving up - given up' complex.** Engel GL.

Eur J Pharmacol. 1983; 91(4):493-9, **A lipopolysaccharide and concanavalin A induce variations of serotonin levels in mouse tissues,** Endo Y.

Ciba Found Symp 1972, 8:57-75, **Conservation-withdrawal: a primary regulatory process for organismic homeostasis,** Engel GL, Schmale AH.

Am J Epidemiol. 2002 Jul 15;156(2):148-57. **Association of dairy products, lactose, and calcium with the risk of ovarian cancer.** Goodman MT, Wu AH, Tung KH, McDuffie K, Kolonel LN, Nomura AM, Terada K, Wilkens LR, Murphy S, Hankin JH.

Med Hypotheses. 2004;62(2):169-72. **Biophotons, microtubules and CNS, is our brain a "holographic computer"?** Grass F, Klima H, Kasper S. "Several experiments show that there is a cell to cell communication by light in different cell types. This article describes theoretical mechanisms and subcellular structures that could be involved in this phenomenon." "Neurons are large colourless cells with wide arborisations, have an active metabolism generating photons, contain little pigment, and have a prominent cytoskeleton consisting of hollow microtubules." "Fluorescent and absorbing substances should interfere with such a communication system. Of all biogenic amines nature has chosen the ones with the strongest fluorescence as neurotransmitters for mood reactions: serotonin, dopamine and norepinephrine. If these mechanisms are of relevance our brain would have to be looked upon as a 'holographic computer'."

Asia Pac J Clin Nutr. 2003;12 Suppl:S12. **Anxiety following increased hind-gut fermentation.** Hanstock TL, Claytons EH, Mallet PE. "Fermentation

and lactic acid production in the caecum and colon can lead to detrimental effects in several animal models, including adverse behaviour in horses." **"Conclusions - Fermentation of carbohydrate leading to increased concentrations of lactic acid in the caecum of rats was associated with increased anxiety in rats."**

Asia Pac J Clin Nutr. 2003;12 Suppl:S12. **Anxiety following increased hind-gut fermentation.** Hanstock TL, Claytons EH, Mallet PE.

Physiol Behav. 2004 Sep 15;82(2-3):357-68. **Anxiety and aggression associated with the fermentation of carbohydrates in the hindgut of rats.** Hanstock TL, Clayton EH, Li KM, Mallet PE. "Lactic acid accumulation in the caecum and colon resulting from the fermentation of carbohydrates can lead to deleterious effects in ruminant and monogastric animals, including humans. In the present study, we examined the behavioural effects of two types of commonly consumed foods: soluble and fermentable carbohydrates (FCs)." "Results showed that anxiety (spending less time in the light compartment during the light/dark emergence test) and aggression (increased fighting during the social interaction test) were increased following raised concentrations of fermentation end products, such as lactic acid and volatile fatty acids (VFAs) in the caecum of rats. These associations occurred regardless of dopamine and 5-HT concentrations in the prefrontal cortex (PFC) and provide evidence supporting a general effect of FCs on behaviour."

Proc Soc Exp Biol Med 1986 Dec;183(3):299-310. **Relationship between dietary fiber and cancer: metabolic, physiologic, and cellular mechanisms.** Jacobs LR. "Experiments in laboratory animals, using chemical induction of large bowel cancer, have in general shown a protective effect with supplements of poorly fermentable fibers such as wheat bran or cellulose. **In contrast, a number of fermentable fiber supplements including pectin, corn bran, oat bran, undegraded carageenan, agar, psyllium, guar gum, and alfalfa have been shown to enhance tumor development.**" "Fermentation of fibers within the large bowel results in the production of short chain fatty acids, which in vivo stimulate cell proliferation, while butyrate appears to be antineoplastic in vitro. Evidence suggests that if dietary fibers stimulate cell proliferation during the stage of initiation, then this may lead to tumor enhancement. Fermentation also lowers luminal pH, which in turn modifies colonic microbial metabolic acidity, and is associated with increased epithelial cell proliferation and colon carcinogenesis."

Biull Eksp Biol Med. 1989 Nov;108(11):545-7. **[Effects of serotonin and stress on the state of the gastric mucosa in rats with the intact and transected vagus nerve].** [Article in Russian] Karevina TG, Khokholia VP, Shevchuk IM.

Gen Pharmacol. 1994 Oct;25(6):1257-62. **Serotonin-induced decrease in brain ATP, stimulation of brain anaerobic glycolysis and elevation of plasma hemoglobin; the protective action of calmodulin antagonists.** Koren-Schwartz N, Chen-Zion M, Ben-Porat H, Beitner R. 1. Injection of serotonin (5-hydroxytryptamine) to rats, induced a dramatic fall in brain ATP level, accompanied by an increase in P(i). Concomitant to these changes, the activity of cytosolic phosphofructokinase, the rate-limiting enzyme of glycolysis, was significantly enhanced. Stimulation of anaerobic glycolysis was also reflected by a marked increase in lactate content in brain. 2. Brain glucose 1,6-bisphosphate level was decreased, whereas fructose 2,6-bisphosphate was unaffected by serotonin. 3. All these serotonin-induced changes in brain, which are characteristic for cerebral ischemia, were prevented by treatment with the calmodulin (CaM) antagonists, trifluoperazine or thioridazine. 4. Injection of serotonin also induced a marked elevation of plasma hemoglobin, reflecting lysed erythrocytes, which was also prevented by treatment with the CaM antagonists. 5. The present results suggest that CaM antagonists may be effective drugs in treatment of many pathological conditions and diseases in which plasma serotonin levels are known to increase.

J Psychopharmacol. 2001 Dec;15(4):219-30. **Effects of tryptophan loading on verbal, spatial and affective working memory functions in healthy adults.** Luciana M, Burgund ED, Berman M, Hanson KL.

J Nutr. 1989 Feb;119(2):235-41. **Independent effects of fiber and protein on colonic luminal ammonia concentration.** Lupton JR, Marchant LJ. "...clearly demonstrates that luminal ammonia concentration is dependent upon both protein level and fiber type, and that a fermentable fiber (pectin), rather than decreasing colonic ammonia concentrations, actually increases them several-fold."

J Neurosci. 2004 Apr 7;24(14):3655-62. **Cognitive correlates of white matter growth and stress hormones in female squirrel monkey adults.** Lyons DM, Yang C, Eliez S, Reiss AL, Schatzberg AF. **"Impaired cognitive response inhibition correlated with the expansion of white matter volume statistically controlling for age, stress hormones, gray matter, and CSF volumes.** These results indicate that

instead of enhancing cognitive control during midlife adulthood, white matter volume expansion contributes to aspects of cognitive decline."

Br Poult Sci. 2001 Mar;42(1):33-42. **Development of pecking damage in layer pullets in relation to dietary protein source.** McKeegan DE, Savory CJ, MacLeod MG, Mitchell MA.

Prev Med 1987 Jul;16(4):540-4. **Fiber, stool bulk, and bile acid output: implications for colon cancer risk.** McPherson-Kay R. "Whereas wheat bran reduces fecal bile acid concentration, pectin, lignin, and oat bran do not. These three fibers significantly increase total bile acid output. Bile acids act as promoters of colonic tumors in mutagenesis assay systems and in various animal models."

Hum Psychopharmacol. 2002 Jun;17 Suppl 1:S13-6. **New directions in the development of antidepressants: the interface of neurobiology and psychiatry.** Nemeroff CB.

J Neural Transm. 2000;107(12):1393-401. **Lipopolysaccharide administration produces time-dependent and region-specific alterations in tryptophan and tyrosine hydroxylase activities in the rat brain,** Nolan Y, Connor TJ, Kelly JP, Leonard BE.

Physiol. Behav. 1996, 60(3):1039-42. **Short-term consumption of a diet rich in fat decreases anxiety response in adult male rats.,** Prasad A, Prasad C.

Fundamental & Clinical Pharmacology 2009;23, 303-309. **Effect of central administration of ondansetron, a 5-hydroxytryptamine-3 receptor antagonist on gastric and duodena ulcers,** Ramesh ST, Asad M, Dhamanigi SS, Prasad VS.

Brain Behav Immun. 2006 Nov;20(6):517-26. Epub 2006 May 2. **Immunity and emotions: lipopolysaccharide increases defensive behaviours and potentiates despair in mice.** Renault J, Aubert A.

Naturwissenschaften, 1965, 12, 519. **Eine durch Serotonin erzeugte Muskeldystrophie,** Selye, H.

Am J. Pathol. 48(2): 299-303. **Production and prevention of stercoral ulcers in the rat,** Selye H, Winandy G, Gabbiani G.

Horm Behav. 2004 Apr;45(4):278-84. **Increased aggressive behavior and decreased affiliative behavior in adult male monkeys after long-term consumption of diets rich in soy protein and isoflavones.** Simon NG, Kaplan JR, Hu S, Register TC, Adams MR.

J Surg Oncol 1993 Feb;52(2):77-82. **The effect of the fiber components cellulose and lignin on experimental colon neoplasia.** Sloan DA, Fleiszer DM, Richards GK, Murray D, Brown RA. "One diet

contained "low-fiber" (3.8% crude fiber); the others contained "high fiber" (28.7% crude fiber) composed of either cellulose or lignin. Although both "high fiber" diets had similar stool bulking effects, only the cellulose diet was associated with a reduction in 1,2-dimethylhydrazine (DMH)-induced colon neoplasms."

Psychoneuroendocrinology. 2004 Oct;29(9):1119-28. **Enhanced expression of cytokines and chemokines by blood monocytes to in vitro lipopolysaccharide stimulation are associated with hostility and severity of depressive symptoms in healthy women.** Suarez EC, Lewis JG, Krishnan RR, Young KH.

Brain Cogn. 2009; 71(3):427-36. **Serotonin: Modulator of a drive to withdraw.** Tops M, Russo S, Boksem MA, Tucker DM.

Food Chem Toxicol 1984 Jul;22(7):573-8. **Effect of Metamucil on tumour formation by 1,2-dimethylhydrazine dihydrochloride in mice.** Toth B. "It was expected that a high amount of dietary fibre would inhibit carcinogenesis in the large intestine. Instead, metamucil increased the incidence of colon tumours induced by 1,2-DMH, although only in males."

Med Hypotheses. 1990 Jan;31(1):7-15. **The major hallucinogens and the central cytoskeleton: an association beyond coincidence? Towards sub-cellular mechanisms in schizophrenia.** van Woerkom AE.

J Nutr 1996 Aug;126(8):1979-91. **Dietary guar gum alters colonic microbial fermentation in azoxymethane-treated rats.** Weaver GA, Tangel C, Krause JA, Alpern HD, Jenkins PL, Parfitt MM, Stragand JJ. "Fecal in vitro fermentation rates were higher for guar-fed rats than for control rats [three-way ANOVA (diet, time, in vitro substrates), $P = 0.002$]."

Immunopharmacology, 1983;6(2):75-86. **Investigation of adrenergic and prostaglandin influences in the endotoxin alteration of hepatic heme oxygenase, microsomal mixed-function oxidase, and glucocorticoid-induced tryptophan oxygenase activities,** Williams JF, Szentifanyi A. (LPS inhibits glucocort. induction of TO activity)

Coll Antropol. 2001 Dec;25(2):485-92. **Depressive disorder as possible risk factor of osteoporosis.** Vrkljan M, Thaller V, Lovricevic I, Gacina P, Resetic J, Bekic M, Sonicki Z.
