

Growth hormone and Klotho

Christoph Schmid¹, Marian C Neidert², Oliver Tschopp¹, Lisa Sze^{1,3}
and René L Bernays^{2,4}

¹Division of Endocrinology and Diabetology, University Hospital Zurich, Rämistrasse 100, CH-8091 Zurich, Switzerland

²Department of Neurosurgery, University Hospital Zurich, Frauenklinikstrasse 10, 8091 Zurich, Switzerland

³Division of Endocrinology and Diabetology, Kantonsspital St Gallen, 9007 St Gallen, Switzerland

⁴Department of Neurosurgery, Clinic Hirslanden, Witellikerstrasse 40, 8032 Zurich, Switzerland

Correspondence
should be addressed
to C Schmid
Email
christoph.schmid@usz.ch

Abstract

Acromegaly is characterized by excessively high GH and IGF1 levels. Recent data suggest that soluble Klotho (sKlotho) is also elevated in patients with active acromegaly. sKlotho decreases towards normal following removal of the GH-producing pituitary adenoma. The *Klotho* gene was identified in mice following its accidental disruption by ectopic DNA. It is an ageing suppressor gene of restricted expression (mainly in kidneys, brain, and parathyroid and pituitary glands) encoding a transmembrane protein, mKlotho. mKlotho serves as a co-receptor in fibroblast growth factor 23 (FGF23) signalling. FGF23 promotes urinary phosphate excretion and inhibits the synthesis of calcitriol. The ectodomain of mKlotho is enzymatically released to result in a humoral factor, sKlotho, which exerts systemic effects (on ion channels and signalling pathways), possibly by working as an enzyme that modifies glycans of cell surface glycoproteins. GH enhances renal phosphate reabsorption and calcitriol production, i.e. exerts effects in the proximal tubule opposing those attributed to mKlotho, and attenuates calciuria in the distal tubule similar to sKlotho. sKlotho can be measured in extracellular fluids (serum, urine and cerebrospinal fluid (CSF)) by an ELISA. In line with predominant expression of Klotho in kidneys and choroid plexus, concentrations of sKlotho are particularly high in urine and CSF. Determination of sKlotho in serum and urine (both presumably reflecting GH action on the kidneys) could be used as a supplementary tool in the diagnosis and follow-up of patients with acromegaly. The question arises whether GH exerts selected actions via modifying activities of Klotho.

Key Words

- Klotho
- growth hormone
- acromegaly
- FGF23
- IGF1

Journal of Endocrinology
(2013) 219, R37–R57

Introduction

This review on growth hormone (GH) and Klotho covers novel findings on a well-known hormone, the first to be extracted and characterized from the pituitary gland, and a more recently, accidentally discovered multifunctional protein. To enhance comprehension of the relationship between the two, we also discuss insulin-like growth factor 1 (IGF1), the first growth factor isolated and characterized from human serum, and fibroblast growth factor 23 (FGF23), currently the last member of the FGF family to be cloned. IGF1 is important for the

apprehension of GH actions, and FGF23 is important for the understanding of the action of Klotho. These two growth factors illustrate the continuum between locally acting growth factors and systemically active hormones. Klotho works both locally and as a circulating humoral factor. Klotho was recently renamed α -Klotho, after the discovery of new Klotho family members; however, β -Klotho will be mentioned only briefly, although GH probably interacts with both α - and β -Klotho endocrine communication lines.

Acromegaly and GH

Acromegaly is a characteristic and disfiguring disease, described as a clinical entity in the late 19th century, which was later found to be caused by an eosinophil pituitary adenoma. GH excess produced by an adenoma of the pituitary gland is the major cause for acromegaly with an incidence of approximately four cases per 1 million persons per year. Clinical features develop insidiously and progressively over many years. It includes changes in the patient's appearance, mainly soft tissue swelling and cartilage and bone growth, resulting in typical acral enlargement and coarse facial features. Thus, with increasing typical signs and symptoms, diagnosis of acromegaly is generally straightforward. However, due to limited awareness of a rare disease, the correct diagnosis is often considerably delayed and is therefore associated with increased mortality even after curative therapy. For patients with acromegaly, (transsphenoidal) surgery remains the most promising therapeutic option, as it offers the best chance for long-term cure and restoration of pituitary function, including normalized GH secretion. Currently, GH and IGF1 (a marker and mediator of GH actions) are the classical biochemical markers of disease activity in acromegaly. Increased levels of GH (i.e. not suppressible to a nadir $<1 \mu\text{g/l}$ during oral glucose tolerance testing, or $<0.3 \mu\text{g/l}$, with more modern assays) and IGF1 on blood testing support the diagnosis. Likewise, to define cure, biochemical assessment is superior to MRI, as tiny tumour remnants are not reliably depicted.

The healthy human pituitary contains several milligrams of GH. Pituitary extracts were found to cause gigantism and pituitary hormones were found to be diabetogenic. GH was first isolated in 1944. In 1957, extraction of human GH from cadaver pituitaries succeeded, and in 1958, use of the extracted hormone to treat pituitary dwarfs was reported. During childhood and adolescence, when the epiphyseal lines have not fused, GH stimulates longitudinal bone growth, and GH excess from adenomas causes gigantism. In 1963, a RIA for GH was developed, and by 1970, Li succeeded with determination of the amino acid sequence and synthesis of GH. In 1985, several cases of Jakob Creutzfeldt disease were reported, and in the same year, recombinant human GH (rhGH) was introduced for therapeutical use in children. GH is effective in promoting growth of children with GH deficiency. However, in the rare condition of GH insensitivity (type Laron-Dwarfism), e.g. due to GH receptor (GHR) deficiency ('GHR-k.o.'), endogenous GH is high, and there is no response to rhGH administration (Godowski *et al.* 1989).

The wide availability of rhGH led to an increasing number of studies in adults with hypopituitarism including GH deficiency, and rhGH was reimbursed in 1996 for use in adulthood. It has never been established that isolated GH deficiency is associated with an increase in morbidity and mortality; however, GH excess is related to an increase in standard mortality rate.

GH is a single-chain polypeptide hormone (with two disulphide bridges) synthesized, stored and secreted by the somatotrophic cells of the pituitary gland. GH consists of 191 amino acids, corresponding to the main circulating form, which is 22 kDa in size and mainly bound to a GH-binding protein (GHBP); the latter is derived from the GHR by proteolytic cleavage. GH is secreted in a pulsatile manner, mainly regulated by hypothalamic neuroendocrine mediators such as (inhibitory) somatostatin and GH-releasing hormone, but also by signals from the periphery such as (stimulatory) ghrelin and by feedback inhibition via IGF1. The effects of GH are mediated by the GHR. The human GHR gene was characterized by Wood and colleagues in 1989 (Godowski *et al.* 1989). It is a member of a large cytokine receptor family that also includes erythropoietin and IL6. Activation of the GHR by GH binding results in a conformational change of the receptor as well as in an activation of a JAK system and several downstream pathways, including STAT members and MAPK. The extracellular fragment of the GHR is cleaved from the cell surface and circulates as a GHBP. GHBP prolongs plasma half-life of circulating GH, which is estimated to be 2–5 min in the free form and 15–20 min in the GHBP-bound form. GH is cleared from the circulation via tissues that are also responsive to the hormone, especially liver and kidneys.

Activities of GH and IGF1

GH promotes (longitudinal bone) growth. It is anabolic, increasing protein synthesis and net nitrogen retention, and insulin is required for these effects. Some of the properties of GH have been fairly well characterized, e.g. on liver (increasing IGF1 production, synergistic with insulin), on cartilage (growth) and on adipose tissue (increasing lipolysis, antagonistic to insulin). However, other actions of GH are still poorly understood, such as the action on kidney function and phosphate homeostasis, or on endothelial function. Some of the effects of GH are direct, e.g. on liver, adipose tissue and striated muscle (i.e. classical insulin target tissues) whereas others are mediated by IGF1 (and possibly other mediators).

Human blood contains several milligrams of IGFs. IGFs were discovered by three independent groups (Salmon & Daughaday 1957, Froesch *et al.* 1963, Pierson & Temin 1972), which were interested in biological activities exerted by serum: i) stimulation of proteoglycan synthesis (sulphate incorporation) in cartilage by a GH-dependent 'sulfation factor' (somatomedin), ii) insulin-like activity (non-suppressible by anti-insulin antibodies) on adipose tissue and iii) mitogenic activity on fibroblasts *in vitro* (multiplication-stimulating activity (MSA)). IGF1 works as a somatomedin in animals and in man (Schoenle *et al.* 1982, Laron *et al.* 1992).

Tests assessing the biological activities of GH and insulin were difficult 60 years ago and required *in vivo* bioassays (measuring the increase in body weight or in epiphyseal width in hypophysectomized rats, or the extent of blood sugar decrease in experimental animals); in search of improved methods (before RIAs were introduced), interesting discoveries were made.

Serum stimulates incorporation of $^{35}\text{SO}_4$ into cartilage *in vitro*. In 1957, Salmon and Daughaday reported that serum of hypophysectomized rats lacked such an activity. It could be reconstituted by administration of GH to hypophysectomized rats, but not by addition of GH to cartilage cultured in serum of hypophysectomized rats. These observations led to the concept that GH does not stimulate growth directly, but rather induces formation of factors that mediate action of GH. These factors were first called sulfation factors and later somatomedins; a GH-dependent plasma factor stimulates the incorporation of sulphate into chondroitin sulphate in cartilage (Salmon & Daughaday 1957, Daughaday *et al.* 1972).

Serum contains insulin-like activity. Froesch *et al.* observed that most of the insulin-like activity in serum could not be suppressed by the addition of antiserum neutralizing insulin to the incubation medium. Therefore, 'non-suppressible insulin-like activity (NSILA)' was different from insulin but could mimic its effects on adipose tissue (Froesch *et al.* 1963). Later, molecules (7.6 kDa in size) responsible for this insulin-like activity were purified from serum and characterized (Rinderknecht & Humbel 1978). The designation IGFs appeared justified as IGFs are functionally and structurally closely related to insulin. As the C-domain of IGFs is not removed during prohormone processing, mature IGF peptides are single-chain polypeptides. IGF1 consists of 70 amino acids and displays 43% sequence identity to the insulin A- and B-chain (Rinderknecht & Humbel 1978). IGFs bound to IGF binding proteins (IGFBPs) cannot interact with insulin

receptors; moreover, the affinity of IGFs to insulin receptors is low.

Serum provides components required for optimal growth of most cells in culture, including growth factors. Factors could be extracted from serum with MSA, i.e. factors that stimulate replication of cells in culture (Pierson & Temin 1972). Subsequently, it was found that MSA was also produced by cultured liver cells (Dulak & Temin 1973).

IGFs (IGF1 and IGF2) signal through type 1 IGF receptors. Insulin and IGF have their own distinct cell surface receptors whose structures have been deduced from analysis of isolated cDNA clones. Type 1 IGF and insulin receptor are homologous (Ullrich *et al.* 1986); both consist of $\alpha_2\beta_2$ 'hetero-tetrameric' structures. The α -subunits lie entirely extracellular and contain a ligand-binding domain, while the β -subunits are transmembrane polypeptides that contain a tyrosine kinase domain in their intracellular portion. Signal transduction by type 1 and insulin receptors involves autophosphorylation at tyrosine residues and phosphorylation of other substrates.

Radiolabelled IGFs had been used to characterize specific IGF binding sites (receptors) on cells. In addition, they allowed the discovery of specific IGFBPs in extracellular fluids. IGFBPs in the strict sense are entirely unrelated to IGF receptors and bind IGFs with comparable or even higher affinity than the type 1 IGF receptor. IGFBP3 is by far the most abundant IGFBP in adult human serum, and its production (like that of IGF1) is dependent on GH both *in vivo* and *in vitro*. Intact IGFBP3 prevents access of IGF1 to its receptors, but proteolysis of IGFBP3 activates IGF1 signalling (Schmid *et al.* 1991). GH (but not IGF1) induces an IGFBP3-proteolytic activity (Rutishauser *et al.* 1993).

IGFBPs appear to affect distribution volumes and half-lives of IGFs; they contribute to the large pool of IGFs present in most extracellular spaces, and they modify their biological activity. IGFBP3 serves as the major circulating carrier of IGFs, most of which are associated with a GH-dependent ternary complex of 140 kDa, comprising an acid-labile 85 kDa glycoprotein, IGFBP3, and IGF1 or IGF2. The apparent half-life of IGFs is about 10 min in their free form and 10–15 h when in the ternary complex; the large molecular weight complex accounts for the slow turnover of serum IGFs (Guler *et al.* 1989).

IGF1 is produced continuously, especially in the liver that is estimated to be responsible for 70–80% of circulating IGF1. Apart from the 'endocrine' IGFs, IGFs are also produced in most tissues where they can act in a paracrine or autocrine manner. Serum IGF1 is a good hepatic marker of GH action. Owing to its long plasma half-life, levels are fairly stable throughout the day,

and timing of blood sampling for diagnostic purposes is not critical.

Until now, GH and especially GH activity, as reflected (in part) by IGF1 concentrations, have been the 'classical' biomarkers for diagnosing and monitoring disease activity during the treatment of patients with acromegaly. Their normalization has been linked to decreased mortality. However, it has been recognized that both parameters entail various shortcomings, both analytical and biological (Clemmons 2011). GH levels are age dependent, and serum levels of IGF1 are influenced not only by GH status but also by age, gender (oestrogens), race, liver function, nutritional status, portal insulin, thyroid hormones and by concomitant inflammatory disease. Unfortunately, measuring IGF1 remains notoriously difficult, especially since IGF1s interfere with immunoassay analysis. For the classical and time-consuming RIAs, IGF carrier proteins are removed before the samples are incubated with the antibodies (Zapf *et al.* 1980, 1981). Ideally, IGF1 should normalize and glucose-suppressed GH should be low after surgery for acromegaly. Most often, 1 µg/l (or lower) has been used as a cut-off value (Giustina *et al.* 2010).

Activities of FGF23

FGF23 may be most easily understood by the recognition of tumours by which it is secreted in excessive amounts. Similar to acromegaly, tumour-induced osteomalacia (TIO) is an acquired condition that can resolve after successful removal of the tumour, the source of excessive FGF23. In 1957, Prader reported the case of an 11½-year-old girl who presented with an acquired disorder of hypophosphataemia, renal phosphate wasting and rickets. He attributed the phenotype of the disease to a circulating factor, a 'rachitogenic substance' secreted by the tumour causing renal phosphate wasting ('phosphate diabetes'), as it disappeared after surgical excision of a benign mixed connective tissue tumour from the rib (Prader *et al.* 1959). The phenotype described corresponds to what we would nowadays designate a case of FGF23-mediated TIO. FGF23 was later cloned as a causative factor from tumours that result in osteomalacia (Shimada *et al.* 2001).

Two other groups, independently, had previously cloned FGF23, by homology as an additional FGF family member, preferentially expressed in the ventrolateral thalamic nucleus of the brain, and by genetic analysis of autosomal dominant hypophosphataemic rickets (ADHR; ADHR Consortium 2000, Yamashita *et al.* 2000).

Shimada reported in 2004 that overexpression of FGF23 in rodents essentially replicated the phenotype of

TIO and ADHR. Independent groups confirmed that FGF23 overexpression in transgenic animals or its administration to rodents resulted in renal phosphate loss, hypophosphataemia, impaired production of 1,25-(OH)₂D and rickets (Bai *et al.* 2004, Larsson *et al.* 2004, Shimada *et al.* 2004c).

FGF23, a 32 kDa glycoprotein with very weak heparin-binding affinity, is produced mainly in bone by osteoblasts/osteocytes; its production is stimulated by decreased glomerular filtration rate (GFR)/low nephron number as with chronic kidney disease (CKD) and ageing and dietary phosphate overload. The 24 amino acid hydrophobic signal sequence is removed and the protein O-glycosylated by GALNT3 to produce the mature, biologically active ²⁵FGF23²⁵¹ form in the circulation, with an estimated half-life of about 50 min (Khosravi *et al.* 2007). FGF23 acts predominantly on kidneys to decrease sodium (Na)-dependent phosphate transporters NaPi2a and NaPi2c (*SLC34a1* and *SLC34a3*) activity and thereby increase phosphate excretion ('phosphatonin') (Shimada *et al.* 2004c, Segawa *et al.* 2007, Gattineni *et al.* 2009). It inhibits vitamin D 1α-hydroxylase (CYP27B1) and calcitriol generation, and it enhances 24α-hydroxylase (CYP24A1) and calcitriol degradation (Saito *et al.* 2003, Shimada *et al.* 2004a). Therefore, excessive FGF23 production (by tumours) or inadequate inactivation (in ADHR) of FGF23 results in hypophosphataemia and rickets (in children) or osteomalacia (in adults). This happens due to renal phosphate loss ('acquired phosphate diabetes'), inhibition of vitamin D 1α-hydroxylase (impaired calcitriol production) and disturbed mineralization. In patients with progressive fatigue, muscle weakness and bone pain, persistent hypophosphataemia due to renal phosphate wasting and an inappropriately low 1,25-(OH)₂D level are the biochemical hallmarks of TIO (Bauer *et al.* 2010).

To understand the physiological roles of FGF23, mice lacking FGF23 were generated by targeted gene disruption. Defective production results in hyperphosphataemia and activation of vitamin D 1α-hydroxylase, severe calcifications (subcutaneous, periarticular, kidneys, lungs and vessels) and premature ageing. These *Fgf23*^{-/-} mice have, reminiscent of *Klotho*^{-/-} mice, hyperphosphataemia and extremely high serum calcitriol levels (Yoshida *et al.* 2002, Shimada *et al.* 2004b). It was realized that *Fgf23*^{-/-} mice (lacking the hormone) and *Klotho*^{-/-} mice (lacking functioning receptors) had similar phenotypes (see below).

FGFs signal through tyrosine kinase receptors and ERK. Among the 22 FGF members, most act in an

autocrine or paracrine fashion. The FGF19 (endocrine) subfamily includes FGF15 (in mice)/FGF19 (in humans), FGF21 and FGF23 (Itoh & Ornitz 2011). These hormone-like FGFs appear to be vertebrate specific and characteristically lack a heparin-binding domain. Therefore, they are not retained by the nearby extracellular matrix and reach the circulation. Moreover, endocrine FGFs do not require heparan sulphate proteoglycans for efficient FGFR tyrosine kinase activation; instead, they need transmembrane Klotho(s) (mKlotho) as co-receptors (see section Klotho, a transmembrane co-receptor and a multifunctional humoral factor in extracellular spaces). FGF15/FGF19 is mainly produced by the small intestine and predominantly targets the liver (FGFR4 and β -Klotho coexpression) to regulate cholesterol and (suppress) bile acid synthesis. FGF21 is mainly secreted from the liver and predominantly acts on white adipose tissue (FGFR1c and β -Klotho coexpression) to regulate glucose and lipid metabolism. This corresponds to FGF23, which is mainly produced by bone and predominantly targets the kidneys to regulate phosphate and vitamin D metabolism. β -Klotho belongs to the receptor complex of the former two endocrine FGFs and α -Klotho to the receptor complex of FGF23. In the kidneys, FGFR1 may be the most relevant receptor to mediate FGF23 actions (Gattineni *et al.* 2009). An activating mutation of FGFR1 has been reported to cause osteoglophonic dysplasia and hypophosphataemia (White *et al.* 2005). Antibodies activating FGFR1 also cause hypophosphataemia (Wu *et al.* 2013).

Klotho, a transmembrane co-receptor and a multifunctional humoral factor in extracellular spaces

While studying the phenotype of transgenic mice overexpressing the rabbit type I sodium proton exchanger, insertion of ectopic DNA happened to occur into the 5' promoter region of the α -Klotho gene; α -Klotho protein was hardly expressed. The *Klotho* gene was thereby identified serendipitously in this mouse model in 1997 after recognition that its disruption caused a phenotype of accelerated ageing (Kuro-o *et al.* 1997). Later, it was found that ageing suppression and lifespan extension were achievable in mice by overexpression of Klotho; these mice lived about 20% longer than WT mice and were more resistant to oxidative stress (Kurosaki *et al.* 2005). The name Klotho stems from the goddess of fate, which is 'spinning the thread of life' in Greek mythology. Mice with deficient *Klotho* expression have a syndrome that resembles human ageing, including shorter lifespan (on average,

only 2 months instead of 2–3 years as for WT mice), premature skin atrophy, osteopenia/osteoporosis, dysfunction of the pituitary gland (GH deficiency), growth retardation, hypogonadotropic hypogonadism (infertility), atrophy of genital organs, thymus and muscle, arteriosclerosis, ectopic (soft tissue and media-) calcifications, pulmonary emphysema and neurodegenerative disorders (hypokinesia, gait disturbances and hearing disorder). Klotho-deficient mice were subsequently characterized in more detail; e.g. the brain phenotype (distinct from the aged human brain with senile plaque deposition) includes memory retention deficits, a reduction in synapses in the hippocampus, disturbed axonal transport and hippocampus degeneration and impaired myelin production in specific areas of the brain (Nagai *et al.* 2003, Li *et al.* 2004, Shiozaki *et al.* 2008, Chen *et al.* 2013a). A recent study suggested that decreased activation of JAK2/STAT3 signalling and cholinergic pathways in hippocampus may play an important role in cognitive impairment observed in Klotho mutant mice (Park *et al.* 2013). Biochemically, Klotho deficiency results in hyperphosphataemia with increased 1,25-(OH)₂ vitamin D₃ and high FGF23 in keeping with FGF23 resistance, moreover in hypoglycaemia with low insulin (insulin hypersensitivity).

The *Klotho* gene product was proposed to function as a part of a signalling pathway that regulates ageing *in vivo* and morbidity in age-related diseases. The *Klotho* mouse created by unintentional insertion is not a strictly null strain, but a *Klotho*^{-/-} mouse was later intentionally generated, and this 'classical' knockout mouse was found to exhibit essentially the same phenotype as the original *Klotho* mouse, including hyperphosphataemia and disturbed vitamin D regulation (Tsujikawa *et al.* 2003). Over the past decade, several aspects reported in the original 1997 paper were described in more detail, including neurodegeneration, hearing loss, sarcopenia, ageing lung/emphysema and arteriosclerosis. However, discussing all of them would be beyond the scope of the current review. Several excellent reviews discuss specific aspects of Klotho in more detail (Nabeshima & Imura 2008, Kuro-o 2010, Manya *et al.* 2010, Martin *et al.* 2012, Hu *et al.* 2013).

The *Klotho* gene encodes a single-pass type I transmembrane protein (1014 amino acids in the mouse and 1012 in humans) with a short cytoplasmic domain and an extracellular domain composed of two β -galactosidase/glycosidase-like tandem repeats (KL1 and KL2) with β -glucuronidase and sialidase activity. It is predominantly expressed in kidneys, brain (choroid plexus (CP), neurons in hippocampus, Purkinje cells and inner ear), the

sinoatrial node of the heart and in several endocrine (pituitary, parathyroid and pancreas) and reproductive organs (gonads and placenta) (Kuro-o *et al.* 1997, Kato *et al.* 2000, Li *et al.* 2004, Takeshita *et al.* 2004, German *et al.* 2012). Although primarily in distal convoluted tubules, Klotho is expressed in all tubular segments along the nephron (Hu *et al.* 2010, Zhou *et al.* 2013). More details on the mouse *Klotho* gene and the identification of the human (96% homology) as well as the rat *Klotho* gene were reported 1 year later (Matsumura *et al.* 1998, Ohyama *et al.* 1998, Shiraki-Iida *et al.* 1998); two distinct transcripts encoding membrane and secreted Klotho protein were characterized. Klotho has been identified not only in the zebrafish, *Danio rerio*, but two *Klotho*-like genes, *klo1* and *klo2*, have also been identified in the nematode *Caenorhabditis elegans* (Polanska *et al.* 2011, Sugano & Lardelli 2011, Mangos *et al.* 2012). The corresponding KLO1 and KLO2 proteins lack a transmembrane domain. The ancestral 'truncated' Klotho forms in nematodes contain only one KL domain and lack the second internal repeat (whereas the full-length transmembrane vertebral Klothos are composed of two KL domains). Therefore, they correspond more closely to the alternatively spliced form for the shorter, directly secreted isoform in men (Polanska *et al.* 2011).

The function of mKlotho protein remained an enigma for years. It was first recognized that FGF23 deficiency and Klotho deficiency (with FGF23 resistance) result in closely resembling phenotypes, namely with hyperphosphataemia by up-regulated NaPi cotransporters, grossly elevated 1,25-(OH)₂ D₃, due to up-regulated 1 α -vitamin D hydroxylase, vascular calcification and premature ageing (Yoshida *et al.* 2002). In 2006, two independent groups reported that mKlotho served as a co-receptor for FGF23 (Kurosu *et al.* 2006, Urakawa *et al.* 2006). mKlotho forms a constitutive binary complex with the FGFR1c to create a *de novo* high-affinity binding site for FGF23. This Klotho–FGFR complex binds to FGF23 with much higher affinity than FGFR or Klotho alone; Klotho greatly enhances the ability of FGF23 to induce phosphorylation of FGFR substrate and ERK and is thus considered an essential cofactor for activation of FGF23 signalling by FGFR23. FGF23 elicits ERK phosphorylation and up-regulation of the expression of early growth-responsive (Egr) mRNA in Klotho-expressing tissues and cells, in the kidneys, parathyroids and pituitary, but not in liver, heart, bone and arterial walls (Kurosu *et al.* 2006, Urakawa *et al.* 2006, Ben-Dov *et al.* 2007, Krajisnik *et al.* 2007, Andrukhova *et al.* 2012, Lindberg *et al.* 2013). These (renal) FGF23 activity-enhancing properties of mKlotho are

particularly important as reducing phosphate (by depleting the diet in vitamin D or phosphate or by depleting NaPi cotransporters) and abolishing vitamin D activities rescue most phenotypes of *Klotho*^{−/−} mice (Morishita *et al.* 2001, Tsujikawa *et al.* 2003, Hesse *et al.* 2007, Nakatani *et al.* 2009, Ohnishi *et al.* 2009).

Klotho protein also exists in a soluble form, which can arise either from a distinct transcript or from ectodomain shedding of mKlotho (Fig. 1). The large extracellular domain of the membrane-bound form can be enzymatically cleaved (by α - and β -secretases) and released as 'secreted' Klotho into blood, urine and cerebrospinal fluid (CSF). The cleaved 130 kDa extracellular domain may be the preferably detected Klotho form in extracellular fluids, as indicated (by the green bars) in Figs 1 and 2 (Imura *et al.* 2004, Yamazaki *et al.* 2010). Soluble Klotho (sKlotho) released into the extracellular spaces can reach and affect a number of target tissues and processes, including regulation of hormone and growth factor signalling (e.g. inhibition of insulin/IGF1) (Kurosu *et al.* 2005, Wolf *et al.* 2008), and the regulation of plasma membrane amount and activity of ion channels and transporters, e.g. to





Figure 1

Transmembrane and soluble α -Klotho. A plasma cell membrane is shown into which mKlotho (single-pass transmembrane, very short cytoplasmic tail) is inserted. mKlotho associates with an FGFR; the latter can transmit (FGF23-induced) signals through its tyrosine kinase domains and ERK activation. sKlotho can be derived from mKlotho when its large extracellular part (130 kDa) is clipped off by enzymes; apart from this large sKlotho form, there may be a smaller, directly secreted sKlotho form (65 kDa), resulting from differential splicing. mKlotho (cofactor for FGF23 binding to FGFR) expression is restricted to a limited number of tissues while sKlotho is present in most extracellular spaces and can regulate many processes throughout the body, including calcium handling, nitric oxide synthesis, insulin and (e.g. insulin-like) growth factor signalling, as discussed in the text.

**Figure 2**

Effect of GH-secreting pituitary adenomas on circulating levels of IGF1 and sKlotho. (A) In the preoperative condition, there is a GH-secreting pituitary tumour in the pituitary gland. Excessive GH directly acts on the liver (predominant source of IGF1) and the increased production of IGF1 results in markedly increased serum IGF1 levels (mean \pm s.e.m., 0.58 ± 0.03 mg/l). Excessive GH also results in markedly increased serum sKlotho levels (4.1 ± 0.4 ng/ml); sKlotho is most likely of renal origin, but the mechanism (direct/indirect) by which GH excess results in increased release of kidney-derived sKlotho has not yet been elucidated. sKlotho particles as detected by the Yamazaki-ELISA (most likely the 130 kDa form) possibly represent only a tiny fraction of the total amount of sKlotho in the serum. The number of both IGF1 particles (small yellow circles) and sKlotho particles (green bars, as in Fig. 1) is dramatically increased, by factors of about 3 and 5 respectively. (B) Postoperatively, the GH-secreting adenoma has been removed, and GH action is normal; IGF1 (mean \pm s.e.m., 0.20 ± 0.01 mg/l) and sKlotho (0.8 ± 0.1 ng/ml) levels returned towards normal (data from Kohler *et al.* 2013).

attenuate calciuria. There is an ongoing debate on whether FGF23 can exert physiologically relevant effects on cells that do not express functional mKlotho, or do so only to a level below detection limits for the protein. Experimental data have suggested that at high doses, FGF23 and 130 kDa sKlotho, when combined, could allow FGF23 signalling, e.g. in chondrocytes and osteoblastic cells (Shalhoub *et al.* 2011, Kawai *et al.* 2013). Concerning vascular function and calcification, more recent data in mice and humans indicate that arterial Klotho expression was low or absent and did not mediate vascular FGF23 signalling (Lindberg *et al.* 2013, Scialla *et al.* 2013). Currently, sKlotho should not be labelled as a hormone as cognate receptors have not been identified so far, in contrast to those known for GH, IGF1, FGFs and for erythropoietin, $1,25\text{-(OH)}_2\text{D}$ and angiotensin II. In this regard, sKlotho (like renin) could be considered a renal enzyme rather than a hormone. Circulating sKlotho can regulate biological processes, e.g. by (enzymatic) glycan modification. Illustrating an action on the nephron, sKlotho modifies *N*-glycan chains of the epithelial Ca^{2+} channel transient receptor potential cation channel, subfamily V, member 5 (TRPV5) in the distal tubule. As a consequence, TRPV5 is retained on the cell surface and the activity of the ion channel is increased, and renal calcium loss is decreased (Chang *et al.* 2005, Cha *et al.* 2008, Alexander *et al.* 2009, Huang 2012, Olauson *et al.* 2012). α -Klotho has been found to be a central player in calcium (particularly sKlotho) and phosphate (predominantly mKlotho, FGF23 dependent) homeostasis, mainly by acting on the kidneys and the parathyroids (Imura *et al.* 2007, Nabeshima & Imura 2008, Kuro-o 2010). FGF23 suppresses production of PTH via mKlotho/FGFR-ERK1/2 (Ben-Dov *et al.* 2007, Krajisnik *et al.* 2007). Considering its effects on renal phosphate and calcium handling, α -Klotho shares phosphaturic effects (proximal tubule) and calcium-retaining effects (distal tubule) with PTH; however, their effects on generation of calcitriol are opposite to each other (stimulation by PTH, inhibition by Klotho). In the proximal tubule, FGF23 inhibits phosphate reabsorption by activating FGFR1 in a mKlotho-dependent fashion. sKlotho could also act as an enzyme (β -glucuronidase-like) to induce phosphaturia in an FGF23-independent manner; however, the direct enzyme target (potentially NaPi2a and NaPi2c) has been less well defined than in the case of TRPV5 (Hu *et al.* 2010). Disruption of the Klotho gene not only causes growth arrest, hyperphosphataemia and high calcitriol but also causes low glucose and insulin levels; however, whether loss of Klotho directly improves insulin sensitivity is still unclear (Utsugi *et al.* 2000, Hesse *et al.* 2007, Lorenzi *et al.* 2010). The mice are lean and have

almost no white adipose tissue (Klotho promotes adipocyte differentiation), lower body temperature and less energy expenditure (Mori *et al.* 2000, Chihara *et al.* 2006, Razzaque 2012). Another important role for Klotho, possibly related to inhibition of insulin/IGF1 signalling, is increased resistance to oxidative stress, which may contribute to the anti-ageing properties of Klotho (Yamamoto *et al.* 2005). Along this line, Klotho protein exerts protective effects, e.g. Klotho protein treatment attenuated cisplatin-induced reactive oxygen species generation. Thereby, it possibly helped to prevent reduction in cell viability in response to this ototoxic drug in an auditory cell line (Park *et al.* 2012). On the other hand, Klotho sensitized lung cancer cells to cisplatin, apparently via the PI3K/PKB pathway (Wang *et al.* 2013).

CP epithelial cells, besides renal tubular cells, express the highest levels of Klotho but they have been studied less often than kidney cells. Imura *et al.* (2007) were the first to address potential roles of Klotho in this tissue and reported on the action of α -Klotho on Na^+/K^+ ATPase activity and transcellular calcium transport. It appears likely that Klotho in CP cells contributes to regulation of composition and homeostasis of CSF similarly as the kidneys contribute to the homeostasis and regulation of plasma composition; therefore, CP has been considered as the 'kidney' of the brain. Recently, a molecular characterization employing a combination of transcriptomics and high-resolution tandem mass spectroscopy-based proteomics methods was carried out (in rats) to gain insight into CP function (Sathyanesan *et al.* 2012). A validation of the findings at the mRNA and protein level was done by *in situ* hybridization and immunohistochemistry. The close relationship between the transcriptome and proteome profiles from the CP and the kidney tubules was a main finding; physiological parallels between CP and renal tubular cells include high energy (and high blood flow) requirement and highly regulated pumps and transporter activities in both tissues. In addition, strong expression of Klotho was confirmed, and co-localization with transthyretin (an excellent CP marker) in CP was demonstrated (Kuro-o *et al.* 1997, Li *et al.* 2004, Sathyanesan *et al.* 2012). Chronic unpredictable stress reduced expression of Klotho in CP of their rats, as Ohyama found that acute inflammatory stress decreased renal Klotho expression (Ohyama *et al.* 1998, Sathyanesan *et al.* 2012).

Soon after discovery of the Klotho-deficient mouse, it was found that Klotho protein protects against endothelial dysfunction (Saito *et al.* 1998, 2000). Klotho contributes to maintenance of endothelial function and NO production via cAMP and calcium channel activation, upregulation of

the mitogen-activated kinase pathway, and increases stress tolerance, decreases oxidative damage, apoptosis and senescence of endothelial cells (Ikushima *et al.* 2006, Rakugi *et al.* 2007, Kusaba *et al.* 2010, Maekawa *et al.* 2011, Carracedo *et al.* 2012).

Germ line mutations or polymorphisms in Klotho do not obviously predispose to the development of neoplasms, and there is no evidence for an increased incidence of malignancies in Klotho-deficient mice. Nevertheless, Klotho can be viewed as a potential tumour suppressor, e.g. in breast, pancreatic, lung and renal cancer (Wolf *et al.* 2008, Chen *et al.* 2010, Abramovitz *et al.* 2011, Zhu *et al.* 2013). Intratumoral Klotho decline correlates with cancer progression, tumour size, TNM stage and nuclear grade in renal cell carcinoma, and with enhanced activation of the PI3K/PKB pathway (Zhu *et al.* 2013). Functional loss of Klotho due to epigenetic silencing was found to promote tumour progression in a variety of cancers.

Inhibition of Wnt signalling may play a role in the senescence-suppressing and renal fibrosis inhibitory effects of Klotho (Liu *et al.* 2007, Doi *et al.* 2011, Satoh *et al.* 2012, Sugiura *et al.* 2012, Zhou *et al.* 2013); TGF β appears to activate β -catenin via suppression of Klotho. sKlotho via its K11 domain blocks Wnt-triggered activation and nuclear translocation of β -catenin and thereby protects the kidneys from developing fibrosis in response to injury (Zhou *et al.* 2013). sKlotho attenuates IGF1 signals in the compensatory renal hypertrophy following uninephrectomy (Nagasu *et al.* 2011). Decreased serum sKlotho was associated with enhanced cyst growth in patients with autosomal dominant polycystic kidney disease (Pavik *et al.* 2012). As described for IGF1/PI3K/PKB signalling, sKlotho may also contribute to apoptosis induction and cell growth inhibition by decreasing the activity of the Wnt- β -catenin signalling pathway (Chen *et al.* 2012).

Gene delivery of Klotho could influence several pathophysiological phenotypes of Klotho mice, including improvement of vascular calcification (Shiraki-Iida *et al.* 2000), protection against endothelial dysfunction (Saito *et al.* 2000) and preventing progression of angiotensin II-induced or hypertensive renal damage (Mitani *et al.* 2002, Wang & Sun 2009). More recently, administration of recombinant sKlotho protein to Klotho^{-/-} mice was found to suppress accelerated ageing (Chen *et al.* 2013b).

Human pathologies due to deficient or excessive Klotho?

Years before immunoassays became available, genetic studies were performed. Arking *et al.* (2002) reported an association of human ageing with a functional variant of

Klotho. Humans with the *Klotho*-VS polymorphism were found to have reduced cognitive abilities (Deary *et al.* 2005). More recently, a functional variant of *Klotho* was found to be associated with early-onset ischaemic stroke (Majumdar *et al.* 2010). A 13-year-old girl with a remarkable phenotype of hyperphosphataemia, and soft tissue and vascular calcification, 'tumoral' calcinosis, was found to harbour a homozygous missense (H193R) mutation in the *Klotho* gene. Hyperphosphataemia was consistent with renal FGF23 resistance, as *Klotho* mutations are expected to prevent complex formation with FGF23 (functionally defective receptors), and lead to decreased FGF23 signalling: FGF23 was indeed markedly elevated (familial 'tumoral' calcinosis was a known phenotype caused by inactivating mutations in FGF23 or GALNT3, i.e. diseases caused by defective ligand had been described before). Hyperparathyroidism was proposed to be caused by a lack of an α -*Klotho*-dependent inhibitory effect of FGF23 on PTH production, although it should be noted that α -*Klotho* could also affect PTH secretion directly (without FGF23), by increasing cell surface Na^+/K^+ -ATPase activity (Ichikawa *et al.* 2007, Imura *et al.* 2007). Another exceptional case was reported, with a translocation adjacent to α -*Klotho* gene, causing markedly increased circulating α -*Klotho* (and FGF23) level, resulting in hypophosphataemia (with rickets) and hyperparathyroidism (Brownstein *et al.* 2008). In animals, circulating α -*Klotho* *per se* may induce FGF23 production and phosphaturia (Smith *et al.* 2012), and so may antibodies activating FGFR1 (Wu *et al.* 2013).

sKlotho is released into the circulation, and its plasma half-life has been estimated to be about 20–30 min by Imura. sKlotho can be detected by a recently developed and commercially available sandwich ELISA (Yamazaki *et al.* 2010; Table 1). Levels of sKlotho decrease with age in healthy subjects and are inversely related to mortality in the elderly (Semba *et al.* 2011). High levels of sKlotho were found in human umbilical cord blood (Ohata *et al.* 2011, Siahianidou *et al.* 2012). The placenta is known to produce sKlotho and may account for these high levels. However, except for the neonatal period, the kidneys appear to be the major source of sKlotho in sera of human subjects. Accordingly, serum sKlotho decreased by about 40% following nephrectomy in living donors (Akimoto *et al.* 2013). So far, there is no well-established reference range for serum sKlotho. In our Zürich area, the median value of healthy adult subjects is around 0.6 ng/ml and, therefore, somewhat lower than previously reported by Japanese and Italian authors using the same assay (Yamazaki *et al.* 2010, Semba *et al.* 2011, Akimoto *et al.*

2012, Neidert *et al.* 2013, Pavik *et al.* 2013). Interestingly, the Italian authors found that low circulating sKlotho levels in elderly adults are not only associated with increased mortality but also associated with poor muscle strength and disability in activities of daily living (Semba *et al.* 2011, 2012, Crasto *et al.* 2012).

It should be mentioned that many uncertainties still exist when assessing sKlotho levels, as the identity of the detected molecules is not always clear (see also Figs 1 and 2). In fact, in 2004, Xiao *et al.* reported on a polyclonal antibody raised in rabbits against the C-terminal of human secreted *Klotho* protein (amino acids 290–549). It could be used for western blotting to detect mainly a 60 kDa protein and for an ELISA to measure sKlotho in human serum; sKlotho was found to decrease with age with this particular assay (Xiao *et al.* 2004).

The ImmunoBiological Laboratories (IBL, Fujioka, Japan) ELISA as developed by Yamazaki and the time-resolved fluorescence immunoassay (Cusabio, Wuhan, China) assays have recently been directly compared in a Danish study where considerable differences became evident (Pedersen *et al.* 2013). The IBL ELISA utilizes two monoclonal antibodies that may detect and measure one larger form of ('intact', full-length shedded) *Klotho*. On the other hand, the Cusabio assay (using mouse monoclonal antibody-precoated plates and goat polyclonal antibodies of unknown epitope specificity for detection) may recognize additional, shorter forms (65 kDa and possibly other forms) that may be much more abundant (Matsumura *et al.* 1998). As mentioned, the IBL assay has been used quite often lately (Table 1), whereas the Cusabio-ELISA has been validated just recently in studies performed in the USA and Denmark (Devaraj *et al.* 2012, Pedersen *et al.* 2013). In their review, Hu *et al.* (2013) estimated plasma concentrations of sKlotho in the range of 10–50 nM, which would be much higher than what is measured by the Yamazaki assay (Yamazaki *et al.* 2010).

CKD can be viewed as a state of *Klotho* deficiency. As the kidney is the primary source and a major target of *Klotho*, the focus of *Klotho* research has been on nephrology, apart from basic and ageing research. Although renal expression of the *Klotho* gene and urinary *Klotho* levels have been found to be decreased in renal disease (Koh *et al.* 2001, Hu *et al.* 2011), it currently remains controversial whether plasma sKlotho is a useful biomarker for kidney disease (Sugiura *et al.* 2011, Pavik *et al.* 2012, 2013, Kim *et al.* 2013, Seiler *et al.* 2013). FGF23 may give a much better readout than sKlotho, as the hormone released by bone in response to kidney disease appears to quite reliably reflect renal resistance to FGF23.

Table 1 Soluble serum Klotho levels in healthy human subjects as assessed by IBL ELISA

Author/year	n	Subjects	Age (years)	Sex (m/f)	Klotho levels (pg/ml)	Comments
Akimoto <i>et al.</i> (2013)	10	Living kidney donors	64 ± 9	4/6	910 (median); 755–1132 (IQR)	The kidneys seem to be the major source of serum sKlotho
Amitani <i>et al.</i> (2013)	11	Healthy	21 ± 1 (S.E.M.)	0/11	1392 ± 145 (S.E.M. ± S.E.)	Klotho levels are lower in patients with anorexia nervosa and patients with obesity than in normal weight controls
Crasto <i>et al.</i> (2012)	802	Elderly	> 65	357/445	689 (mean); 238 (S.D.)	Low serum Klotho is associated with ADL disability
Neidert <i>et al.</i> (2013)	26	Healthy	39 ± 8	15/11	596 (median); 506–734 (IQR)	Serum Klotho is excessively elevated in patients with untreated acromegaly
Pedersen <i>et al.</i> (2013)	120	Healthy	40 (19–66) median (range)	58/62	472 ± 137 (mean ± S.D.)	Different immunoassays recognize distinct forms of sKlotho
Pavik <i>et al.</i> (2012)	20	Healthy	32 ± 6	12/8	1200 ± 385 (mean ± S.D.)	Low serum sKlotho is associated with cyst growth and low phosphaturic activity of FGF23 in ADPKD
Pavik <i>et al.</i> (2013)	21	Healthy	48 ± 8	9/12	600 (median); 429–862 (IQR)	Serum Klotho decreases in advanced CKD
Semba <i>et al.</i> (2011)	1023	Elderly	73 (69–79) median (IQR)	564/459	676 (median); 530–819 (IQR)	Klotho is inversely related to CVD risk
Semba <i>et al.</i> (2011)	804	Elderly	> 65	354/450	697 ± 325 (mean ± S.D.)	Serum sKlotho is suggested as an independent predictor of all-cause mortality
Sugiura <i>et al.</i> (2011)	10	Healthy	20–44	NA	404 ± 87 (mean ± S.D.)	Increased serum sKlotho is suggested as a renoprotective factor in CKD
Yamazaki <i>et al.</i> (2010)	181	142 healthy adults (39 children)	61 ± 19 (7 ± 8)	66/76 (23/16)	562 ± 146 (mean ± S.D.) 952 ± 282 (mean ± S.D.)	Serum Klotho is inversely related to age and creatinine

ADL, activities of daily living; ADPKD, autosomal dominant polycystic kidney disease; CKD, chronic kidney disease; CVD, cardiovascular disease; eGFR, estimated glomerular filtration rate; FGF23, fibroblast growth factor 23; IBL, ImmunoBiological Laboratories Co. Ltd.; IQR, interquartile range.

In this context, it may be worth mentioning potential off (classical) target effects of hormones adapting to a resistance induced by Klotho deficiency: FGF23 excess may (not result in hypophosphataemia but) induce left ventricular hypertrophy in a Klotho-independent manner (Faul *et al.* 2011).

As reported by Imura *et al.* (2004), sKlotho can be found not only in serum and urine but also in CSF. Interestingly, levels of sKlotho are quite high in CSF and in urine; CP and kidney as Klotho-expressing sites may account for these findings. CSF Klotho levels as measured by immunoassay have not been reported in the literature, but data for urine are available and suggest that urine Klotho is linked to renal function (Akimoto *et al.* 2012). Over the past 6 months (January–June 2013), we have also started to assess CSF and urine samples for sKlotho. According to these preliminary data, concentrations of

sKlotho are similar or slightly higher in CSF than in serum: sKlotho was 888 (595–1319) pg/ml (median (range)) in CSF ($n=9$) vs 557 (255–1017) pg/ml in the corresponding sera. This is remarkable, given that the total protein concentration is two orders of magnitude lower in CSF than in plasma and serum. Despite samples obtained from quite distinct patients (none of them, with acromegaly) for CSF analysis, we found a narrow concentration range in CSF when compared with serum. This could suggest that Klotho levels are maintained within a certain limit within this compartment. Klotho may be biologically active and play an important role in brain and regulating functions of CSF (Imura *et al.* 2004, 2007). It is currently unknown whether changes in CSF Klotho reflect and/or contribute to neurological and neurosurgical disorders.

As mentioned earlier, urinary Klotho concentrations have been measured in patients with renal disease. We had

a special interest to compare serum and urine sKlotho levels in patients with acromegaly.

Urinary Klotho concentrations remained below the detection limit in unconcentrated spot urine samples of nine patients cured from acromegaly and of six patients with non-functioning pituitary adenomas but detectable in nine patients with newly diagnosed acromegaly, which suggests that increased urinary sKlotho could have a high positive predictive value for the diagnosis of the disease (Fig. 3). However, technical issues need to be addressed and the sensitivity of the analysis refined before urinary Klotho can be considered a useful marker for estimating GH excess in individual patients.

Based on sequence similarity with the *Klotho* gene, β -Klotho was identified, and the former (the founder of the family) occasionally renamed α -Klotho (Ito *et al.* 2000). γ -Klotho was identified as another, functionally more remote member of the family, which is expressed particularly in kidneys, adipose tissue and eyes (Ito *et al.* 2002). α -Klotho protein shares 41% amino acid sequence identity with β -Klotho. β -Klotho is expressed predominantly in the liver and white adipose tissue, i.e. in tissues that are targets of FGF19 and FGF21 (endocrine FGFs, like

FGF23), and β -Klotho was found to work as co-receptor for FGF19 and FGF21 signalling. In mammals, Klothos (α and β) are type 1 transmembrane proteins that function as obligatory cofactors for signalling of all known, vertebrate specific, hormone-like ('endocrine') FGFs: FGF15/FGF19 (enterocyte), 21 (hepatocyte) and 23 (osteocyte). The (FGF) ligand, the (FGF) receptor and its (Klotho) co-receptor usually define the specificity of the endocrine actions (Tomiya *et al.* 2010); however, in artificial experimental settings, promiscuity of signalling is observed, for instance by FGF23 mutants with an ability to activate FGFR signalling through both α - and β -Klotho (Wu *et al.* 2012), or with antibodies activating FGFR1, which may act in a FGF21-like manner (requiring β -Klotho on target cells) or in a FGF23-like manner (requiring α -Klotho on target cells), to induce hypophosphataemia (Wu *et al.* 2013).

Role for α -Klotho in acromegaly?

Luft and colleagues recognized that acromegaly had a marked impact on renal function (Ikko *et al.* 1956). Administration of GH increased renal GFR and tubular phosphate reabsorption (Corvilain *et al.* 1962, Bianda *et al.* 1997). The latter effect of GH cannot be explained by a deficiency of known classical phosphaturic hormones such as PTH and FGF23. Reviewing old data on GH, IGF1 and phosphate transport, it is important to realize that stimulatory effects of GH on renal phosphate reabsorption in adult humans have been repeatedly found *in vivo*. A stimulatory effect of IGF1 on phosphate uptake can be documented in most cultured cells, but the finding of phosphate uptake stimulation by IGF1 *in vitro* has not been shared by GH, and the claim that IGF1 is mediating this effect of GH may have been premature (Quigley & Baum 1991, Zoidis *et al.* 2004). Maximal tubular phosphate reabsorption per GFR was raised by GH but not by IGF1 compared with saline; thus, IGF1 does not mimic (and possibly not mediate) the TmP/GFR raising effect of GH on renal phosphate reabsorption despite comparable effects on GFR (Bianda *et al.* 1997, Schmid & Meili 2000).

Acromegaly has unique biochemical and endocrine features; it leads to high serum phosphate and FGF23 levels despite enhanced GFR, as well as to insulin resistance (IR) and hyperglycaemia (Table 2). These characteristics are distinct from those observed in more common conditions such as renal failure and type 2 diabetes: high phosphate levels cannot be attributed to decreased GFR, and IR and hyperglycaemia are not caused by increased visceral fat. Although levels of phosphate and FGF23 are high in




Figure 3

Concentrations of soluble Klotho (sKlotho) in serum and urine of patients with acromegaly. Concentrations of Klotho were assessed by ELISA in serum and urine samples (spots, obtained the same time point as blood) from patients with pituitary adenomas (between 1/2013 and 6/2013). Each point represents an individual; filled circles, active (newly diagnosed, untreated) acromegaly (nine patients, three females); empty circles, successfully (by transphenoidal surgery) treated acromegaly (as confirmed by normal IGF1 and GH suppressible to <1 ng/ml during oGTT; nine patients, three females) and filled triangles, clinically non-functioning pituitary adenoma (six patients, three females). Note that both the x-axis (serum sKlotho (pg/ml)) and y-axis (urine Klotho, expressed in pg sKlotho/mmol creatinine) are drawn on logarithmic scales. Patients with active acromegaly have higher serum and urinary sKlotho levels than patients without active acromegaly; in urine, sKlotho was readily detectable (concentration above the detection limit of the ELISA) only in samples from patients with acromegaly.

Table 2 Parameters and phenotypes characterizing GH excess, Klotho deficiency and CKD

Parameter	GH excess	Klotho deficiency	CKD
Growth	↑↑	↓↓	↓
Lifespan	↓	↓↓↓	↓↓
Energy expenditure	↑	↓	↔↓
Glucose, IR	↑↑	↓↓	↔↑
Creatinine	↓	↑	↑↑↑
eGFR	↑	↓	↓↓↓
1,25-(OH) ₂ vitamin D	↑↑	↑↑↑	↓↓
Phosphate	↑	↑↑↑	↑↑
FGF23	↑	↑↑↑	↑↑↑
Calcium	↔↑	↑↑	↔↓
PTH	↔	↔↓	↑↑

CKD, chronic kidney disease; eGFR, estimated glomerular filtration rate; IR, insulin resistance.

acromegaly, calcitriol is also increased in this disease. FGF23 (usually inhibiting renal phosphate reabsorption and calcitriol production) and phosphate tend to be high despite increased GFR, in keeping with a condition of renal FGF23 resistance (Table 2; Ito *et al.* 2007).

Longer lasting excessive secretion of GH by a pituitary adenoma results in premature death, mainly due to an increased risk for cardiovascular disease. In human adults, GH (but not IGF1) increases the set points of plasma glucose and phosphate, and both IR and elevated serum phosphate are associated with increased mortality in the general population.

As Klotho levels may relate to FGF23 resistance and IR (or high phosphate and glucose respectively) in acromegaly, we measured sKlotho in sera of patients before and after transsphenoidal surgery. Removal of the pituitary adenoma led to the expected decrease in GH and IGF1. sKlotho was markedly increased in relation to GH excess and declined towards normal levels after surgery (Sze *et al.* 2012); differences in sKlotho before and after surgery were at least as pronounced as IGF1 (Sze *et al.* 2012, Kohler *et al.* 2013; Fig. 2). While sKlotho was elevated in sera of patients with active acromegaly, this did not hold true for patients with clinically non-functioning pituitary adenomas (Neidert *et al.* 2013). In the latter study, it was shown that sKlotho fell at least as quickly as IGF1 after surgery. Moreover, the pituitary adenomas were analysed for Klotho by immunohistochemical staining. It was found that the rise in serum α -Klotho is not explained by increased pituitary (adenoma) Klotho expression, but rather by increased pituitary GH secretion and possibly due to a systemic action of GH. However, renal biopsies to check for changes in amounts of mKlotho were not performed.

As sKlotho may reflect activity of GH-producing adenomas, it could serve as an additional serum marker in the long-term follow-up of acromegalic patients. Overall, when assessing patients with GH-producing adenomas, we observed concomitant and parallel changes in serum IGF1 and sKlotho over time in a given individual patient, and levels of sKlotho and IGF1 appeared to be similarly dependent on GH (Sze *et al.* 2012, Kohler *et al.* 2013). A remarkable difference was found with regard to gender; for a given GH (excess) status, IGF1 tended to be higher in males whereas sKlotho levels were higher in females; more detailed analysis revealed that oestrogens accounted for lower IGF1 in (premenopausal) women, but not for the higher sKlotho; overall, GH was by far the strongest predictor of both sKlotho and IGF1 (L Sze, MC Neidert, RL Bernays & C Schmid, 2013, unpublished observations).

In human adults, GH deficiency results in a more or less characteristic but much less obvious phenotype than in children (dwarfism); hypoglycaemia and hypophosphataemia are not obvious features. In the study by Neidert, sKlotho did not appear to be particularly low in patients with GH deficiency. It remains to be tested whether sKlotho levels are decreased in patients with GH deficiency, and whether GH treatment results in increasing serum sKlotho levels. Considering sKlotho as a GH-dependent serum protein, a response to GH replacement might be anticipated, but in view of kidney function and phosphate homeostasis, differences in sKlotho are possibly smaller between healthy adults and patients with GH deficiency when compared with those with GH excess.

GH and Klotho

As mentioned, *Klotho* mutant (*Klotho*^{-/-}) mice exhibit growth retardation after weaning and premature death. Electron microscopic examination of GH-producing cells in pituitary glands revealed a reduction in GH granules. Growth retardation was described in the original 1997 paper (Kuro-o *et al.* 1997). As it has not been known whether growth retardation in *Klotho* mutant mice is related to loss of GH function, Kashimada *et al.* examined whether treatment with GH could rescue retardation of growth. At the end of 3 weeks of treatment with human GH, body weight was not increased in *Klotho* mutant (*Klotho*^{-/-}) mice in contrast to WT mice. Growth retardation in *Klotho* mutant mice is therefore resistant against GH treatment, and GH deficiency is not the cause of growth arrest in *Klotho*^{-/-} mice (Kashimada *et al.* 2002).

There are data suggesting a negative impact on lifespan by an excess in the anabolic peptide hormones GH, IGF1 and insulin. These data are difficult to accept not only by people abusing such hormones but also for the medical community, including endocrinologists. With regard to life expectancy, apparent contradictions concerning the action of GH should be considered. GH has generally been associated with positive attributes of adolescence and youth, such as growth, health, muscle mass and well-being, yet the bulk of scientific evidence suggests that signalling through GH, IGF1 and insulin receptors is related to a shortened lifespan in adults (Berryman *et al.* 2008, Junnila *et al.* 2013). It is currently difficult to estimate whether high sKlotho levels can attenuate insulin and IGF1 receptor signalling *in vivo*, e.g. in patients with acromegaly. In their overviews, Kopchick and colleagues have underlined the importance of GH, IGF1 and insulin signalling in regulating lifespan; according to our view, not only the GH-IGF1 axis and IR but also FGF23 resistance and hyperphosphataemia in the context of the GH-Klotho axis deserve more attention.

Concerning the renal effects of GH, several aspects of Klotho can be considered (Table 2). Apparently, GH clearly opposes the effects attributed to mKlotho and its essential role in mediating FGF23-induced renal phosphate loss and FGF23-mediated inhibition of calcitriol production (Bianda *et al.* 1997). Hypervitaminosis D due to over-expression of CYP27B1 is central to the phenotypes of α -mKlotho and FGF23 deficiency, and hypovitaminosis D due to suppression of CYP27B1 is a central phenotype of FGF23 transgenic mice. With regard to vitamin D metabolism, GH increases vitamin D activation in humans and experimental animals and can, therefore, be considered as an antagonist of FGF23 action mediated via the FGFR/ α -mKlotho complex (Bianda *et al.* 1997, 1998, Zoidis *et al.* 2002). GH stimulates 25-hydroxy-vitamin D 1 α -hydroxylase (CYP27B1) mRNA and markedly down regulates 24-hydroxylase (CYP24A1) mRNA expression; the former enzyme favours activation and the latter degradation of active vitamin D (Zoidis *et al.* 2002). As mentioned in the previous section, the high levels of phosphate and calcitriol together with increased FGF23 suggest a state of renal FGF23 resistance in patients with acromegaly (Ito *et al.* 2007, Sze *et al.* 2012). It is, therefore, conceivable that GH induces functional impairment of the FGF23-mKlotho signalling pathway in proximal tubular cells.

On the other hand, GH shares the effects of sKlotho on the distal convoluted tubule to attenuate calciuria. However, as calcium, calcitriol and GFR tend to be

increased in patients with acromegaly, increased calcium reabsorption in the distal convoluted tubule may be masked (acromegalic patients tend to have fasting hypercalciuria). sKlotho may not only account for an increased calcium loss when deficient (as in diabetic nephropathy) but also for attenuation of renal calcium loss or when present in excess (as in acromegaly) (Asai *et al.* 2012, Kamenicky *et al.* 2012).

Concerning parathyroid function, findings reported from patients with acromegaly have been variable, especially in studies of drug-treated patients (White *et al.* 2006, Kamenicky *et al.* 2012). More consistent data from patients with adenectomy-treated acromegaly suggest that PTH levels remain essentially unchanged when the preoperative condition is compared with the postoperative condition (Lund *et al.* 1981, Takamoto *et al.* 1985). Likewise, high-dose short-term GH treatment in a more controlled setting of healthy volunteers or GH-deficient patients did not alter PTH levels, despite significant increases in 1,25-(OH)₂ D and a trend for increased calcium levels, as observed in acromegaly (Lund *et al.* 1981, Takamoto *et al.* 1985, Bianda *et al.* 1997, Bianda *et al.* 1998). Given the fact that in acromegalic patients, consistently calcium tends to be high in parallel with increased calcitriol and FGF23 values, but in the presence of unchanged PTH, it is likely that inappropriate feedback inhibition of PTH occurs (Table 2). One potential explanation would be that FGF23 action on PTH secretion is impaired possibly by a functional mKlotho deficiency (as proposed for kidneys; Ben-Dov *et al.* 2007, Krajisnik *et al.* 2007).

Considering effects of GH on the endothelium, it appears that some are potentially favourable for cardiovascular health whereas others such as IR, hyperglycaemia and hyperphosphataemia negatively impact on endothelial function. Ageing is associated with a decrease in GH, endothelial function (nitric oxide synthase activity and abundance of eNOS), cardiovascular and renal health. GH deficiency has been associated with impaired endothelial function, and GH replacement could improve endothelial function (Boger *et al.* 1996, Evans *et al.* 2000). As mentioned earlier, Klotho protein protects against endothelial dysfunction and exerts several beneficial (anti-senescence) effects on endothelial cells, including some mediated via the MAPK pathway (Saito *et al.* 1998, 2000, Ikushima *et al.* 2006, Kusaba *et al.* 2010, Maekawa *et al.* 2011, Carracedo *et al.* 2012). However, it is currently unknown whether Klotho is involved (a downstream effector or mediator) in the effects of GH on the endothelium.

In the context of acromegaly, it has been well recognized that hypertension and hyperglycaemia/diabetes contribute to the increased cardiovascular disease risk in these patients. Phosphate has been considered less as a risk factor, although more recent data suggest that hyperphosphataemia and possibly phosphate intake are associated with increased cardiovascular risk, even in the absence of established renal disease (Tonelli *et al.* 2005, Dhingra *et al.* 2007, Foley 2009, Shuto *et al.* 2009, Ellam & Chico 2012, Osuka & Razzaque 2012, Yamamoto *et al.* 2013). Excessive phosphate not merely favours ectopic (including media) calcification but also impairs endothelial function (e.g. NO synthesis, annexin II regulation, etc.) (Lau *et al.* 2011, Di Marco *et al.* 2013). In acromegaly, it may be that potentially beneficial effects of GH and Klotho on endothelial function and NO synthesis are outweighed by IR and hyperphosphataemia, so that there is a net increase in cardiovascular risk. GH-induced high phosphate levels may make sense in the context of growth, but it appears likely that they are harmful in adulthood.

The findings of remarkably high serum sKlotho levels in patients with active acromegaly are difficult to interpret, and additional data are required. The biological meaning remains unclear, and currently, it is uncertain whether changes in Klotho turnover account for the high threshold of renal phosphate reabsorption and FGF23 resistance with GH excess; moreover, the mechanisms by which acromegaly leads to excess sKlotho remain to be elucidated. One can speculate about several mechanisms; GH may increase Klotho gene expression in kidneys (and other sites). Soluble forms of α -Klotho could be generated by alternative splicing of its transcript but they could also result from proteolytic cleavage of the transmembrane form by secretases into various body fluids. Two members of the 'A Disintegrin and Metalloproteinase' (ADAM) family, ADAM10 and ADAM17 (α -secretases), and β -site amyloid precursor protein-cleaving enzyme 1 (BACE1) (β -secretase), have been suggested as responsible enzymes (Chen *et al.* 2007, Bloch *et al.* 2009, Saftig & Reiss 2011). The high sKlotho levels in acromegalic patients do not necessarily reflect a high level of cell mKlotho (in the kidneys, the parathyroids and elsewhere). A discrepancy between renal mKlotho and sKlotho has been discussed for experimental settings, and it was considered that increased sKlotho could be caused by accelerated shedding of Klotho without corresponding changes in mKlotho expression (Lau *et al.* 2012). In acromegaly, FGF23 resistance cannot be explained by high sKlotho *per se*, unless sKlotho is elevated at the expense of mKlotho, which

serves as a co-receptor for FGF23. Formally, resistance to FGF23 may be caused by decreased expression of the Klotho–FGFR1 complex. This should be investigated at the level of the kidneys and the parathyroids, and this has indeed been addressed in patients with CKD where parathyroids were studied to find the basis for parathyroid resistance to FGF23 (Galitzer *et al.* 2010, Komaba *et al.* 2010, Krajisnik *et al.* 2010). Similar investigations have not been performed with tissues from patients with acromegaly. According to our hypothesis, the activity of secretases (shedding the ectodomain from the integral mKlotho) may be increased in acromegaly, either directly by GH or indirectly by factors or by proteolytic activity induced by GH. As previously mentioned, GH has a proteolytic activity on other systems, as it induced an IGFBP3-proteolytic activity *in vivo* in rats; moreover, GH stimulates the mRNA expression not only of the cytochrome P450 enzyme CYP27B1 involved in the production of active vitamin D (calcitriol) but also the phex mRNA that encodes for an endopeptidase important in phosphate homeostasis (Rutishauser *et al.* 1993, Zoidis *et al.* 2002). Increased mKlotho-proteolytic activity could result in deficient mKlotho and impaired FGF23 signalling, thereby explaining FGF23 resistance; the enhanced enzymatic clipping of the extracellular Klotho domain would lead to an excess in sKlotho contributing to channel and transporter regulation, to attenuation of calciuria, growth factor signalling and IR.

Thus far, it has not been studied whether GH stimulates the production of Klotho. It will be of interest to see whether GH increases serum levels of sKlotho in GH-deficient subjects and to see whether GH influences expression of Klotho at the mRNA level and whether it regulates the abundance of mKlotho in tissues and sKlotho in extracellular spaces. Such studies have been performed in experimental animals with calcium- and phosphate-regulating hormones (PTH, calcitriol, FGF23; increasing renal expression and serum levels in rats) but not with GH (Takenaka *et al.* 2013).

As mentioned earlier, β -Klotho is important as a co-receptor for FGF19 and FGF21 signalling. GH induces hepatic production of FGF21, apparently through a mechanism dependent on lipolysis in adipocytes. In this context, it is interesting to know that GH stimulates lipolysis especially in the state of fasting (when insulin is low). FGF21, in turn, inhibits GH action on chondrocytes directly at the growth plate; FGF21 may also mediate GH resistance in anorexia nervosa as illustrated by high GH and low IGF1. Taken together, there are not only multiple interacting targets between GH and α -Klotho in calcium

and phosphate mineral handling but it is likely that GH and β -Klotho pathways are closely linked in order to regulate fuel metabolism, at both afferent and efferent parts of endocrine loops. So far, due to lack of appropriate assays to measure β -Klotho and α -Klotho, data lag far behind basic research in this field.

Conclusions

There is increasing interest in the field of GH and Klotho and their mutual interaction. Many questions can be addressed by both basic and clinical studies. Both IGF1 and sKlotho are relatively abundant in the circulation; both of them are regulated by GH (Sze *et al.* 2012, Hu *et al.* 2013, Neidert *et al.* 2013). In the case of IGF1, there is fairly good evidence that it mediates some actions of GH and that there is a negative feedback at the pituitary level. α -Klotho expression can also be documented in the pituitary gland (Kuro-o *et al.* 1997, Li *et al.* 2004, Neidert *et al.* 2013), and FGF23 administration to rats increased early growth response-1 (*Egr1*) mRNA levels in the pituitary as it did in the kidney and parathyroids (Urakawa *et al.* 2006); however, a feedback action of FGF23 and Klotho on the pituitary GH secretion has not been described so far. There are no data available that show how FGF23 or Klotho may mediate GH actions.

In acromegaly, IGF1 (predominantly liver-derived), a marker and mediator of GH action, is increased, as is sKlotho (predominantly kidney-derived), a marker of GH-producing pituitary adenomas. Further observations and experiments should allow more insight, e.g. as to whether generation of sKlotho could mediate selected actions of GH, causing FGF23 resistance, hyperphosphataemia, possibly contributing to IR and hyperglycaemia, contributing to improved endothelial function, increased energy expenditure and decreased life expectancy.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the review reported.

Funding

This review did not receive any specific grant from any funding agency in the public, commercial or not-for-profit sector.

Acknowledgements

The authors would like to thank the patients participating in our studies and physicians referring them to us: Drs M Brändle, Ch Geiss, S Hofer,

E Keller, S Kohler, D Möller, L Regli, U Schwarz, St Suter and P Wiesli. They also thank Cornelia Zwimpfer for performing the α -Klotho assays and Dr E Christ for critical reading and valuable suggestions.

References

- Abramovitz L, Rubinek T, Ligumsky H, Bose S, Barshack I, Avivi C, Kaufman B & Wolf I 2011 KL1 internal repeat mediates Klotho tumor suppressor activities and inhibits bFGF and IGF-I signaling in pancreatic cancer. *Clinical Cancer Research* **17** 4254–4266. (doi:10.1158/1078-0432.CCR-10-2749)
- ADHR Consortium 2000 Autosomal dominant hypophosphataemic rickets is associated with mutations in FGF23. *Nature Genetics* **26** 345–348. (doi:10.1038/81664)
- Akimoto T, Yoshizawa H, Watanabe Y, Numata A, Yamazaki T, Takeshima E, Iwazu K, Komada T, Otani N, Morishita Y *et al.* 2012 Characteristics of urinary and serum soluble Klotho protein in patients with different degrees of chronic kidney disease. *BMC Nephrology* **13** 155. (doi:10.1186/1471-2369-13-155)
- Akimoto T, Kimura T, Watanabe Y, Ishikawa N, Iwazu Y, Saito O, Muto S, Yagisawa T & Kusano E 2013 The impact of nephrectomy and renal transplantation on serum levels of soluble Klotho protein. *Transplantation Proceedings* **45** 134–136. (doi:10.1016/j.transproceed.2012.07.150)
- Alexander RT, Woudenberg-Vrenken TE, Buurman J, Dijkman H, van der Eerden BC, van Leeuwen JP, Bindels RJ & Hoenderop JG 2009 Klotho prevents renal calcium loss. *Journal of the American Society of Nephrology* **20** 2371–2379. (doi:10.1681/ASN.2008.121273)
- Amitani M, Asakawa A, Amitani H, Kaimoto K, Sameshima N, Koyama KI, Haruta I, Tsai M, Nakahara T, Ushikai M *et al.* 2013 Plasma klotho levels decrease in both anorexia nervosa and obesity. *Nutrition* **29** 1106–1109. (doi:10.1016/j.nut.2013.02.005)
- Andrukhova O, Zeitz U, Goetz R, Mohammadi M, Lanske B & Erben RG 2012 FGF23 acts directly on renal proximal tubules to induce phosphaturia through activation of the ERK1/2-SGK1 signaling pathway. *Bone* **51** 621–628. (doi:10.1016/j.bone.2012.05.015)
- Arking DE, Krebsova A, Macek M Sr, Macek M Jr, Arking A, Mian IS, Fried L, Hamosh A, Dey S, McIntosh I *et al.* 2002 Association of human aging with a functional variant of Klotho. *PNAS* **99** 856–861. (doi:10.1073/pnas.022484299)
- Asai O, Nakatani K, Tanaka T, Sakan H, Imura A, Yoshimoto S, Samejima K, Yamaguchi Y, Matsui M, Akai Y *et al.* 2012 Decreased renal α -Klotho expression in early diabetic nephropathy in humans and mice and its possible role in urinary calcium excretion. *Kidney International* **81** 539–547. (doi:10.1038/ki.2011.423)
- Bai X, Miao D, Li J, Goltzman D & Karaplis AC 2004 Transgenic mice overexpressing human fibroblast growth factor 23 (R176Q) delineate a putative role for parathyroid hormone in renal phosphate wasting disorders. *Endocrinology* **145** 5269–5279. (doi:10.1210/en.2004-0233)
- Bauer C, Brucker R, Butzberger S & Schmid C 2010 Hypophosphataemia-inducing mesenchymal tumour in the foot. *BMJ Case Reports*. (doi:10.1136/bcr.06.2009.2034)
- Ben-Dov IZ, Galitzer H, Lavi-Moshayoff V, Goetz R, Kuro-o M, Mohammadi M, Sirkis R, Naveh-Manly T & Silver J 2007 The parathyroid is a target organ for FGF23 in rats. *Journal of Clinical Investigation* **117** 4003–4008. (doi:10.1172/JCI32409)
- Berryman DE, Christiansen JS, Johannsson G, Thorner MO & Kopchick JJ 2008 Role of the GH/IGF-1 axis in lifespan and healthspan: lessons from animal models. *Growth Hormone & IGF Research* **18** 455–471. (doi:10.1016/j.ghir.2008.05.005)
- Bianda T, Hussain MA, Glatz Y, Bouillon R, Froesch ER & Schmid C 1997 Effects of short-term insulin-like growth factor-I or growth hormone treatment on bone turnover, renal phosphate reabsorption and 1,25 dihydroxyvitamin D3 production in healthy man. *Journal of Internal Medicine* **241** 143–150. (doi:10.1046/j.1365-2796.1997.94101000.x)

- Bianda T, Glatz Y, Bouillon R, Froesch ER & Schmid C 1998 Effects of short-term insulin-like growth factor-I (IGF-I) or growth hormone (GH) treatment on bone metabolism and on production of 1,25-dihydroxy-cholecalciferol in GH-deficient adults. *Journal of Clinical Endocrinology and Metabolism* **83** 81–87. (doi:10.1210/jc.83.1.81)
- Bloch L, Sineshchekova O, Reichenbach D, Reiss K, Saftig P, Kuro-o M & Kaether C 2009 Klotho is a substrate for α -, β - and γ -secretase. *FEBS Letters* **583** 3221–3224. (doi:10.1016/j.febslet.2009.09.009)
- Boger RH, Skamira C, Bode-Boger SM, Brabant G, von zur Muhlen A & Frolich JC 1996 Nitric oxide may mediate the hemodynamic effects of recombinant growth hormone in patients with acquired growth hormone deficiency. A double-blind, placebo-controlled study. *Journal of Clinical Investigation* **98** 2706–2713. (doi:10.1172/JCI119095)
- Brownstein CA, Adler F, Nelson-Williams C, Iijima J, Li P, Imura A, Nabeshima Y, Reyes-Mugica M, Carpenter TO & Lifton RP 2008 A translocation causing increased α -Klotho level results in hypophosphatemic rickets and hyperparathyroidism. *PNAS* **105** 3455–3460. (doi:10.1073/pnas.0712361105)
- Carracedo J, Buendia P, Merino A, Madueno JA, Peralbo E, Ortiz A, Martin-Malo A, Aljama P, Rodriguez M & Ramirez R 2012 Klotho modulates the stress response in human senescent endothelial cells. *Mechanisms of Ageing and Development* **133** 647–654. (doi:10.1016/j.mad.2012.09.002)
- Cha SK, Ortega B, Kurosu H, Rosenblatt KP, Kuro OM & Huang CL 2008 Removal of sialic acid involving Klotho causes cell-surface retention of TRPV5 channel via binding to galectin-1. *PNAS* **105** 9805–9810. (doi:10.1073/pnas.0803223105)
- Chang Q, Hoefs S, van der Kemp AW, Topala CN, Bindels RJ & Hoenderop JG 2005 The β -glucuronidase Klotho hydrolyzes and activates the TRPV5 channel. *Science* **310** 490–493. (doi:10.1126/science.1114245)
- Chen CD, Podvin S, Gillespie E, Leeman SE & Abraham CR 2007 Insulin stimulates the cleavage and release of the extracellular domain of Klotho by ADAM10 and ADAM17. *PNAS* **104** 19796–19801. (doi:10.1073/pnas.0709805104)
- Chen B, Wang X, Zhao W & Wu J 2010 Klotho inhibits growth and promotes apoptosis in human lung cancer cell line A549. *Journal of Experimental & Clinical Cancer Research* **29** 99. (doi:10.1186/1756-9966-29-99)
- Chen B, Ma X, Liu S, Zhao W & Wu J 2012 Inhibition of lung cancer cells growth, motility and induction of apoptosis by Klotho, a novel secreted Wnt antagonist, in a dose-dependent manner. *Cancer Biology & Therapy* **13** 1221–1228. (doi:10.4161/cbt.21420)
- Chen CD, Sloane JA, Li H, Aytan N, Giannaris EL, Zeldich E, Hinman JD, Dedeoglu A, Rosene DL, Bansal R *et al.* 2013a The antiaging protein Klotho enhances oligodendrocyte maturation and myelination of the CNS. *Journal of Neuroscience* **33** 1927–1939. (doi:10.1523/JNEUROSCI.2080-12.2013)
- Chen TH, Kuro OM, Chen CH, Sue YM, Chen YC, Wu HH & Cheng CY 2013b The secreted Klotho protein restores phosphate retention and suppresses accelerated aging in Klotho mutant mice. *European Journal of Pharmacology* **698** 67–73. (doi:10.1016/j.ejphar.2012.09.032)
- Chihara Y, Rakugi H, Ishikawa K, Ikushima M, Maekawa Y, Ohta J, Kida I & Ogihara T 2006 Klotho protein promotes adipocyte differentiation. *Endocrinology* **147** 3835–3842. (doi:10.1210/en.2005.1529)
- Clemmons DR 2011 Consensus statement on the standardization and evaluation of growth hormone and insulin-like growth factor assays. *Clinical Chemistry* **57** 555–559. (doi:10.1373/clinchem.2010.150631)
- Corvilain J, Abramow M & Bergans A 1962 Some effects of human growth hormone on renal hemodynamics and on tubular phosphate transport in man. *Journal of Clinical Investigation* **41** 1230–1235. (doi:10.1172/JCI104584)
- Crasto CL, Semba RD, Sun K, Cappola AR, Bandinelli S & Ferrucci L 2012 Relationship of low-circulating “anti-aging” Klotho hormone with disability in activities of daily living among older community-dwelling adults. *Rejuvenation Research* **15** 295–301. (doi:10.1089/rej.2011.1268)
- Daughaday WH, Hall K, Raben MS, Salmon WD Jr, van den Brande JL & van Wyk JJ 1972 Somatomedin: proposed designation for sulphation factor. *Nature* **235** 107. (doi:10.1038/235107a0)
- Deary IJ, Harris SE, Fox HC, Hayward C, Wright AF, Starr JM & Whalley LJ 2005 KLOTHO genotype and cognitive ability in childhood and old age in the same individuals. *Neuroscience Letters* **378** 22–27. (doi:10.1016/j.neulet.2004.12.005)
- Devaraj S, Syed B, Chien A & Jialal I 2012 Validation of an immunoassay for soluble Klotho protein: decreased levels in diabetes and increased levels in chronic kidney disease. *American Journal of Clinical Pathology* **137** 479–485. (doi:10.1309/AJCPGPMF7SFRBO4)
- Dhingra R, Sullivan LM, Fox CS, Wang TJ, D'Agostino RB Sr, Gaziano JM & Vasan RS 2007 Relations of serum phosphorus and calcium levels to the incidence of cardiovascular disease in the community. *Archives of Internal Medicine* **167** 879–885. (doi:10.1001/archinte.167.9.879)
- Di Marco GS, König M, Stock C, Wiesinger A, Hillebrand U, Reiermann S, Reuter S, Amler S, Kohler G, Buck F *et al.* 2013 High phosphate directly affects endothelial function by downregulating annexin II. *Kidney International* **83** 213–222. (doi:10.1038/ki.2012.300)
- Doi S, Zou Y, Togao O, Pastor JV, John GB, Wang L, Shiizaki K, Gotschall R, Schiavi S, Yorioka N *et al.* 2011 Klotho inhibits transforming growth factor- β 1 (TGF- β 1) signaling and suppresses renal fibrosis and cancer metastasis in mice. *Journal of Biological Chemistry* **286** 8655–8665. (doi:10.1074/jbc.M110.174037)
- Dulak NC & Temin HM 1973 A partially purified polypeptide fraction from rat liver cell conditioned medium with multiplication-stimulating activity for embryo fibroblasts. *Journal of Cellular Physiology* **81** 153–160. (doi:10.1002/jcp.1040810204)
- Ellam TJ & Chico TJ 2012 Phosphate: the new cholesterol? The role of the phosphate axis in non-uremic vascular disease *Atherosclerosis* **220** 310–318. (doi:10.1016/j.atherosclerosis.2011.09.002)
- Evans LM, Davies JS, Anderson RA, Ellis GR, Jackson SK, Lewis MJ, Frenneaux MP, Rees A & Scanlon MF 2000 The effect of GH replacement therapy on endothelial function and oxidative stress in adult growth hormone deficiency. *European Journal of Endocrinology* **142** 254–262. (doi:10.1530/eje.0.1420254)
- Faul C, Amaral AP, Oskouei B, Hu MC, Sloan A, Isakova T, Gutierrez OM, Aguillon-Prada R, Lincoln J, Hare JM *et al.* 2011 FGF23 induces left ventricular hypertrophy. *Journal of Clinical Investigation* **121** 4393–4408. (doi:10.1172/JCI46122)
- Foley RN 2009 Phosphate levels and cardiovascular disease in the general population. *Clinical Journal of the American Society of Nephrology* **4** 1136–1139. (doi:10.2215/CJN.01660309)
- Froesch ER, Buergi H, Ramseier EB, Bally P & Labhart A 1963 Antibody-suppressible and nonsuppressible insulin-like activities in human serum and their physiologic significance. an insulin assay with adipose tissue of increased precision and specificity. *Journal of Clinical Investigation* **42** 1816–1834. (doi:10.1172/JCI104866)
- Galitzer H, Ben-Dov IZ, Silver J & Naveh-Manny T 2010 Parathyroid cell resistance to fibroblast growth factor 23 in secondary hyperparathyroidism of chronic kidney disease. *Kidney International* **77** 211–218. (doi:10.1038/ki.2009.464)
- Gattineni J, Bates C, Twombly K, Dwarakanath V, Robinson ML, Goetz R, Mohammadi M & Baum M 2009 FGF23 decreases renal NaPi-2a and NaPi-2c expression and induces hypophosphatemia in vivo predominantly via FGF receptor 1. *American Journal of Physiology. Renal Physiology* **297** F282–F291. (doi:10.1152/ajprenal.90742.2008)
- German DC, Khobahy I, Pastor J, Kuro OM & Liu X 2012 Nuclear localization of Klotho in brain: an anti-aging protein. *Neurobiology of Aging* **33** e1425–e1430. (doi:10.1016/j.neurobiolaging.2011.12.018)
- Giustina A, Chanson P, Bronstein MD, Klibanski A, Lamberts S, Casanueva FF, Trainer P, Ghigo E, Ho K & Melmed S 2010 A consensus on criteria for cure of acromegaly. *Journal of Clinical Endocrinology and Metabolism* **95** 3141–3148. (doi:10.1210/jc.2009-2670)
- Godowski PJ, Leung DW, Meacham LR, Galgani JP, Hellmiss R, Keret R, Rotwein PS, Parks JS, Laron Z & Wood WI 1989 Characterization of the

- human growth hormone receptor gene and demonstration of a partial gene deletion in two patients with Laron-type dwarfism. *PNAS* **86** 8083–8087. (doi:10.1073/pnas.86.20.8083)
- Guler HP, Zapf J, Schmid C & Froesch ER 1989 Insulin-like growth factors I and II in healthy man. Estimations of half-lives and production rates. *Acta Endocrinologica* **121** 753–758. (doi:10.1530/acta.0.1210753)
- Hesse M, Frohlich LF, Zeitl U, Lanske B & Erben RG 2007 Ablation of vitamin D signaling rescues bone, mineral, and glucose homeostasis in Fgf-23 deficient mice. *Matrix Biology* **26** 75–84. (doi:10.1016/j.matbio.2006.10.003)
- Hu MC, Shi M, Zhang J, Pastor J, Nakatani T, Lanske B, Razzaque MS, Rosenblatt KP, Baum MG, Kuro-o M *et al.* 2010 Klotho: a novel phosphaturic substance acting as an autocrine enzyme in the renal proximal tubule. *FASEB Journal* **24** 3438–3450. (doi:10.1096/fj.10-154765)
- Hu MC, Shi M, Zhang J, Quinones H, Griffith C, Kuro-o M & Moe OW 2011 Klotho deficiency causes vascular calcification in chronic kidney disease. *Journal of the American Society of Nephrology* **22** 124–136. (doi:10.1681/ASN.2009121311)
- Hu MC, Shiizaki K, Kuro-o M & Moe OW 2013 Fibroblast growth factor 23 and Klotho: physiology and pathophysiology of an endocrine network of mineral metabolism. *Annual Review of Physiology* **75** 503–533. (doi:10.1146/annurev-physiol-030212-183727)
- Huang CL 2012 Regulation of ion channels by secreted Klotho. *Advances in Experimental Medicine and Biology* **728** 100–106. (doi:10.1007/978-1-4614-0887-1_7)
- Ichikawa S, Imel EA, Kreiter ML, Yu X, Mackenzie DS, Sorenson AH, Goetz R, Mohammadi M, White KE & Econs MJ 2007 A homozygous missense mutation in human KLOTHO causes severe tumoral calcinosis. *Journal of Clinical Investigation* **117** 2684–2691. (doi:10.1172/JCI31330)
- Ikkos D, Ljunggren H & Luft R 1956 Glomerular filtration rate and renal plasma flow in acromegaly. *Acta Endocrinologica* **21** 226–236. (doi:10.1530/acta.0.0210226)
- Ikushima M, Rakugi H, Ishikawa K, Maekawa Y, Yamamoto K, Ohta J, Chihara Y, Kida I & Ogihara T 2006 Anti-apoptotic and anti-senescence effects of Klotho on vascular endothelial cells. *Biochemical and Biophysical Research Communications* **339** 827–832. (doi:10.1016/j.bbrc.2005.11.094)
- Imura A, Iwano A, Tohyama O, Tsuji Y, Nozaki K, Hashimoto N, Fujimori T & Nabeshima Y 2004 Secreted Klotho protein in sera and CSF: implication for post-translational cleavage in release of Klotho protein from cell membrane. *FEBS Letters* **565** 143–147. (doi:10.1016/j.febslet.2004.03.090)
- Imura A, Tsuji Y, Murata M, Maeda R, Kubota K, Iwano A, Obuse C, Togashi K, Tominaga M, Kita N *et al.* 2007 α -Klotho as a regulator of calcium homeostasis. *Science* **316** 1615–1618. (doi:10.1126/science.1135901)
- Ito S, Kinoshita S, Shiraiishi N, Nakagawa S, Sekine S, Fujimori T & Nabeshima YI 2000 Molecular cloning and expression analyses of mouse β Klotho, which encodes a novel Klotho family protein. *Mechanisms of Development* **98** 115–119. (doi:10.1016/S0925-4773(00)00439-1)
- Ito S, Fujimori T, Hayashizaki Y & Nabeshima Y 2002 Identification of a novel mouse membrane-bound family 1 glycosidase-like protein, which carries an atypical active site structure. *Biochimica et Biophysica Acta* **1576** 341–345. (doi:10.1016/S0167-4781(02)00281-6)
- Ito N, Fukumoto S, Taguchi M, Takeshita A, Takeuchi Y, Yamada S & Fujita T 2007 Fibroblast growth factor (FGF)23 in patients with acromegaly. *Endocrine Journal* **54** 481–484. (doi:10.1507/endocrj.K06-217)
- Itoh N & Ornitz DM 2011 Fibroblast growth factors: from molecular evolution to roles in development, metabolism and disease. *Journal of Biochemistry* **149** 121–130. (doi:10.1093/jb/mvq121)
- Junnala RK, List EO, Berryman DE, Murrey JW & Kopchick JJ 2013 The GH/IGF-1 axis in ageing and longevity. *Nature Reviews. Endocrinology* **9** 366–376. (doi:10.1038/nrendo.2013.67)
- Kamenicky P, Blanchard A, Gauci C, Salenave S, Letierce A, Lombes M, Brailly-Tabard S, Azizi M, Prie D, Souberbielle JC *et al.* 2012 Pathophysiology of renal calcium handling in acromegaly: what lies behind hypercalciuria? *Journal of Clinical Endocrinology and Metabolism* **97** 2124–2133. (doi:10.1210/jc.2011-3188)
- Kashimada K, Yamashita T, Tsuji K, Nifuji A, Mizutani S, Nabeshima Y & Noda M 2002 Defects in growth and bone metabolism in Klotho mutant mice are resistant to GH treatment. *Journal of Endocrinology* **174** 403–410. (doi:10.1677/joe.0.1740403)
- Kato Y, Arakawa E, Kinoshita S, Shirai A, Furuya A, Yamano K, Nakamura K, Iida A, Anazawa H, Koh N *et al.* 2000 Establishment of the anti-Klotho monoclonal antibodies and detection of Klotho protein in kidneys. *Biochemical and Biophysical Research Communications* **267** 597–602. (doi:10.1006/bbrc.1999.2009)
- Kawai M, Kinoshita S, Kimoto A, Hasegawa Y, Miyagawa K, Yamazaki M, Ohata Y, Ozono K & Michigami T 2013 FGF23 suppresses chondrocyte proliferation in the presence of soluble α -Klotho both *in vitro* and *in vivo*. *Journal of Biological Chemistry* **288** 2414–2427. (doi:10.1074/jbc.M112.410043)
- Khosravi A, Cutler CM, Kelly MH, Chang R, Royal RE, Sherry RM, Wodajo FM, Fedarko NS & Collins MT 2007 Determination of the elimination half-life of fibroblast growth factor-23. *Journal of Clinical Endocrinology and Metabolism* **92** 2374–2377. (doi:10.1210/jc.2006-2865)
- Kim HR, Nam BY, Kim DW, Kang MW, Han JH, Lee MJ, Shin DH, Doh FM, Koo HM, Ko KI *et al.* 2013 Circulating α -Klotho levels in CKD and relationship to progression. *American Journal of Kidney Diseases* **61** 899–909. (doi:10.1053/ajkd.2013.01.024)
- Koh N, Fujimori T, Nishiguchi S, Tamori A, Shiomi S, Nakatani T, Sugimura K, Kishimoto T, Kinoshita S, Kuroki T *et al.* 2001 Severely reduced production of Klotho in human chronic renal failure kidney. *Biochemical and Biophysical Research Communications* **280** 1015–1020. (doi:10.1006/bbrc.2000.4226)
- Kohler S, Tschopp O, Sze L, Neidert M, Bernays RL, Spanaus KS, Wiesli P & Schmid C 2013 Monitoring for potential residual disease activity by serum insulin-like growth factor 1 and soluble Klotho in patients with acromegaly after pituitary surgery: is there an impact of the genomic deletion of exon 3 in the growth hormone receptor (d3-GHR) gene on “safe” GH cut-off values? *General and Comparative Endocrinology* **188** 282–287. (doi:10.1016/j.ygcen.2013.04.024)
- Komaba H, Goto S, Fujii H, Hamada Y, Kobayashi A, Shibuya K, Tominaga Y, Otsuki N, Nibu K, Nakagawa K *et al.* 2010 Depressed expression of Klotho and FGF receptor 1 in hyperplastic parathyroid glands from uremic patients. *Kidney International* **77** 232–238. (doi:10.1038/ki.2009.414)
- Krajisnik T, Bjorklund P, Marsel R, Ljunggren O, Akerstrom G, Jonsson KB, Westin G & Larsson TE 2007 Fibroblast growth factor-23 regulates parathyroid hormone and 1 α -hydroxylase expression in cultured bovine parathyroid cells. *Journal of Endocrinology* **195** 125–131. (doi:10.1677/JOE-07-0267)
- Krajisnik T, Olsson H, Mirza MA, Hellman P, Akerstrom G, Westin G, Larsson TE & Bjorklund P 2010 Parathyroid Klotho and FGF-receptor 1 expression decline with renal function in hyperparathyroid patients with chronic kidney disease and kidney transplant recipients. *Kidney International* **78** 1024–1032. (doi:10.1038/ki.2010.260)
- Kuro-o M 2010 Klotho. *Pflügers Archiv* **459** 333–343. (doi:10.1007/s00424-009-0722-7)
- Kuro-o M, Matsumura Y, Aizawa H, Kawaguchi H, Suga T, Utsugi T, Ohya Y, Kurabayashi M, Kaname T, Kume E *et al.* 1997 Mutation of the mouse klotho gene leads to a syndrome resembling ageing. *Nature* **390** 45–51. (doi:10.1038/36285)
- Kurosu H, Yamamoto M, Clark JD, Pastor JV, Nandi A, Gurnani P, McGuinness OP, Chikuda H, Yamaguchi M, Kawaguchi H *et al.* 2005 Suppression of aging in mice by the hormone Klotho. *Science* **309** 1829–1833. (doi:10.1126/science.1112766)

- Kurosu H, Ogawa Y, Miyoshi M, Yamamoto M, Nandi A, Rosenblatt KP, Baum MG, Schiavi S, Hu MC, Moe OW *et al.* 2006 Regulation of fibroblast growth factor-23 signaling by Klotho. *Journal of Biological Chemistry* **281** 6120–6123. (doi:10.1074/jbc.C500457200)
- Kusaba T, Okigaki M, Matui A, Murakami M, Ishikawa K, Kimura T, Sonomura K, Adachi Y, Shibuya M, Shirayama T *et al.* 2010 Klotho is associated with VEGF receptor-2 and the transient receptor potential canonical-1 Ca^{2+} channel to maintain endothelial integrity. *PNAS* **107** 19308–19313. (doi:10.1073/pnas.1008544107)
- Laron Z, Anin S, Klipper-Aurbach Y & Klinger B 1992 Effects of insulin-like growth factor on linear growth, head circumference, and body fat in patients with Laron-type dwarfism. *Lancet* **339** 1258–1261. (doi:10.1016/0140-6736(92)91594-X)
- Larsson T, Marsell R, Schipani E, Ohlsson C, Ljunggren O, Tenenhouse HS, Juppner H & Jonsson KB 2004 Transgenic mice expressing fibroblast growth factor 23 under the control of the $\alpha 1(\text{I})$ collagen promoter exhibit growth retardation, osteomalacia, and disturbed phosphate homeostasis. *Endocrinology* **145** 3087–3094. (doi:10.1210/en.2003-1768)
- Lau WL, Pai A, Moe SM & Giachelli CM 2011 Direct effects of phosphate on vascular cell function. *Advances in Chronic Kidney Disease* **18** 105–112. (doi:10.1053/j.ackd.2010.12.002)
- Lau WL, Leaf EM, Hu MC, Takeno MM, Kuro-o M, Moe OW & Giachelli CM 2012 Vitamin D receptor agonists increase Klotho and osteopontin while decreasing aortic calcification in mice with chronic kidney disease fed a high phosphate diet. *Kidney International* **82** 1261–1270. (doi:10.1038/ki.2012.322)
- Li SA, Watanabe M, Yamada H, Nagai A, Kinuta M & Takei K 2004 Immunohistochemical localization of Klotho protein in brain, kidney, and reproductive organs of mice. *Cell Structure and Function* **29** 91–99. (doi:10.1247/csf.29.91)
- Lindberg K, Olauson H, Amin R, Ponnusamy A, Goetz R, Taylor RF, Mohammadi M, Canfield A, Kublickiene K & Larsson TE 2013 Arterial Klotho expression and FGF23 effects on vascular calcification and function. *PLoS ONE* **8** e60658. (doi:10.1371/journal.pone.0060658)
- Liu H, Fergusson MM, Castilho RM, Liu J, Cao L, Chen J, Malide D, Rovira II, Schimel D, Kuo CJ *et al.* 2007 Augmented Wnt signaling in a mammalian model of accelerated aging. *Science* **317** 803–806. (doi:10.1126/science.1143578)
- Lorenzi O, Veyrat-Durebex C, Wollheim CB, Villemin P, Rohner-Jeanrenaud F, Zanchi A & Vischer UM 2010 Evidence against a direct role of Klotho in insulin resistance. *Pflügers Archiv* **459** 465–473. (doi:10.1007/s00424-009-0735-2)
- Lund B, Eskildsen PC, Norman AW & Sorensen OH 1981 Calcium and vitamin D metabolism in acromegaly. *Acta Endocrinologica* **96** 444–450. (doi:10.1530/acta.0.0960444)
- Maekawa Y, Ohishi M, Ikushima M, Yamamoto K, Yasuda O, Oguro R, Yamamoto-Hanasaki H, Tatara Y, Takeya Y & Rakugi H 2011 Klotho protein diminishes endothelial apoptosis and senescence via a mitogen-activated kinase pathway. *Geriatrics & Gerontology International* **11** 510–516. (doi:10.1111/j.1447-0594.2011.00699.x)
- Majumdar V, Nagaraja D & Christopher R 2010 Association of the functional KL-VS variant of Klotho gene with early-onset ischemic stroke. *Biochemical and Biophysical Research Communications* **403** 412–416. (doi:10.1016/j.bbrc.2010.11.045)
- Mangos S, Amaral AP, Faul C, Juppner H, Reiser J & Wolf M 2012 Expression of fgf23 and α Klotho in developing embryonic tissues and adult kidney of the zebrafish, *Danio rerio*. *Nephrology, Dialysis, Transplantation* **27** 4314–4322. (doi:10.1093/ndt/gfs335)
- Manya H, Akasaka-Manya K & Endo T 2010 Klotho protein deficiency and aging. *Geriatrics & Gerontology International* **10**(Suppl 1) S80–S87. (doi:10.1111/j.1447-0594.2010.00596.x)
- Martin A, David V & Quarles LD 2012 Regulation and function of the FGF23/Klotho endocrine pathways. *Physiological Reviews* **92** 131–155. (doi:10.1152/physrev.00002.2011)
- Matsumura Y, Aizawa H, Shiraki-Iida T, Nagai R, Kuro-o M & Nabeshima Y 1998 Identification of the human klotho gene and its two transcripts encoding membrane and secreted Klotho protein. *Biochemical and Biophysical Research Communications* **242** 626–630. (doi:10.1006/bbrc.1997.8019)
- Mitani H, Ishizaka N, Aizawa T, Ohno M, Usui S, Suzuki T, Amaki T, Mori I, Nakamura Y, Sato M *et al.* 2002 *In vivo* klotho gene transfer ameliorates angiotensin II-induced renal damage. *Hypertension* **39** 838–843. (doi:10.1161/01.HYP.0000013734.33441.EA)
- Mori K, Yahata K, Mukoyama M, Suganami T, Makino H, Nagae T, Masuzaki H, Ogawa Y, Sugawara A, Nabeshima Y *et al.* 2000 Disruption of klotho gene causes an abnormal energy homeostasis in mice. *Biochemical and Biophysical Research Communications* **278** 665–670. (doi:10.1006/bbrc.2000.3864)
- Morishita K, Shirai A, Kubota M, Katakura Y, Nabeshima Y, Takeshige K & Kamiya T 2001 The progression of aging in klotho mutant mice can be modified by dietary phosphorus and zinc. *Journal of Nutrition* **131** 3182–3188.
- Nabeshima Y & Imura H 2008 α -Klotho: a regulator that integrates calcium homeostasis. *American Journal of Nephrology* **28** 455–464. (doi:10.1159/000112824)
- Nagai T, Yamada K, Kim HC, Kim YS, Noda Y, Imura A, Nabeshima Y & Nabeshima T 2003 Cognition impairment in the genetic model of aging klotho gene mutant mice: a role of oxidative stress. *FASEB Journal* **17** 50–52. (doi:10.1096/fj.02-0448fje)
- Nagasu H, Satoh M, Kuwabara A, Yorimitsu D, Kidokoro K, Nishi Y, Tomita N, Sasaki T & Kashihara N 2011 Overexpression of Klotho protein modulates uninephrectomy-induced compensatory renal hypertrophy by suppressing IGF-I signals. *Biochemical and Biophysical Research Communications* **407** 39–43. (doi:10.1016/j.bbrc.2011.02.089)
- Nakatani T, Sarraj B, Ohnishi M, Densmore MJ, Taguchi T, Goetz R, Mohammadi M, Lanske B & Razzaque MS 2009 *In vivo* genetic evidence for Klotho-dependent, fibroblast growth factor 23 (Fgf23)-mediated regulation of systemic phosphate homeostasis. *FASEB Journal* **23** 433–441. (doi:10.1096/fj.08-114397)
- Neider MC, Sze L, Zwimpfer C, Samthein J, Seifert B, Frei K, Leske H, Rushing EJ, Schmid C & Bernays RL 2013 Soluble α -Klotho: a novel serum biomarker for the activity of GH-producing pituitary adenomas. *European Journal of Endocrinology* **168** 575–583. (doi:10.1530/EJE-12-1045)
- Ohata Y, Arahori H, Namba N, Kitaoka T, Hirai H, Wada K, Nakayama M, Michigami T, Imura A, Nabeshima Y *et al.* 2011 Circulating levels of soluble α -Klotho are markedly elevated in human umbilical cord blood. *Journal of Clinical Endocrinology and Metabolism* **96** E943–E947. (doi:10.1210/jc.2010-2357)
- Ohnishi M, Nakatani T, Lanske B & Razzaque MS 2009 Reversal of mineral ion homeostasis and soft-tissue calcification of klotho knockout mice by deletion of vitamin D 1α -hydroxylase. *Kidney International* **75** 1166–1172. (doi:10.1038/ki.2009.24)
- Ohyama Y, Kurabayashi M, Masuda H, Nakamura T, Aihara Y, Kaname T, Suga T, Arai M, Aizawa H, Matsumura Y *et al.* 1998 Molecular cloning of rat klotho cDNA: markedly decreased expression of klotho by acute inflammatory stress. *Biochemical and Biophysical Research Communications* **251** 920–925. (doi:10.1006/bbrc.1998.9576)
- Olauson H, Lindberg K, Amin R, Jia T, Wernerson A, Andersson G & Larsson TE 2012 Targeted deletion of Klotho in kidney distal tubule disrupts mineral metabolism. *Journal of the American Society of Nephrology* **23** 1641–1651. (doi:10.1681/ASN.2012010048)
- Osuka S & Razzaque MS 2012 Can features of phosphate toxicity appear in normophosphatemia? *Journal of Bone and Mineral Metabolism* **30** 10–18. (doi:10.1007/s00774-011-0343-z)
- Park SJ, Park SH, Chang JW, Choi J, Jung HH & Im GJ 2012 Protective effect of klotho protein against cisplatin ototoxicity in an auditory cell line. *Journal of Laryngology and Otology* **126** 1003–1009. (doi:10.1017/S0022215112001715)
- Park SJ, Shin EJ, Min SS, An J, Li Z, Hee Chung Y, Hoon Jeong J, Bach JH, Nah SY, Kim WK *et al.* 2013 Inactivation of JAK2/STAT3 signaling axis and downregulation of M1 mAChR cause cognitive impairment

- in klotho mutant mice, a genetic model of aging. *Neuropsychopharmacology* **38** 1426–1437. (doi:10.1038/npp.2013.39)
- Pavik I, Jaeger P, Ebner L, Poster D, Krauer F, Kistler AD, Rentsch K, Andreisek G, Wagner CA, Devuyst O *et al.* 2012 Soluble klotho and autosomal dominant polycystic kidney disease. *Clinical Journal of the American Society of Nephrology* **7** 248–257. (doi:10.2215/CJN.09020911)
- Pavik I, Jaeger P, Ebner L, Wagner CA, Petzold K, Spichtig D, Poster D, Wuthrich RP, Russmann S & Serra AL 2013 Secreted Klotho and FGF23 in chronic kidney disease stage 1 to 5: a sequence suggested from a cross-sectional study. *Nephrology, Dialysis, Transplantation* **28** 352–359. (doi:10.1093/ndt/gfs460)
- Pedersen L, Pedersen SM, Brasen CL & Rasmussen LM 2013 Soluble serum Klotho levels in healthy subjects. Comparison of two different immunoassays. *Clinical Biochemistry* **46** 1079–1083. (doi:10.1016/j.clinbiochem.2013.05.046)
- Pierson RW Jr & Temin HM 1972 The partial purification from calf serum of a fraction with multiplication-stimulating activity for chicken fibroblasts in cell culture and with non-suppressible insulin-like activity. *Journal of Cellular Physiology* **79** 319–330. (doi:10.1002/jcp.1040790302)
- Polanska UM, Edwards E, Fernig DG & Kinnunen TK 2011 The cooperation of FGF receptor and Klotho is involved in excretory canal development and regulation of metabolic homeostasis in *Caenorhabditis elegans*. *Journal of Biological Chemistry* **286** 5657–5666. (doi:10.1074/jbc.M110.173039)
- Prader A, Illig R, Uehlinger E & Stalder G 1959 Rachitis infolge Knochentumors [Rickets following bone tumor]. *Helvetica Paediatrica Acta* **14** 554–565.
- Quigley R & Baum M 1991 Effects of growth hormone and insulin-like growth factor I on rabbit proximal convoluted tubule transport. *Journal of Clinical Investigation* **88** 368–374. (doi:10.1172/JCI115312)
- Rakugi H, Matsukawa N, Ishikawa K, Yang J, Imai M, Ikushima M, Maekawa Y, Kida I, Miyazaki J & Ogihara T 2007 Anti-oxidative effect of Klotho on endothelial cells through cAMP activation. *Endocrine* **31** 82–87. (doi:10.1007/s12020-007-0016-9)
- Razzaque MS 2012 The role of Klotho in energy metabolism. *Nature Reviews. Endocrinology* **8** 579–587. (doi:10.1038/nrendo.2012.75)
- Rinderknecht E & Humbel RE 1978 The amino acid sequence of human insulin-like growth factor I and its structural homology with proinsulin. *Journal of Biological Chemistry* **253** 2769–2776.
- Rutishauser J, Schmid C, Hauri C, Froesch ER & Zapf J 1993 Growth hormone, but not insulin-like growth factor I, induces a serum protease activity for insulin-like growth factor binding protein-3 in hypophysectomized rats *in vivo*. *FEBS Letters* **334** 23–26. (doi:10.1016/0014-5793(93)81672-M)
- Saftig P & Reiss K 2011 The “A Disintegrin And Metalloproteases” ADAM10 and ADAM17: novel drug targets with therapeutic potential? *European Journal of Cell Biology* **90** 527–535. (doi:10.1016/j.ejcb.2010.11.005)
- Saito Y, Yamagishi T, Nakamura T, Ohyama Y, Aizawa H, Suga T, Matsumura Y, Masuda H, Kurabayashi M, Kuro-o M *et al.* 1998 Klotho protein protects against endothelial dysfunction. *Biochemical and Biophysical Research Communications* **248** 324–329. (doi:10.1006/bbrc.1998.8943)
- Saito Y, Nakamura T, Ohyama Y, Suzuki T, Iida A, Shiraki-Iida T, Kuro-o M, Nabeshima Y, Kurabayashi M & Nagai R 2000 *In vivo* klotho gene delivery protects against endothelial dysfunction in multiple risk factor syndrome. *Biochemical and Biophysical Research Communications* **276** 767–772. (doi:10.1006/bbrc.2000.3470)
- Saito H, Kusano K, Kinoshita M, Ito H, Hirata M, Segawa H, Miyamoto K & Fukushima N 2003 Human fibroblast growth factor-23 mutants suppress Na⁺-dependent phosphate co-transport activity and 1 α ,25-dihydroxyvitamin D3 production. *Journal of Biological Chemistry* **278** 2206–2211. (doi:10.1074/jbc.M207872200)
- Salmon WD Jr & Daughaday WH 1957 A hormonally controlled serum factor which stimulates sulfate incorporation by cartilage *in vitro*. *Journal of Laboratory and Clinical Medicine* **49** 825–836.
- Sathyanesan M, Girgenti MJ, Banasr M, Stone K, Bruce C, Guilchick E, Wilczak-Havill K, Nairn A, Williams K, Sass S *et al.* 2012 A molecular characterization of the choroid plexus and stress-induced gene regulation. *Translational Psychiatry* **2** e139. (doi:10.1038/tp.2012.64)
- Satoh M, Nagasu H, Morita Y, Yamaguchi TP, Kanwar YS & Kashihara N 2012 Klotho protects against mouse renal fibrosis by inhibiting Wnt signaling. *American Journal of Physiology. Renal Physiology* **303** F1641–F1651. (doi:10.1152/ajprenal.00460.2012)
- Schmid C & Meili E 2000 Acromegaly in a hemophilic patient infected with the hepatitis C and human immunodeficiency viruses. *American Journal of Medicine* **108** 757–758. (doi:10.1016/S0002-9343(00)00421-6)
- Schmid C, Rutishauser J, Schlapfer I, Froesch ER & Zapf J 1991 Intact but not truncated insulin-like growth factor binding protein-3 (IGFBP-3) blocks IGF I-induced stimulation of osteoblasts: control of IGF signalling to bone cells by IGFBP-3-specific proteolysis? *Biochemical and Biophysical Research Communications* **179** 579–585. (doi:10.1016/0006-291X(91)91410-E)
- Schoenle E, Zapf J, Humbel RE & Froesch ER 1982 Insulin-like growth factor I stimulates growth in hypophysectomized rats. *Nature* **296** 252–253. (doi:10.1038/296252a0)
- Scialla JJ, Lau WL, Reilly MP, Isakova T, Yang HY, Crouthamel MH, Chavkin NW, Rahman M, Wahl P, Amaral AP *et al.* 2013 Fibroblast growth factor 23 is not associated with and does not induce arterial calcification. *Kidney International* **83** 1159–1168. (doi:10.1038/ki.2013.3)
- Segawa H, Yamanaka S, Ohno Y, Onitsuka A, Shiozawa K, Aranami F, Furutani J, Tomoe Y, Ito M, Kuwahata M *et al.* 2007 Correlation between hyperphosphatemia and type II Na-Pi cotransporter activity in klotho mice. *American Journal of Physiology. Renal Physiology* **292** F769–F779. (doi:10.1152/ajprenal.00248.2006)
- Seiler S, Wen M, Roth HJ, Fehrenz M, Flugge F, Herath E, Weihrauch A, Fliser D & Heine GH 2013 Plasma Klotho is not related to kidney function and does not predict adverse outcome in patients with chronic kidney disease. *Kidney International* **83** 121–128. (doi:10.1038/ki.2012.288)
- Semba RD, Cappola AR, Sun K, Bandinelli S, Dalal M, Crasto C, Guralnik JM & Ferrucci L 2011 Plasma klotho and mortality risk in older community-dwelling adults. *Journals of Gerontology. Series A, Biological Sciences and Medical Sciences* **66** 794–800. (doi:10.1093/gerona/glr058)
- Semba RD, Cappola AR, Sun K, Bandinelli S, Dalal M, Crasto C, Guralnik JM & Ferrucci L 2012 Relationship of low plasma klotho with poor grip strength in older community-dwelling adults: the INCHIANTI study. *European Journal of Applied Physiology* **112** 1215–1220. (doi:10.1007/s00421-011-2072-3)
- Shalhoub V, Ward SC, Sun B, Stevens J, Renshaw L, Hawkins N & Richards WG 2011 Fibroblast growth factor 23 (FGF23) and α -Klotho stimulate osteoblastic MC3T3.E1 cell proliferation and inhibit mineralization. *Calcified Tissue International* **89** 140–150. (doi:10.1007/s00223-011-9501-5)
- Shimada T, Mizutani S, Muto T, Yoneya T, Hino R, Takeda S, Takeuchi Y, Fujita T, Fukumoto S & Yamashita T 2001 Cloning and characterization of FGF23 as a causative factor of tumor-induced osteomalacia. *PNAS* **98** 6500–6505. (doi:10.1073/pnas.101545198)
- Shimada T, Hasegawa H, Yamazaki Y, Muto T, Hino R, Takeuchi Y, Fujita T, Nakahara K, Fukumoto S & Yamashita T 2004a FGF-23 is a potent regulator of vitamin D metabolism and phosphate homeostasis. *Journal of Bone and Mineral Research* **19** 429–435. (doi:10.1359/JBMR.0301264)
- Shimada T, Kakitani M, Yamazaki Y, Hasegawa H, Takeuchi Y, Fujita T, Fukumoto S, Tomizuka K & Yamashita T 2004b Targeted ablation of Fgf23 demonstrates an essential physiological role of FGF23 in phosphate and vitamin D metabolism. *Journal of Clinical Investigation* **113** 561–568. (doi:10.1172/JCI200419081)
- Shimada T, Urakawa I, Yamazaki Y, Hasegawa H, Hino R, Yoneya T, Takeuchi Y, Fujita T, Fukumoto S & Yamashita T 2004c FGF-23 transgenic mice demonstrate hypophosphatemic rickets with reduced expression of sodium phosphate cotransporter type IIa. *Biochemical and*

- Biophysical Research Communications* **314** 409–414. (doi:10.1016/j.bbrc.2003.12.102)
- Shiozaki M, Yoshimura K, Shibata M, Koike M, Matsuura N, Uchiyama Y & Gotow T 2008 Morphological and biochemical signs of age-related neurodegenerative changes in klotho mutant mice. *Neuroscience* **152** 924–941. (doi:10.1016/j.neuroscience.2008.01.032)
- Shiraki-Iida T, Aizawa H, Matsumura Y, Sekine S, Iida A, Anazawa H, Nagai R, Kuro-o M & Nabeshima Y 1998 Structure of the mouse klotho gene and its two transcripts encoding membrane and secreted protein. *FEBS Letters* **424** 6–10. (doi:10.1016/S0014-5793(98)00127-6)
- Shiraki-Iida T, Iida A, Nabeshima Y, Anazawa H, Nishikawa S, Noda M & Kuro-o M 2000 Improvement of multiple pathophysiological phenotypes of klotho (kl/kl) mice by adenovirus-mediated expression of the klotho gene. *Journal of Gene Medicine* **2** 233–242. (doi:10.1002/1521-2254(200007/08)2:4<233::AID-JGM110>3.0.CO;2-5)
- Shuto E, Taketani Y, Tanaka R, Harada N, Isshiki M, Sato M, Nashiki K, Amo K, Yamamoto H, Higashi Y *et al.* 2009 Dietary phosphorus acutely impairs endothelial function. *Journal of the American Society of Nephrology* **20** 1504–1512. (doi:10.1681/ASN.2008101106)
- Siahanidou T, Garatzioti M, Lazaropoulou C, Kourlaba G, Papassotiropoulos I, Kino T, Imura A, Nabeshima Y & Chrousos G 2012 Plasma soluble α -Klotho protein levels in premature and term neonates: correlations with growth and metabolic parameters. *European Journal of Endocrinology* **167** 433–440. (doi:10.1530/EJE-12-0476)
- Smith RC, O'Bryan LM, Farrow EG, Summers LJ, Clinkenberg EL, Roberts JL, Cass TA, Saha J, Broderick C, Ma YL *et al.* 2012 Circulating α -Klotho influences phosphate handling by controlling FGF23 production. *Journal of Clinical Investigation* **122** 4710–4715. (doi:10.1172/JCI64986)
- Sugano Y & Lardelli M 2011 Identification and expression analysis of the zebrafish orthologue of Klotho. *Development Genes and Evolution* **221** 179–186. (doi:10.1007/s00427-011-0367-3)
- Sugiura H, Tsuchiya K & Nitta K 2011 Circulating levels of soluble α -Klotho in patients with chronic kidney disease. *Clinical and Experimental Nephrology* **15** 795–796. (doi:10.1007/s10157-011-0511-4)
- Sugiura H, Yoshida T, Shiohira S, Kohei J, Mitobe M, Kurosu H, Kuro-o M, Nitta K & Tsuchiya K 2012 Reduced Klotho expression level in kidney aggravates renal interstitial fibrosis. *American Journal of Physiology. Renal Physiology* **302** F1252–F1264. (doi:10.1152/ajprenal.00294.2011)
- Sze L, Bernays RL, Zwimpfer C, Wiesli P, Brandle M & Schmid C 2012 Excessively high soluble Klotho in patients with acromegaly. *Journal of Internal Medicine* **272** 93–97. (doi:10.1111/j.1365-2796.2012.02542.x)
- Takamoto S, Tsuchiya H, Onishi T, Morimoto S, Imanaka S, Mori S, Seino Y, Uozumi T & Kumahara Y 1985 Changes in calcium homeostasis in acromegaly treated by pituitary adenectomy. *Journal of Clinical Endocrinology and Metabolism* **61** 7–11. (doi:10.1210/jcem-61-1-7)
- Takenaka T, Watanabe Y, Inoue T, Miyazaki T & Suzuki H 2013 Fibroblast growth factor 23 enhances renal klotho abundance. *Pflügers Archiv* **465** 935–943. (doi:10.1007/s00424-013-1226-z)
- Takeshita K, Fujimori T, Kurotaki Y, Honjo H, Tsujikawa H, Yasui K, Lee JK, Kamiya K, Kitaichi K, Yamamoto K *et al.* 2004 Sinoatrial node dysfunction and early unexpected death of mice with a defect of klotho gene expression. *Circulation* **109** 1776–1782. (doi:10.1161/01.CIR.0000124224.48962.32)
- Tomiyama K, Maeda R, Urakawa I, Yamazaki Y, Tanaka T, Ito S, Nabeshima Y, Tomita T, Odori S, Hosoda K *et al.* 2010 Relevant use of Klotho in FGF19 subfamily signaling system *in vivo*. *PNAS* **107** 1666–1671. (doi:10.1073/pnas.0913986107)
- Tonelli M, Sacks F, Pfeffer M, Gao Z & Curhan G 2005 Relation between serum phosphate level and cardiovascular event rate in people with coronary disease. *Circulation* **112** 2627–2633. (doi:10.1161/CIRCULATIONAHA.105.553198)
- Tsujikawa H, Kurotaki Y, Fujimori T, Fukuda K & Nabeshima Y 2003 Klotho, a gene related to a syndrome resembling human premature aging, functions in a negative regulatory circuit of vitamin D endocrine system. *Molecular Endocrinology* **17** 2393–2403. (doi:10.1210/me.2003-0048)
- Ullrich A, Gray A, Tam AW, Yang-Feng T, Tsubokawa M, Collins C, Henzel W, Le Bon T, Kathuria S, Chen E *et al.* 1986 Insulin-like growth factor I receptor primary structure: comparison with insulin receptor suggests structural determinants that define functional specificity. *EMBO Journal* **5** 2503–2512.
- Urakawa I, Yamazaki Y, Shimada T, Iijima K, Hasegawa H, Okawa K, Fujita T, Fukumoto S & Yamashita T 2006 Klotho converts canonical FGF receptor into a specific receptor for FGF23. *Nature* **444** 770–774. (doi:10.1038/nature05315)
- Utsugi T, Ohno T, Ohyama Y, Uchiyama T, Saito Y, Matsumura Y, Aizawa H, Itoh H, Kurabayashi M, Kawazu S *et al.* 2000 Decreased insulin production and increased insulin sensitivity in the klotho mutant mouse, a novel animal model for human aging. *Metabolism* **49** 1118–1123. (doi:10.1053/meta.2000.8606)
- Wang Y & Sun Z 2009 Klotho gene delivery prevents the progression of spontaneous hypertension and renal damage. *Hypertension* **54** 810–817. (doi:10.1161/HYPERTENSIONAHA.109.134320)
- Wang Y, Chen L, Huang G, He D, He J, Xu W, Zou C, Zong F, Li Y, Chen B *et al.* 2013 Klotho sensitizes human lung cancer cell line to cisplatin via PI3k/Akt pathway. *PLoS ONE* **8** e57391. (doi:10.1371/journal.pone.0057391)
- White KE, Cabral JM, Davis SI, Fishburn T, Evans WE, Ichikawa S, Fields J, Yu X, Shaw NJ, McLellan NJ *et al.* 2005 Mutations that cause osteoglophonic dysplasia define novel roles for FGFR1 in bone elongation. *American Journal of Human Genetics* **76** 361–367. (doi:10.1086/427956)
- White HD, Ahmad AM, Durham BH, Chandran S, Patwala A, Fraser WD & Vora JP 2006 Effect of active acromegaly and its treatment on parathyroid circadian rhythmicity and parathyroid target-organ sensitivity. *Journal of Clinical Endocrinology and Metabolism* **91** 913–919. (doi:10.1210/jc.2005-1602)
- Wolf I, Levanon-Cohen S, Bose S, Ligumsky H, Sredni B, Kanety H, Kuro-o M, Karlan B, Kaufman B, Koeffler HP *et al.* 2008 Klotho: a tumor suppressor and a modulator of the IGF-1 and FGF pathways in human breast cancer. *Oncogene* **27** 7094–7105. (doi:10.1038/onc.2008.292)
- Wu X, Weiszmam J, Ge H, Baribault H, Stevens J, Hawkins N, Vonderfecht S, Gardner J, Gupta J, Sheng J *et al.* 2012 A unique FGF23 with the ability to activate FGFR signaling through both α -Klotho and β -Klotho. *Journal of Molecular Biology* **418** 82–89. (doi:10.1016/j.jmb.2012.02.027)
- Wu AL, Feng B, Chen MZ, Kolumam G, Zavala-Solorio J, Wyatt SK, Gandham VD, Carano RA & Sonoda J 2013 Antibody-mediated activation of FGFR1 induces FGF23 production and hypophosphatemia. *PLoS ONE* **8** e57322. (doi:10.1371/journal.pone.0057322)
- Xiao NM, Zhang YM, Zheng Q & Gu J 2004 Klotho is a serum factor related to human aging. *Chinese Medical Journal* **117** 742–747.
- Yamamoto M, Clark JD, Pastor JV, Gurnani P, Nandi A, Kurosu H, Miyoshi M, Ogawa Y, Castrillon DH, Rosenblatt KP *et al.* 2005 Regulation of oxidative stress by the anti-aging hormone klotho. *Journal of Biological Chemistry* **280** 38029–38034. (doi:10.1074/jbc.M509039200)
- Yamamoto KT, Robinson-Cohen C, de Oliveira MC, Kostina A, Nettleton JA, Ix JH, Nguyen H, Eng J, Lima JA, Siscovick DS *et al.* 2013 Dietary phosphorus is associated with greater left ventricular mass. *Kidney International* **83** 707–714. (doi:10.1038/ki.2012.303)
- Yamashita T, Yoshioka M & Itoh N 2000 Identification of a novel fibroblast growth factor, FGF-23, preferentially expressed in the ventrolateral thalamic nucleus of the brain. *Biochemical and Biophysical Research Communications* **277** 494–498. (doi:10.1006/bbrc.2000.3696)
- Yamazaki Y, Imura A, Urakawa I, Shimada T, Murakami J, Aono Y, Hasegawa H, Yamashita T, Nakatani K, Saito Y *et al.* 2010 Establishment of sandwich ELISA for soluble α -Klotho measurement: age-dependent change of soluble α -Klotho levels in healthy subjects. *Biochemical and Biophysical Research Communications* **398** 513–518. (doi:10.1016/j.bbrc.2010.06.110)

- Yoshida T, Fujimori T & Nabeshima Y 2002 Mediation of unusually high concentrations of 1,25-dihydroxyvitamin D in homozygous klotho mutant mice by increased expression of renal 1 α -hydroxylase gene. *Endocrinology* **143** 683–689. (doi:10.1210/en.143.2.683)
- Zapf J, Morell B, Walter H, Laron Z & Froesch ER 1980 Serum levels of insulin-like growth factor (IGF) and its carrier protein in various metabolic disorders. *Acta Endocrinologica* **95** 505–517. (doi:10.1530/acta.0.0950505)
- Zapf J, Walter H & Froesch ER 1981 Radioimmunological determination of insulin-like growth factors I and II in normal subjects and in patients with growth disorders and extrapancreatic tumor hypoglycemia. *Journal of Clinical Investigation* **68** 1321–1330. (doi:10.1172/JCI110379)
- Zhou L, Li Y, Zhou D, Tan RJ & Liu Y 2013 Loss of Klotho contributes to kidney injury by derepression of Wnt/ β -catenin signaling. *Journal of the American Society of Nephrology* **24** 771–785. (doi:10.1681/ASN.2012080865)
- Zhu Y, Xu L, Zhang J, Xu W, Liu Y, Yin H, Lv T, An H, Liu L, He H *et al.* 2013 Klotho suppresses tumor progression via inhibiting PI3K/Akt/GSK3 β /Snail signaling in renal cell carcinoma. *Cancer Science* **104** 663–671. (doi:10.1111/cas.12134)
- Zoidis E, Gosteli-Peter M, Ghirlanda-Keller C, Meinel L, Zapf J & Schmid C 2002 IGF-I and GH stimulate Phex mRNA expression in lungs and bones and 1,25-dihydroxyvitamin D(3) production in hypophysectomized rats. *European Journal of Endocrinology* **146** 97–105. (doi:10.1530/eje.0.1460097)
- Zoidis E, Ghirlanda-Keller C, Gosteli-Peter M, Zapf J & Schmid C 2004 Regulation of phosphate (Pi) transport and NaPi-III transporter (Pit-1) mRNA in rat osteoblasts. *Journal of Endocrinology* **181** 531–540. (doi:10.1677/joe.0.1810531)

Received in final form 12 July 2013

Accepted 4 September 2013

Accepted Preprint published online 20 August 2013