



# Endothelin receptor antagonists: Potential in Alzheimer's disease

Jennifer Palmer\*, Seth Love

Dementia Research Group, Institute of Clinical Neurosciences, School of Clinical Sciences, University of Bristol, Frenchay Hospital, Bristol BS16 1LE, United Kingdom

## ARTICLE INFO

### Article history:

Received 13 October 2010

Received in revised form

17 December 2010

Accepted 17 December 2010

### Keywords:

Alzheimer's disease

Endothelin

Endothelin-converting enzyme-2

Cerebral blood flow

Beta-amyloid

## ABSTRACT

Alzheimer's disease (AD) is believed to be initiated by the accumulation of neurotoxic forms of A $\beta$  peptide within the brain. AD patients show reduction of cerebral blood flow (CBF), the extent of the reduction correlating with the impairment of cognition. There is evidence that cerebral hypoperfusion precedes and may even trigger the onset of dementia in AD. Cerebral hypoperfusion impairs neuronal function, reduces the clearance of A $\beta$  peptide and other toxic metabolites from the brain, and upregulates A $\beta$  production. Studies in animal models of AD have shown the reduction in CBF to be more than would be expected for the reduction in neuronal metabolic activity. A $\beta$  may contribute to the reduction in CBF in AD, as both A $\beta$ <sub>1–40</sub> and A $\beta$ <sub>1–42</sub> induce cerebrovascular dysfunction. A $\beta$ <sub>1–40</sub> acts directly on cerebral arteries to cause cerebral smooth muscle cell contraction. A $\beta$ <sub>1–42</sub> causes increased neuronal production and release of endothelin-1 (ET-1), a potent vasoconstrictor, and upregulation of endothelin-converting enzyme-2 (ECE-2), the enzyme which cleaves ET-1 from its inactive precursor. ET-1 and ECE-2 are also elevated in AD, making it likely that upregulation of the ECE-2–ET-1 axis by A $\beta$ <sub>1–42</sub> contributes to the chronic reduction of CBF in AD. At present, only a few symptomatic treatment options exist for AD. The involvement of ET-1 in the pathogenesis of endothelial dysfunction associated with elevated A $\beta$  indicates the potential for endothelin receptor antagonists in the treatment of AD. It has already been demonstrated that the endothelin receptor antagonist bosentan, preserves aortic and carotid endothelial function in Tg2576 mice, and our findings suggest that endothelin receptor antagonists may be beneficial in maintaining CBF in AD.

© 2011 Elsevier Ltd. All rights reserved.

## Contents

1. Introduction.....	525
2. Pathological findings in AD.....	526
3. The cerebral vasculature in AD.....	526
4. A $\beta$ acts on the endothelin system to cause reduced CBF.....	527
5. Does the endothelin system hold therapeutic potential in AD?.....	528
6. Enhancement of ECE activity for treatment of AD: need to evaluate risk of vascular dysfunction.....	528
7. Conclusions.....	528
Acknowledgments.....	529
References.....	529

## 1. Introduction

Alzheimer's disease (AD) is a progressive, degenerative disease of the brain that leads to deficits in cognitive function. Manifestations can include amnesia, aphasia, apraxia, agnosia and psychiatric symptoms such as depression, personality change, delusions and

hallucinations. Sufferers experience difficulties in activities of daily living; as the disease progresses these include such basic tasks as dressing, feeding and toileting [1]. AD is the most common of the diseases that cause dementia, of which the estimated global prevalence in 2005 was 24.3 million people [2]. In the absence of effective prevention or curative treatments, this number is predicted to double every 20 years as populations age, to 81.1 million by 2040. About 43% of patients are estimated to need a high level of care equivalent to that in a nursing home [3]. There is clearly a pressing need to find interventions to delay disease onset and progression.

\* Corresponding author at: University of Bristol, Dementia Research Group, John James Laboratories, The Day Hospital Frenchay Hospital, Bristol, United Kingdom.  
E-mail address: [jen.palmer@bristol.ac.uk](mailto:jen.palmer@bristol.ac.uk) (J. Palmer).

## 2. Pathological findings in AD

The brain undergoes atrophy, reflecting a decrease in the volume of the cerebrocortical ribbon and cerebral white matter, resulting from shrinkage or death of neurons, loss of the dendrites and degeneration of nerve fibres [4–6]. In advanced disease there is neuronal loss from the hippocampus, entorhinal regions and neocortex [7]; the cornu ammonis (CA) can be as much as 66% smaller than normal in end-stage AD with over 80% loss of neurons [8]. Loss of neocortical synapses is a relatively early abnormality that progresses with AD severity [7,9–12].

The two 'hallmark' pathological lesions in the AD brain are the extracellular parenchymal deposits of A $\beta$  peptide that form plaques and the intracellular aggregates of hyperphosphorylated tau protein that form neurofibrillary tangles (NFTs) [13]. Hyperphosphorylated tau also aggregates within nerve cell processes, as neurofibrillary threads, and in most AD patients A $\beta$  also accumulates in the walls of cerebral blood vessels, causing amyloid angiopathy (CAA) – present to some extent in over 90% of patients with AD and about one-third of elderly people who do not have AD [14]. After cleavage from its precursor, A $\beta$  initially takes the form of a coiled single peptide but this tends to aggregate, resulting in a dynamic equilibrium of monomeric, dimeric, oligomeric, protofibrillar (soluble) and fibrillar (insoluble) states [15]. Although a diagnosis of AD requires, amongst other pathological abnormalities, the presence of A $\beta$  plaques, which are predominantly made up of the insoluble fibrillar form, there is poor correlation between A $\beta$  plaque load and cognitive impairment or neuronal loss [16,17]. The correlation between soluble A $\beta$  and cognitive impairment is much stronger, suggesting that soluble species of A $\beta$  are more toxic than fibrillar forms [17]. There is recent evidence from *in vitro* and *in vivo* studies that the major toxic species of A $\beta$  are the soluble oligomeric forms [18–21]. Oligomeric A $\beta$  is increased in brain tissue from AD patients [22–26], and correlates with cognitive impairment [23,25,27]. The size or conformation of oligomer may be critical in determining toxicity, in that soluble A $\beta$  aggregates from AD cases were reported to be more toxic than soluble A $\beta$  aggregates from age-matched controls [28]. Both A $\beta$  and tau are normal constituents of the brain but have an abnormal insoluble fibrillary structure in AD. Physiological levels of A $\beta$  probably have important physiological roles [29] but AD is believed to be initiated by the abnormal accumulation of A $\beta$  peptide within the brain [30]. The amount of A $\beta$  in the brain is determined by the balance between its production and clearance. Familial autosomal dominant forms of AD, which either increase overall production of A $\beta$  or increase the ratio of A $\beta$ <sub>1–42</sub>:A $\beta$ <sub>1–40</sub>, account for fewer than 5% of cases. The mechanisms by which A $\beta$  accumulates in late-onset sporadic AD (LOAD), the major form of the disease, are unknown. There is evidence of increased activity of the rate-limiting enzyme in A $\beta$  synthesis,  $\beta$ -secretase (BACE-1) in LOAD [31–33] but animal studies suggest that this may, at least in part, simply be a reaction to the accumulation of A $\beta$  [34]. Another possible explanation for the accumulation of A $\beta$  in LOAD is an impairment of A $\beta$  clearance. Several potential pathways of A $\beta$  clearance have been identified. These include perivascular drainage [35,36], receptor-mediated transport across the blood–brain barrier [37,38] and proteolytic degradation of A $\beta$  within the brain [39–41].

## 3. The cerebral vasculature in AD

The brain does not have the capacity to store significant energy reserves, thus its viability is dependent on a continuous supply of oxygen and glucose, delivered wholly by the blood. Blood flow is directed to the cerebral circulation at the expense of other regions when required [42]. Cerebrovascular autoregulation ensures that cerebral blood flow (CBF) remains independent of mean arterial

pressure within a range of approximately 60–150 mm Hg [43]. This autoregulation is mediated by contraction of smooth muscle cells of cerebral vessels when intravascular pressure increases and relaxation when it decreases [44,45]. The distribution of CBF within the brain is regulated according to the activity of different brain regions. Neural activity increases the energy demands of the brain tissue, increasing the need for nutrient delivery and removal of toxic by-products of metabolism and heat by the circulating blood [46]. This is accomplished by increases in CBF which are highly restricted to the activated region and occur within seconds of the increase in neural activity [47–49], a phenomenon termed 'functional hyperaemia' [50] and requires neurons, astrocytes and vascular cells to work together as a neurovascular unit. Neurons that have axons or dendrites in contact with blood vessels, particularly interneurons, can rapidly communicate the need for increased blood flow by releasing vasoactive factors [51]. A direct link between functional hyperaemia and the activity of interneurons was shown by Iadecola et al. [52] in a study of blood flow in the cerebellar cortex. The increase in flow produced by stimulation of somatosensory fibres was markedly attenuated by inhibition or genetic deletion of neuronal nitric oxide synthase (NOS), present in high levels in interneurons [53,54]. The increase in CBF produced by activation of somatosensory fibres was also less in *cyclin D2*-null mice, which lack satellite interneurons [55].

There is evidence that vascular dysfunction and lowered CBF could be key factors in the development of AD, and may even trigger the onset of dementia [56]. AD patients have reduced CBF [57–62], the extent of the reduction correlating with the impairment of performance on mini-mental state examination [63–65]. Most cardiovascular risk factors (such as diabetes, hypertension, high cholesterol levels, atherosclerosis and obesity) are also risk factors for AD, suggesting an interrelationship between vascular and neurodegenerative pathology [66–75], and the onset of dementia [56,74,76]. In a large study of CBF velocity in people without clinical dementia (the Rotterdam Study, 1730 participants aged 55 years and older), those who had greater CBF velocity were less likely to go on to develop dementia [77], suggesting that cerebral hypoperfusion precedes and possibly contributes to the development of dementia. Higher CBF velocity was associated with significantly less cognitive decline, and larger hippocampus and amygdala volumes [77].

Patients with Down's syndrome have similar functional vascular abnormalities to those in AD, even before the formation of A $\beta$  plaques and NFTs although soluble forms of A $\beta$  may be elevated [78]. The earliest abnormality in transgenic mice that have AD-related *APP* mutations is a profound alteration in the regulation of CBF at 2–3 months of age [79–81]. Under normal circumstances there is a close relationship between CBF and cerebral glucose utilisation (CGU), which reflects neural metabolic activity [82,83]. AD patients show early reductions in CGU in temporal, parietal, and posterior cingulate cortices in a pattern that is distinct from that in normal ageing and other dementias [84–89]. Although reduction in CBF has long been known to be an early event in AD, it has been unclear whether this is simply a consequence of diminished metabolic demand because of neuronal dysfunction. The Rotterdam study suggested that cerebral hypoperfusion precedes and possibly contributes to the dementia but more convincing evidence came from studies of the relationship between CBF and CGU, which can be measured by positron emission tomography scanning using the glucose metabolic tracer [<sup>18</sup>F]-fluorodeoxyglucose. The Tg2576 mouse model of AD, transgenic for human *APP* containing the Swedish double mutation, was found to show reduction in CBF and CGU before the formation of A $\beta$  plaques or neurodegeneration [90,91]. However, although the reductions in CBF and CGU were coupled, the coupling differed between Tg2576 and wild-type mice: for any measured level of CGU, CBF was about 30% lower in Tg2576 than wild-type mice, indicating that the reduction in blood

flow in Tg2576 mice was not simply a reflection of lower metabolic activity.


#### 4. A $\beta$ acts on the endothelin system to cause reduced CBF

There is evidence that A $\beta$  may contribute to the reduction in CBF in AD: CBF was reduced by 20–40% in mice that over-express APP [81,92] and regulation of CBF was most impaired in transgenic mice with highest intracerebral levels of A $\beta$  [81]. Several experimental studies have shown A $\beta_{1-42}$  and A $\beta_{1-40}$  to induce endothelial dysfunction (defined as impaired endothelium-dependent relaxation and enhanced endothelium-dependent vasoconstriction) *in vivo* and *in vitro* [93–101]. For example, exposure of isolated vessels to A $\beta_{1-40}$  or A $\beta_{25-35}$  for 24–48 h altered endothelial function without changing cell morphology or inducing apoptosis [102]. In these studies of isolated vessels, A $\beta$  peptides act directly on cerebral arteries to cause cerebral smooth muscle cell contraction (Fig. 1). This direct action of A $\beta$  was associated with A $\beta_{1-40}$ , not A $\beta_{1-42}$  or the highly neurotoxic fragment A $\beta_{25-35}$ , suggesting that the factors responsible for direct mediation of A $\beta$  vasoactivity are distinct from those mediating A $\beta$  cytotoxicity. The precise mechanism of enhanced vasoconstriction is not entirely clear. It may be partly mediated through the production of free radicals [93,95,98,99,103]: interaction of A $\beta$  with endothelial cells of rat aorta produces an excess of superoxide radicals, which inactivate endothelium-derived relaxing factor [104], leading to enhanced vasoconstriction [98]. However, in some studies vasoconstriction was reported not to be caused by enhanced production or biological activity of free radicals during A $\beta$  incubation [100,102,105]. A pro-inflammatory pathway involving upregulation of COX-2 has also been suggested [106–108].

Our own research has focused on the contribution of the endothelin system to vascular dysfunction in AD. Endothelin-converting enzymes-1 and -2 (ECE-1 and -2) expressed, respectively, by endothelial cells [109] and neurons [110] within the CNS [111], convert the inactive precursor ‘big-endothelin’ to the potent

vasoconstrictor endothelin-1 (ET-1). ECE-1 and -2 are also capable of degrading A $\beta$  [112], and *ECE-1* heterozygous and *ECE-2* homozygous knock-out mice [113] have significantly elevated levels of A $\beta_{1-40}$  and A $\beta_{1-42}$  [114], attributed to decreased A $\beta$  degradation. In addition, *ECE-2* homozygous knock-out mice are deficient in learning and memory [115]. However, we found expression of *ECE-1* to be unchanged [109] and *ECE-2* to be significantly elevated [110] at both the mRNA and protein level in post-mortem (PM) brain tissue from patients with AD compared to that from normal controls and patients with vascular dementia (VaD). The fact that *ECE-2* expression was unchanged in VaD despite neuronal damage and gliosis, suggested that the elevated expression in AD might be secondary to the accumulation of A $\beta$ . 24 h exposure of SH-SY5Y cells to monomeric or oligomeric A $\beta_{1-42}$  caused marked upregulation of *ECE-2* mRNA and protein [110], supporting this hypothesis. The upregulation of *ECE-2* may be a physiological feedback or protective mechanism that serves to increase degradation of A $\beta$  and reduce its accumulation. Immunostaining of ET-1 was found to be elevated in frontal, temporal, parietal and occipital cortex in AD compared to control brains [116] and ET-1 expression was elevated in the cerebral vasculature of mice that had been infused with A $\beta$  [117]. We have recently extended these observations and shown that the levels of both *EDN1* mRNA and of ET-1 itself are significantly elevated in temporal neocortex in AD, and that A $\beta_{1-42}$  induces the production and release of ET-1 by neuroblastoma cells *in vitro* [118,119]. ET-1 was shown to be present in perivascular nerves surrounding cerebral vessels [120], providing further support for the idea that release of neuronally derived ET-1 is a cause of vasoconstriction [121]. We suggest that the cerebrovascular dysfunction in AD reflects not only a direct action of A $\beta_{1-40}$  on cerebral blood vessels but also the indirect effects of A $\beta_{1-42}$ , mediated through upregulation of neuronal *ECE-2* and increased production and release of ET-1, causing a chronic reduction in CBF.

It should be noted that there are also other ways in which A $\beta$  may cause cerebral vasoconstriction in AD. A $\beta_{1-42}$  upregulates neuronal production of angiotensin-converting enzyme (ACE) [122], the activity of which is increased in the brain in AD. ACE is another



**Fig. 1.** Two of the mechanisms by which A $\beta$  is thought to reduce cerebral blood flow in AD. A $\beta_{1-40}$  acts directly on cerebral arteries to cause cerebrovascular smooth muscle cell contraction. A $\beta_{1-42}$  acts on neurons to upregulate *ECE-2* and *EDN1* expression, increase cleavage of big ET-1 and release of ET-1. High levels of ET-1 cause chronic vasoconstriction by acting on ET<sub>A</sub> receptors on cerebral smooth muscle cells (represented by asterisks \*).

example of an enzyme that is capable of degrading A $\beta$  [123,124] but may also reduce CBF, through the cleavage of angiotensin I to form the vasoconstrictors angiotensin II and III. This is a further mechanism through which A $\beta$  may reduce CBF, potentially treatable through the use of ACE inhibitors and angiotensin receptor blockers [125].

## 5. Does the endothelin system hold therapeutic potential in AD?

At present, only a few symptomatic treatment options exist for AD. Acetylcholinesterase inhibitors (which prolong availability of ACh at synapses) and glutamate receptor (*N*-methyl-D-aspartate receptor) antagonists (which are thought to reduce glutamate-mediated neurotoxicity) are the only treatments approved by the National Institute for Health and Clinical Excellence for AD [126–131]. Many other therapies have been proposed and investigated, including anti-inflammatory drugs, antioxidants, selegiline, *G. biloba*, oestrogen replacement, statins, and compounds designed to inhibit aggregation or toxicity of A $\beta$  or tau but trial data for all of these are lacking, negative or inconclusive [132].

Pathological changes and neuronal degeneration in AD begin many years before a clinical diagnosis can be made [133–136]. Once the disease has reached the middle to late stages, a cure for AD seems unlikely because of the degradation and loss of neuronal networks that control cognitive functions. A $\beta$  immunisation (Elan AN1792 trial) resulted in significant plaque removal [137,138] in patients with AD. However, the relationship between neurodegeneration and plaque load is tenuous [139] and there was no improvement in survival or delayed progression to severe dementia in the AN1792 compared to the placebo group [138]. It has been suggested that drugs aimed at reducing the amount of A $\beta$  in the brain need to be administered before significant neurodegeneration has occurred. However, the concentration of soluble A $\beta$  in the brain is considerably higher in early than later life [140] which may reflect a physiological role in neuronal development and suggests a need for caution in attempting to lower A $\beta$  level too aggressively at too early a stage. Modulation of other risk factors may be a more realistic strategy. de la Torre [141] proposed screening subjects at mid-life for vascular risk factors associated with AD, such as atherosclerosis, hypertension and cardiovascular disease, and treating those conditions before they become symptomatic; yet this should constitute normal good clinical practice for the prevention of stroke and other adverse vascular events irrespective of the potential beneficial impact of such screening on development of AD.

As noted above, one of the earliest abnormalities in AD is a reduction in CBF [79–81,85] and both A $\beta_{1-40}$  and A $\beta_{1-42}$  induce endothelial dysfunction [103]. The cerebral hypoperfusion observed in AD is likely to impair neuronal function and reduce the clearance of A $\beta$  and other solutes from the brain – whether by perivascular drainage or by transport across vessel walls into the circulation. Furthermore, hypoperfusion results in upregulation of BACE1 and so increases A $\beta$  production [142,143]. Cerebral hypoperfusion may also lead to cerebral capillary damage, further reducing blood flow [144]. Intervention at an early stage may be of critical importance in preserving vascular function, neuronal metabolism and cognitive function [145,146].

Outside of the CNS several diseases are associated with abnormal activation of the endothelin system. These include pulmonary hypertension, some forms of renal disease, systemic arterial hypertension, heart failure, allograft rejection and diabetes/insulin resistance [147]. Substantial physiological and clinical benefit derives from the use of ET receptor antagonists, particularly selective ET<sub>A</sub> receptor blockers, in these disorders [148–155]. ET<sub>A</sub> receptor antagonists improve endothelium-dependent, NO-

mediated vasodilatation [156], which is blocked by selective ET<sub>B</sub> receptor antagonists [157].

In keeping with our own observations on the upregulation of the ECE-2-ET-1 axis by A $\beta$ , Elesber et al. [158] demonstrated that bosentan (Tracleer®), a dual ET<sub>A</sub>/ET<sub>B</sub> receptor antagonist, preserved aortic and carotid endothelial function in Tg2576 mice, which overexpress *APP*. This provides additional evidence of the involvement of ET-1 in the pathogenesis of endothelial dysfunction associated with elevated A $\beta$  and demonstrates the therapeutic potential of drugs such as bosentan in this context. Several selective ET<sub>A</sub> receptor antagonists have been used for the treatment of pulmonary hypertension, including ambrisentan (Letairis®, Volibris®), darusentan and sixaxsantan (Thelin®) [147]. Experimental and clinical studies have shown beneficial effects of treating pulmonary hypertension with both ET<sub>A</sub> receptor-selective and non-selective ET<sub>A</sub>/ET<sub>B</sub> receptor antagonists [159] but not selective ET<sub>B</sub> receptor antagonists [148,150–153]. In view of the opposing functions of the ET<sub>A</sub> and ET<sub>B</sub> receptor in the maintenance of vascular tone, and the probable role that endothelial ET<sub>B</sub> receptors play in the clearance of excess ET-1 from the circulation [159], it seems likely that selective ET<sub>A</sub> receptor inhibitors would be most beneficial in maintaining CBF in AD.

## 6. Enhancement of ECE activity for treatment of AD: need to evaluate risk of vascular dysfunction

Reduction of A $\beta$  accumulation is a major objective of treatment in AD, and enhancement of the activity of A $\beta$ -degrading enzymes is a potential therapeutic approach [160]. The ECEs have A $\beta$ -degrading potential [39,161–163]. Intracranial administration of serotype 5 recombinant adeno-associated viral vector containing the *ECE-1* synthetic gene resulted in an ~50% reduction in the total A $\beta$  load in the cortex and hippocampus in transgenic mice overexpressing *APP* and *PSEN1* [162]. This led to the suggestion that increasing the expression of the ECEs is a promising therapeutic approach for the treatment of AD [162,163]. In these studies, the A $\beta$  load was assessed by immunohistochemistry of PM brain tissue, and no measurements of blood pressure, CBF or cognitive function were made antemortem, so it remains to be investigated whether measures that increase ECE activity affect the production of ET-1, cerebral perfusion or cognition.

The corollary of the concern that enhanced ECE activity might reduce CBF is that the use of ECE inhibitors to improve CBF might exacerbate the accumulation of A $\beta$ . The advantage of using ET-1 receptor antagonists rather than ECE inhibitors is that the former would be expected to prevent the cerebrovascular dysfunction caused by excess ET-1 without the potentially deleterious effect on ECE-2-mediated degradation of A $\beta$ .

## 7. Conclusions

Many patients with AD have concomitant white matter pathology, attributed to ischaemia caused by cerebral hypoperfusion [164,165]. Current data indicate that increased levels of A $\beta$  in the brain may have a significant influence on CBF but the precise mechanism of A $\beta$ -mediated vasoconstriction remains to be established. Little is known about the normal function of A $\beta$ , which is present in the normal brain and is produced throughout life but accumulates in excess in AD [166]. A $\beta$  starts to accumulate decades before the onset of AD and is likely to influence cerebral perfusion from as early as mid-life in a much wider context than solely in patients with AD. This influence could extend to conditions such as neurogenic hypertension and all of its systemic consequences [167,168] including stroke and cerebral ischaemic damage.

The relationship between vascular risk factors and AD is difficult to interpret from review of the scientific literature, as the pres-



ence of significant cerebrovascular disease was long considered an exclusion criterion for the diagnosis of AD. However, recent experimental studies have strengthened the 'vascular hypothesis' of AD, in which dysfunction of the cerebral microvasculature contributes directly to the development of AD [74]. There is also a growing body of evidence that A $\beta$  contributes to cerebrovascular insufficiency, by both direct and indirect actions on the cerebral vasculature, the latter mediated through the influences of A $\beta$  on neuronal function and signalling.

## Acknowledgments

Some of the research described in this article was supported by grants from Alzheimer's Research Trust, the British Heart Foundation, and BRACE (Bristol Research into Alzheimer's and Care of the Elderly).

## References

- Burns A, Byrne EJ, Maurer K. Alzheimer's disease. *Lancet* 2002;360:163–5.
- Ferri CP, Prince M, Brayne C, Brodaty H, Fratiglioni L, Ganguli M, Hall K, Hasegawa K, Hendrie H, Huang Y, Jorm A, Mathers C, Menezes PR, Rimmer E, Scazufca M. Alzheimer's disease I. Global prevalence of dementia: a delphi consensus study. *Lancet* 2005;366:2112–7.
- Brookmeyer R, Johnson E, Ziegler-Graham K, Arrighi HM. Forecasting the global burden of alzheimer's disease. *Alzheimer's Dement* 2007;3:186–91.
- Smith AD. Imaging the progression of alzheimer pathology through the brain. *Proc Natl Acad Sci U S A* 2002;99:4135–7.
- Duyckaerts C, Hauw JJ, Piette F, Rainsard C, Poulain V, Berthaux P, Escourolle R. Cortical atrophy in senile dementia of the alzheimer type is mainly due to a decrease in cortical length. *Acta Neuropathol (Berl)* 1985;66:72–4.
- Lerch JP, Pruessner J, Zijdenbos AP, Collins DL, Teipel SJ, Hampel H, Evans AC. Automated cortical thickness measurements from mri can accurately separate alzheimer's patients from normal elderly controls. *Neurobiol Aging* 2008;29:23–30.
- Terry RD, Peck A, Deteresa R, Schechter R, Horoupian DS. Some morphometric aspects of the brain in senile dementia of the alzheimer type. *Ann Neurol* 1981;10:184–92.
- Bobinski M, Wegiel J, Wisniewski HM, Tarnawski M, Reisberg B, DeLeon MJ, Miller DC. Neurofibrillary pathology – correlation with hippocampal formation atrophy in alzheimer disease. *Neurobiol Aging* 1996;17:909–19.
- Rapoport SI. In vivo pet imaging and postmortem studies suggest potentially reversible and irreversible stages of brain metabolic failure in alzheimer's disease. *Eur Arch Psychiatry Clin Neurosci* 1999;249:46–55.
- Terry RD, Masliah E, Salmon DP, Butters N, Deteresa R, Hill R, Hansen LA, Katzman R. Physical basis of cognitive alterations in alzheimer's disease – synapse loss is the major correlate of cognitive impairment. *Ann Neurol* 1991;30:572–80.
- Masliah E, Mallory M, Alford M, DeTeresa R, Hansen LA, McKeel DW, Morris JC. Altered expression of synaptic proteins occurs early during progression of alzheimer's disease. *Neurology* 2001;56:127–9.
- Masliah E, Mallory M, Hansen L, Richard D, Alford M, Terry R. Synaptic and neuritic alterations during the progression of alzheimer's disease. *Neurosci Lett* 1994;174:67–72.
- Braak H, Braak E. Neuropathological staging of alzheimer-related changes. *Acta Neuropathol (Berl)* 1991;82:239–59.
- Love S. Contribution of cerebral amyloid angiopathy to alzheimer's disease. *J Neurol Neurosurg Psychiatry* 2004;75:1–4.
- Montalto MC, Farrar G, Hehir CT. Fibrillar and oligomeric  $\beta$ -amyloid as distinct local biomarkers for alzheimer's disease. *Ann NY Acad Sci* 2007;1097:239–58.
- Crystal H, Dickson D, Fuld P, Masur D, Scott R, Mehler M, Masdeu J, Kawas C, Aronson M, Wolfson L. Clinicopathologic studies in dementia – nondemented subjects with pathologically confirmed alzheimer's disease. *Neurology* 1988;38:1682–7.
- Glabe CG. Common mechanisms of amyloid oligomer pathogenesis in degenerative disease. *Neurobiol Aging* 2006;27:570–5.
- Dahlgren KN, Manelli AN, Stine WB, Baker LK, Krafft GA, LaDu MJ. Oligomeric and fibrillar species of amyloid- $\beta$  peptides differentially affect neuronal viability. *J Biol Chem* 2002;277:32046–53.
- Lesne S, Koh MT, Kotilinek L, Kaye R, Glabe CG, Yang A, Gallagher M, Ashe KH. A specific amyloid- $\beta$  protein assembly in the brain impairs memory. *Nature* 2006;440:352–7.
- Townsend M, Shankar GM, Mehta T, Walsh DM, Selkoe DJ. Effects of secreted oligomers of amyloid  $\beta$ -protein on hippocampal synaptic plasticity: a potent role for trimers. *J Physiol-Lond* 2006;572:477–92.
- Shankar GM, Li SM, Mehta TH, Garcia-Munoz A, Shepardson NE, Smith I, Brett FM, Farrell MA, Rowan MJ, Lemere CA, Regan CM, Walsh DM, Sabatini BL, Selkoe DJ. Amyloid- $\beta$  protein dimers isolated directly from alzheimer's brains impair synaptic plasticity and memory. *Nat Med* 2008;14:837–42.
- van Helmond Z, Heesom K, Love S. Characterisation of two antibodies to oligomeric  $\beta$  and their use in elisas on human brain tissue homogenates. *J Neurosci Methods* 2009;176:206–12.
- Lue LF, Kuo YM, Roher AE, Brachova L, Shen Y, Sue L, Beach T, Kurth JH, Rydel RE, Rogers J. Soluble amyloid- $\beta$  peptide concentration as a predictor of synaptic change in alzheimer's disease. *Am J Pathol* 1999;155:853–62.
- Kuo YM, Emmerling MR, Vigo-Pelfrey C, Kasunic TC, Kirkpatrick JB, Murdoch GH, Ball MJ, Roher AE. Water-soluble  $\beta$  (n-40, n-42) oligomers in normal and alzheimer disease brains. *J Biol Chem* 1996;271:4077–81.
- Wang J, Dickson DW, Trojanowski JQ, Lee VMY. The levels of soluble versus insoluble brain  $\beta$  distinguish alzheimer's disease from normal and pathologic aging. *Exp Neurol* 1999;158:328–37.
- McLean CA, Cherny RA, Fraser FW, Fuller SJ, Smith MJ, Beyreuther K, Bush AI, Masters CL. Soluble pool of  $\beta$  amyloid as a determinant of severity of neurodegeneration in alzheimer's disease. *Ann Neurol* 1999;46:860–6.
- Gong YS, Chang L, Viola KL, Lacor PN, Lambert MP, Finch CE, Krafft GA, Klein WL. Alzheimer's disease-affected brain: presence of oligomeric  $\beta$  ligands (addls) suggests a molecular basis for reversible memory loss. *Proc Natl Acad Sci U S A* 2003;100:10417–22.
- Piccini A, Russo C, Gliazzi A, Relini A, Vitali A, Borghi R, Giliberto L, Armirotti A, D'Arrigo C, Bachi A, Cattaneo A, Canale C, Torrasa S, Saido TC, Markesbery W, Gambetti P, Tabaton M. B-amyloid is different in normal aging and in alzheimer disease. *J Biol Chem* 2005;280:34186–92.
- Pearson HA, Peers C. Physiological roles for amyloid  $\beta$  peptides. *J Physiol-Lond* 2006;575:5–10.
- Tanzi RE, Bertram L. Twenty years of the alzheimer's disease amyloid hypothesis: a genetic perspective. *Cell* 2005;120:545–55.
- Li R, Lindholm K, Yang LB, Yue X, Citron M, Yao RQ, Beach T, Sue L, Sabbagh M, Cai HB, Wong P, Price D, Shen Y. Amyloid  $\beta$  peptide load is correlated with increased  $\beta$ -secretase activity in sporadic alzheimer's disease patients. *Proc Natl Acad Sci U S A* 2004;101:3632–7.
- Yang LB, Lindholm K, Yan RQ, Citron M, Xia WM, Yang XL, Beach T, Sue L, Wong P, Price D, Li R, Shen Y. Elevated  $\beta$ -secretase expression and enzymatic activity detected in sporadic alzheimer disease. *Nat Med* 2003;9:3–4.
- Tyler SJ, Dawbarn D, Wilcock GK, Allen SJ. A- and  $\beta$ -secretase: profound changes in alzheimer's disease. *Biochem Biophys Res Commun* 2002;299:373–6.
- Zhao J, Fu YF, Yasvoina M, Shao PZ, Hitt B, O'Connor T, Logan S, Maus E, Citron M, Berry R, Binder L, Vassar R. B-site amyloid precursor protein cleaving enzyme 1 levels become elevated in neurons around amyloid plaques: implications for alzheimer's disease pathogenesis. *J Neurosci* 2007;27:3639–49.
- Preston SD, Steart PV, Wilkinson A, Nicoll JAR, Weller RO. Capillary and arterial cerebral amyloid angiopathy in alzheimer's disease: defining the perivascular route for the elimination of amyloid  $\beta$  from the human brain. *Neuropathol Appl Neurobiol* 2003;29:106–17.
- Weller RO, Massey A, Newman TA, Hutchings M, Kuo YM, Roher AE. Cerebral amyloid angiopathy – amyloid- $\beta$  accumulates in putative interstitial fluid drainage pathways in alzheimer's disease. *Am J Pathol* 1998;153:725–33.
- Lam FC, Liu RH, Lu PH, Shapiro AB, Renoir JM, Sharom FJ, Reiner PB. B-amyloid efflux mediated by p-glycoprotein. *J Neurochem* 2001;76:1121–8.
- Shibata M, Yamada S, Kumar SR, Calero M, Bading J, Frangione B, Holtzman DM, Miller CA, Strickland DK, Ghiso J, Zlokovic BV. Clearance of alzheimer's amyloid- $\beta_{1-40}$  peptide from brain by ldl receptor-related protein-1 at the blood–brain barrier. *J Clin Invest* 2000;106:1489–99.
- Eckman EA, Eckman CB.  $\beta$ -degrading enzymes: modulators of alzheimer's disease pathogenesis and targets for therapeutic intervention. *Biochem Soc Trans* 2005;33:1101–5.
- Jacobsen JS, Comery TA, Martone RL, Elokda H, Crandall DL, Oganessian A, Aschmies J, Kirksey Y, Gonzales C, Xu J, Zhou H, Atchison K, Wagner E, Zaleska MM, Das I, Arias RL, Bard J, Riddell D, Gardell SJ, Abou-Gharbia M, Robichaud A, Magolda R, Vlasuk GP, Bjornsson T, Reinhart PH, Pangalos MN. Enhanced clearance of  $\beta$  in brain by sustaining the plasmin proteolysis cascade. *Proc Natl Acad Sci U S A* 2008;105:8754–9.
- Jiang Q, Lee CYD, Mandrekar S, Wilkinson B, Cramer P, Zelcer N, Mann K, Lamb B, Willson TM, Collins JL, Richardson JC, Smith JD, Comery TA, Riddell D, Holtzman DM, Tontonoz P, Landreth GE. ApoE promotes the proteolytic degradation of  $\beta$ . *Neuron* 2008;58:681–93.
- van Lieshout JJ, Wieling W, Karemaker JM, Secher NH. Syncope, cerebral perfusion, and oxygenation. *J Appl Physiol* 2003;94:833–48.
- Cipolla MJ. Cerebrovascular function in pregnancy and eclampsia. *Hypertension* 2007;50:14–24.
- Hill MA, Sun Z, Martinez-Lemus L, Meininger GA. New technologies for dissecting the arteriolar myogenic response. *Trends Pharmacol Sci* 2007;28:308–15.
- Bayliss WM. On the local reactions of the arterial wall to changes of internal pressure. *J Physiol-Lond* 1902;28:220–31.
- Iadecola C, Nedergaard M. Glial regulation of the cerebral microvasculature. *Nat Neurosci* 2007;10:1369–76.
- Cox SB, Woolsey TA, Rovainen CM. Localized dynamic changes in cortical blood-flow with whisker stimulation corresponds to matched vascular and neuronal architecture of rat barrels. *J Cereb Blood Flow Metab* 1993;13:899–913.
- Ngai AC, Ko KR, Morii S, Winn HR. Effect of sciatic-nerve stimulation on pial arterioles in rats. *Am J Physiol* 1988;254:H133–9.
- Silva AC, Lee SP, Iadecola C, Kim SG. Early temporal characteristics of cerebral blood flow and deoxyhemoglobin changes during somatosensory stimulation. *J Cereb Blood Flow Metab* 2000;20:201–6.
- Iadecola C. Neurovascular regulation in the normal brain and in alzheimer's disease. *Nat Rev Neurosci* 2004;5:347–60.
- Freund TF. Interneuron diversity series: rhythm and mood in perisomatic inhibition. *Trends Neurosci* 2003;26:489–95.

- [53] Yang G, Chen G, Ebner TJ, Iadecola C. Nitric oxide is the predominant mediator of cerebellar hyperemia during somatosensory activation in rats. *Am J Physiol-Regul Integr Comp Physiol* 1999;277:R1760–70.
- [54] Yang G, Zhang Y, Ross ME, Iadecola C. Attenuation of activity-induced increases in cerebellar blood flow in mice lacking neuronal nitric oxide synthase. *Am J Physiol-Heart Circul Physiol* 2003;285:H298–304.
- [55] Yang G, Huard JMT, Beitz AJ, Ross ME, Iadecola C. Stellate neurons mediate functional hyperemia in the cerebellar molecular layer. *J Neurosci* 2000;20:6968–73.
- [56] de la Torre JC. Is alzheimer's disease preceded by neurodegeneration or cerebral hypoperfusion? *Ann Neurol* 2005;57:783–4.
- [57] Jagust WJ. Neuroimaging in dementia. *Neurol Clin* 2000;18:885–901.
- [58] Sharp P, Gemmell H, Cherryman G, Besson J, Crawford J, Smith F. Application of i-123 labeled isopropylamphetamine imaging to the study of dementia. *J Nucl Med* 1986;27:761–8.
- [59] Jagust WJ, Budinger TF, Reed BR. The diagnosis of dementia with single photon-emission computed-tomography. *Arch Neurol* 1987;44:258–62.
- [60] Pakrasi S, O'Brien JT. Emission tomography in dementia. *Nucl Med Commun* 2005;26:189–96.
- [61] Schuff N, Matsumoto S, Kmiecik J, Studholme C, Du A, Ezekiel F, Miller BL, Kramer JH, Jagust WJ, Chui HC, Weiner MW. Cerebral blood flow in ischemic vascular dementia and alzheimer's disease, measured by arterial spin-labeling magnetic resonance imaging. *Alzheimer's Dement* 2009;5:454–62.
- [62] O'Brien JT. Role of imaging techniques in the diagnosis of dementia. *Br J Radiol* 2007;80:S71–77.
- [63] Dekosky ST, Shih WJ, Schmitt FA, Coupal J, Kirkpatrick C. Assessing utility of single photon emission computed tomography spect scan in alzheimer disease correlation with cognitive severity. *Alzheimer Dis Assoc Disord* 1990;4:14–23.
- [64] Imran MB, Kawashima R, Awata S, Sato K, Kinomura S, Ono S, Sato M, Fukuda H. Tc-99m hmpao spect in the evaluation of alzheimer's disease: correlation between neuropsychiatric evaluation and cbf images. *J Neurol Neurosurg Psychiatry* 1999;66:228–32.
- [65] Shiraishi H, Chang CC, Kanno H, Yamamoto I. The relationship between cerebral blood flow and cognitive function in patients with brain insult of various etiology. *J Clin Neurosci* 2004;11:138–41.
- [66] Qiu CX, Winblad B, Fratiglioni L. The age-dependent relation of blood pressure to cognitive function and dementia. *Lancet Neurol* 2005;4:487–99.
- [67] Skoog I, Gustafson D. Update on hypertension and alzheimer's disease. *Neurol Res* 2006;28:605–11.
- [68] Bellew KM, Pigeon JG, Stang PE, Fleischman W, Gardner RM, Baker WW. Hypertension and the rate of cognitive decline in patients with dementia of the alzheimer type. *Alzheimer Dis Assoc Disord* 2004;18:208–13.
- [69] Petrovitch H, White LR, Izmirlian G, Ross GW, Havlik RJ, Markesbery W, Nelson J, Davis DG, Hardman J, Foley DJ, Launer LJ. Midlife blood pressure and neurotic plaques, neurofibrillary tangles, and brain weight at death: the haas. *Neurobiol Aging* 2000;21:57–62.
- [70] Sparks DL, Scheff SW, Liu HC, Landers TM, Coyne CM, Hunsaker JC. Increased incidence of neurofibrillary tangles (nft) in nondemented individuals with hypertension. *J Neurol Sci* 1995;131:162–9.
- [71] Korf ESC, White LR, Scheltens P, Launer LJ. Midlife blood pressure and the risk of hippocampal atrophy – the honolulu asia aging study. *Hypertension* 2004;44:29–34.
- [72] den Heijer T, Launer LJ, Prins ND, van Dijk EJ, Vermeer SE, Hofman A, Koudstaal PJ, Breteler MMB. Association between blood pressure, white matter lesions, and atrophy of the medial temporal lobe. *Neurology* 2005;64:263–7.
- [73] Kivipelto M, Helkala EL, Laakso MP, Hanninen T, Hallikainen M, Alhainen K, Soininen H, Tuomilehto J, Nissien A. Midlife vascular risk factors and alzheimer's disease in later life: longitudinal, population based study. *Br Med J* 2001;322:1447–51.
- [74] Dickstein DL, Walsh J, Brautigam H, Stockton SD, Gandy S, Hof PR. Role of vascular risk factors and vascular dysfunction in alzheimer's disease. *Mt Sinai J Med* 2010;77:82–102.
- [75] Ott A, Stolk RP, van Harskamp F, Pols HAP, Hofman A, Breteler MMB. Diabetes mellitus and the risk of dementia – the rotterdam study. *Neurology* 1999;53:1937–42.
- [76] Snowden DA, Greiner LH, Mortimer JA, Riley KP, Greiner PA, Markesbery WR. Brain infarction and the clinical expression of alzheimer disease – the nun study. *JAMA-J Am Med Assoc* 1997;277:813–7.
- [77] Ruitenberg A, Heijer Td, Bakker SLM, Swieten JCV, Koudstaal PJ, Hofman A, Breteler MMB. Cerebral hypoperfusion and clinical onset of dementia: the rotterdam study. *Ann Neurol* 2005;57:789–94.
- [78] Buee L, Hof PR, Bouras C, Delacourte A, Perl DP, Morrison JH, Fillit HM. Pathological alterations of the cerebral microvasculature in alzheimer's-disease and related dementing disorders. *Acta Neuropathol (Berl)* 1994;87:469–80.
- [79] Iadecola C, Zhang FY, Niwa K, Eckman C, Turner SK, Fischer E, Yonkin S, Borchelt DR, Hsiao KK, Carlson GA. Sod1 rescues cerebral endothelial dysfunction in mice overexpressing amyloid precursor protein. *Nat Neurosci* 1999;2:157–61.
- [80] Niwa K, Kazama K, Yonkin L, Yonkin SG, Carlson GA, Iadecola C. Cerebrovascular autoregulation is profoundly impaired in mice overexpressing amyloid precursor protein. *Am J Physiol-Heart Circul Physiol* 2002;283:H315–23.
- [81] Niwa K, Kazama K, Yonkin SG, Carlson GA, Iadecola C. Alterations in cerebral blood flow and glucose utilization in mice overexpressing the amyloid precursor protein. *Neurobiol Dis* 2002;9:61–8.
- [82] Reivich M. Blood flow metabolism couple in brain. *Res Publ Assoc Res Nerv Ment Dis* 1974;53:125–40.
- [83] Sokoloff L, Reivich M, Kennedy C, Desrosiers MH, Patlak CS, Pettigrew KD, Sakurada O, Shinohara M. Deoxyglucose-c-14 method for measurement of local cerebral glucose-utilization – theory, procedure, and normal values in conscious and anesthetized albino-rat. *J Neurochem* 1977;28:897–916.
- [84] Chetelat G, Desgranges B, de la Sayette V, Viader F, Eustache F, Baron JC. Mild cognitive impairment – can fdg-pet predict who is to rapidly convert to alzheimer's disease? *Neurology* 2003;60:1374–7.
- [85] Jagust W, Gitcho A, Sun F, Kuczynski B, Mungas D, Haan M. Brain imaging evidence of preclinical alzheimer's disease in normal aging. *Ann Neurol* 2006;59:673–81.
- [86] Silverman DHS, Small GW, Chang CY, Lu CS, de Aburto MAK, Chen W, Czernin J, Rapoport SI, Pietrini P, Alexander GE, Schapiro MB, Jagust WJ, Hoffman JM, Welsh-Bohmer KA, Alavi A, Clark CM, Salmon E, de Leon MJ, Mielke R, Cummings JL, Kowell AP, Gambhir SS, Hoh CK, Phelps ME. Positron emission tomography in evaluation of dementia – regional brain metabolism and long-term outcome. *JAMA-J Am Med Assoc* 2001;286:2120–7.
- [87] Minoshima S, Giordani B, Berent S, Frey KA, Foster NL, Kuhl DE. Metabolic reduction in the posterior cingulate cortex in very early alzheimer's disease. *Ann Neurol* 1997;42:85–94.
- [88] Herholz K. Pet studies in dementia. *Ann Nucl Med* 2003;17:79–89.
- [89] Jagust W, Reed B, Mungas D, Ellis W, DeCarli C. What does fluorodeoxyglucose pet imaging add to a clinical diagnosis of dementia? *Neurology* 2007;69:871–7.
- [90] Hsiao K, Chapman P, Nilsen S, Eckman C, Harigaya Y, Yonkin S, Yang FS, Cole G. Correlative memory deficits, a $\beta$  elevation, and amyloid plaques in transgenic mice. *Science* 1996;274:99–102.
- [91] Hsiao KK, Borchelt DR, Olson K, Johannsdottir R, Kitt C, Yunis W, Xu S, Eckman C, Yonkin S, Price D, Iadecola C, Clark HB, Carlson G. Age-related cns disorder and early death in transgenic fvb/n mice overexpressing alzheimer amyloid precursor proteins. *Neuron* 1995;15:1203–18.
- [92] Bourasset F, Ouellet M, Tremblay C, Julien C, Do TM, Oddo S, LaFerla F, Calon F. Reduction of the cerebrovascular volume in a transgenic mouse model of alzheimer's disease. *Neuropharmacology* 2009;56:808–13.
- [93] Crawford F, Suo ZM, Fang CH, Mullan M. Characteristics of the in vitro vasoactivity of  $\beta$ -amyloid peptides. *Exp Neurol* 1998;150:159–68.
- [94] Price JM, Chi XD, Hellermann G, Sutton ET. Physiological levels of  $\beta$ -amyloid induce cerebral vessel dysfunction and reduce endothelial nitric oxide production. *Neurol Res* 2001;23:506–12.
- [95] Price JM, Sutton ET, Hellermann A, Thomas T. B-amyloid induces cerebrovascular endothelial dysfunction in the rat brain. *Neurol Res* 1997;19:534–8.
- [96] Niwa K, Carlson GA, Iadecola C. Exogenous a $\beta_{1-40}$  reproduces cerebrovascular alterations resulting from amyloid precursor protein overexpression in mice. *J Cereb Blood Flow Metab* 2000;20:1659–68.
- [97] Niwa K, Porter VA, Kazama K, Cornfield D, Carlson GA, Iadecola C. A $\beta$ -peptides enhance vasoconstriction in cerebral circulation. *Am J Physiol-Heart Circul Physiol* 2001;281:H2417–24.
- [98] Thomas T, Thomas G, McLendon C, Sutton T, Mullan M. B-amyloid-mediated vasoactivity and vascular endothelial damage. *Nature* 1996;380:168–71.
- [99] Thomas T, McLendon C, Sutton ET, Thomas G. Cerebrovascular endothelial dysfunction mediated by  $\beta$ -amyloid. *Neuroreport* 1997;8:1387–91.
- [100] Gentile MT, Vecchione C, Maffei A, Aretini A, Marino G, Poulet R, Capobianco L, Selvetella G, Lembo G. Mechanisms of soluble  $\beta$ -amyloid impairment of endothelial function. *J Biol Chem* 2004;279:48135–42.
- [101] Luo F, Seifert TR, Edalji R, Loebbert RW, Hradil VP, Harlan J, Schmidt M, Nim-mirich V, Cox BF, Fox GB. Non-invasive characterization of  $\beta$ -amyloid(1–40) vasoactivity by functional magnetic resonance imaging in mice. *Neuroscience* 2008;155:263–9.
- [102] Chisari M, Merlo S, Sortino MA, Salomone S. Long-term incubation with  $\beta$ -amyloid peptides impairs endothelium-dependent vasodilatation in isolated rat basilar artery. *Pharmacol Res* 2010;61:157–61.
- [103] Dietrich H, Xiang C, Han B, Zipfel G, Holtzman D. Soluble amyloid- $\beta$ , effect on cerebral arteriolar regulation and vascular cells. *Molecular Neurodegeneration* 2010;5:15.
- [104] Rubanyi GM, Vanhoutte PM. Superoxide anions and hyperoxia inactivate endothelium-derived relaxing factor. *Am J Physiol* 1986;250:H822–7.
- [105] Paris D, Parker TA, Town T, Suo ZM, Fang CH, Humphrey J, Crawford F, Mullan M. Role of peroxynitrite in the vasoactive and cytotoxic effects of alzheimer's  $\beta$ -amyloid(1–40) peptide. *Exp Neurol* 1998;152:116–22.
- [106] Townsend KP, Obregon D, Quadros A, Patel N, Volmar C, Paris D, Mullan M. Proinflammatory and vasoactive effects of a beta in the cerebrovasculature. *Alzheimer's Dis: Vasc Etiol Pathol* 2002;97:65–76.
- [107] Paris D, Town T, Mori T, Parker TA, Humphrey J, Mullan M. Soluble  $\beta$ -amyloid peptides mediate vasoactivity via activation of a pro-inflammatory pathway. *Neurobiol Aging* 2000;21:183–97.
- [108] Paris D, Humphrey J, Quadros A, Patel N, Crescentini R, Crawford F, Mullan M. Vasoactive effects of a $\beta$  in isolated human cerebrovessels and in a transgenic mouse model of alzheimer's disease: role of inflammation. *Neurol Res* 2003;25:642–51.
- [109] Palmer JC, Kehoe PG, Love S. Endothelin-converting enzyme-1 in alzheimer's disease and vascular dementia. *Neuropathol Appl Neurobiol* 2010;36:487–97.
- [110] Palmer JC, Baig S, Kehoe PG, Love S. Endothelin-converting enzyme-2 is increased in alzheimer's disease and up-regulated by a $\beta$ . *Am J Pathol* 2009;175:262–70.
- [111] Emoto N, Yanagisawa M. Endothelin-converting enzyme-2 is a membrane-bound, phosphoramidon-sensitive metalloprotease with acidic pH optimum. *J Biol Chem* 1995;270:15262–8.

- [112] Eckman EA, Reed DK, Eckman CB. Degradation of the alzheimer's amyloid- $\beta$  peptide by endothelin-converting enzyme. *J Biol Chem* 2001;276:24540–8.
- [113] Yanagisawa H, Hammer RE, Richardson JA, Emoto N, Williams SC, Takeda S, Clouthier DE, Yanagisawa M. Disruption of ECE-1 and ECE-2 reveals a role for endothelin-converting enzyme-2 in murine cardiac development. *J Clin Invest* 2000;105:1373–82.
- [114] Eckman EA, Watson M, Marlow L, Sambamurti K, Eckman CB. Alzheimer's disease  $\beta$ -amyloid peptide is increased in mice deficient in endothelin-converting enzyme. *J Biol Chem* 2003;278:2081–4.
- [115] Rodriguiz RM, Gadnidez K, Ragnauth A, Dorr N, Yanagisawa M, Wetsel WC, Devi LA. Animals lacking endothelin-converting enzyme-2 are deficient in learning and memory. *Genes Brain Behav* 2008;7:418–26.
- [116] Minami M, Kimura M, Iwamoto N, Arai H. Endothelin-1-like immunoreactivity in cerebral-cortex of alzheimer-type dementia. *Prog Neuropsychopharmacol Biol Psychiatry* 1995;19:509–13.
- [117] Deane R, Yan SD, Subramanyam RK, LaRue B, Jovanovic S, Hogg E, Welch D, Manness L, Lin C, Yu J, Zhu H, Ghiso J, Frangione B, Stern A, Schmidt AM, Armstrong DL, Arnold B, Liliensiek B, Nawroth P, Hofman F, Kindy M, Stern D, Zlokovic B. Rage mediates amyloid- $\beta$  peptide transport across the blood–brain barrier and accumulation in brain. *Nat Med* 2003;9:907–13.
- [118] Palmer J, Kehoe P, Love S. A $\beta$  induces endothelin-1 production by sh-sy5y neuroblastoma cells. *Brain Pathol* 2010;20(suppl 1):18.
- [119] Palmer J, Kehoe P, Love S. Et-1 is elevated in the brain in ad and its neuronal production and release are upregulated by a $\beta$ 1–42. *Neuropathol Appl Neurobiol* 2011;37(Suppl. 1):34.
- [120] Loesch A. Localisation of endothelin-1 and its receptors in vascular tissue as seen at the electron microscopic level. *Curr Vasc Pharmacol* 2005;3:381–92.
- [121] Milner P, Loesch A, Burnstock G. Neural endothelin in hypertension: increased expression in ganglia and nerves to cerebral arteries of the spontaneously hypertensive rat. *J Vasc Res* 2000;37:39–49.
- [122] Miners JS, Ashby EL, Baig S, Harrison R, Tayler H, Speedy E, Prince JA, Love S, Kehoe PG. Angiotensin-converting enzyme levels and activity in alzheimer's disease: differences in brain and csf ace and association with ace1 genotypes. *Am J Transl Res* 2009;1:163–77.
- [123] Zou K, Maeda T, Watanabe A, Liu J, Liu S, Oba R, Satoh Y-I, Komano H, Michikawa M. A $\beta$ 42-to-a $\beta$ 40-and angiotensin-converting activities in different domains of angiotensin-converting enzyme. *J Biol Chem* 2009;284:31914–20.
- [124] Hemming ML, Selkoe DJ. Amyloid  $\beta$ -protein is degraded by cellular angiotensin-converting enzyme (ace) and elevated by an ace inhibitor. *J Biol Chem* 2005;280:37644–50.
- [125] Kehoe PG, Miners S, Love S. Angiotensins in alzheimer's disease – friend or foe? *Trends Neurosci* 2009;32:619–28.
- [126] Farlow M. A clinical overview of cholinesterase inhibitors in alzheimer's disease. *Int Psychogeriatr* 2002;14:93–126.
- [127] Cummings JL. Use of cholinesterase inhibitors in clinical practice – evidence-based recommendations. *Am J Geriatr Psychiatry* 2003;11:131–45.
- [128] Birks J. Cholinesterase inhibitors for alzheimer's disease. In: *Cochrane Database of Systematic Reviews*; 2006.
- [129] Burns A, O'Brien J. Clinical practice with anti-dementia drugs: a consensus statement from british association for psychopharmacology. *J Psychopharmacol (Oxf)* 2006;20:732–55.
- [130] Farlow MR. Utilizing combination therapy in the treatment of alzheimer's disease. *Expert Rev Neurother* 2004;4:799–808.
- [131] McShane R, Sastre AA, Minakaran N. Memantine for dementia. In: *Cochrane Database of Systematic Reviews*; 2006.
- [132] Farlow MR, Miller ML, Pejovic V. Treatment options in alzheimer's disease: maximizing benefit, managing expectations. *Dement Geriatr Cogn Disord* 2008;25:408–22.
- [133] Mondadori CRA, Buchmann A, Mustovic H, Schmidt CF, Boesiger P, Nitsch RM, Hock C, Streffer J, Henke K. Enhanced brain activity may precede the diagnosis of alzheimer's disease by 30 years. *Brain* 2006;129:2908–22.
- [134] Gomez-Isla T, Price JL, McKeel DW, Morris JC, Growdon JH, Hyman BT. Profound loss of layer ii entorhinal cortex neurons occurs in very mild alzheimer's disease. *J Neurosci* 1996;16:4491–500.
- [135] Morris JC, Storandt M, McKeel DW, Rubin EH, Price JL, Grant EA, Berg L. Cerebral amyloid deposition and diffuse plaques in “normal” aging: evidence for presymptomatic and very mild alzheimer's disease. *Neurology* 1996;46:707–19.
- [136] Ingelsson M, Fukumoto H, Newell KL, Growdon JH, Hedley-Whyte ET, Frosch MP, Albert MS, Hyman BT, Irizarry MC. Early a $\beta$  accumulation and progressive synaptic loss, gliosis, and tangle formation in ad brain. *Neurology* 2004;62:925–31.
- [137] Boche D, Denham N, Holmes C, Nicoll JAR. Neuropathology after active a $\beta$ 42 immunotherapy: implications for alzheimer's disease pathogenesis. *Acta Neuropathol (Berl)* 2010;120.
- [138] Holmes C, Boche D, Wilkinson D, Yadegarfar G, Hopkins V, Bayer A, Jones RW, Bullock R, Love S, Neal JW, Zotova E, Nicoll JAR. Long-term effects of a $\beta$ 42 immunisation in alzheimer's disease: follow-up of a randomised, placebo-controlled phase i trial. *Lancet* 2008;372:216–23.
- [139] Josephs KA, Whitwell JL, Ahmed Z, Shiung MM, Weigand SD, Knopman DS, Boeve BF, Parisi JE, Petersen RC, Dickson DW, Jack CR. B-amyloid burden is not associated with rates of brain atrophy. *Ann Neurol* 2008;63:204–12.
- [140] van Helmond Z, Miners JS, Kehoe PG, Love S. Higher soluble amyloid- $\beta$  concentration in frontal cortex of young adults than in normal elderly or alzheimer's disease. *Brain Pathol* 2010:9999.
- [141] de la Torre JC. Vascular risk factor detection and control may prevent alzheimer's disease. *Ageing Res Rev* 2010;9:218–25.
- [142] Sun X, He G, Qing H, Zhou W, Dobie F, Cai F, Staufenbiel M, Huang LE, Song W. Hypoxia facilitates alzheimer's disease pathogenesis by up-regulating bace1 gene expression. *Proc Natl Acad Sci U S A* 2006;103:18727–32.
- [143] Zhang X, Zhou K, Wang RS, Cui JK, Lipton SA, Liao FF, Xu HX, Zhang YW. Hypoxia-inducible factor 1  $\alpha$  (hif-1  $\alpha$ )-mediated hypoxia increases bace1 expression and  $\beta$ -amyloid generation. *J Biol Chem* 2007;282:10873–80.
- [144] De Jong GI, Farkas E, Stienstra CM, Plass JRM, Keijser JN, de la Torre JC, Luiten PGM. Cerebral hypoperfusion yields capillary damage in the hippocampal ca1 area that correlates with spatial memory impairment. *Neuroscience* 1999;91:203–10.
- [145] DeKosky S. Early intervention is key to successful management of alzheimer disease. *Alzheimer Dis Assoc Disord* 2003;17:S99–104.
- [146] Jagust W. Molecular neuroimaging in alzheimer's disease. *NeuroRx* 2004;1:206–12.
- [147] Barton M, Yanagisawa M. Endothelin: 20 years from discovery to therapy. *Can J Physiol Pharmacol* 2008;86:485–98.
- [148] Wenzel RR, Fleisch M, Shaw S, Noll G, Kaufmann U, Schmitt R, Jones CR, Clozel M, Meier B, Luscher TF. Hemodynamic and coronary effects of the endothelin antagonist bosentan in patients with coronary artery disease. *Circulation* 1998;98:2235–40.
- [149] Kohan DE. Endothelin, hypertension and chronic kidney disease: new insights. *Curr Opin Nephrol Hypertens* 2010;19:134–9.
- [150] Givertz MM, Colucci WS, Lejemtel TH, Gottlieb SS, Hare JM, Slawsky MT, Leier CV, Loh E, Nicklas JM, Lewis BE. Study I. Acute endothelin a receptor blockade causes selective pulmonary vasodilation in patients with chronic heart failure. *Circulation* 2000;101:2922–7.
- [151] Kato T, Kassab S, Wilkins FC, Kirchner KA, Keiser J, Granger JP. Endothelin antagonists improve renal-function in spontaneously hypertensive rats. *Hypertension* 1995;25:883–7.
- [152] Ohlstein EH, Douglas SA, Ezekiel M, Gellai M. Antihypertensive effects of the endothelin receptor antagonist bq-123 in conscious spontaneously hypertensive rats. *J Cardiovasc Pharmacol* 1993;22:S321–4.
- [153] Williamson DJ, Wallman LL, Jones R, Keogh AM, Scroope F, Penny R, Weber C, Macdonald PS. Hemodynamic effects of bosentan, an endothelin receptor antagonist, in patients with pulmonary hypertension. *Circulation* 2000;102:411–8.
- [154] Chen SJ, Chen YF, Meng QC, Durand J, Dicarlo VS, Oparil S. Endothelin-receptor antagonist bosentan prevents and reverses hypoxic pulmonary hypertension in rats. *J Appl Physiol* 1995;79:2122–31.
- [155] Galie N, Rubin L, Hoeper M, Jansa P, Al-Hit H, Meyer G, Chiossi E, Kusic-Pajic A, Simonneau G. Treatment of patients with mildly symptomatic pulmonary arterial hypertension with bosentan (early study): a double-blind, randomised controlled trial [see comment]. *Lancet* 2008;371:2093–100.
- [156] Barton M. Endothelial dysfunction and atherosclerosis: endothelin receptor antagonists as novel therapeutics. *Curr Hypertens Rep* 2000;2:84–91.
- [157] Verhaar MC, Strachan FE, Newby DE, Cruden NL, Koomans HA, Rabelink TJ, Webb DJ. Endothelin-a receptor antagonist-mediated vasodilatation is attenuated by inhibition of nitric oxide synthesis and by endothelin-b receptor blockade. *Circulation* 1998;97:752–6.
- [158] Elesber AA, Bonetti PO, Woodrum JE, Zhu XY, Lerman LO, Younkin SG, Lerman A. Bosentan preserves endothelial function in mice overexpressing app. *Neurobiol Aging* 2006;27:446–50.
- [159] Davie NJ, Schermuly RT, Weissmann N, Grimminger F, Ghofrani HA. The science of endothelin-1 and endothelin receptor antagonists in the management of pulmonary arterial hypertension: current understanding and future studies. *Eur J Clin Invest* 2009;39:38–49.
- [160] Bates KA, Verdile G, Li QX, Ames D, Hudson P, Masters CL, Martins RN. Clearance mechanisms of alzheimer's amyloid- $\beta$  peptide: implications for therapeutic design and diagnostic tests. *Mol Psychiatry* 2009;14:469–86.
- [161] Nalivaeva NN, Fisk LR, Belyaev ND, Turner AJ. Amyloid-degrading enzymes as therapeutic targets in alzheimer's disease. *Curr Alzheimer Res* 2008;5:212–24.
- [162] Carty NC, Nash K, Lee D, Mercer M, Gottschall PE, Meyers C, Muzyczka N, Gordon MN, Morgan D. Adeno-associated viral (aav) serotype 5 vector mediated gene delivery of endothelin-converting enzyme reduces a beta deposits in app+ps1 transgenic mice. *Mol Ther* 2008;16:1580–6.
- [163] Nilsson P, Iwata N, Muramatsu S-i, Tjernberg LO, Winblad B, Saido TC. Gene therapy in alzheimer's disease – potential for disease modification. *J Cell Mol Med* 2010;14:741–57.
- [164] Zhang FY, Eckman C, Younkin S, Hsiao KK, Iadecola C. Increased susceptibility to ischemic brain damage in transgenic mice overexpressing the amyloid precursor protein. *J Neurosci* 1997;17:7655–61.
- [165] Suter OC, Sunthorn T, Kraftsik R, Straubel J, Darekar P, Khalili K, Miklossy J. Cerebral hypoperfusion generates cortical watershed microinfarcts in alzheimer disease. *Stroke* 2002;33:1986–92.
- [166] Selkoe DJ. Alzheimer's disease: genes, proteins, and therapy. *Physiol Rev* 2001;81:741–66.
- [167] Paton JFR, Dickinson CJ, Mitchell G. Harvey cushing and the regulation of blood pressure in giraffe, rat and man: introducing ‘cushing's mechanism’. *Exp Physiol* 2009;94:11–7.
- [168] Paton JFR, Waki H, Abdala APL, Dickinson J, Kasparov S. Vascular-brain signaling in hypertension: role of angiotensin ii and nitric oxide. *Curr Hypertens Rep* 2007;9:242–7.