

# Concomitant diagnosis of primary Sjögren's syndrome and systemic AL amyloidosis

I Delèvaux, M André, Z Amoura, J-L Kémény, J-C Piette, O Aumaitre

## Abstract

A 48 year old woman was referred to hospital for buccal discomfort. Physical examination showed a macroglossia and features of xerostomia. She was diagnosed as having primary Sjögren's syndrome according to the criteria proposed by the European Community study group in 1993. Furthermore, a lower lip salivary gland biopsy showed amyloid deposits that were also seen in the stomach and in the bone marrow. Echocardiography was consistent with cardiac amyloidosis. Serum immunofixation identified a monoclonal IgG $\lambda$ . As far as is known, this is the first report of systemic primary amyloidosis associated with Sjögren's syndrome. The relation between these two disorders is discussed.

(Ann Rheum Dis 2001;60:694-695)

Primary Sjögren's syndrome is an autoimmune disorder that exposes a patient to the development of lymphoproliferative disorders. The occurrence of amyloidosis in primary Sjögren's syndrome has rarely been reported: only a few cases of localised amyloidosis either with or without lymphoma are mentioned. We describe herein the case of a patient who presented with both primary Sjögren's syndrome and systemic AL amyloidosis.

## Case report

A 48 year old woman was admitted to our department for a tongue discomfort of three weeks' duration. She complained of xerostomia, xerophthalmia, and trigeminal dysaesthesia. Physical examination showed a macroglossia and mouth dryness.

Her white and red blood cell count was normal. Serum creatinine, C reactive protein, thyroid stimulating hormone, and  $\beta_2$  microglobulin were within the normal range. Serum concentrations of immunoglobulins were 23 g/l for IgG (normal 6.6-12.3), 0.7 g/l for IgA (normal 0.85-2.7), and 1.18 g/l for IgM (normal 0.62-1.74). Serum immunofixation showed a monoclonal IgG $\lambda$ . Anti-Ro/SSA antibodies were positive by enzyme linked immunosorbent assay (ELISA) (ENADOT bmd). Searches for cryoglobulin (home method) and markers for hepatitis C virus were negative.<sup>1</sup> A Schirmer test was positive and the break up time was abnormal. Histological examination of a minor salivary gland biopsy specimen (fig 1) showed an infiltration of glandular interstitium by lymphocytes and plasma cells. There were three inflammatory foci containing more than 50 cells in six lobules examined. According to the Chisholm scale, the patient had a biopsy grade 4. There were also amorphous deposits in the interstitium and in

Department of  
Internal Medicine, G  
Montpied hospital, BP  
69, 63003  
Clermont-Ferrand,  
France  
I Delèvaux  
M André  
O Aumaitre

Department of  
Pathology, G Montpied  
hospital  
J-L Kémény

Department of  
Internal Medicine,  
Pitié-Salpêtrière  
hospital, 47-83  
boulevard de l'Hôpital,  
75651 Paris cedex 13,  
France  
Z Amoura  
J-C Piette

Correspondence to:  
Professor O Aumaitre  
oaumaitre@  
chu-clermontferrand.fr

Accepted 15 November 2000




Figure 1 Labial salivary gland biopsy showing (A) two foci with more than 50 lymphocytes/mm<sup>2</sup> (haematoxylin and eosin, original magnification  $\times 80$ ); (B) amyloid deposits around the salivary ducts with green birefringence under polarised light microscopy after exposure to potassium permanganate (Congo red, original magnification  $\times 40$ ).

the vessel walls. After pretreatment with potassium permanganate, they were stained with Congo red and so were consistent with AL amyloid protein deposits. Immunohistochemical stains showed positive intracytoplasmic immunoreactivity of plasma cells for anti-IgG, IgM, and sometimes IgA with  $\kappa$  chain restriction. Bone marrow biopsy showed 7% of normal plasma cells; Congo red staining demonstrated amyloid deposits within vessel walls. Immunohistochemical studies showed that the cytoplasm of the plasma cells was stained with anti- $\kappa$  and anti- $\lambda$  light chains and with anti-IgA heavy chain antibodies. Thoracoabdominal computed tomography (CT) was normal. Endoscopic examination of the upper gastrointestinal tract was unremarkable, but amyloid deposits were found near the fundus glands on biopsy specimens. An electromyogram was normal. An electrocardiogram showed pseudo infarct Q waves on V5 to V6 leads and echocardiography an abnormal diastolic function. No lytic bone lesion was observed on radiographs.

Systemic AL amyloidosis associated with Sjögren's syndrome was diagnosed. High dose chemotherapy with cyclophosphamide followed by autologous stem cell transplantation was given to the patient. Three months after transplantation she improved, but the gammopathy and anti-Ro/SSA antibodies persisted.

## Discussion

Our patient complained of xerophthalmia and xerostomia. The diagnosis of primary Sjögren's syndrome was supported by a Schirmer test, the break up time, the positivity of anti-Ro/SSA antibodies, and the lower lip salivary gland biopsy. The presence of a macroglossia led us to search for an associated amyloidosis. The immunohistochemical examination of several tissues confirmed the presence of AL amyloid deposits. Salivary gland amyloid deposition may cause sicca syndrome, and minor salivary gland biopsy has been said to be a reliable test to diagnose amyloidosis.<sup>2</sup> Our patient had both true primary Sjögren's syndrome and primary systemic amyloidosis.<sup>3-4</sup> Amyloid deposits were found not only in the salivary gland but also in stomach and bone marrow specimens, and were probably responsible for macroglossia and heart abnormalities. An IgG $\lambda$  gammopathy was found by immunofixation, but the diagnosis of multiple myeloma was not allowed because bone marrow infiltration by plasma cells was lower than 10%. A hypothetical underlying lymphoma was ruled out by normal thoracoabdominal CT scan and bone marrow biopsy.

Unlike the well known association of Sjögren's syndrome with low grade marginal zone B cell lymphoma,<sup>5-6</sup> the coexistence of primary amyloidosis and Sjögren's syndrome is rare. Only eight cases have been previously reported in the English literature. In all these published cases, amyloidosis was localised or limited to dermis,<sup>7-9</sup> lung,<sup>10-11</sup> or tongue.<sup>12</sup> In some cases, plasma cells were found surrounding the amyloid deposits. Monotypic or polypotypic light


chains, thought to be the precursors of amyloidosis, were discovered in these cells and in amyloid deposits.<sup>12-13</sup> In one case a localised cutaneous amyloidosis with a monoclonal gammopathy was found, but there was no multiple myeloma.<sup>7</sup>

To our knowledge, primary systemic amyloidosis has not been reported so far in association with Sjögren's syndrome. One case of primary nodal plasmacytoma with Sjögren's syndrome has been described but without amyloidosis.<sup>14</sup> Wong *et al* reported on a patient with secondary amyloidosis (amyloid protein AA) of the lung during Sjögren's syndrome.<sup>15</sup>

Our patient differed from these cases because of the presence of systemic amyloidosis and the slight plasma cell infiltration on bone marrow biopsy. Even if extensive investigations have not been performed in previous reports of patients with localised amyloidosis and Sjögren's syndrome, it should be noted that none of them developed systemic amyloid involvement or plasma cell proliferation.

Although a chance association cannot be firmly excluded, this case report shows that the spectrum of lymphoid proliferations associated with Sjögren's syndrome extends beyond the classical B cell lymphoma. Staining for amyloidosis should be performed in patient with Sjögren's syndrome and unexplained findings, such as a macroglossia.

- Romaszko JP, Tridon A, Jouanel P, Subtil E, Dibet P, Betail G. Détection et analyse des cryoglobulines: étude comparative d'une population de malades et d'une population de témoins sains. *Pathol Biol* 1993;41:525-9.
- Hachulla E, Janin A, Flipo RM, Saile R, Facon T, Bataille D, *et al*. Labial salivary gland biopsy is a reliable test for the diagnosis of primary and secondary amyloidosis. A prospective clinical and immunohistologic study in 59 patients. *Arthritis Rheum* 1993;36:691-7.
- Myssiorek D, Alvi A, Bhuiya T. Primary salivary gland amyloidosis causing sicca syndrome. *Ann Otol Rhinol Laryngol* 1992;101:487-90.
- Schlesinger I. Multiple myeloma and AL amyloidosis mimicking Sjögren syndrome. *South Med J* 1993;86:568-9.
- Mariette X. Lymphomas in patients with Sjögren's syndrome: review of the literature and physiopathologic hypothesis. *Leuk Lymphoma* 1999;33:93-9.
- Royer B, Cazals-Hatem D, Sibilia J, Agbalika F, Cayuela JM, Soussi T, *et al*. Lymphomas in patients with Sjögren's syndrome are marginal zone B-cell neoplasms, arise in diverse extranodal and nodal sites, and are not associated with viruses. *Blood* 1997;7:66-75.
- Kitajima Y, Seno J, Aoki S, Tada S, Yaoita H. Nodular primary cutaneous amyloidosis. *Arch Dermatol* 1986;122:1425-30.
- Pablos JL, Cogolludo V, Pinedo F, Carreira PE. Subcutaneous nodular amyloidosis in Sjögren's syndrome. *Scand J Rheumatol* 1993;22:250-1.
- Inazumi T, Hakuno M, Yamada H, Tanaka M, Naka W, Tajima S, *et al*. Characterisation of the amyloid fibril from primary localized cutaneous nodular amyloidosis associated with Sjögren's syndrome. *Dermatology* 1994;189:125-8.
- Kobayashi H, Matsuoka R, Kitamura S, Tsunoda N, Saito K. Sjögren's syndrome with multiple bullae and pulmonary nodular amyloidosis. *Chest* 1988;94:438-40.
- Milburn JM, Kay D, Ridpat C. Pulmonary nodular amyloidosis in a patient with Sjögren's syndrome diagnosed by transthoracic biopsy. *J La State Med Soc* 1994;146:395-8.
- Haraguchi H, Ohashi K, Yamada M, Hasegawa M, Maeda S, Komatsuzaki A. Primary localized nodular tongue amyloidosis associated with Sjögren's syndrome. *ORL J Otorhinolaryngol Relat Spec* 1997;1:60-3.
- Ishihara T, Takahashi M, Koga M, Yokota T, Yamashita Y, Uchino F. Amyloid fibril formation in the rough endoplasmic reticulum of plasma cells from a patient with localized AL amyloidosis. *Lab Invest* 1991;64:265-71.
- Watanabe K, Yatabe Y, Narita M, Ogawa T, Mori N. An autopsy case of primary nodal plasmacytoma associated with Sjögren's syndrome. *Pathol Int* 1999;49:577-9.
- Wong BC, Wong KL, Ip MS, Wang EP, Chan KW, Cheng LC. Sjögren's syndrome with amyloid A presenting as multiple pulmonary nodules. *J Rheumatol* 1994;21:165-7.



# Concomitant diagnosis of primary Sjögren's syndrome and systemic AL amyloidosis

I Delèvaux, M André, Z Amoura, J-L Kémény, J-C Piette and O Aumaître

*Ann Rheum Dis* 2001 60: 694-695  
doi: 10.1136/ard.60.7.694

---

Updated information and services can be found at:  
<http://ard.bmj.com/content/60/7/694>

---

	<i>These include:</i>
<b>References</b>	This article cites 15 articles, 1 of which you can access for free at: <a href="http://ard.bmj.com/content/60/7/694#BIBL">http://ard.bmj.com/content/60/7/694#BIBL</a>
<b>Email alerting service</b>	Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

---

## Topic Collections

Articles on similar topics can be found in the following collections

[Genetics](#) (896)  
[Pathology](#) (413)  
[Clinical diagnostic tests](#) (1194)  
[Immunology \(including allergy\)](#) (4673)  
[Radiology](#) (1040)  
[Radiology \(diagnostics\)](#) (704)  
[Surgical diagnostic tests](#) (402)

---

## Notes

---

To request permissions go to:  
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:  
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:  
<http://group.bmj.com/subscribe/>