

Identical But Not the Same: The Value of Discordant Monozygotic Twins in Genetic Research

Petra J.G. Zwijsenburg,^{1*} Hanne Meijers-Heijboer,¹ and Dorret I. Boomsma²

¹Department of Clinical Genetics, VU University Medical Center, Amsterdam, the Netherlands

²Department of Biological Psychology, VU University, Amsterdam, the Netherlands

Received 6 November 2008; Accepted 4 March 2010

Monozygotic (MZ) twins show remarkable resemblance in many aspects of behavior, health, and disease. Until recently, MZ twins were usually called “genetically identical”; however, evidence for genetic and epigenetic differences within rare MZ twin pairs has accumulated. Here, we summarize the literature on MZ twins discordant for Mendelian inherited disorders and chromosomal abnormalities. A systematic literature search for English articles on discordant MZ twin pairs was performed in Web of Science and PubMed. A total number of 2,016 publications were retrieved and reviewed and 439 reports were retained. Discordant MZ twin pairs are informative in respect to variability of phenotypic expression, pathogenetic mechanisms, epigenetics, and post-zygotic mutagenesis and may serve as a model for research on genetic defects. The analysis of single discordant MZ twin pairs may represent an elegant approach to identify genes in inherited disorders. © 2010 Wiley-Liss, Inc.

Key words: twins; monozygotic; twin studies; discordant; chromosome disorders; genetic diseases; inborn

TWINNING

Twins and multiples always have intrigued humans, as illustrated by numerous legends, myths, tales, and anecdotes in which twins play a principal part.

Twins can result from a single ovum, fertilized by one sperm, and are then called monozygotic (MZ) or identical twins. Dizygotic (DZ) or fraternal twins result from two different ova, fertilized by two different sperm. Hence, MZ twins originate from one zygote and are considered to be genetically identical, whereas DZ twins originate from two zygotes and share on average half of the genetic variance present in unrelated individuals (like non-twin sibling pairs). Triplets and higher order multiples can consist of any combination of MZ individuals (including none) together with “additional siblings.”

The prevalence of twin births varies between populations, in particular due to differences in the DZ twinning rate. The highest prevalence of twinning is found in Nigeria, where 1 in every 12 persons is a member of a twin pair, and the lowest twinning rates are found in Asia (8/1,000 births in China) [Gan et al., 2007; Hoekstra et al., 2008]. In Europe and the USA, the incidence of twinning is roughly 1 twin pair in every 60 births [Hall, 2003;

How to Cite this Article:

Zwijsenburg PJG, Meijers-Heijboer HEJ, Boomsma DI. 2010. Identical But Not the Same: The Value of Discordant Monozygotic Twins in Genetic Research.

Am J Med Genet Part B 153B:1134–1149.

Hoekstra et al., 2008]. Within Europe and the United States, twinning rates also can vary considerably across time and place [Hoekstra et al., 2008]. As MZ twinning generally occurs at a constant rate of 3–4 per 1,000 maternities around the world [Tong et al., 1997; Hall, 2003], the variation in twinning rates is generally accepted as being the result of variation in DZ twinning rates. The rates of DZ, and possibly also MZ twinning, have increased the past decades, in part due to assisted reproductive technologies (ART) [Chang et al., 2009] and due to increases in maternal age.

The causes of MZ twinning in humans are unknown. Until recently, the process of MZ twinning was considered to represent a random process. However, new data indicate that (epi)genetic mechanisms may be involved in the phenomenon of splitting of the zygote [Shur, 2009]. An excess of females over males has been observed in spontaneous, surviving, MZ twins. This female predominance is observed in particular in monochorionic diamniotic MZ twins, and even more pronounced among monochorionic monoamniotic twins and conjoined twins, suggesting a relationship with the timing of twinning [James, 1975, 1980; Hall, 2003]. Skewed X-inactivation has been suggested to contribute to the process of twinning and as such to the observation of female predominance [Hall, 2003].

Familial MZ twinning has long been recognized [Machin, 2009a] and the MZ twinning rate in ART-induced pregnancies seems to be

*Correspondence to:

Dr. Petra J.G. Zwijsenburg, Department of Clinical Genetics, VU University Medical Center, de Boelelaan 1117, 1081 HV Amsterdam, the Netherlands. E-mail: p.zwijsenburg@vumc.nl

Published online 28 April 2010 in Wiley Online Library (wileyonlinelibrary.com).

DOI 10.1002/ajmg.b.31091

elevated [Chang et al., 2009; Shur, 2009]. Damage to the inner cell mass has been suggested to contribute to twinning in human beings and both familial twinning and ART may be associated with an increased risk of damage to the inner cell mass [Hall, 2003].

Furthermore, and of interest in the context of this review, it has been suggested that genetic differences between cells of the early blastomere may trigger division of the zygote and initiate the process of twinning [Hall, 1996a,b, 2003]. As such, a post-zygotic genetic event may trigger the origin of an MZ twin. In line with this assumption is the observation of an increased prevalence of certain genetic syndromes among MZ twins as compared to singletons, such as Beckwith–Wiedemann syndrome (BWS) [Weksberg et al., 2002; Blik et al., 2009; Shur, 2009]. Although MZ twin concordance for BWS has been described, most MZ twin pairs are discordant. It has been suggested that unequal splitting of the inner cell mass results in differential methylation between the two cell masses or alternatively that a lack of methylation maintenance in the early zygote results in splitting of the zygote. In both cases a difference in methylation between the two MZ twins may account for phenotypic discordance. Thus, the role of DNA methylation in MZ twinning needs further elucidation.

In the past, zygosity in same-sex twin pairs was mainly determined by blood group and HLA typing, dermatoglyphic studies, and comparison of physical characteristics. Nowadays, DNA polymorphisms are increasingly used (microsatellite markers and/or other polymorphisms such as single-nucleotide polymorphisms (SNPs)) [Hall, 1996b, 2003], although in large-scale epidemiological projects, zygosity still often is reliably determined on a series of questionnaire items [Rietveld et al., 2000]. Correct determination of zygosity is important for individual twin pairs with respect to genetic counseling, preventive medical strategies, or surveillance when one twin has a disease, and to offer optimal guidance in case of organ transplantation [Machin, 2009b]. Obviously, correct determination of zygosity is crucial in research projects involving twins. Methods of zygosity testing are extensively reviewed elsewhere [Machin, 2009b].

TWIN RESEARCH

From the late 19th century onwards, twin studies played an important role in genetics. In classical twin studies, comparison of the resemblance (concordance rates or correlations) for a disorder or a quantitative trait between MZ and DZ twins facilitates research into the etiology of population variation. The heritability (standardized estimate of the contribution of genetic effects of a (complex) trait) can be estimated through fitting quantitative genetic models to twin data. As such, the results of twin studies are applicable to the whole population the twins originated from and have contributed enormously to current genetic insights. For example, multiple sclerosis (MS) was thought to be a non-genetic disorder until a large study was done in Canadian twins [Martin et al., 1997].

DISCORDANT MZ TWINS

Most MZ twins are remarkably similar and MZ twin pairs are in general expected to be concordant for congenital malformations,

chromosomal abnormalities, and Mendelian disorders. However, an increasing number of reports are being published on discordant twin pairs. We performed a systematic literature search in Web of Science and PubMed for English language articles on discordant MZ twin pairs, published before November 1, 2009. The terms used for the PubMed search were: “Monozyg* OR Twins[mesh] OR “Twin Studies as Topic”[Mesh] AND Discordan*[tiab]” and the criteria used for Web of Science were topic = (monozyg*) AND topic = (discordan*). A total of 2,016 publications were primarily screened by P.Z. Four hundred thirty-nine papers concerned reports on individual discordant MZ twin pairs. References were also obtained from articles found through the literature search. It should be noted that some older publications lack the appropriate criteria for establishing monozygosity. Determination of zygosity may be limited in older publications, especially in single discordant pairs [Machin, 2009b]; however, for the purpose of this review we generally assumed that the zygosity status was correctly described.

In this journal, 81 publications on individual discordant MZ twin pairs have been published over the last years. The MZ twin pairs in these reports are informative either because both twins are affected, with a different phenotype in each twin, or because only one twin from an MZ twin pair is affected. Our fascination for twins and the educational value of discordant MZ twins is illustrated by three reports that appear to be based on a one single MZ twin pair [Michalski et al., 1978; Schlegelberger et al., 1984, 1986] discordant for Klinefelter syndrome.

DISCORDANT PHENOTYPES IN MZ TWINS

Tables I and II provide an overview of the publications, which report a difference in phenotype between members from MZ twin pairs. Numerous monogenetic and chromosomal disorders are characterized by significant variability that can be observed between unrelated individuals, sibs, and twins. Since affected sibs and DZ twins share on average only 50% of their segregating genes,

TABLE I. Overview of Publications on Monozygotic Twin Pairs With a Chromosomal Abnormality That Report a Difference in Phenotype Between Both Twins

| Chromosomal abnormality | Refs. |
|-------------------------|---------------------------------------------------------------------|
| del 7q | Tsukamoto et al. [1993] |
| del 8p23.1 | Wat et al. [2009] |
| dup 11p mosaicism | Marcus-Soekarman et al. [2004] |
| dup 4q + del 1p36 | Angle et al. [2002] |
| dup 4q28.3-qter | Celle et al. [2000] |
| inv dup(15) | Peng et al. [2007] |
| Pseudic 13 | Lynch et al. [1995] |
| Trisomy 13 | Loevy et al. [1985], Naor et al. [1987] |
| Trisomy 18 | Lee et al. [2004], Mulder et al. [1989], Schlessel et al. [1990] |
| Trisomy 21 | Beattie et al. [1993], Grynberg et al. [2007] |

del, deletion; dup, duplication; inv dup, inverted duplication; psu dic, pseudodicentric chromosome.

TABLE II. Overview of Publications on Monozygotic Twin Pairs With a Monogenetic Disorder That Report a Difference in Phenotype Between Both Twins

| Disorder | MIM number ^a | Refs. |
|-----------------------------------------------|---------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------|
| 22q11-deletion syndrome | #611867, #192430, #188400 | Fryer [1996], Goodship et al. [1995], Hillebrand et al. [2000], Rauch et al. [1998], Singh et al. [2002], Vincent et al. [1999], Yamagishi et al. [1998] |
| D-2-hydroxyglutaric aciduria | #600721 | Misra et al. [2005] |
| Alagille syndrome | #118450 | Kamath et al. [2002] |
| Alport syndrome | #301050 | Matsukura et al. [2004] |
| Anophthalmia-esophageal atresia | #206900 | Zenteno et al. [2006] |
| Cystic fibrosis | #219700 | Mekus et al. [2000], Picci et al. [2007] |
| Crouzon syndrome | #123500 | Lajeunie et al. [2000], Sher et al. [2008] |
| Dravet syndrome | #607208 | Miyama et al. [2008] |
| Familial amyloid polyneuropathy | +176300 | Holmgren et al. [2004], Munar-Ques et al. [1999], Saporta et al. [2009] |
| Fragile X syndrome | #300624 | Helderman-van den Enden et al. [1999], Kruger et al. [1994] |
| Frijns syndrome | %229850 | Vargas et al. [2000] |
| Facioscapulohumeral muscular dystrophy (FSHD) | %158900 | Tawil et al. [1993] |
| G syndrome | %145410 | Young et al. [1988] |
| Gaucher disease | #231000 | Lachmann et al. [2004] |
| Gerstmann–Straussler disease | #137440 | Webb et al. [2009] |
| Huntington disease | #143100 | Friedman et al. [2005], Panas et al. [2008], Anca et al. [2004], Georgiou et al. [1999] |
| Joubert syndrome | %213300 | Raynes et al. [1999] |
| Kallmann syndrome | +308700 | Hipkin et al. [1990] |
| Keratoconus | #148300 | McMahon et al. [1999], Weed et al. [2006] |
| Langerhans histiocytosis | 604856 | Chen et al. [2004] |
| LEOPARD syndrome | #151100 | Rudolph et al. [2001] |
| Lymphedema-distichiasis | #153400 | Kumar et al. [2007] |
| McCune–Albright syndrome | #174800 | Endo et al. [1991] |
| Neurofibromatosis type 1 | #162200 | Kelly et al. [1998], Detjen et al. [2007] |
| Neurofibromatosis type 2 | #101000 | Baser et al. [1996] |
| Goldenhar syndrome | %164210 | Ryan et al. [1988], Satoh et al. [1995] |
| Pai syndrome | 155145 | Al-Mazrou et al. [2001], Guion-Almeida et al. [2007] |
| Photosensitive epilepsy | %609569 | de Haan et al. [2005] |
| Primary ciliary dyskinesia | #242650 | Noone et al. [1999] |
| Severe combined immunodeficiency (SCID) | #102700+ | Niehues et al. [1996] |
| Sickle cell anemia | #603903 | Amin et al. [1991] |
| Thanatophoric dysplasia | #187601/#187600 | Corsello et al. [1992], Horton et al. [1983], Horton et al. [1983] |
| Tibial hemimelia with ectrodactyly | %119100 | Dayer et al. [2007] |
| Tricho-rhino-phalangeal syndrome | #190350 | Naselli et al. [1998] |
| Tuberous sclerosis | #190350/#191100 | Humphrey et al. [2004], Kondo et al. [1991], Martin et al. [2003] |
| WAGR syndrome | #194072 | Bremond-Gignac et al. [2005] |
| Williams syndrome | #194050 | Castorina et al. [1997], Maurer et al. [1979], Murphy et al. [1990], Oorthuys [1984], Pankau et al. [2001] |
| Wilson disease | #277900 | Czlonkowska et al. [2009] |
| (X-linked) adrenoleukodystrophy | #300100 | Di et al. [2001], Sobue et al. [1994] |

Symbols preceding a MIM number: #, a descriptive phenotype, not necessarily representing a unique locus; +, description of a gene of known sequence and a phenotype; %, a confirmed Mendelian phenotype or phenotypic locus for which the underlying molecular basis is not known; no symbol before a MIM number generally indicates a description of a phenotype for which the Mendelian basis, although suspected, has not been clearly established.^aMendelian Inheritance in Man (MIM) numbers correspond to Online Mendelian Inheritance in Man, OMIM (TM). McKusick-Nathans Institute of Genetic Medicine, Johns Hopkins University (Baltimore, MD) and National Center for Biotechnology Information, National Library of Medicine (Bethesda, MD) (November 1, 2009). World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>.

phenotypic differences between them may be caused by non-shared genetic variation. Therefore, studying the variability of genetic disorders in MZ twins can be very valuable. Discordant MZ twin pairs have highlighted the possible variability of some diseases, sometimes making us aware of the previously unrecognized ends of the spectrum of disease, such as in Frijns syndrome [Vargas et al., 2000]. An MZ twin pair concordant for a number of dysmorphic features, but discordant for a diaphragmatic defect was described. Twin A came to clinical attention in the context of his twin brother's more obvious clinical characteristics that led to the diagnosis of Frys syndrome in both twins. At 1 year of age Twin A had mild developmental delay, whereas his twin brother died on day 5. This MZ twin pair indicated that clinically significant diaphragmatic defects are not invariably present, and developmental outcome can be mild in Frijns syndrome, suggesting that the clinical spectrum of this disorder is broader than previously suspected.

A considerable part of research in human genetics is focused on finding explanations for variability in genetically determined diseases, and discordant MZ twin pairs are a valuable source of information in this respect. Numerous reports on MZ twins with 22q11 deletion syndrome, Joubert syndrome, neurofibromatosis, Williams syndrome, and many other diseases with discordant phenotypes emphasize that non-genetic factors may affect the phenotypic variability (see Table II).

A single phenotypically discordant MZ twin pair can contribute to our understanding of the pathogenesis of a disease. Recently, by studying one MZ twin pair, one child with preleukemic and the other one with frank childhood precursor B-cell acute lymphoblastic leukemia, the lineal relationship between several events in carcinogenesis was established [Hong et al., 2008].

Furthermore, discordant MZ twins can guide us toward different manifestations of a genetically determined syndrome. Germline mutations of the gene encoding succinate dehydrogenase subunit B (*SDHB*) are associated with inherited paraganglioma. Boccon-Gibod et al. [2004] and Pasini et al. [2008] described MZ twin brothers of whom one underwent surgery for a paraganglioma and the other for a stomach gastrointestinal stromal tumor (GIST). They both carried a germline mutation of the *SDHB* gene and subsequent studies in other patients confirmed an association between germline mutations in *SDHB* and GISTs. A careful medical and family history regarding gastrointestinal symptoms and/or tumors is now suggested in patients with germline mutations in *SDHB*.

Finally, MZ twin pairs concordant for a disease, but discordant for prescribed therapy [Davin et al., 2006] or discordant for compliance [Verougstraete et al., 1999] may serve as a model for the effect of therapy. Comparison of the course of the disease in such twin pairs is informative since many other factors that determine the course of the disease are shared by MZ twins.

There have been a number of reports on MZ twin pairs with a chromosomal abnormality with a significant difference in phenotype. Some of these articles describe severely unbalanced chromosomal patterns in blood from healthy children or adults [Celle et al., 2000; Marcus-Soekarman et al., 2004; O'Donnell et al., 2004; Bourthoumieu et al., 2005]. Chimerism, a condition in which an individual is composed of two genetically different types of cells, originating from more than one zygote, has been demonstrated

extensively in DZ twins [van Dijk et al., 1996; Abuelo, 2009]. Blood and buccal DNA likely reflects (in part) "donor" DNA in case of chimerism [Thiede et al., 2000; Hong et al., 2007]. The finding of unbalanced chromosomal patterns in healthy MZ co-twins of affected individuals suggests hematopoietic chimerism between the affected and unaffected twin, as in some cases mentioned by the authors.

MZ TWINS DISCORDANT FOR CHROMOSOMAL ABNORMALITIES AND MENDELIAN DISORDERS

Tables III and IV summarize studies of MZ twin pairs where one twin is considered to have a (possibly) Mendelian disorder or chromosomal abnormality and the other twin is unaffected. In the past, MZ twins were considered to be genetically identical. However, it has become increasingly clear that some MZ twins are not fully genetically identical. Figure 1 summarizes possible post-zygotic genetic effects that may account for MZ twin discordance. Obviously, it is important to realize that discordant MZ twins may become concordant over time. In 1993, an MZ twin pair discordant for Leber's hereditary optic neuropathy was reported [Johns et al., 1993]. However, a few years later in 1998 the unaffected twin had become affected as well [Lam, 1998].

MZ twins discordant for aneuploidy have been recognized for several decades; discordant MZ twins have been reported for monosomy X, trisomy 1, trisomy 13, and trisomy 21 [Rogers et al., 1982; Weiss et al., 1982; Gilgenkrantz and Janot, 1983; Uchida et al., 1983; Reiss et al., 1993; Nieuwint et al., 1999; O'Donnell et al., 2004; Rohrer et al., 2004; Cheng et al., 2006; Sethupathy et al., 2007; Dahoun et al., 2008; Gentilin et al., 2008; Taylor et al., 2008]. Moreover, unbalanced chromosomal abnormalities have been described in one member from an MZ twin pair [Marcus-Soekarman et al., 2004; Bourthoumieu et al., 2005].

TABLE III. Overview of Publications on Monozygotic Twin Pairs of Whom Only One Twin Is Considered to Have a Chromosomal Abnormality

| Chromosomal abnormality | Refs. |
|-------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 45,X | Gentilin et al. [2008], Nieuwint et al. [1999], Reiss et al. [1993], Rohrer et al. [2004], Uchida et al. [1983], Weiss et al. [1982] |
| del(10) | Juberg et al. [1981] |
| der(11) | Bourthoumieu et al. [2005] |
| Trisomy 1 (partial) | Watson et al. [1990] |
| Trisomy 13 | Taylor et al. [2008] |
| Trisomy 21 | Cheng et al. [2006], Dahoun et al. [2008], Gilgenkrantz and Janot [1983], Nieuwint et al. [1999], O'Donnell et al. [2004], Rogers et al. [1982], Sethupathy et al. [2007] |

del, deletion; der, derivative chromosome.

TABLE IV. Overview of Publications on Monozygotic Twin Pairs of Whom Only One Twin Is Considered to Have a (Possibly) Mendelian Disorder

| Disorder | MIM ^a | Refs. |
|-----------------------------------------------|-------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Aglossia-adactylia | 103300 | Robinow et al. [1978] |
| Aicardi syndrome | %304050 | Costa et al. [1997] |
| Asplenia syndrome | %208530 | Hwang et al. [2006], Wilkinson et al. [1979] |
| Bladder exstrophy | %600057 | Bugge [1981], Reutter et al. [2003] |
| Body stalk anomaly | 230750 | Daskalakis and Nicolaides [2002], Vidaeff et al. [2005] |
| Beckwith—Wiedemann syndrome | #122470 | Berry et al. [1980], Bose et al. [1985], Chien et al. [1990], Clayton-Smith et al. [1992], Leonard et al. [1996], Litz et al. [1988], Olney et al. [1988], Orstavik et al. [1995], Weksberg et al. [2002], Bliek et al. [2009] |
| Caudal duplication anomaly | #607864 | Kroes et al. [2002], Oates et al. [2006] |
| Congenital hypothyroidism | #275200/#218700 | Rettig et al. [1980] |
| Colorblindness | +303800 | Jorgensen et al. [1992] |
| Complete congenital heartblock | 234700 | Cooley et al. [1997] |
| Congenital bilateral perisylvian syndrome | #300388 | Lenti and Triulzi [1996] |
| Cornelia de Lange syndrome | #122470 | Carakushansky and Berthier [1976], Carakushansky et al. [1996] |
| Darrier's disease | #124200 | Sakuntabhai et al. [1999] |
| Duane's retraction | #604356, +%126800 | Kaufman et al. [1989], Rosenbaum and Weiss [1978] |
| Duchenne muscular dystrophy | #310200 | Burn et al. [1986], Chutkow et al. [1987], Gomez et al. [1977], Lupski et al. [1991], Pena et al. [1987], Richards et al. [1990], Tremblay et al. [1993], Zneimer et al. [1993], Bonilla et al. [1990] |
| Epilepsy with structural brain abnormalities | | Briellmann et al. [1998], Brodtkorb et al. [2000], Kuzniecky et al. [1995], Sisodiya et al. [1999], Supprian et al. [2000] |
| Fabry disease | #301500 | Redonnet-Vernhet et al. [1996] |
| Fragile X syndrome | #300624 | Kruger et al. [1994], Tuckerman et al. [1985] |
| Frontonasal dysplasia | 136760 | Mohammed et al. [2004], Wu et al. [2007] |
| Facioscapulohumeral muscular dystrophy (FSHD) | %158900 | Griggs et al. [1995], Tawil et al. [1993] |
| Goldenhar syndrome | %164210 | Boles et al. [1987], Stoll et al. [1984], Touliatou et al. [2006], Verona et al. [2006], Wieczorek et al. [2007] |
| Growth hormone deficiency | +139250 | Simpson et al. [1999] |
| Hemihypertrophy | %235000 | West et al. [2003] |
| Hemophilia A | +306700 | Bennett et al. [2008] |
| Hemophilia B | #306900 | Kitchens [1987], Revesz et al. [1972] |
| Holoprosencephaly | %236100 | Machin et al. [1985], Peng et al. [2007] |
| Humero-radial synostosis | %236400/143050 | McCredie [1975] |
| Hunter syndrome | +309900 | Winchester et al. [1992] |
| Idiopathic torsion dystonia (ITD) | #128100/%602124 | Chan and Tsui [1997] |
| Kabuki syndrome | %147920 | Shotelersuk et al. [2002] |
| Kearns—Sayre syndrome | #530000 | Blakely et al. [2004] |
| Klippel—Feil syndrome | %118100 | Toyoshima et al. [2006] |
| Klippel—Trenaunay syndrome | %149000 | Hofer et al. [2005], Oduber et al. [2010] |
| Landau—Kleffner syndrome | 245570 | Feekery et al. [1993] |
| Leber's hereditary optic atrophy | #535000 | Biousse et al. [1997], Johns et al. [1993] |
| Lesch—Nyhan syndrome | #300322 | De et al. [2005] |
| Mayer Rokitansky Küster syndrome | %277000 | Heidenreich et al. [1977], Duru and Laufer [2009] |
| McCune—Albright syndrome | #174800 | Peleg et al. [2009] |
| Ocular cicatricial pemphigoid | 164185 | Bhol et al. [1995] |
| Oculo-oto-radial (IVIC) syndrome | #147750 | Elcioglu and Berry [1997] |
| Oto-palato-digital syndrome | #311300 | Robertson et al. [2006] |
| Oral—facial—digital syndrome | #311200 | Shotelersuk et al. [1999] |
| Primary lateral sclerosis | %611637 | Sorenson [2006] |
| Primary progressive aphasia | #607485 | Doran and Larner [2004] |
| Proteus syndrome | %176920 | Brockmann et al. [2008] |
| Retinitis pigmentosa | #180100 + others | Bernstein and Aptsiauri [2003] |
| Rett syndrome | #312750 | Carter et al. [2008], Migeon et al. [1995], Subramaniam et al. [1997] |
| Rubinstein—Taybi syndrome | #180849 | Kajii et al. [1981] |
| Say syndrome | 181180 | Ashton-Prolla and Felix [1997] |

(Continued)

TABLE IV. (Continued)

| Disorder | MIM ^a | Refs. |
|------------------------------------------|------------------|---------------------------------------------------------------------------------------|
| Schimmelpenning–Feuerstein–Mims syndrome | %163200 | Schworm et al. [1996] |
| Silver-Russell syndrome | %180860 | Bailey et al. [1995], Sagot et al. [1996], Samn et al. [1990], Yamazawa et al. [2008] |
| Sotos syndrome | #117550 | Brown et al. [1998] |
| Spondylocostal dysostosis | #277300 | Van Thienen and Van der Auwera [1994] |
| Thumb polydactyly | #174400 | Peterson and Rayan [2004] |
| VACTERL | 192350 | Becker et al. [2005], Camacho et al. [2008] |
| Van der Woude syndrome | #119300 | Kondo et al. [2002] |
| X-linked adrenoleukodystrophy | #300100 | Korenke et al. [1996] |
| X-linked hypophosphataemic rickets | #307800 | Owen et al. [2009] |

Symbols preceding a MIM number: #, a descriptive phenotype, not necessarily representing a unique locus; +, description of a gene of known sequence and a phenotype; %, a confirmed Mendelian phenotype or phenotypic locus for which the underlying molecular basis is not known; no symbol before a MIM number generally indicates a description of a phenotype for which the Mendelian basis, although suspected, has not been clearly established.^aMendelian Inheritance in Man (MIM) numbers correspond to Online Mendelian Inheritance in Man, OMIM (TM). McKusick-Nathans Institute of Genetic Medicine, Johns Hopkins University (Baltimore, MD) and National Center for Biotechnology Information, National Library of Medicine (Bethesda, MD) [November 1, 2009]. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>.

Discordant genotypes due to point mutations in one twin have been identified in a few MZ twin pairs. First, in a pair of MZ twins discordant for Darier's disease, a point mutation in *ATP2A2* was identified in the affected twin only [Sakuntabhai et al., 1999]. Furthermore, a mutation in *IRF6* was found in the affected twin from an MZ twin pair discordant for Van der Woude syndrome [Kondo et al., 2002] and discordance for a point mutation in *FLNA* explains the occurrence of otopalatodigital syndrome in one twin from an MZ twin pair [Robertson et al., 2006]. Mosaicism of a mutation in the *COL4A5* gene in the affected twin was suggested to cause discordance of Alport syndrome in an MZ twin pair [Matsukura et al., 2004]. These reports highlight the potential value of discordant MZ twins in identifying new genes [Mansilla et al., 2005]. For complex traits, this approach has not yet seen large-scale application for gene-finding so far.

It has been hypothesized that the inheritance of a single recessive mutated allele of a gene followed by a somatic mutation in the normal allele during critical periods of development in the affected twin may result in discordance for complex diseases in MZ twins [Guidry and Kent, 1999; Singh et al., 2009].

Discordance for many other genetic abnormalities has been demonstrated in the past years. Uniparental disomy (UPD) may occur in only twin from an MZ pair [West et al., 2003;

Smith et al., 2006]. In these cases, a post-zygotic mitotic recombination error is believed to result in exclusively paternal origin of a segment of chromosome 11p, resulting in BWS or hemihypertrophy. To our knowledge, no reports are available on discordance for whole chromosome UPD within MZ twin pairs.

Discordance for a triplet repeat expansion length may cause a difference in expression, as illustrated for fragile-X syndrome [Kruyer et al., 1994; Helderman-van den Enden et al., 1999]. Discordance for a phenotype may be the result of different levels of (unrecognized) mosaicism between the two twins. This phenomenon has been demonstrated for monogenetic diseases [Norremolle et al., 2004] and chromosomal abnormalities (45,X). Different rates of mosaicism in 45,X/46,XY or 46,XX/46,XY mosaicism may even result in MZ twins of opposite sex [Schmidt et al., 1976; Schmid et al., 2000; Dallapiccola et al., 1985; Reindollar et al., 1987; Fujimoto et al., 1991; Kurosawa et al., 1992; Costa et al., 1998; Somkuti et al., 2000; Wachtel et al., 2000; Tho et al., 2007; Zech et al., 2008]. Likewise, 45,X mosaicism may result in discordance for symptomatic X-linked disorders, such as Duchenne muscular dystrophy, in women [Bonilla et al., 1990].

In diseases caused by mitochondrial DNA mutations, different levels of heteroplasmy are suggested to be the cause of discordance for Leber's hereditary optic atrophy [Biousse et al., 1997] and




FIG. 1. Possible triggers for splitting of the zygote [left] and [epi]genetic mechanisms that may cause MZ twin discordance [right].

chronic progressive external ophthalmoplegia (CPEO) [Blakely et al., 2004].

The presence of copy-number variations (CNVs) has been demonstrated both within concordant and discordant MZ twin pairs. Bruder et al. studied a total of 19 pairs of MZ twins; 9 of these MZ twin pairs were discordant for probable or possible Parkinson disease, Parkinsonism, or Lewy body dementia and 10 MZ twin pairs were phenotypically unselected. Comparisons within the discordant twin pairs revealed a considerable number of loci suggestive of putative CNV. Analysis of the concordant normal MZ twin pairs also showed discordances in CNVs, suggesting putative de novo somatic CNV events, in addition to the concordant genotypes, including CNVs shared by both twins of a pair [Bruder et al., 2008]. The implications of these findings may be far reaching and the study of CNVs in discordant MZ twin pairs is likely to be an important field of further research.

Telomere length and reduction of telomere length are partly genetically determined, nonetheless significant differences can be found within MZ twin pairs [Slagboom et al., 1994; Andrew et al., 2006; Bakaysa et al., 2007]. The within-pair correlation of telomere length in MZ twins decreases with increasing age [Bakaysa et al., 2007]. The analysis of telomere lengths in lymphocytes from an opposite-sex DZ twin pair with chimerism suggested that telomere length attrition is partly determined by the extracellular factors [Bruderlein et al., 2008]. Twin studies have contributed to our insight into the role of telomere length reduction in the process of aging and the development of age-related disease [Bakaysa et al., 2007; Gilley et al., 2008; Kimura et al., 2008; Valdes et al., 2008].

Finally, few differences in SNPs [Montgomery et al., 2005] have been recognized within MZ twin pairs. The significance of this finding remains to be elucidated.

Thus, genetic differences between MZ twins may reflect a post-zygotic mutation. In singletons, post-zygotic mutations result in (somatic) mosaicism. Somatic mosaicism is an increasingly recognized phenomenon [Gripp et al., 2006] and is believed to be involved in diseases in which the presumed causative mutation is hypothesized to be lethal in the non-mosaic state, such as Proteus syndrome and McCune–Albright syndrome [Endo et al., 1991; Peleg et al., 2009]. Moreover, somatic mosaicism has been associated with various non-Mendelian disorders and is considered to play a key role in carcinogenesis, ageing and possibly autoimmunity [Yousoufian and Pyeritz, 2002]. Hence, genetic differences between MZ twins may provide insight into early post-zygotic mutagenesis.

Discordant epigenetic phenomena also have been shown to contribute to phenotypic differences in MZ twins. Significant epigenetic differences have been demonstrated in MZ twins [Kato et al., 2005; Wong et al., 2005; Heijmans et al., 2007; Kaminsky et al., 2009]. Fraga et al. found remarkable (age-dependent) discordance within MZ twin pairs for DNA methylation and histone acetylation in one-third of investigated MZ twin pairs [Fraga et al., 2005; Machin, 2009b].

An increased incidence of BWS has been reported in MZ (female) twins [Weksberg et al., 2002; Smith et al., 2006; Blik et al., 2009]. In most cases the twin pair is discordant for the disease [Weksberg et al., 2002], due to discordance for loss of methylation at imprinting control region 2 (IC2) at chromosome 11p15 [Gaston et al.,

2001; Weksberg et al., 2002]. In line with these data, a discordant methylation pattern of differentially methylated region in the upstream of *H19* (H19-DMR) was found in an MZ twin pair discordant for Silver–Russell syndrome [Yamazawa et al., 2008]. In the affected twin from an MZ twin pair discordant for a caudal duplication anomaly, hypermethylation was shown at the promoter region of the *AXIN1* gene [Oates et al., 2006].

Discordance for a monogenetic disease in women may be the result of a difference in randomization of X-inactivation [Lubinsky and Hall, 1991; Tiberio, 1994; Watkiss et al., 1994; Goodship et al., 1996]; this mechanism has been described for hemophilia [Revesz et al., 1972; Kitchens, 1987; Valleix et al., 2002; Bennett et al., 2008], Fabry disease [Redonnet-Vernhet et al., 1996], colorblindness [Jorgensen et al., 1992], Hunter syndrome [Winchester et al., 1992], Fragile-X syndrome [Tuckerman et al., 1985; Kruyer et al., 1994], and Duchenne muscular dystrophy [Gomez et al., 1977; Burn et al., 1986; Chutkow et al., 1987; Pena et al., 1987; Richards et al., 1990; Lupski et al., 1991; Tremblay et al., 1993; Zneimer et al., 1993]. Apparently, non-shared factors can determine randomization of X-inactivation.

The process of MZ twinning may coincide the timing of X-inactivation, randomly resulting in clones with different X-inactivation patterns and unequal blastomere allocation may result in skewed X-inactivation in the smaller clone [Machin, 2009b].

MZ twin pairs discordant for a genetic abnormality elegantly serve as an in vivo model for the genetic defect. In case reports or series on singletons with a (relatively) uncommon disorder the effects related to the genetic aberration may be difficult to extract. Nevertheless, in a single discordant MZ twin pair, sharing many environmental factors and the genetic abnormality being the sole genetic difference, the comparison between the two twins is very illustrative and provides valuable information. The report of an MZ twin pair, of whom one twin has a 45,X genotype and the other twin a 46,XX genotype, increased our insight into the net effect of missing one X-chromosome in females [Rohrer et al., 2004]. Follow-up and comparison of the physical features, growth, and hormone production of both girls, presumably being otherwise genetically identical, indicates the pure effect of the absence of a second X-chromosome on these parameters over time. Likewise, an MZ twin pair discordant for a malformation sequence clearly illustrates the sequential effects of the primary defect. The urinary production from the unaffected twin from an MZ twin pair discordant for renal agenesis prevented the development of sequential abnormalities occurring in Potter sequence in the affected twin [Perez-Brayfield et al., 2004].

Discordant MZ twin pairs may increase our knowledge on a disease upon detailed phenotyping. A thorough analysis of discordant MZ twin pairs may elucidate differences and highlight mechanisms that play a key role in the disease. For example, magnetic resonance imaging (MRI) analysis of MZ twin pairs, discordant for Rett syndrome [Carter et al., 2008], Asperger syndrome [Belmonte and Carper, 2006], ADHD [van 't et al., 2007], and obsessive-compulsive symptoms [den Braber et al., 2008] revealed significant differences in brain structure or functioning.

The intrauterine environment is not necessarily equal for both twins from an MZ twin pair because of the number of cells allocated

to each twin, the timing of the twinning process, and the vascular distribution from the placenta [Machin, 1996]. An increased prevalence of deformations is observed in MZ twins compared to DZ twins and singletons. A substantial part of these deformations is present in only one twin [Hall, 1996b, 2003]. Evidently, discordance for congenital deformations or malformations in an MZ twin pair may also reflect environmental origins [van der Knaap and Barth, 1994; Mohammed et al., 2004].

In addition to descriptions of discordance for genotypes or phenotypes in Mendelian and chromosomal disorders, discordant MZ twin pairs also have been investigated for aspects of multifactorial traits and disorders, by comparing the affected (high-scoring) and unaffected (low-scoring) twin. For example, the immunologic responses in MZ twins discordant for rheumatoid arthritis [Gulwani-Akolkar et al., 1995] and Crohn's disease [Gulwani-Akolkar et al., 1994] were compared and cerebral glucose metabolism was studied in MZ twins discordant for Alzheimer disease [Virta et al., 2009]. Similarly, cytokine profiles in MZ twins discordant for myasthenia gravis differed significantly, suggesting a role of these mediators in the pathogenesis of the disease [Kakoulidou et al., 2004]. The amount of exercise in a pair of MZ female twins discordant for adolescent idiopathic scoliosis (AIS) was evaluated by Potoupnis et al. [2008], and this case report questioned the "cause and effect" relationship between AIS and exercising, as both girls were elite athletes twins.

Discordant MZ twin pairs also comprise a unique source of information on the effects of acquired diseases such as HIV [Agrati et al., 2006] or lifestyle factors [e.g., smoking, Kaprio and Koskenvuo, 1989; van Leeuwen et al., 2007 and drug use, Ferguson et al., 2007]. In population-based case-control studies, it is often difficult to distinguish the cause and the effect of a disorder (or, e.g., of medication use) on gene expression. MZ twins, discordant for the disorder, may be very valuable in the analysis of the effect of the disease on gene expression, as demonstrated for acquired obesity [Pietilainen et al., 2008]. In these pairs, mtDNA copy number was reduced by 47% in the obese twin's fat and pathway analyses of the adipose tissue transcription profiles showed significant down-regulation of mitochondrial branched-chain amino acid (BCAA) catabolism. These findings pointed to a substantial role of mitochondrial energy and amino acid metabolism in obesity.

MZ twins, both exposed to toxics may suggest variability of the effect that is probably not genetically determined, as published, for example, for diethylstilbestrol (DES), [Sandberg and Christian, 1980], thalidomide [Jorgensen et al., 1970], and alcohol [Streissguth and Dehaene, 1993] exposure.

The Human Genome Project and its associated technologies have opened up many new opportunities for gene finding and positional cloning has been successful in the identification of genes involved in many human disorders [Khoury and Yang, 1998; Landegren, 2000; Collins and McKusick, 2001]. However, the genetic basis of many heritable disorders remains unclear and the identification of a causative mutation responsible for a phenotype still represents a very demanding task. With today's whole-genome sequencing technologies and increasing evidence for genetic and epigenetic differences in some MZ twins, twin research provides new opportunities. The genetic analysis of discordant MZ twin pairs may identify new genes, as demonstrated in

Darier's disease and Van der Woude syndrome [Sakuntabhai et al., 1999; Kondo et al., 2002]. The occurrence of a possibly monogenetic disorder in only one twin from an MZ twin pair has been used as an argument against Mendelian inheritance. In 1992, an MZ twin pair discordant for Fuchs' heterochromic uveitis (FHU) persuaded the authors to state that "regular Mendelian inheritance is now proved to be impossible" [Jones and Read, 1992]. Recognizing numerous reports on discordance for genetic abnormalities in MZ twins, such a conclusion would nowadays probably be drawn more carefully. Also in prenatal testing, awareness of the possibility of genetic or phenotypic discordance is justified.

The exact extent of genetic differences between (unselected) MZ twins still remains to be elucidated. Based on the very first results of studies in which both members of MZ twin pairs were genotyped, such differences seem rare in unselected samples. It is therefore unlikely that such differences would significantly bias the results of large-scale epidemiological studies that analyze data from twins to estimate heritability. However, the precise effect of this possible variance in twin studies remains to be established based on empirical data.

Genetic difference within a discordant MZ twin pair may guide us toward causative genes in monogenetic diseases, without the confounding effect of polymorphisms [Kruglyak and Nickerson, 2001]. As such, hunting for genetic differences between discordant MZ twin pairs appears to be an encouraging strategy for future twin research. In the era of large-scale genome-wide association studies including thousands of patients in each study, single discordant MZ twin pairs may serve as an alternative to increase our knowledge on the spectrum of diseases, factors contributing to variability, pathogenic mechanisms, and post-zygotic mutagenesis. The search for differences in genotypes within discordant MZ twin pairs seems to be a promising approach in gene-finding. With future possibilities for human genome sequencing of large numbers of individuals [e.g., Drmanac et al., 2010] it may even become feasible to turn this strategy around and sequence large numbers of unselected MZ twin pairs, after which differences within pairs can be correlated with differences in phenotypes or transcription profiles.

REFERENCES

- Abuelo D. 2009. Clinical significance of chimerism. *Am J Med Genet Part C* 151C:148–151.
- Agrati C, Gioia C, Soldani F, Martini F, Antinori A, Poccia F. 2006. T cell selection and differentiation in AIDS disease: The model of HIV-discordant monozygotic twins. *J Biol Regul Homeost Agents* 20:24–28.
- Al-Mazrou KA, Al-Rekabi A, Alorainy IA, Al-Kharfi T, Al-Serhani AM. 2001. Pai syndrome: A report of a case and review of the literature. *Int J Pediatr Otorhinolaryngol* 61:149–153.
- Amin BR, Bauersachs RM, Meiselman HJ, Mohandas N, Heibel RP, Bowen PE, Schlegel RA, Williamson P, Westerman MP. 1991. Monozygotic twins with sickle cell anemia and discordant clinical courses: Clinical and laboratory studies. *Hemoglobin* 15:247–256.
- Anca MH, Gazit E, Loewenthal R, Ostrovsky O, Frydman M, Giladi N. 2004. Different phenotypic expression in monozygotic twins with Huntington disease. *Am J Med Genet Part A* 124A:89–91.
- Andrew T, Aviv A, Falchi M, Surdulescu GL, Gardner JP, Lu X, Kimura M, Kato BS, Valdes AM, Spector TD. 2006. Mapping genetic loci that

- determine leukocyte telomere length in a large sample of unselected female sibling pairs. *Am J Hum Genet* 78:480–486.
- Angle B, Yen F, Hersh JH, Gowans G, Barch M. 2002. Partial duplication 4q and deletion 1p36 in monozygotic twins with discordant phenotypes. *Am J Med Genet* 111:307–312.
- Ashton-Prolla P, Felix TM. 1997. Say syndrome: A new case with cystic renal dysplasia in discordant monozygotic twins. *Am J Med Genet* 70:353–356.
- Bailey W, Popovich B, Jones KL. 1995. Monozygotic twins discordant for the Russell-Silver syndrome. *Am J Med Genet* 58:101–105.
- Bakaysa SL, Mucci LA, Slagboom PE, Boomsma DI, McClearn GE, Johansson B, Pedersen NL. 2007. Telomere length predicts survival independent of genetic influences. *Aging Cell* 6:769–774.
- Baser ME, Ragge NK, Riccardi VM, Janus T, Gantz B, Pulst SM. 1996. Phenotypic variability in monozygotic twins with neurofibromatosis 2. *Am J Med Genet* 64:563–567.
- Beattie RB, Manson IW, Whittle MJ. 1993. Identical twins with trisomy 21 discordant for exomphalos. *Prenat Diagn* 13:1067–1070.
- Becker J, Hernandez A, Dipietro M, Coran AG. 2005. Identical twins concordant for pulmonary sequestration communicating with the esophagus and discordant for the VACTERL association. *Pediatr Surg Int* 21:541–546.
- Belmonte MK, Carper RA. 2006. Monozygotic twins with Asperger syndrome: Differences in behaviour reflect variations in brain structure and function. *Brain Cogn* 61:110–121.
- Bennett CM, Boye E, Neufeld EJ. 2008. Female monozygotic twins discordant for hemophilia A due to nonrandom X-chromosome inactivation. *Am J Hematol* 83:778–780.
- Bernstein PS, Aptsiauri N. 2003. Discordant retinitis pigmentosa in monozygotic twins. *Arch Ophthalmol* 121:1059–1062.
- Berry AC, Belton EM, Chantler C. 1980. Monozygotic twins discordant for Wiedemann–Beckwith syndrome and the implications for genetic counselling. *J Med Genet* 17:136–138.
- Bhol K, Udell I, Haider N, Yunis JJ, Mohimen A, Neuman R, Grasso C, Ahmed AR, Foster S. 1995. Ocular cicatricial pemphigoid. A case report of monozygotic twins discordant for the disease. *Arch Ophthalmol* 113:202–207.
- Biousse V, Brown MD, Newman NJ, Allen JC, Rosenfeld J, Meola G, Wallace DC. 1997. De novo 14484 mitochondrial DNA mutation in monozygotic twins discordant for Leber's hereditary optic neuropathy. *Neurology* 49:1136–1138.
- Blakely EL, He L, Taylor RW, Chinnery PF, Lightowlers RN, Schaefer AM, Turnbull DM. 2004. Mitochondrial DNA deletion in "identical" twin brothers. *J Med Genet* 41:e19.
- Blik J, Alders M, Maas SM, Oostra RJ, Mackay DM, van der LK, Callaway JL, Brooks A, van 't PS, Westerveld A, Leschot NJ, Mannens MM. 2009. Lessons from BWS twins: Complex maternal and paternal hypomethylation and a common source of haematopoietic stem cells. *Eur J Hum Genet* 17:1625–1634.
- Boccon-Gibod L, Boman F, Boudjemaa S, Fabre M, Leverger G, Carney AJ. 2004. Separate occurrence of extra-adrenal paraganglioma and gastrointestinal stromal tumor in monozygotic twins: Probable familial Carney syndrome. *Pediatr Dev Pathol* 7:380–384.
- Boles DJ, Bodurtha J, Nance WE. 1987. Goldenhar complex in discordant monozygotic twins: A case report and review of the literature. *Am J Med Genet* 28:103–109.
- Bonilla E, Younger DS, Chang HW, Tantravahi U, Miranda AF, Medori R, DiMauro S, Warburton D, Rowland LP. 1990. Partial dystrophin deficiency in monozygous twin carriers of the Duchenne gene discordant for clinical myopathy. *Neurology* 40:1267–1270.
- Bose B, Wilkie RA, Madlom M, Forsyth JS, Faed MJ. 1985. Wiedemann–Beckwith syndrome in one of monozygotic twins. *Arch Dis Child* 60:1191–1192.
- Bourthoumieu S, Yardin C, Terro F, Gilbert B, Laroche C, Saura R, Vincent MC, Esclaire F. 2005. Monozygotic twins concordant for blood karyotype, but phenotypically discordant: A case of "mosaic chimerism." *Am J Med Genet Part A* 135A:190–194.
- Bremond-Gignac D, Gerard-Blanluet M, Copin H, Bitoun P, Baumann C, Crolla JA, Benzacken B, Verloes A. 2005. Three patients with hallucal polydactyly and WAGR syndrome, including discordant expression of Wilms tumor in MZ twins. *Am J Med Genet Part A* 134A:422–425.
- Briellmann RS, Jackson GD, Torn-Broers Y, Berkovic SF. 1998. Twins with different temporal lobe malformations: Schizencephaly and arachnoid cyst. *Neuropediatrics* 29:284–288.
- Brockmann K, Happel R, Oeffner F, Konig A. 2008. Monozygotic twins discordant for Proteus syndrome. *Am J Med Genet Part A* 146A:2122–2125.
- Brodtkorb E, Myhr G, Gimse R. 2000. Is monochorionic twinning a risk factor for focal cortical dysgenesis? *Acta Neurol Scand* 102:53–59.
- Brown WT, Wisniewski KE, Sudhalter V, Keogh M, Tsiouris J, Miezieski C, Schaefer GB. 1998. Identical twins discordant for Sotos syndrome. *Am J Med Genet* 79:329–333.
- Bruder CE, Piotrowski A, Gijsbers AA, Andersson R, Erickson S, de Stahl TD, Menzel U, Sandgren J, von TD, Poplawski A, Crowley M, Crasto C, Partridge EC, Tiwari H, Allison DB, Komorowski J, van Ommen GJ, Boomsma DI, Pedersen NL, den Dunnen JT, Wirdefeldt K, Dumanski JP. 2008. Phenotypically concordant and discordant monozygotic twins display different DNA copy-number-variation profiles. *Am J Hum Genet* 82:763–771.
- Bruderlein S, Muller K, Melzner J, Hogel J, Wiegand P, Moller P. 2008. Different rates of telomere attrition in peripheral lymphocytes in a pair of dizygotic twins with hematopoietic chimerism. *Aging Cell* 7:663–666.
- Bugge M. 1981. Monozygotic twins discordant for exstrophy of the urinary bladder. *J Med Genet* 18:139–141.
- Burn J, Povey S, Boyd Y, Munro EA, West L, Harper K, Thomas D. 1986. Duchenne muscular dystrophy in one of monozygotic twin girls. *J Med Genet* 23:494–500.
- Camacho AF, Schneider A, Dorsainville D, Schutzman DL. 2008. Monozygotic twins discordant for VACTERL association. *Prenat Diagn* 28:366–368.
- Carakushansky G, Berthier C. 1976. The de Lange syndrome in one of twins. *J Med Genet* 13:404–406.
- Carakushansky G, Aguiar MB, Goncalves MR, Berthier CO, Kahn E, Carakushansky M, Pena SD. 1996. Identical twin discordance for the Brachmann-de Lange syndrome revisited. *Am J Med Genet* 63:458–460.
- Carter JC, Lanham DC, Pham D, Bibat G, Naidu S, Kaufmann WE. 2008. Selective cerebral volume reduction in Rett syndrome: A multiple-approach MR imaging study. *AJNR Am J Neuroradiol* 29:436–441.
- Castorina P, Selicorni A, Bedeschi F, Dalpra L, Larizza L. 1997. Genotype–phenotype correlation in two sets of monozygotic twins with Williams syndrome. *Am J Med Genet* 69:107–111.
- Celle L, Lee L, Rintoul N, Savani RC, Long W, Mennuti MT, Krantz ID. 2000. Duplication of chromosome region 4q28.3-qter in monozygotic twins with discordant phenotypes. *Am J Med Genet* 94:125–140.
- Chan JH, Tsui JK. 1997. Idiopathic cervical dystonia: Report of discordance in a pair of monozygotic twins. *Parkinsonism Relat Disord* 3:167–169.

- Chang HJ, Lee JR, Jee BC, Suh CS, Kim SH. 2009. Impact of blastocyst transfer on offspring sex ratio and the monozygotic twinning rate: A systematic review and meta-analysis. *Fertil Steril* 91:2381–2390.
- Chen CJ, Ho TY, Lu JJ, Sheu LF, Lee SY, Tien CH, Cheng SN. 2004. Identical twin brothers concordant for Langerhans' cell histiocytosis and discordant for Epstein-Barr virus-associated haemophagocytic syndrome. *Eur J Pediatr* 163:536–539.
- Cheng PJ, Shaw SW, Shih JC, Soong YK. 2006. Monozygotic twins discordant for monosomy 21 detected by first-trimester nuchal translucency screening. *Obstet Gynecol* 107:538–541.
- Chien CH, Lee JS, Tsai WY, Wang TR. 1990. Wiedemann–Beckwith syndrome with congenital central hypothyroidism in one of monozygotic twins. *J Formos Med Assoc* 89:132–136.
- Chutkow JG, Hyser CL, Edwards JA, Heffner RR Jr, Czyrny JJ. 1987. Monozygotic female twin carriers discordant for the clinical manifestations of Duchenne muscular dystrophy. *Neurology* 37:1147–1151.
- Clayton-Smith J, Read AP, Donnai D. 1992. Monozygotic twinning and Wiedemann–Beckwith syndrome. *Am J Med Genet* 42:633–637.
- Collins FS, McKusick VA. 2001. Implications of the Human Genome Project for medical science. *J Am Med Assoc* 285:540–544.
- Cooley HM, Keech CL, Melny BJ, Menahem S, Morahan G, Kay TW. 1997. Monozygotic twins discordant for congenital complete heart block. *Arthritis Rheum* 40:381–384.
- Corsello G, Maresi E, Rossi C, Giuffrè L, Cittadini E. 1992. Thanatophoric dysplasia in monozygotic twins discordant for cloverleaf skull: Prenatal diagnosis, clinical and pathological findings. *Am J Med Genet* 42:122–126.
- Costa T, Greer W, Rysiecki G, Buncic JR, Ray PN. 1997. Monozygotic twins discordant for Aicardi syndrome. *J Med Genet* 34:688–691.
- Costa T, Lambert M, Teshima I, Ray PN, Richer CL, Dallaire L. 1998. Monozygotic twins with 45,X/46,XY mosaicism discordant for phenotypic sex. *Am J Med Genet* 75:40–44.
- Czlonkowska A, Gromadzka G, Chabik G. 2009. Monozygotic female twins discordant for phenotype of Wilson's disease. *Mov Disord* 24:1066–1069.
- Dahoun S, Gagos S, Gagnebin M, Gehrig C, Burgi C, Simon F, Vieux C, Extermann P, Lyle R, Morris MA, Antonarakis SE, Bena F, Blouin JL. 2008. Monozygotic twins discordant for trisomy 21 and maternal 21q inheritance: A complex series of events. *Am J Med Genet Part A* 146A:2086–2093.
- Dallapiccola B, Stomeo C, Ferranti G, Di LA, Purpura M. 1985. Discordant sex in one of three monozygotic triplets. *J Med Genet* 22:6–11.
- Daskalakis GJ, Nicolaides KH. 2002. Monozygotic twins discordant for body stalk anomaly. *Ultrasound Obstet Gynecol* 20:79–81.
- Davin JC, Olie KH, Verlaak R, Horuz F, Florquin S, Weening JJ, Groothoff JW, Strain L, Goodship TH. 2006. Complement factor H-associated atypical hemolytic uremic syndrome in monozygotic twins: Concordant presentation, discordant response to treatment. *Am J Kidney Dis* 47:e27–e30.
- Dayer R, Ceroni D, Bottani A, Kaelin A. 2007. Tibial aplasia-hypoplasia and ectrodactyly in monozygotic twins with a discordant phenotype. *J Pediatr Orthop* 27:266–269.
- De GL, Jinnah HA, Harris JC, Nyhan WL, Schretlen DJ, Trombley LM, O'Neill JP. 2005. Lesch-Nyhan disease in a female with a clinically normal monozygotic twin. *Mol Genet Metab* 85:70–77.
- de Haan GJ, Trenite DK, Stroink H, Parra J, Voskuyl R, van KM, Lindhout D, Bertram E. 2005. Monozygous twin brothers discordant for photosensitive epilepsy: First report of possible visual priming in humans. *Epilepsia* 46:1545–1549.
- den Braber A, Ent D, Blokland GA, van Grootheest DS, Cath DC, Veltman DJ, de Ruiter MB, Boomsma DI. 2008. An fMRI study in monozygotic twins discordant for obsessive-compulsive symptoms. *Biol Psychol* 79:91–102.
- Detjen AK, Tinschert S, Kaufmann D, Algermissen B, Nurnberg P, Schuelke M. 2007. Analysis of mitochondrial DNA in discordant monozygotic twins with neurofibromatosis type 1. *Twin Res Hum Genet* 10:486–495.
- Di RM, Doria-Lamba L, Caruso U. 2001. Monozygotic twins with X-linked adrenoleukodystrophy and different phenotypes. *Ann Neurol* 50:424.
- Doran M, Lerner AJ. 2004. Monozygotic twins discordant for primary progressive aphasia. *Alzheimer Dis Assoc Disord* 18:48–49.
- Drmanac R, Sparks AB, Callow MJ, Halpern AL, Burns NL, Kermani BG, Carnevali P, Nazarenko I, Nilsen GB, Yeung G, Dahl F, Fernandez A, Staker B, Pant KP, Baccash J, Borchertding AP, Brownley A, Cedeno R, Chen L, Chernikoff D, Cheung A, Chirita R, Curson B, Ebert JC, Hacker CR, Hartlage R, Hauser B, Huang S, Jiang Y, Karpinchyk V, Koenig M, Kong C, Landers T, Le C, Liu J, McBride CE, Morenzoni M, Morey RE, Mutch K, Perazich H, Perry K, Peters BA, Peterson J, Pethiyagoda CL, Pothuraju K, Richter C, Rosenbaum AM, Roy S, Shafto J, Sharanovich U, Shannon KW, Sheppy CG, Sun M, Thakuria JV, Tran A, Vu D, Zaranek AW, Wu X, Drmanac S, Oliphant AR, Banyai WC, Martin B, Ballinger DG, Church GM, Reid CA. 2010. Human genome sequencing using unchained base reads on self-assembling DNA nanoarrays. *Science* 327:78–81.
- Duru UA, Laufer MR. 2009. Discordance in Mayer-von Rokitansky-Kuster-Hauser syndrome noted in monozygotic twins. *J Pediatr Adolesc Gynecol* 22:e73–e75.
- Elcioglu N, Berry AC. 1997. Monozygotic twins discordant for the oculo-oto-radial syndrome (IVIC syndrome). *Genet Couns* 8:201–206.
- Endo M, Yamada Y, Matsuura N, Niikawa N. 1991. Monozygotic twins discordant for the major signs of McCune–Albright syndrome. *Am J Med Genet* 41:216–220.
- Feekery CJ, Parry-Fielder B, Hopkins JJ. 1993. Landau-Kleffner syndrome: Six patients including discordant monozygotic twins. *Pediatr Neurol* 9:49–53.
- Ferguson RJ, McDonald BC, Saykin AJ, Ahles TA. 2007. Brain structure and function differences in monozygotic twins: Possible effects of breast cancer chemotherapy. *J Clin Oncol* 25:3866–3870.
- Fraga MF, Ballestar E, Paz MF, Ropero S, Setien F, Ballestar ML, Heine-Suner D, Cigudosa JC, Urioste M, Benitez J, Boix-Chornet M, Sanchez-Aguilera A, Ling C, Carlsson E, Poulsen P, Vaag A, Stephan Z, Spector TD, Wu YZ, Plass C, Esteller M. 2005. Epigenetic differences arise during the lifetime of monozygotic twins. *Proc Natl Acad Sci USA* 102:10604–10609.
- Friedman JH, Trieschmann ME, Myers RH, Fernandez HH. 2005. Monozygotic twins discordant for Huntington disease after 7 years. *Arch Neurol* 62:995–997.
- Fryer A. 1996. Monozygotic twins with 22q11 deletion and discordant phenotypes. *J Med Genet* 33:173.
- Fujimoto A, Boelter WD, Sparkes RS, Lin MS, Battersby K. 1991. Monozygotic twins of discordant sex both with 45,X/46,X,idic(Y) mosaicism. *Am J Med Genet* 41:239–245.
- Gan JP, Wu ZH, Tu ZM, Zheng J. 2007. The comparison of twinning rates between urban and rural areas in China. *Twin Res Hum Genet* 10:633–637.
- Gaston V, Le BY, Soupre V, Burglen L, Donadieu J, Oro H, Audry G, Vazquez MP, Gicquel C. 2001. Analysis of the methylation status of the KCNQ1OT and H19 genes in leukocyte DNA for the diagnosis and prognosis of Beckwith–Wiedemann syndrome. *Eur J Hum Genet* 9:409–418.

- Gentilin B, Gueneri S, Bianchi V, Natacci F, Colombo A, Fogliani R, Fortuna R, Coviello DA, Curcio C, Lalatta F. 2008. Discordant prenatal phenotype and karyotype of monozygotic twins characterized by the unequal distribution of two cell lines investigated by different methods: A review. *Twin Res Hum Genet* 11:352–356.
- Georgiou N, Bradshaw JL, Chiu E, Tudor A, O’Gorman L, Phillips JG. 1999. Differential clinical and motor control function in a pair of monozygotic twins with Huntington’s disease. *Mov Disord* 14:320–325.
- Gilgenkrantz S, Janot C. 1983. Monozygotic twins discordant for trisomy 21 or chimeric dizygotic twins? *Am J Med Genet* 15:159–160.
- Gilley D, Herbert BS, Huda N, Tanaka H, Reed T. 2008. Factors impacting human telomere homeostasis and age-related disease. *Mech Ageing Dev* 129:27–34.
- Gomez MR, Engel AG, Dewald G, Peterson HA. 1977. Failure of inactivation of Duchenne dystrophy X-chromosome in one of female identical twins. *Neurology* 27:537–541.
- Goodship J, Cross I, Scambler P, Burn J. 1995. Monozygotic twins with chromosome 22q11 deletion and discordant phenotype. *J Med Genet* 32:746–748.
- Goodship J, Carter J, Burn J. 1996. X-inactivation patterns in monozygotic and dizygotic female twins. *Am J Med Genet* 61:205–208.
- Griggs RC, Tawil R, McDermott M, Forrester J, Figlewicz D, Weiffenbach B. 1995. Monozygotic twins with facioscapulohumeral dystrophy (FSHD): Implications for genotype/phenotype correlation. FSH-DY Group. *Muscle Nerve* 2:S50–S55.
- Gripp KW, Stabley DL, Nicholson L, Hoffman JD, Sol-Church K. 2006. Somatic mosaicism for an HRAS mutation causes Costello syndrome. *Am J Med Genet Part A* 140A:2163–2169.
- Grynberg M, Graesslin O, Teyssedre J, Quereux C, Gaillard D, Carre-Pigeon F. 2007. Prenatal diagnosis in monozygotic twins with Down syndrome who had different phenotypes. *Prenat Diagn* 27:552–554.
- Guidry J, Kent TA. 1999. New genetic hypothesis of schizophrenia. *Med Hypotheses* 52:69–75.
- Guion-Almeida ML, Mellado C, Beltran C, Richieri-Costa A. 2007. Pai syndrome: Report of seven South American patients. *Am J Med Genet Part A* 143A:3273–3279.
- Gulwani-Akolkar B, Shalon L, Akolkar PN, Fisher SE, Silver J. 1994. Analysis of the peripheral blood T-cell receptor (TCR) repertoire in monozygotic twins discordant for Crohn’s disease. *Autoimmunity* 17:241–248.
- Gulwani-Akolkar B, Akolkar PN, Gregersen PK, Silver J. 1995. Analysis of the peripheral blood T-cell receptor repertoire in monozygotic twins discordant for rheumatoid arthritis. *Ann N Y Acad Sci* 756:176–178.
- Hall JG. 1996a. Twinning: Mechanisms and genetic implications. *Curr Opin Genet Dev* 6:343–347.
- Hall JG. 1996b. Twins and twinning. *Am J Med Genet* 61:202–204.
- Hall JG. 2003. Twinning. *Lancet* 362:735–743.
- Heidenreich W, Pfeiffer A, Kumbhani HK, Scholz W, Zeuner W. 1977. Discordant monozygotic twins with Mayer Rokitansky Kutser syndrome (author’s transl). *Geburtshilfe Frauenheilkd* 37:221–223.
- Heijmans BT, Kremer D, Tobi EW, Boomsma DI, Slagboom PE. 2007. Heritable rather than age-related environmental and stochastic factors dominate variation in DNA methylation of the human IGF2/H19 locus. *Hum Mol Genet* 16:547–554.
- Helderman-van den Enden AT, Maaswinkel-Mooij PD, Hoogendoorn E, Willemsen R, Maat-Kievit JA, Losekoot M, Oostra BA. 1999. Monozygotic twin brothers with the fragile X syndrome: Different CGG repeats and different mental capacities. *J Med Genet* 36:253–257.
- Hillebrand G, Siebert R, Simeoni E, Santer R. 2000. DiGeorge syndrome with discordant phenotype in monozygotic twins. *J Med Genet* 37:E23.
- Hipkin LJ, Casson IF, Davis JC. 1990. Identical twins discordant for Kallmann’s syndrome. *J Med Genet* 27:198–199.
- Hoekstra C, Zhao ZZ, Lambalk CB, Willemsen G, Martin NG, Boomsma DI, Montgomery GW. 2008. Dizygotic twinning. *Hum Reprod Update* 14:37–47.
- Hofer T, Frank J, Itin PH. 2005. Klippel-Trenaunay syndrome in a monozygotic male twin: Supportive evidence for the concept of para-dominant inheritance. *Eur J Dermatol* 15:341–343.
- Holmgren G, Wikstrom L, Lundgren HE, Suhr OB. 2004. Discordant penetrance of the trait for familial amyloidotic polyneuropathy in two pairs of monozygotic twins. *J Intern Med* 256:453–456.
- Hong YC, Liu HM, Chen PS, Chen YJ, Lyuu JY, Hu HY, Yi MF, Lin JS, Tzeng CH. 2007. Hair follicle: A reliable source of recipient origin after allogeneic hematopoietic stem cell transplantation. *Bone Marrow Transplant* 40:871–874.
- Hong D, Gupta R, Ancliff P, Atzberger A, Brown J, Soneji S, Green J, Colman S, Piacibello W, Buckle V, Tsuzuki S, Greaves M, Enver T. 2008. Initiating and cancer-propagating cells in TEL-AML1-associated childhood leukemia. *Science* 319:336–339.
- Horton WA, Harris DJ, Collins DL. 1983. Discordance for the Kleeblatt-schadel anomaly in monozygotic twins with thanatophoric dysplasia. *Am J Med Genet* 15:97–101.
- Humphrey A, Higgins JN, Yates JR, Bolton PF. 2004. Monozygotic twins with tuberous sclerosis discordant for the severity of developmental deficits. *Neurology* 62:795–798.
- Hwang MS, Su WJ, Lin JL. 2006. Asplenia syndrome in a pair of monozygotic twins. *Acta Paediatr* 95:500–501.
- James WH. 1975. Sex ratio in twin births. *Ann Hum Biol* 2:365–378.
- James WH. 1980. Secular changes in twinning rates in England and Wales. *Ann Hum Biol* 7:485–487.
- Johns DR, Smith KH, Miller NR, Sulewski ME, Bias WB. 1993. Identical twins who are discordant for Leber’s hereditary optic neuropathy. *Arch Ophthalmol* 111:1491–1494.
- Jones NP, Read AP. 1992. Is there a genetic basis for Fuchs’ heterochromic uveitis? Discordance in monozygotic twins. *Br J Ophthalmol* 76:22–24.
- Jorgensen G, Lenz W, Pfeiffer RA, Schaafhausen C. 1970. Thalidomide-embryopathy in twins: A collaborative study. *Acta Genet Med Gemellol (Roma)* 19:205–210.
- Jorgensen AL, Philip J, Raskind WH, Matsushita M, Christensen B, Dreyer V, Motulsky AG. 1992. Different patterns of X inactivation in MZ twins discordant for red-green color-vision deficiency. *Am J Hum Genet* 51:291–298.
- Juberg RC, Stallard R, Straughen WJ, Avotri KJ, Washington JW. 1981. Clinicopathological conference: A newborn monozygotic twin with abnormal facial appearance and respiratory insufficiency. *Am J Med Genet* 10:193–200.
- Kajii T, Hagiwara K, Tsukahara M, Nakajima H, Fukuda Y. 1981. Monozygotic twins discordant for Rubinstein-Taybi syndrome. *J Med Genet* 18:312–314.
- Kakoulidou M, Ahlberg R, Yi Q, Giscombe R, Pirskanen R, Lefvert AK. 2004. The autoimmune T and B cell repertoires in monozygotic twins discordant for myasthenia gravis. *J Neuroimmunol* 148:183–191.
- Kamath BM, Krantz ID, Spinner NB, Heubi JE, Piccoli DA. 2002. Monozygotic twins with a severe form of Alagille syndrome and phenotypic discordance. *Am J Med Genet* 112:194–197.

- Kaminsky ZA, Tang T, Wang SC, Ptak C, Oh GH, Wong AH, Feldcamp LA, Virtanen C, Halfvarson J, Tysk C, McRae AF, Visscher PM, Montgomery GW, Gottesman II, Martin NG, Petronis A. 2009. DNA methylation profiles in monozygotic and dizygotic twins. *Nat Genet* 41:240–245.
- Kaprio J, Koskenvuo M. 1989. Twins, smoking and mortality: A 12-year prospective study of smoking-discordant twin pairs. *Soc Sci Med* 29:1083–1089.
- Kato T, Iwamoto K, Kakiuchi C, Kuratomi G, Okazaki Y. 2005. Genetic or epigenetic difference causing discordance between monozygotic twins as a clue to molecular basis of mental disorders. *Mol Psychiatry* 10:622–630.
- Kaufman LW, Folk ER, Miller MT. 1989. Monozygotic twins discordant for Duane's retraction syndrome. Case report. *Arch Ophthalmol* 107:324–325.
- Kelly TE, Sproul GT, Huerta MG, Rogol AD. 1998. Discordant puberty in monozygotic twin sisters with neurofibromatosis type 1 (NF1). *Clin Pediatr (Phila)* 37:301–304.
- Khoury MJ, Yang Q. 1998. The future of genetic studies of complex human diseases: An epidemiologic perspective. *Epidemiology* 9:350–354.
- Kimura M, Hjelmberg JV, Gardner JP, Bathum L, Brimacombe M, Lu X, Christiansen L, Vaupel JW, Aviv A, Christensen K. 2008. Telomere length and mortality: A study of leukocytes in elderly Danish twins. *Am J Epidemiol* 167:799–806.
- Kitchens CS. 1987. Discordance in a pair of identical twin carriers of factor IX deficiency. *Am J Hematol* 24:225–228.
- Kondo S, Yamashina U, Sato N, Aso K. 1991. Discordant expression of tuberous sclerosis in monozygotic twins. *J Dermatol* 18:178–180.
- Kondo S, Schutte BC, Richardson RJ, Bjork BC, Knight AS, Watanabe Y, Howard E, de Lima RL, Daack-Hirsch S, Sander A, Donald-McGinn DM, Zackai EH, Lammer EJ, Aylsworth AS, Ardinger HH, Lidral AC, Pober BR, Moreno L, Arcos-Burgos M, Valencia C, Houdayer C, Bahuau M, Moretti-Ferreira D, Richieri-Costa A, Dixon MJ, Murray JC. 2002. Mutations in IRF6 cause Van der Woude and popliteal pterygium syndromes. *Nat Genet* 32:285–289.
- Korenke GC, Fuchs S, Krasemann E, Doerr HG, Wilichowski E, Hunneman DH, Hanefeld F. 1996. Cerebral adrenoleukodystrophy (ALD) in only one of monozygotic twins with an identical ALD genotype. *Ann Neurol* 40:254–257.
- Kroes HY, Takahashi M, Zijlstra RJ, Baert JA, Kooi KA, Hofstra RM, van Essen AJ. 2002. Two cases of the caudal duplication anomaly including a discordant monozygotic twin. *Am J Med Genet* 112:390–393.
- Kruglyak L, Nickerson DA. 2001. Variation is the spice of life. *Nat Genet* 27:234–236.
- Kruyer H, Mila M, Glover G, Carbonell P, Ballesta F, Estivill X. 1994. Fragile X syndrome and the (CGG)_n mutation: Two families with discordant MZ twins. *Am J Hum Genet* 54:437–442.
- Kumar S, Carver C, McCall S, Brice G, Ostergaard P, Mortimer P, Jeffery S. 2007. A family with lymphoedema-distichiasis where identical twins have a discordant phenotype. *Clin Genet* 71:285–287.
- Kurosawa K, Kuromaru R, Imaizumi K, Nakamura Y, Ishikawa F, Ueda K, Kuroki Y. 1992. Monozygotic twins with discordant sex. *Acta Genet Med Gemellol (Roma)* 41:301–310.
- Kuzniecky R, Gilliam F, Faught E. 1995. Discordant occurrence of cerebral unilateral heterotopia and epilepsy in monozygotic twins. *Epilepsia* 36:1155–1157.
- Lachmann RH, Grant IR, Halsall D, Cox TM. 2004. Twin pairs showing discordance of phenotype in adult Gaucher's disease. *Q J Med* 97:199–204.
- Lajeunie E, Bonaventure J, El Ghouzzi V, Catala M, Renier D. 2000. Monozygotic twins with Crouzon syndrome: Concordance for cranio-synostosis and discordance for thumb duplication. *Am J Med Genet* 91:159–160.
- Lam BL. 1998. Identical twins no longer discordant for Leber's hereditary optic neuropathy. *Arch Ophthalmol* 116:956–957.
- Landegren U. 2000. Consulting the source code: Prospects for gene-based medical diagnostics. *J Intern Med* 248:271–276.
- Lee JT, Chou HC, Tsao PN, Hsieh WS, Hwu WL. 2004. Trisomy 18 in monozygotic twins with discordant phenotypes. *J Formos Med Assoc* 103:314–316.
- Lenti C, Triulzi F. 1996. Discordant clinical and neuroradiological features of congenital bilateral perisylvian syndrome in monozygotic female twins. *Ital J Neurol Sci* 17:287–290.
- Leonard NJ, Bernier FP, Rudd N, Machin GA, Bamforth F, Bamforth S, Grundy P, Johnson C. 1996. Two pairs of male monozygotic twins discordant for Wiedemann-Beckwith syndrome. *Am J Med Genet* 61:253–257.
- Litz CE, Taylor KA, Qiu JS, Pescovitz OH, de MB. 1988. Absence of detectable chromosomal and molecular abnormalities in monozygotic twins discordant for the Wiedemann-Beckwith syndrome. *Am J Med Genet* 30:821–833.
- Loevey HT, Miller M, Rosenthal IM. 1985. Discordant monozygotic twins with trisomy 13. *Acta Genet Med Gemellol (Roma)* 34:185–188.
- Lubinsky MS, Hall JG. 1991. Genomic imprinting, monozygous twinning, and X inactivation. *Lancet* 337:1288.
- Lupski JR, Garcia CA, Zoghbi HY, Hoffman EP, Fenwick RG. 1991. Discordance of muscular dystrophy in monozygotic female twins: Evidence supporting asymmetric splitting of the inner cell mass in a manifesting carrier of Duchenne dystrophy. *Am J Med Genet* 40:354–364.
- Lynch SA, Ashcroft KA, Zwolinski S, Clarke C, Burn J. 1995. Kabuki syndrome-like features in monozygotic twin boys with a pseudodicentric chromosome 13. *J Med Genet* 32:227–230.
- Machin GA. 1996. Some causes of genotypic and phenotypic discordance in monozygotic twin pairs. *Am J Med Genet* 61:216–228.
- Machin G. 2009a. Familial monozygotic twinning: A report of seven pedigrees. *Am J Med Genet Part C* 151C:152–154.
- Machin G. 2009b. Non-identical monozygotic twins, intermediate twin types, zygosity testing, and the non-random nature of monozygotic twinning: A review. *Am J Med Genet Part C* 151C:110–127.
- Machin GA, Sperber GH, Wootliffe J. 1985. Monozygotic twin aborted fetuses discordant for holoprosencephaly/synotia. *Teratology* 31:203–215.
- Mansilla MA, Kimani J, Mitchell LE, Christensen K, Boomsma DI, Daack-Hirsch S, Nepomucena B, Wyszynski DF, Felix TM, Martin NG, Murray JC. 2005. Discordant MZ twins with cleft lip and palate: A model for identifying genes in complex traits. *Twin Res Hum Genet* 8:39–46.
- Marcus-Soekarman D, Hamers G, Velzeboer S, Nijhuis J, Loneus WH, Herbergs J, de Die-Smulders C, Schrander-Stumpel C, Engelen J. 2004. Mosaic trisomy 11p in monozygotic twins with discordant clinical phenotypes. *Am J Med Genet Part A* 124A:288–291.
- Martin N, Boomsma D, Machin G. 1997. A twin-pronged attack on complex traits. *Nat Genet* 17:387–392.
- Martin N, Zugge K, Brandt R, Friebe D, Janssen B, Zimmerhackl LB. 2003. Discordant clinical manifestations in monozygotic twins with the identical mutation in the TSC2 gene. *Clin Genet* 63:427–430.
- Matsukura H, Higuchi A, Ieki A, Miyawaki T. 2004. Discordant phenotypic expression of Alport syndrome in monozygotic twins. *Clin Nephrol* 62:313–318.

- Maurer HS, Pendergrass TW, Borges W, Honig GR. 1979. The role of genetic factors in the etiology of Wilms' tumor: Two pairs of monozygous twins with congenital abnormalities (aniridia; hemihypertrophy) and discordance for Wilms' tumor. *Cancer* 43:205–208.
- McCredie J. 1975. Congenital fusion of bones: Radiology, embryology and pathogenesis. *Clin Radiol* 26:47–51.
- McMahon TT, Shin JA, Newlin A, Edrington TB, Sugar J, Zadnik K. 1999. Discordance for keratoconus in two pairs of monozygotic twins. *Cornea* 18:444–451.
- Mekus F, Ballmann M, Bronsveld I, Bijman J, Veeze H, Tummler B. 2000. Categories of deltaF508 homozygous cystic fibrosis twin and sibling pairs with distinct phenotypic characteristics. *Twin Res* 3:277–293.
- Michalski JP, Snyder SM, McLeod RL, Talal N. 1978. Monozygotic twins with Klinefelter's syndrome discordant for systemic lupus erythematosus and symptomatic myasthenia gravis. *Arthritis Rheum* 21:306–309.
- Migeon BR, Dunn MA, Thomas G, Schmeckpeper BJ, Naidu S. 1995. Studies of X inactivation and isodisomy in twins provide further evidence that the X chromosome is not involved in Rett syndrome. *Am J Hum Genet* 56:647–653.
- Misra VK, Struys EA, O'Brien W, Salomons GS, Glover T, Jakobs C, Innis JW. 2005. Phenotypic heterogeneity in the presentation of D-2-hydroxyglutaric aciduria in monozygotic twins. *Mol Genet Metab* 86:200–205.
- Miyama S, Goto T, Inoue Y, Yamakawa K. 2008. Monozygotic twins with severe myoclonic epilepsy in infancy discordant for clinical features. *Pediatr Neurol* 39:120–122.
- Mohammed SN, Swan MC, Wall SA, Wilkie AO. 2004. Monozygotic twins discordant for frontonasal malformation. *Am J Med Genet Part A* 130A:384–388.
- Montgomery GW, Campbell MJ, Dickson P, Herbert S, Siemering K, Ewen-White KR, Visscher PM, Martin NG. 2005. Estimation of the rate of SNP genotyping errors from DNA extracted from different tissues. *Twin Res Hum Genet* 8:346–352.
- Mulder AF, van EJ, Groenendaal F, Wladimiroff JW. 1989. Trisomy 18 in monozygotic twins. *Hum Genet* 83:300–301.
- Munar-Ques M, Pedrosa JL, Coelho T, Gusmao L, Seruca R, Amorim A, Sequeiros J. 1999. Two pairs of proven monozygotic twins discordant for familial amyloid neuropathy (FAP) TTR Met 30. *J Med Genet* 36:629–632.
- Murphy MB, Greenberg F, Wilson G, Hughes M, DiLiberti J. 1990. Williams syndrome in twins. *Am J Med Genet Suppl* 6:97–99.
- Naor N, Amir Y, Cohen T, Davidson S. 1987. Trisomy 13 in monozygotic twins discordant for major congenital anomalies. *J Med Genet* 24:500–502.
- Naselli A, Vignolo M, Di BE, Papale V, Aicardi G, Becchetti S, Toma P. 1998. Trichorhinophalangeal syndrome type I in monozygotic twins discordant for hip pathology. Report on the morphological evolution of cone-shaped epiphyses and the unusual pattern of skeletal maturation. *Pediatr Radiol* 28:851–855.
- Niehues T, Schwarz K, Schneider M, Schrotten H, Schroder E, Stephan V, Wahn V. 1996. Severe combined immunodeficiency (SCID) associated neutropenia: A lesson from monozygotic twins. *Arch Dis Child* 74:340–342.
- Nieuwint A, Van Zalen-Sprock R, Hummel P, Pals G, Van VJ, Van Der HH, Heins Y, Madan K. 1999. 'Identical' twins with discordant karyotypes. *Prenat Diagn* 19:72–76.
- Noone PG, Bali D, Carson JL, Sannuti A, Gipson CL, Ostrowski LE, Bromberg PA, Boucher RC, Knowles MR. 1999. Discordant organ laterality in monozygotic twins with primary ciliary dyskinesia. *Am J Med Genet* 82:155–160.
- Norremolle A, Hasholt L, Petersen CB, Eiberg H, Hasselbalch SG, Gideon P, Nielsen JE, Sorensen SA. 2004. Mosaicism of the CAG repeat sequence in the Huntington disease gene in a pair of monozygotic twins. *Am J Med Genet Part A* 130A:154–159.
- Oates NA, van VJ, Duffy DL, Kroes HY, Martin NG, Boomsma DI, Campbell M, Coulthard MG, Whitelaw E, Chong S. 2006. Increased DNA methylation at the AXIN1 gene in a monozygotic twin from a pair discordant for a caudal duplication anomaly. *Am J Hum Genet* 79:155–162.
- O'Donnell CP, Pertile MD, Sheffield LJ, Sampson A. 2004. Monozygotic twins with discordant karyotypes: A case report. *J Pediatr* 145:406–408.
- Oduber CE, Blik J, van der Horst CM, van Steensel MA, Hennekam RC. 2010. Monozygotic twins discordant for vascular malformations and dysregulated growth. *Eur J Med Genet*. 53:14–18.
- Olney AH, Buehler BA, Waziri M. 1988. Wiedemann–Beckwith syndrome in apparently discordant monozygotic twins. *Am J Med Genet* 29:491–499.
- Oorthuys JW. 1984. Monozygotic twins with Williams-Beuren or 'elfenface' syndrome. *Tijdschr Kindergeneesk* 52:197–200.
- Orstavik RE, Tommerup N, Eiklid K, Orstavik KH. 1995. Non-random X chromosome inactivation in an affected twin in a monozygotic twin pair discordant for Wiedemann–Beckwith syndrome. *Am J Med Genet* 56:210–214.
- Owen CJ, Habeb A, Pearce SH, Wright M, Ichikawa S, Sorenson AH, Econs MJ, Cheetham TD. 2009. Discordance for X-linked hypophosphataemic rickets in identical twin girls. *Horm Res* 71:237–244.
- Panas M, Karadima G, Markianos M, Kalfakis N, Vassilopoulos D. 2008. Phenotypic discordance in a pair of monozygotic twins with Huntington's disease. *Clin Genet* 74:291–292.
- Pankau R, Siebert R, Kautza M, Schneppenheim R, Gosch A, Wessel A, Partsch CJ. 2001. Familial Williams-Beuren syndrome showing varying clinical expression. *Am J Med Genet* 98:324–329.
- Pasini B, McWhinney SR, Bei T, Matyakhina L, Stergiopoulos S, Muchow M, Boikos SA, Ferrando B, Pacak K, Assie G, Baudin E, Chompret A, Ellison JW, Briere JJ, Rustin P, Gimenez-Roqueplo AP, Eng C, Carney JA, Stratakis CA. 2008. Clinical and molecular genetics of patients with the Carney-Stratakis syndrome and germline mutations of the genes coding for the succinate dehydrogenase subunits SDHB, SDHC, and SDHD. *Eur J Hum Genet* 16:79–88.
- Peleg R, Luba A, Eliakim A, Israeli-Shani L, Manor E, Birk R, Parvari R. 2009. McCune–Albright syndrome in a discordant monozygotic twin. *Isr Med Assoc J* 11:343–347.
- Pena SD, Karpati G, Carpenter S, Fraser FC. 1987. The clinical consequences of X-chromosome inactivation: Duchenne muscular dystrophy in one of monozygotic twins. *J Neurol Sci* 79:337–344.
- Peng HH, Kuo PL, Chao AS, Wang TH, Chang YL, Soong YK, Chang SD. 2007. Discordant semilobar holoprosencephaly in monozygotic twins with de novo inv dup(15) marker chromosome and de novo mutation on SHH gene. *Fetal Diagn Ther* 22:389–393.
- Perez-Brayfield MR, Kirsch AJ, Smith EA. 2004. Monoamniotic twin discordant for bilateral renal agenesis with normal pulmonary function. *Urology* 64:589.
- Peterson SL, Rayan GM. 2004. Monozygotic twins discordant for thumb polydactyly. *Plast Reconstr Surg* 113:449–451.
- Picci L, Cameran M, Scarpa M, Pradal U, Melotti P, Assael BM, Castellani C. 2007. TG15 T5 allele in clinically discordant monozygotic twins with cystic fibrosis. *Am J Med Genet Part A* 143A:1936–1937.
- Pietilainen KH, Naukkarinen J, Rissanen A, Saharinen J, Ellonen P, Keranen H, Suomalainen A, Gotz A, Suortti T, Yki-Jarvinen H, Oresic

- M, Kaprio J, Peltonen L. 2008. Global transcript profiles of fat in monozygotic twins discordant for BMI: Pathways behind acquired obesity. *PLoS Med* 5:e51.
- Potoupnis ME, Kenanidis E, Papavasiliou KA, Kapetanios GA. 2008. The role of exercising in a pair of female monozygotic (high-class athletes) twins discordant for adolescent idiopathic scoliosis. *Spine (Phila Pa 1976)* 33:E607–E610.
- Rauch A, Hofbeck M, Bähring S, Leipold G, Trautmann U, Singer H, Pfeiffer RA. 1998. Monozygotic twins concordant for Cayler syndrome. *Am J Med Genet* 75:113–117.
- Raynes HR, Shanske A, Goldberg S, Burde R, Rapin I. 1999. Joubert syndrome: Monozygotic twins with discordant phenotypes. *J Child Neurol* 14:649–654.
- Redonnet-Vernhet I, Ploos van Amstel JK, Jansen RP, Wevers RA, Salvayre R, Levade T. 1996. Uneven X inactivation in a female monozygotic twin pair with Fabry disease and discordant expression of a novel mutation in the alpha-galactosidase A gene. *J Med Genet* 33:682–688.
- Reindollar RH, Byrd JR, Hahn DH, Haseltine FP, McDonough PG. 1987. A cytogenetic and endocrinologic study of a set of monozygotic isokaryotic 45,X/46,XY twins discordant for phenotypic sex: Mosaicism versus chimerism. *Fertil Steril* 47:626–633.
- Reiss AL, Freund L, Plotnick L, Baumgardner T, Green K, Sozer AC, Reader M, Boehm C, Denckla MB. 1993. The effects of X monosomy on brain development: Monozygotic twins discordant for Turner's syndrome. *Ann Neurol* 34:95–107.
- Rettig KR, Vargas A, Reiter E, Root AW. 1980. Discordance for congenital athyrotic hypothyroidism in one monozygotic twin. *Clin Pediatr (Phila)* 19:63–65.
- Reutter H, Shapiro E, Gruen JR. 2003. Seven new cases of familial isolated bladder exstrophy and epispadias complex (BEEC) and review of the literature. *Am J Med Genet Part A* 120A:215–221.
- Revesz T, Schuler D, Goldschmidt B, Elodi S. 1972. Christmas disease in one of a pair of monozygotic twin girls, possibly the effect of lyonization. *J Med Genet* 9:396–400.
- Richards CS, Watkins SC, Hoffman EP, Schneider NR, Milsark IW, Katz KS, Cook JD, Kunkel LM, Cortada JM. 1990. Skewed X inactivation in a female MZ twin results in Duchenne muscular dystrophy. *Am J Hum Genet* 46:672–681.
- Rietveld MJ, Der Valk JC, Bongers IL, Stroet TM, Slagboom PE, Boomsma DI. 2000. Zygosity diagnosis in young twins by parental report. *Twin Res* 3:134–141.
- Robertson SP, Thompson S, Morgan T, Holder-Espinasse M, Martinot-Duquenoy V, Wilkie AO, Manouvrier-Hanu S. 2006. Postzygotic mutation and germline mosaicism in the otopalatodigital syndrome spectrum disorders. *Eur J Hum Genet* 14:549–554.
- Robinow M, Marsh JL, Edgerton MT, Sabio H, Johnson GF. 1978. Discordance in monozygotic twins for aglossia-adactylia, and possible clues to the pathogenesis of the syndrome. *Birth Defects Orig Artic Ser* 14:223–230.
- Rogers JG, Voullaire L, Gold H. 1982. Monozygotic twins discordant for trisomy 21. *Am J Med Genet* 11:143–146.
- Rohrer TR, Gassmann KF, Rauch A, Pfeiffer RA, Doerr HG. 2004. Growth of heterokaryotic monozygotic twins discordant for Ullrich-Turner syndrome during the first years of life. *Am J Med Genet Part A* 126A:78–83.
- Rosenbaum AL, Weiss SJ. 1978. Monozygotic twins discordant for Duane's retraction syndrome. *J Pediatr Ophthalmol Strabismus* 15:359–361.
- Rudolph G, Haritoglou C, Kalpadakis P, Boergen KP, Meitinger T. 2001. LEOPARD syndrome with iris-retina-choroid coloboma. Discordant findings in monozygotic twins (MIM # 151 100). *Ophthalmologie* 98:1101–1103.
- Ryan CA, Finer NN, Ives E. 1988. Discordance of signs in monozygotic twins concordant for the Goldenhar anomaly. *Am J Med Genet* 29:755–761.
- Sagot P, David A, Talmant C, Pascal O, Winer N, Boog G. 1996. Russell-Silver syndrome: An explanation for discordant growth in monozygotic twins. *Fetal Diagn Ther* 11:72–78.
- Sakuntabhai A, Ruiz-Perez V, Carter S, Jacobsen N, Burge S, Monk S, Smith M, Munro CS, O'Donovan M, Craddock N, Kucherlapati R, Rees JL, Owen M, Lathrop GM, Monaco AP, Strachan T, Hovnanian A. 1999. Mutations in ATP2A2, encoding a Ca²⁺ pump, cause Darier disease. *Nat Genet* 21:271–277.
- Samn M, Lewis K, Blumberg B. 1990. Monozygotic twins discordant for the Russell-Silver syndrome. *Am J Med Genet* 37:543–545.
- Sandberg EC, Christian JC. 1980. Diethylstilbestrol-exposed monozygotic twins discordant for cervicovaginal clear cell adenocarcinoma. *Am J Obstet Gynecol* 137:220–228.
- Saporta MA, Plante-Bordeneuve V, Misrahi M, Cruz MW. 2009. Discordant expression of familial amyloid polyneuropathy in monozygotic Brazilian twins. *Amyloid* 16:38–41.
- Satoh K, Shibata Y, Tokushige H, Onizuka T. 1995. A mirror image of the first and second branchial arch syndrome associated with cleft lip and palate in monozygotic twins. *Br J Plast Surg* 48:601–605.
- Schlegelberger B, Schlegelberger T, Kekow J, Gross WL, Grote W. 1984. Monozygotic twins with Klinefelter syndrome (concordant) and systemic lupus erythematosus (discordant). *Klin Wochenschr* 62:906–910.
- Schlegelberger T, Kekow J, Gross WL. 1986. Impaired T-cell-independent B-cell maturation in systemic lupus erythematosus: Coculture experiments in monozygotic twins concordant for Klinefelter's syndrome but discordant for systemic lupus erythematosus. *Clin Immunol Immunopathol* 40:365–370.
- Schlessel JS, Brown WT, Lysikiewicz A, Schiff R, Zaslav AL. 1990. Monozygotic twins with trisomy 18: A report of discordant phenotype. *J Med Genet* 27:640–642.
- Schmid O, Trautmann U, Ashour H, Ulmer R, Pfeiffer RA, Beinder E. 2000. Prenatal diagnosis of heterokaryotypic mosaic twins discordant for fetal sex. *Prenat Diagn* 20:999–1003.
- Schmidt R, Sobel EH, Nitowsky HM, Dar H, Allen FH Jr. 1976. Monozygotic twins discordant for sex. *J Med Genet* 13:64–68.
- Schworm HD, Jedeke KB, Holinski E, Hortnagel K, Rudolph G, Boergen KP, Kampik A, Meitinger T. 1996. Discordant monozygotic twins with the Schimmelpenning-Feuerstein-Mims syndrome. *Clin Genet* 50:393–397.
- Sethupathy P, Borel C, Gagnebin M, Grant GR, Deutsch S, Elton TS, Hatzigeorgiou AG, Antonarakis SE. 2007. Human microRNA-155 on chromosome 21 differentially interacts with its polymorphic target in the AGTR1 3' untranslated region: A mechanism for functional single-nucleotide polymorphisms related to phenotypes. *Am J Hum Genet* 81:405–413.
- Sher S, Cole P, Kaufman Y, Hatef DA, Hollier L. 2008. Craniosynostotic variations in syndromic, identical twins. *Ann Plast Surg* 61:290–293.
- Shotelersuk V, Tift CJ, Vacha S, Peters KF, Biesecker LG. 1999. Discordance of oral-facial-digital syndrome type 1 in monozygotic twin girls. *Am J Med Genet* 86:269–273.
- Shotelersuk V, Punyashthiti R, Srivuthana S, Wacharasindhu S. 2002. Kabuki syndrome: Report of six Thai children and further phenotypic and genetic delineation. *Am J Med Genet* 110:384–390.
- Shur N. 2009. The genetics of twinning: From splitting eggs to breaking paradigms. *Am J Med Genet Part C* 151C:105–109.

- Simpson H, Camcho-Hubner C, Sonksen P, Russell-Jones D. 1999. Clinical experience in identical twins discordant for adult-onset growth hormone deficiency. *J Endocrinol Invest* 22:61–63.
- Singh SM, Murphy B, O'Reilly R. 2002. Epigenetic contributors to the discordance of monozygotic twins. *Clin Genet* 62:97–103.
- Singh SM, Castellani CA, O'Reilly RL. 2009. Copy number variation showers in schizophrenia: An emerging hypothesis. *Mol Psychiatry* 14:356–358.
- Sisodiya SM, Marques W Jr, Everitt A, Sander JW. 1999. Male monozygotic twins discordant for periventricular nodular heterotopia and epilepsy. *Epilepsia* 40:248–250.
- Slagboom PE, Droog S, Boomsma DI. 1994. Genetic determination of telomere size in humans: A twin study of three age groups. *Am J Hum Genet* 55:876–882.
- Smith AC, Rubin T, Shuman C, Estabrooks L, Aylsworth AS, McDonald MT, Steele L, Ray PN, Weksberg R. 2006. New chromosome 11p15 epigenotypes identified in male monozygotic twins with Beckwith–Wiedemann syndrome. *Cytogenet Genome Res* 113:313–317.
- Sobue G, Ueno-Natsukari I, Okamoto H, Connell TA, Aizawa I, Mizoguchi K, Honma M, Ishikawa G, Mitsuma T, Natsukari N. 1994. Phenotypic heterogeneity of an adult form of adrenoleukodystrophy in monozygotic twins. *Ann Neurol* 36:912–915.
- Somkuti SG, Wachtel SS, Schinfeld JS, Jackson L, Tharapel AT, DiGeorge AM. 2000. 46, XY monozygotic twins with discordant sex: phenotype. *Fertil Steril* 74:1254–1256.
- Sorenson EJ. 2006. Monozygotic twins discordant for primary lateral sclerosis. *Amyotroph Lateral Scler* 7:123–125.
- Stoll C, Roth MP, Dott B, Bigel P. 1984. Discordance for skeletal and cardiac defect in monozygotic twins. *Acta Genet Med Gemellol (Roma)* 33:501–504.
- Streissguth AP, Dehaene P. 1993. Fetal alcohol syndrome in twins of alcoholic mothers: Concordance of diagnosis and IQ. *Am J Med Genet* 47:857–861.
- Subramaniam B, Naidu S, Reiss AL. 1997. Neuroanatomy in Rett syndrome: Cerebral cortex and posterior fossa. *Neurology* 48:399–407.
- Supprian T, Bengel D, Hofmann E, Fallgatter AJ, Franzek E. 2000. Cavum veli interpositi and psychotic disorder in a monozygotic twin. *Eur Arch Psychiatry Clin Neurosci* 250:76–78.
- Tawil R, Storvick D, Feasby TE, Weiffenbach B, Griggs RC. 1993. Extreme variability of expression in monozygotic twins with FSH muscular dystrophy. *Neurology* 43:345–348.
- Taylor DM, Thum MY, Abdalla H. 2008. Dichorionic triamniotic triplet pregnancy with monozygotic twins discordant for trisomy 13 after preimplantation genetic screening: Case report. *Fertil Steril* 90:2017.e5–e9.
- Thiede C, Prange-Krex G, Freiberg-Richter J, Bornhauser M, Ehninger G. 2000. Buccal swabs but not mouthwash samples can be used to obtain pretransplant DNA fingerprints from recipients of allogeneic bone marrow transplants. *Bone Marrow Transplant* 25:575–577.
- Tho SP, Jackson R, Kulharya AS, Reindollar RH, Layman LC, McDonough PG. 2007. Long-term follow-up and analysis of monozygotic twins concordant for 45,X/46,XY peripheral blood karyotype but discordant for phenotypic sex. *Am J Med Genet Part A* 143A:2616–2622.
- Tiberio G. 1994. MZ female twins discordant for X-linked diseases: A review. *Acta Genet Med Gemellol (Roma)* 43:207–214.
- Tong S, Caddy D, Short RV. 1997. Use of dizygotic to monozygotic twinning ratio as a measure of fertility. *Lancet* 349:843–845.
- Touliatou V, Fryssira H, Mavrou A, Kanavakis E, Kitsiou-Tzeli S. 2006. Clinical manifestations in 17 Greek patients with Goldenhar syndrome. *Genet Couns* 17:359–370.
- Toyoshima M, Maegaki Y, Yuasa I, Ohno K. 2006. Monozygotic twins discordant for Klippel-Feil syndrome. *Pediatr Neurol* 34:76–78.
- Tremblay JP, Bouchard JP, Malouin F, Theau D, Cottrell F, Collin H, Rouche A, Gilgenkrantz S, Abbadi N, Tremblay M. 1993. Myoblast transplantation between monozygotic twin girl carriers of Duchenne muscular dystrophy. *Neuromuscul Disord* 3:583–592.
- Tsukamoto H, Inui K, Taniike M, Kamiyama K, Hori M, Sumi K, Okada S. 1993. Different clinical features in monozygotic twins: A case of 7q-syndrome. *Clin Genet* 43:139–142.
- Tuckerman E, Webb T, Bunday SE. 1985. Frequency and replication status of the fragile X, fra(X)(q27-28), in a pair of monozygotic twins of markedly differing intelligence. *J Med Genet* 22:85–91.
- Uchida IA, deSa DJ, Whelan DT. 1983. 45,X/46,XX mosaicism in discordant: Monozygotic twins. *Pediatrics* 71:413–417.
- Valdes AM, Deary IJ, Gardner J, Kimura M, Lu X, Spector TD, Aviv A, Cherkas LF. 2008. Leukocyte telomere length is associated with cognitive performance in healthy women. *Neurobiol Aging*. DOI: 10.1016/j.neurobiolaging.2008.07.012
- Valleix S, Vinciguerra C, Lavergne JM, Leuer M, Delpech M, Negrier C. 2002. Skewed X-chromosome inactivation in monochorionic diamniotic twin sisters results in severe and mild hemophilia A. *Blood* 100:3034–3036.
- van der Knaap MS, Barth PG. 1994. Discordant infantile encephalopathy with symmetrical thalamic calcifications in identical twins. *Am J Med Genet* 52:218–222.
- van Dijk BA, Boomsma DI, de Man AJ. 1996. Blood group chimerism in human multiple births is not rare. *Am J Med Genet* 61:264–268.
- van Leeuwen DM, van AE, Gottschalk RW, Vlietinck R, Gielen M, van Herwijnen MH, Maas LM, Kleinjans JC, van Delft JH. 2007. Cigarette smoke-induced differential gene expression in blood cells from monozygotic twin pairs. *Carcinogenesis* 28:691–697.
- van 't ED, Lehn H, Derks EM, Hudziak JJ, Van Strien NM, Veltman DJ, de Geus EJ, Todd RD, Boomsma DI. 2007. A structural MRI study in monozygotic twins concordant or discordant for attention/hyperactivity problems: Evidence for genetic and environmental heterogeneity in the developing brain. *Neuroimage* 35:1004–1020.
- Van Thienen MN, Van der Auwera BJ. 1994. Monozygotic twins discordant for spondylocostal dysostosis. *Am J Med Genet* 52:483–486.
- Vargas JE, Cox GF, Korf BR. 2000. Discordant phenotype in monozygotic twins with Fryns syndrome. *Am J Med Genet* 94:42–45.
- Verona LL, Damian NG, Pavarina LP, Ferreira CH, Melo DG. 2006. Monozygotic twins discordant for Goldenhar syndrome. *J Pediatr (Rio J)* 82:75–78.
- Verougstraete CM, Libert JA, Dorchy HR. 1999. Discordant diabetic retinopathy in homozygous twins: The importance of good metabolic control. *J Pediatr* 134:658.
- Vidaeff AC, Delu AN, Silva JB, Yeomans ER. 2005. Monoamniotic twin pregnancy discordant for body stalk anomaly: Case report with nosologic implications. *J Ultrasound Med* 24:1739–1744.
- Vincent MC, Heitz F, Tricoire J, Bourrouillou G, Kuhlein E, Rolland M, Calvas P. 1999. 22q11 deletion in DGS/VCFS monozygotic twins with discordant phenotypes. *Genet Couns* 10:43–49.
- Virta JJ, Aalto S, Jarvenpaa T, Karrasch M, Kaprio J, Koskenvuo M, Raiha I, Viljanen T, Rinne JO. 2009. Voxel-based analysis of cerebral glucose metabolism in mono- and dizygotic twins discordant for Alzheimer disease. *J Neurol Neurosurg Psychiatry* 80:259–266.

- Wachtel SS, Somkuti SG, Schinfeld JS. 2000. Monozygotic twins of opposite sex. *Cytogenet Cell Genet* 91:293–295.
- Wat MJ, Shchelochkov OA, Holder AM, Breman AM, Dagli A, Bacino C, Scaglia F, Zori RT, Cheung SW, Scott DA, Kang SH. 2009. Chromosome 8p23.1 deletions as a cause of complex congenital heart defects and diaphragmatic hernia. *Am J Med Genet Part A* 149A:1661–1677.
- Watkiss E, Webb T, Rysiecki G, Girdler N, Hewett E, Bunday S. 1994. X inactivation patterns in female monozygotic twins and their families. *J Med Genet* 31:754–757.
- Watson WJ, Katz VL, Albright SG, Rao KW, Aylsworth AS. 1990. Monozygotic twins discordant for partial trisomy 1. *Obstet Gynecol* 76:949–951.
- Webb T, Mead S, Beck J, Uphill J, Pal S, Hampson S, Wadsworth JD, Mena ID, O'Malley C, Wroe S, Schapira A, Brandner S, Collinge J. 2009. Seven-year discordance in age at onset in monozygotic twins with inherited prion disease (P102L). *Neuropathol Appl Neurobiol* 35:427–432.
- Weed KH, MacEwen CJ, McGhee CN. 2006. The variable expression of keratoconus within monozygotic twins: Dundee University Scottish Keratoconus Study (DUSKS). *Cont Lens Anterior Eye* 29:123–126.
- Weiss E, Loevy H, Saunders A, Pruzansky S, Rosenthal IM. 1982. Monozygotic twins discordant for Ullrich-Turner syndrome. *Am J Med Genet* 13:389–399.
- Weksberg R, Shuman C, Caluseriu O, Smith AC, Fei YL, Nishikawa J, Stockley TL, Best L, Chitayat D, Olney A, Ives E, Schneider A, Bestor TH, Li M, Sadowski P, Squire J. 2002. Discordant KCNQ1OT1 imprinting in sets of monozygotic twins discordant for Beckwith–Wiedemann syndrome. *Hum Mol Genet* 11:1317–1325.
- West PM, Love DR, Stapleton PM, Winship IM. 2003. Paternal uniparental disomy in monozygotic twins discordant for hemihypertrophy. *J Med Genet* 40:223–226.
- Wieczorek D, Ludwig M, Boehringer S, Jongbloet PH, Gillessen-Kaesbach G, Horsthemke B. 2007. Reproduction abnormalities and twin pregnancies in parents of sporadic patients with oculo-auriculo-vertebral spectrum/Goldenhar syndrome. *Hum Genet* 121:369–376.
- Wilkinson JL, Holt PA, Dickinson DF, Jivani SK. 1979. Asplenia syndrome in one of mono-zygotic twins. *Eur J Cardiol* 10:301–304.
- Winchester B, Young E, Geddes S, Genet S, Hurst J, Middleton-Price H, Williams N, Webb M, Habel A, Malcolm S. 1992. Female twin with Hunter disease due to nonrandom inactivation of the X-chromosome: A consequence of twinning. *Am J Med Genet* 44:834–838.
- Wong AH, Gottesman II, Petronis A. 2005. Phenotypic differences in genetically identical organisms: The epigenetic perspective. *Hum Mol Genet* 14(SpecNo. 1): R11–R18.
- Wu E, Vargevik K, Slavotinek AM. 2007. Subtypes of frontonasal dysplasia are useful in determining clinical prognosis. *Am J Med Genet Part A* 143A:3069–3078.
- Yamagishi H, Ishii C, Maeda J, Kojima Y, Matsuoka R, Kimura M, Takao A, Momma K, Matsuo N. 1998. Phenotypic discordance in monozygotic twins with 22q11.2 deletion. *Am J Med Genet* 78:319–321.
- Yamazawa K, Kagami M, Fukami M, Matsubara K, Ogata T. 2008. Monozygotic female twins discordant for Silver-Russell syndrome and hypomethylation of the H19-DMR. *J Hum Genet* 53:950–955.
- Young ID, Dalgleish R, MacKay EH, MacFadyen UM. 1988. Discordant expression of the G syndrome in monozygotic twins. *Am J Med Genet* 29:863–869.
- Youssefian H, Pyeritz RE. 2002. Mechanisms and consequences of somatic mosaicism in humans. *Nat Rev Genet* 3:748–758.
- Zech NH, Wisser J, Natalucci G, Riegel M, Baumer A, Schinzel A. 2008. Monochorionic-diamniotic twins discordant in gender from a naturally conceived pregnancy through postzygotic sex chromosome loss in a 47,XXY zygote. *Prenat Diagn* 28:759–763.
- Zenteno JC, Perez-Cano HJ, Aguinaga M. 2006. Anophthalmia-esophageal atresia syndrome caused by an SOX2 gene deletion in monozygotic twin brothers with markedly discordant phenotypes. *Am J Med Genet Part A* 140A:1899–1903.
- Zneimer SM, Schneider NR, Richards CS. 1993. In situ hybridization shows direct evidence of skewed X inactivation in one of monozygotic twin females manifesting Duchenne muscular dystrophy. *Am J Med Genet* 45:601–605.