

## **Hemodynamic and hemorheological profiles in women with proteinuric hypertension of pregnancy and in pregnant controls**

L. Heilmann and U. Siekmann

Department of Obstetrics and Gynecology, University of Essen, FRG

**Summary.** We obtained blood samples from 52 patients with pre-eclampsia and from 40 pregnant controls for measurement of plasma urate levels, hematocrit, white cell count and various hemorheological parameters (see Table 3). We also used impedance cardiography to measure cardiac output in both groups and from the results derived values for total peripheral resistance and oxygen transport. Central venous pressure was measured with a superior vena cava catheter in patients with pre-eclampsia but not in controls. Women with pre-eclampsia had significantly lower cardiac output and central venous pressure when compared with a control group. A modest correlation was observed between central venous pressure and cardiac output. The majority of pre-eclamptic patients had significantly raised hematocrit, leucocyte count, uric acid and red cell aggregation. Red cell deformability was significantly decreased in patients with pre-eclampsia. Most patients with severe pre-eclampsia (BP diast. > 100 mmHg) had a low Antithrombin III and colloid osmotic pressure level. The leucocyte count was raised when compared with the women with moderate pre-eclampsia. Oxygen delivery was reduced in patients with pre-eclampsia because of impaired rheological properties of their blood.

**Key words:** Pre-eclampsia – Hemorheology – Cardiac output – Impedance Cardiography

### **Introduction**

Pregnancy-induced proteinuric hypertension (or pre-eclampsia) is associated with exaggerated arterial reactivity (Gant 1973). Although the rate of tissue perfusion is dependent on cardiovascular factors recent studies (Guyton 1963,

Chien 1972, Schmid-Schönbein 1976) have indicated that the rheological properties of blood can also play a significant role in the regulation of blood flow. Proteinuric hypertension may also be associated with a hyperviscosity syndrome (Mathews 1974, Hobbs et al. 1982, Buchan 1982, Oggolter-Siekmann et al. 1984).

The viscosity of normal blood is a function of hematocrit, plasma viscosity, red cell aggregation and red cell deformability with the last two parameters being flow dependent. In small vessels the concentration and adhesiveness of white cells can also exert a significant effect on flow resistance (Chien 1985, Braide 1984). Postcapillary venules, small veins and the intervillous space of the placenta have the lowest shear stress in the circulation and represent the most likely sites of red cell aggregation (Chien 1969, Scholz et al. 1975). With normal cardiovascular and respiratory function, hematocrit and blood viscosity reflect the rate of oxygen delivery (Chien 1987). It would appear that the physiological mechanisms for the control of red cell volume and plasma are coordinated such that the hematocrit stays in the optimum range.

The optimum hematocrit in pregnancy was given by Garn et al. (1981) and by Murphy et al. (1986) as between 30% and 38%. This entails a hemoglobin content of 10.0–13.2 g/dl (Murphy et al. 1986) or of 10.4–13.3 g/dl (Garn et al. 1981). Hemoglobin values above 13.2 g/dl or a hematocrit above 38% were associated with a statistically significant rise in fetal pathology (perinatal death, premature babies, low birth weight) and also with pregnancy-induced hypertension. In severe proteinuric hypertension and fetal pathology Sagen et al. (1984) found hemoconcentration (Hb more than 13 g/dl) in 82% of cases.

The aim of the present study was to examine the relationship between the rheological properties of blood and hemodynamic and clinical data in a group of pre-eclamptic patients.

## Patients and methods

The study was performed on 52 women with pregnancy-induced proteinuric hypertension. No antihypertensive medication was administered before hemodynamic and hemorheological measurements. All patients had a blood pressure which was persistently above 140/85 mmHg and more than 0.1% protein in a 24 h urine sample. Hypertension was classified as severe when the diastolic blood pressure was > 100 mmHg or signs of severe microangiopathy or the HELLP-Syndrome occurred consists of hemolysis, elevated liver enzymes and a low platelet count (Weinstein 1985). Poor perinatal outcome was defined as perinatal death or fetal acidosis ( $\text{pH} < 7.2$ ) and/or intrauterine growth retardation (IUGR). 40 women with uncomplicated pregnancies served as controls and contributed a total 235 samples. Samples were longitudinal after the 24th week of gestation until term.

### *Hemodynamic monitoring*

All pre-eclamptic patients (but no controls) had an intravenous right superior vena cava catheter with continuous venous pressure monitoring. Cardiac output was derived from calculation of stroke volume multiplied by heart rate. Stroke volume was assessed by the formula of Kubicek (1974). Stroke volume was determined from the signals recorded by an impedance cardiograph (Diefenbach GmbH, Frankfurt). In the Kubicek-formula

$$SV = \frac{R \times L^2 \times T \times dz/dt}{(z_0)^2}$$

was R the resistivity of blood, L the distance between these electrodes,  $dz/dt$  the maximum impedance rate changes, T the ventricular ejection time and  $z_0$  the basic chest impedance. Heart rate was derived from an ECG.

#### *Hemorheological parameters*

Venous blood was collected in an EDTA tube from the antecubital vein with minimal occlusion, centrifuged at 3500 for 10 min, plasma was aspirated, buffy coat removed by vacuum suction, the suspension was washed twice in isotonic buffer, pH 7.4 and resuspended in the same medium containing 1 g/100 ml of albumin. The hematocrit was adjusted to 10% and the percentage of residual leucocytes after cotton wool filtration was < 15% (Diepenhorst et al. 1972, Stuart 1985).

#### *Erythrocyte deformability*

was studied by determining the pressure – flow relationship of RBC suspensions passed through polycarbonate sieves with a pore diameter of 5  $\mu\text{m}$ . The analyses of the initial flow curve was performed by a Filtrometer MF 4 as a function of time and the flow curve is automatically received, processed and printed. In order to derive an average deformation behavior of the entire population we measured the initial relative resistance at the beginning of the filtration process.

*Red cell aggregation* was determined at 22°C in the "Rheoaggregometer" Typ MAI (Myrhenne GmbH, Roetgen, FRG) with a corrected hematocrit of 40%. This cone plate system rotates the blood 10 s at shear rate of 600  $\text{s}^{-1}$ , where after during 10 s the average aggregation is determined by the quantity of infrared light measured by photosensors. The data were processed by a microcomputer and the average aggregation registered on a digital display, expressed in arbitrary units. 5 measurements were performed on each blood sample and the average values was assessed.

*Plasma viscosity* was measured with a capillary viscosimeter at 37°C. The calibration factor was calculated with 0.033. Three measurements were performed on each plasma sample.

*Hematocrit, leucocytes and platelets* were measured using a Coulter Counter.

*Serum osmolality* were determined by freezing point depression (Osmometer, Gonotec, Berlin, FRG).

*Colloid osmotic pressure* with the use of a Colloid Osmometer (Knauer AG, Bad Homburg, FRG).

*Plasma fibrinogen* was measured according to the method Clauss (1957).

*Plasma fibronectin* was measured by radial immunodiffusion using antiserum from Behringwerke (Marburg, FRG).

*Antithrombin III* was measured using chromogenic peptide substrates (Boehringer, Mannheim, FRG).

*Plasma urate* was assayed by the uricase-method.

*Systemic oxygen transport capacity* or rate of oxygen delivery to tissues is equal to the product of cardiac output, hematocrit and the Huefner factor (= 1.34  $\text{mlO}_2/\text{gHb}$ ).

Means and standard errors were calculated for the data from each group. Since it was not clear whether the data were normally distributed, Wilcoxon rank-sum testing was used for statistical analysis.

## Results

### *Clinical data*

During the study period 52 Primiparae with proteinuric hypertension (but not the controls) were studied by catheterisation of the superior vena cava. The mean gestational age was  $33.8 \pm 4.2$  weeks (range 25 to 40 weeks) and the mean maternal age was  $33.8 \pm 5.3$  (range 18 to 39 years). The control group consisted of 40 women with uncomplicated pregnancy (Table 1): the mean gestational age was  $38.5 \pm 2.4$  (range 36 to 41 weeks) and the maternal age  $32.4 \pm 4.6$  (17 to 38 years). Seven patients presented signs of microangiopathy (HELLP-Syndrome). The perinatal mortality in the study group was 20.3%. Five patients had an intrauterine fetal death and there were four neonatal deaths. All infants survived in the control group and in the group with the HELLP-Syndrome.

### *Hemodynamic data*

Mean values for arterial blood pressure, central venous pressure, cardiac output and total peripheral resistance are summarized in Table 2. Cardiac output in patients with proteinuric hypertension was low (Mean:  $3.11 \pm 1.5$  l/min) compared with normal pregnancy ( $6.56 \pm 0.79$  l/min). Forty-one percent ( $n = 30$ ) had a Cardiac output  $> 3.0$  l/min compared with 82% ( $n = 32$ ) in normal pregnancy. The mean central venous pressure was  $1.47 \pm 4.0$  cm (range  $-6$  to  $+14$ ) and was similar in patients with moderate and severe hypertension. 38% ( $n = 27$ ) of patients in the study group had hypovolemia with a central

**Table 1.** Clinical data and pregnancy outcome in controls and patients with pre-eclampsia

Parameters	Controls	HELLP-Syndrome	Pre-eclampsia	
			RR diast. $\leq 100$ mmHg	RR diast. $> 100$ mmHg
Number (N)	40	7	31	14
Mean age (years)	$32.4 \pm 4.6$ (17–38)	$29.1 \pm 5.3$ (21–36)	$26.7 \pm 4.8$ (18–38)	$28.1 \pm 6.4$ (20–39)
Mean parity (N)	$2.3 \pm 1.2$ (0–7)	0	0	0
Mean gestational age (weeks)	$38.5 \pm 2.4$ (36–41)	$33.7 \pm 4.4$ (27–38)	$33.8 \pm 4.2$ (25–40)	$34.0 \pm 4.5$ (28–40)
Birthweight (g)	$3181 \pm 441$ (2600–3800)	$2104 \pm 1286$ (710–4450)	$1874 \pm 941$ (500–3700)	$1672 \pm 783$ (820–3180)
pH-cord blood	$7.25 \pm 0.1$	$7.25 \pm 0.04$	$7.21 \pm 0.11$	$7.17 \pm 0.11$
Perinatal mortality (N)	0	0	5	4

**Table 2.** Hemodynamic parameters in controls and patient with pre-eclampsia

Parameters	Controls (A)	HELLP- Syndrome (B)	Pre-eclampsia	
			RR diast. ≤ 100 mm Hg (C)	RR diast. > 100 mm Hg (D)
Number	40	7	31	14
RR syst. (mm Hg)	115.4 ± 8.4	160.7 ± 13.0	156.1 ± 9.2	175.7 ± 19.4
RR diast. (mm Hg)	75.4 ± 8.9	91.4 ± 14.6	93.0 ± 7.8	111.2 ± 5.3
Proteinuria (‰)	0	0.5–12.0	0.1–4.0	0.1–5.0
Cardiac output (l/min)	6.6 ± 0.8	2.4 ± 0.5	3.4 ± 1.8 <sup>a</sup>	2.9 ± 1.1 <sup>a</sup>
Total peripheral resistance (dynes/cm <sup>5</sup> × 10 <sup>3</sup> )	1.1 ± 0.1	4.0 ± 0.9	3.2 ± 1.3 <sup>a</sup>	4.3 ± 1.6 <sup>a</sup>
Central venous pressure (mm Hg)	n. m.	–5 to +8	–2 to +14	–6 to +9
Oxygen transport capacity (l/min)	289.3 ± 78.4	105.5 ± 24.4	173.6 ± 83.0	152.6 ± 67.1 <sup>a</sup>

<sup>a</sup> Avs B;  $P < 0.05$ . n. m.: not measured

venous pressure of < 0 cm. The relationship between central venous pressure and cardiac output was  $r = 0.3$  ( $P < 0.05$ ). Two women had signs of pulmonary oedema with a high CVP, a high cardiac output and a low hematocrit.

The hemodynamic values for normal pregnancy were obtained from a longitudinal study (Siekmann 1985). The technique used in that study was impedance cardiography (Kubicek et al. 1974) and we consider the normal range for cardiac output in the second and third trimesters to be 4.5–8.5 l/min. The total peripheral resistance in normal pregnancy is very low ( $1.05 \pm 0.11$  dynes/cm<sup>5</sup> × 10<sup>3</sup>) when compared with patients with proteinuric hypertension (86.3% above  $2.2$  dynes/cm<sup>5</sup> × 10<sup>3</sup>).

### *Hemorheologic data*

The hemorheological results are presented in Table 3. Red cell aggregation was significantly higher in patients with proteinuric hypertension than in normal pregnancy ( $P < 0.01$ ), but plasma viscosity and serum osmolality were similar. The difference in hematocrit and leucocyte count between patients and controls was significant ( $P < 0.05$ ). COP and erythrocyte flow rate were decreased in proteinuric hypertension but we could not show any significant differences between moderate and severe forms of hypertension. Plasma fibrinogen was only higher in the severe hypertensive group compared with normals or moderate hypertensive patients. Plasma fibronectin and uric acid were raised in proteinuric hypertension ( $P < 0.001$ ).

**Table 3.** Hemorheological data for patients with normal pregnancy, women with HELLP-syndrome and patients with pre-eclampsia

Parameters	Controls (A)	HELLP (B)	Pre-eclampsia		Patients with unsuccessful pregnancy (E)
			(C) RR diast. ≤ 100 mmHg	(D) RR diast. > 100 mmHg	
Number	40	7	31	14	9/52 (20.3%)
Hkt (%)	35.2 ± 3.4	32.9 ± 4.8	38.9 ± 4.3	39.2 ± 4.5 <sup>a</sup>	41.1 ± 5.9 <sup>b</sup>
White cells (/nl)	9.1 ± 1.8	12.6 ± 5.4	11.6 ± 3.4	14.9 ± 5.8 <sup>a</sup>	13.1 ± 2.5
Platelets (mm <sup>3</sup> )	309 ± 85	82 ± 39	223 ± 67	221 ± 83	183 ± 50
Red cell aggre- gation (–)	20.3 ± 4.9	23.7 ± 5.8	28.7 ± 4.7	27.5 ± 8.4 <sup>a</sup>	30.4 ± 5.9
Plasma viscosity (cst)	1.31 ± 0.04	1.30 ± 0.06	1.28 ± 0.1	1.32 ± 0.07	1.32 ± 0.1
Serum osmolality (mmol/kg)	288 ± 6.8	301 ± 9.1	294 ± 10.2	293 ± 5.3	292 ± 10.3
COP (mmHg)	21.0 ± 3.2	16.1 ± 3.6	18.2 ± 3.9	16.5 ± 3.8 <sup>a</sup>	17.4 ± 3.8
Fibrinogen (mg%)	383 ± 57	248 ± 152	393 ± 71	418 ± 100 <sup>a</sup>	366 ± 105
Fibronectin (mg%)	28.5 ± 1.8	44.3 ± 10.7	52.9 ± 14.4	56.9 ± 12.6 <sup>a</sup>	56.2 ± 12.8
AT III (I. U./ml)	11.5 ± 2.3	7.1 ± 4.6	10.8 ± 2.2	8.2 ± 2.2 <sup>a</sup>	9.6 ± 4.3
Red cell deform- ability (μl/s)	10.7 ± 2.7 (4.8–19.7)	8.4 ± 4.1 (2.9–16.3)	7.8 ± 4.3 (0–18.8)	5.1 ± 4.8 <sup>a</sup> (0.08–15.9)	3.9 ± 4.7 <sup>b</sup> (0.08–13.5)
Plasma urate (mg%)	3.37 ± 1.8	8.6 ± 2.9	7.8 ± 1.4	6.9 ± 1.4 <sup>a</sup>	6.8 ± 1.1

<sup>a</sup> Avs D:  $P < 0.05$ ; <sup>b</sup> Avs E:  $P < 0.01$ 

In cases with perinatal loss of the fetus, maternal red cell deformability was very low ( $3.9 \pm 4.7 \mu\text{l/s}$ ) and the mean hematocrit at delivery (at a mean of 29 weeks gestation) was high at  $41.1 \pm 5.9\%$ . The lower hematocrit of normal pregnancy is associated with a decreased oxygen capacity but with an increased oxygen delivery. In cases with proteinuric hypertension a hematocrit above normal is associated with reduced systemic oxygen transport rate. This fall in oxygen availability is due to impendence of the venous return (low CVP and CO) as a result of increased blood viscosity.

We studied 7 patients with hemolysis, elevated liver enzyme levels and low platelet count (HELLP-Syndrome). They had decreased plasma fibrinogen and Antithrombin III-level and their cardiac output was  $2.4 \pm 0.5 \text{ l/min}$  and the calculated vascular resistance  $4.0 \pm 0.9 \text{ dyn/cm}^{-5} \times 10^3$  (Table 2).

## Discussion

Previous hemodynamic investigations have revealed a wide range of cardiac output in cases of severe pre-eclampsia. 59% of our pre-eclampsia patients had a low cardiac output. The difference between our results and those of Cotton et al. (1988) may be due to differences in methodology. Most studies use dye dilution (Lees et al. 1967) or thermodilution (Hankin et al. 1984, Groenendijk et al. 1984, Cotton et al. 1987, Rafferty et al. 1980). More recently (Siekman 1985, Davies et al. 1986) impedance cardiography – a noninvasive technique for the measurement of cardiac stroke volume has been used. Unlike Atkins et al. (1981) and de Swiet (1986) we found that the impedance method permitted repeated measurements to study changes of cardiac output in the same patient during pregnancy (Siekman 1985). The results of Milson et al. (1983) in late pregnancy show a strong correlation ( $r = 0.93$ ) between impedance and dye dilution technique. Groenendijk et al. (1984) and Wallenburg (1988) suggest that pre-eclampsia is associated with a low cardiac output and a reduced blood volume. This would be in agreement with the view that the plasma volume is reduced in pre-eclampsia (Soffronoff et al. 1977, Gallery et al. 1979, 1981). A decrease in venous return can occur as a result of increased peripheral resistance, vasoconstriction, cava compression or increased blood viscosity. The fact that the viscosity plays an important role in determining the cardiac output or peripheral resistance is generally accepted (Heilmann et al. 1981, Messmer 1981, Goslinga 1984). Low flow states lead to an increase in blood viscosity at a given hematocrit. Under these conditions  $O_2$ -delivery is better maintained at a lower hematocrit such as is normal in pregnancy. Experiments have shown that a high hematocrit and viscosity are associated with low oxygen delivery to tissue (Guyton 1963). In agreement with Chien (1972) the oxygen capacity of blood in the low flow areas in pre-eclamptic patients is reduced by 50%. A logical consequence is that at every flow rate, a particular value of hematocrit will be associated with an optimal viscosity and a maximal  $O_2$ -transport.

In pre-eclampsia statistically significant differences from normal pregnancy were observed in erythrocyte aggregation, hematocrit and white cell count. The degree of hemoconcentration we found in our patients is in agreement with previously published results (Sagen 1984, Heilmann 1981, Kiesewetter 1983, Buchan 1982, Thornburn et al. 1982, Siekman et al. 1984). A high hematocrit and a low central venous pressure are the results of inadequate plasma volume expansion. Increased vascular and rheological resistance lead to an endothelial damage with release of mediators and disorders of hemostasis (Saleh et al. 1987, 1988, Oian et al. 1985, Poldre 1987, Weinstein 1985). The release reaction is not only started by the leucocyte (endothelial interaction) but also by the polypeptide, Interleukin-1. Interleukin-1 makes neutrophils and hepatocytes release high molecular weight acute phase proteins, i.e. fibrinogen and procoagulant products (Friedmann 1988, Martin 1988).

Fibronectin is a large glycoprotein of 440000 Daltons, presenting either in soluble form in blood or in insoluble form in connective tissue. The fibronectin rise in pre-eclamptic patients would suggest that an endothelial cell abnormality may occur early in patients with gestational hypertension. The mechanisms are

unclear: changes in rates of synthesis and/or consumption, as well as variation in plasma volume have been discussed as causes (Lazarchick et al. 1986).

Red cell aggregation brought about by changes in the suspension stability of blood (Chien 1972) was thought to be a possible mechanism for precapillary plugging. It is now clear that the forces disrupting aggregation in the fastmoving precapillary stream are greater than the forces holding the aggregate together, so that this possibility appears unlikely (Lowe 1987). However, red cell aggregation in the postcapillary and intervillous vessels with resultant stasis is a real possibility. Plasma viscosity did not seem to be influenced by the severity of the hypertension.

In pre-eclamptic patients "hematological stress" (Dinaretto 1984, Stuart et al. 1983, Stuart 1985) leads to arise in the leucocyte count and fibrinogen level. Both factors may decrease the erythrocyte filtration rate because leucocytes show poor deformability (Bagge et al. 1977) and plasma proteins decrease erythrocyte flow (Stuart 1985) through narrow channels in vitro and in vivo. We used Nucleporefilters in our experiments as these have pores similar in diameter to capillaries. Although red cell filterability in pre-eclamptic and normal pregnancies overlapped poor perinatal outcome was associated with decreased flow. The increased erythrocyte rigidity may result from a disturbance of cellular calcium metabolism (Cunningham et al. 1985) or changes in the lipid composition of cell membrane (Worley et al. 1982).

The increased leucocyte count in patients with pre-eclampsia may occlude small vessels and could be a further factor impairing intervillous flow. Trapping of white blood cells may be due to factors released by earlier white cells after an interaction with endothelial and trophoblast cells. These activated white cells would be expected to have altered mechanical and adhesive properties (Ernst et al. 1987). Factors released in such circumstances might include leucotriens, thromboxanes, platelet activating factor and leucocyte derived free radicals (Harlan 1985).

The rheological changes found in this study would be expected to affect the flow of white cells through the microcirculation. Further studies are needed to test specific markers for activation of white cells and to examine whether white cells abnormalities correlate with the clinical severity of pre-eclampsia.

## References

1. Atkins AF, Watt JM (1981) A longitudinal study of cardiovascular dynamic changes throughout pregnancy. *Eur J Obstet Gynecol Reprod Biol* 12:215–224
2. Bagge U, Stolack R, Attefors R (1977) Granulocyte rheology: experimental studies in an in vitro micro flow-system. *Adv Microcirc* 7:29–48
3. Braide M, Ammundson B, Chien S, Bagge U (1984) Quantitative studies on the influence of leucocytes on the vascular resistance in a skeletal muscle preparation. *Microvasc Res* 27:331–352
4. Buchan PC (1982) Pre-eclampsia a hyperviscosity syndrome. *Am J Obstet Gynecol* 142:111–112
5. Chien S (1969) Blood rheology and its relation to flow resistance and transcapillary exchange with special referece to shock. *Adv Microcirc* 2:89–103
6. Chien S (1972) Present state of blood rheology. In: Messmer K, Schmid-Schönbein H (eds) *Hemodilution. Theoretical basis and clinical application*. Karger, Basel München Paris London New York Sydney



7. Chien S (1985) Role of blood cells in microcirculatory regulation. *Microvasc Res* 29:129–151
8. Chien S (1987) Physiological and pathophysiological significance of hemorheology. In: Chien S, Dormandy J, Ernst E, Matrai A (eds) *Clinical hemorheology*. Martinus Lighoff, pp 125–164
9. Clauss A (1957) Gerinnungsphysiologische Schnellmethode zur Bestimmung des Fibrinogens. *Acta Haemat* 17:237–246
10. Cotton DB, Lee W, Huhto JC, Dorman KF (1988) Hemodynamic profile of severe pregnancy-induced hypertension. *Am J Obstet Gynecol* 158:523–529
11. Cunningham FG, Lowe Th, Guss St, Mason R (1985) Erythrocyte morphology in women with severe pre-eclampsia and eclampsia. *Am J Obstet Gynecol* 153:358–363
12. Davies P, Francis RI, Dockes MF, Watt JM, Crawford JS (1986) Analysis of impedance cardiography longitudinally applied in pregnancy. *Br J Obstet Gynecol* 93:717–720
13. De Swiet M, Talbert DG (1986) The measurement of cardiac output by electrical impedance plethysmography in pregnancy. Are the assumptions valid? *Br J Obstet Gynecol* 93:721–726
14. Diepenhorst P, Sprokholt R, Prius MK (1972) Removal of leucocytes from whole blood and erythrocytes suspensions by filtration through cotton wool. *Vox Sang* 23:308–320
15. Dinorell CA (1984) Interleukin-1 and the pathogenesis of the acute phase response. *N Engl J Med* 311:1413–1418
16. Ernst E, Hammerschmidt DE, Bagge U, Matrai A, Dormandy JA (1987) Leucocytes and the risk of ischemic disease. *JAMA* 257:2318–2324
17. Friedmann StA (1988) Pre-eclampsia: a review of the role of prostaglandins. *Obstet Gynecol* 71:122–137
18. Gallery EDM, Hunyor S, Gyory AZ (1979) Plasma volume contraction: a significant factor in both pregnancy-associated hypertension (pre-eclampsia) and chronic hypertension in pregnancy. *O J Med* 192:593–602
19. Gallery EDM, Delprado W, Gyory AZ (1981) Antihypertensive effect of plasma volume expansion in pregnancy-associated hypertension. *Aust N Z J Med* 11:20–24
20. Gant NF, Daley GL, Chands S, Whalley PJ, Mac Donald PC (1973) A study of angiotension II pressor response throughout primigravid pregnancy. *J Clin Invest* 52:2682–2686
21. Garn SM, Ridella SA, Petzold AS, Falkner F (1981) Maternal hematologic levels and pregnancy outcomes. *Sem Perinat* 5:155–162
22. Goslinga H (1984) Blood viscosity and shock. Springer, Berlin Heidelberg New York
23. Groenendijk R, Trimpos JBJ, Wallenburg HCS (1984) Hemodynamic measurement in pre-eclampsia: preliminary observations. *Am J Obstet Gynecol* 150:232–236
24. Guyton AC (1963) Effect of peripheral resistance and capacitance on venous return. In: Guyton AC (ed) *Cardiac output and its regulation*. WB Saunders, Philadelphia London, pp 209–220
25. Hankin GDV, Wendele GD, Cunningham FG, Leveno KJ (1984) Longitudinal evaluation of hemodynamic changes in eclampsia. *Am J Obstet Gynecol* 150:306–512
26. Harlan JM (1985) Leucocyte-endothelial interactions. *Blood* 65:513–525
27. Haubius GDV, Wendel GP, Cunningham FG, Leveno KJ (1984) Longitudinal evaluation of hemodynamic changes in eclampsia. *Am J Obstet Gynecol* 150:506–512
28. Heilmann L, Siekmann U, Schmid-Schönbein H, Ludwig H (1981) Hemoconcentration and pre-eclampsia. *Arch Gynecol* 231:7–14
29. Hobbs JB, Oats JN, Palmer AA (1982) Whole blood viscosity in pre-eclampsia. *Am J Obstet Gynecol* 142:288–292
30. Kiese Wetter H (1983) Bedeutung haemorheologischer Messverfahren für Diagnostik und Therapie von Durchblutungsstörungen. *Essentialia* 1:151–164
31. Kubicek WG, Notthe FJ, Ramos MU (1974) The minnesota impedance cardiograph – theory and applications. *Biomed Eng* 9:410–416
32. Lazarschick J, Stubbs ThM, Romein L, van Dorsten JP, Loadholt CB (1986) Predictive value of fibronectin levels in normotensive gravid women destined to become pre-eclamptic. *Am J Obstet Gynecol* 154:1050–1052
33. Lies MM, Scott DB, Kerr MG, Taylor SA (1967) The circulatory effects of remember postural change in late-pregnancy. *Clin Sci* 32:453
34. Lowe GDO (1987) Blood rheology in vitro and in vivo. In: Lowe GDO (ed) *Blood rheology and hyperviscosity syndromes*. Bailliere Tindall, London Philadelphia Toronto Sydney Tokyo, pp 597–636

35. Martin M, Resch K (1988) Interleukin 1: more than a mediator between leucocytes trends in pharm. Sciences 9:171–177
36. Mathews JD, Mason TW (1974) Plasma viscosity and pre-eclampsia. *Lancet* I:409
37. Messmer K (1981) Compensatory mechanismus for acute dilutional anemia. In: Schmid-Schönbein H, Messmer K, Rieger H (eds) Hemodilution and flow improvement. Karger, Basel, pp 31–42
38. Milson J, Forssman L, Biber B, Dottari O, Sivertsson R (1983) Measurement od cardiac stroke volume during cesarean section: a comparison between impedance cardiography and the dye dilution technique. *Acta Anaesthesiol Scand* 27:421–426
39. Murphy JF, Riordan JO, Newcombe RG, Coles EC, Pearson JF (1986) Relation of hemoglobin levels in first and second trimesters to outcome or pregnancy. *Lancet* I:992–994
40. Øion P, Omsjo I, Maltau JM, Østernd B (1985) Increased sensitivity to thromboplastic synthesis in blood monocytes from pre-eclamptic patients. *Br J Obstet Gynecol* 92:511–517
41. Oggolter-Siekmann H (1985) Die prognostische Bedeutung hämorrheologischer Parameter zur Früherkennung der Gestose. Thesis Essen
42. Poldre PA (1985) Haptoglobin helps diagnose the HELLP syndrome. *Am J Obstet Gynecol* 157:1267
43. Rafferty TD, Berkowitz RL (1980) Hemodynamics in patient with severe toxemia during labor and delivery. *Am J Obstet Gynecol* 138:263
44. Sagen N, Nilsen ST, Kim HC, Bergsjo P, Koller O (1984) Maternal hemoglobin concentration is closely related to birth weight in normal pregnancies. *Acta Obstet Gynecol Scand* 63:245–248
45. Sagen N, Nilsen ST, Kim HC, Koller O, Bergsjo P (1984) The predictive value of total estriol, HPL and Hb on perinatal outcome in severe pre-eclampsia. *Acta Obstet Gynecol Scand* 63:603–608
46. Saleh AA, Bottoms F, Welch RA, Ali AM, Marione FG, Mammen EF (1987) Pre-eclampsia, delivery and the hemostatic system. *Am J Obstet Gynecol* 157:331–336
47. Saleh AA, Bottoms SF, Norman G, Farag A, Mammen EF (1988) Hemostasis in hypertensive disorders of pregnancy. *Obstet Gynecol* 71:719–722
48. Schmid-Schönbein H (1976) Microrheologys of erythrocytes, blood viscosity and the distribution of blood in the microcirculation. In: Guyton AC, Cowley AW (eds) Cardiovascular. Physiology II, Vol 9. University Park Press, Baltimore, pp 1–62
49. Scholz PM, Karis JH, Gump FE, Kinney FE, Chien S (1973) Correlation of blood rheology with vascular resistance in critically ill patients. *J Appl Physiol* 39:1008–1011
50. Siekmann U, Oggolter-Siekmann H, Heilmann L (1984) The rheological behavior of red blood cells in the course of pregnancy-induced hypertension. In: Heilmann L, Buchan PC (eds) Hemorheological disorders in obstetrics and neonatology. Schattauer, Stuttgart New York, pp 21–28
51. Siekmann U (1985) Die Beziehung zwischen der maternalen Hämodynamik und den Fließeigenschaften des Blutes in der Schwangerschaft. Habilitationsschrift, Essen
52. Soffronoff EC, Kaufmann BM, Connanghton JF (1977) Intravascular volume determination and fetal outcome in hypertensive disease of pregnancy. *Am J Obstet Gynecol* 127:4–9
53. Stuart J, Kenny MW, Inglu TCM (1983) Erythrocyte filterability in normal pregnancy and pre-eclampsia. *Br J Haematol* 53:353–355
54. Stuart J (1985) Erythrocyte rheology. *J Clin Pathol* 38:965–977
55. Stuart J, Marshall J (1985) Therapeutic approaches to abnormalities of blood rheology in vascular disease. In: Das PC (eds) Supportive therapy in hematology. Nijhoff, Boston Dordrecht Lancaster, pp 351–364
56. Thornburn J, Drummond MM, Whigham KA (1982) Blood viscosity and hemostatic in late pregnancy pre-eclampsia and fetal growth retardation. *Br J Obstet Gynecol* 89:117–122
57. Wallenburg HCSC (1988) Hemodynamics in hypertensive pregnancy. In: Rubin PC (eds) Hypertension in pregnancy. Elsevier, Amsterdam New York Oxford, pp 66–101
58. Weinstein L (1985) Pre-eclampsia/eclampsia with hemolysis, elevated liver enzymes and thrombocytopenia. *Obstet Gynecol* 66:657–660
59. Worley RJ, Hentschel WM, Comier C, Nutting S, Pead G, Zelenkiov K, Smith JB, Ash KO, Williams RR (1982) Increased sodium-lithium counter-transport in erythrocytes of pregnant women. *N Engl J Med* 307:412–426