

# Potassium Citrate Prevents Increased Urine Calcium Excretion and Bone Resorption Induced by a High Sodium Chloride Diet

DEBORAH E. SELLMAYER, MONIQUE SCHLOETTER, AND ANTHONY SEBASTIAN

*Division of Endocrinology (D.E.S., A.S.) and the General Clinical Research Center (M.S., A.S.), University of California, San Francisco, California 94105*

The amount of sodium chloride in the diet of industrialized nations far exceeds physiological requirements. The impact of abundant dietary salt on skeletal health has yet to be established, but is potentially detrimental through increased urinary calcium losses. We examined the effect of increased dietary sodium chloride on urine calcium excretion and bone turnover markers in postmenopausal women and, further, whether potassium citrate attenuates the effects of increased dietary salt. Postmenopausal women ( $n = 60$ ) were adapted to a low-salt (87 mmol/d sodium) diet for 3 wk, then randomized to a high-salt (225 mmol/d sodium) diet plus potassium citrate (90 mmol/d) or a high-salt diet plus placebo for 4 wk. Urine calcium, urine N-telopeptide, urine cAMP, serum osteocalcin, and fasting serum PTH were measured at the end of the low- and high-salt diets. On the high salt plus placebo diet, urine calcium increased  $42 \pm 12$  mg/d (mean  $\pm$  SEM), but decreased

$8 \pm 14$  mg/d in the high salt plus potassium citrate group ( $P = 0.008$ , potassium citrate *vs.* placebo, unpaired  $t$  test). N-telopeptide increased  $6.4 \pm 1.4$  nanomoles bone collagen equivalents per millimole creatinine in the high salt plus placebo group and  $2.0 \pm 1.7$  nanomoles bone collagen equivalents per millimole creatinine in the high salt plus potassium citrate group ( $P < 0.05$ , potassium citrate *vs.* placebo, unpaired  $t$  test). Osteocalcin, PTH, and cAMP were not significantly altered. The addition of oral potassium citrate to a high-salt diet prevented the increased excretion of urine calcium and the bone resorption marker caused by a high salt intake. Increased intake of dietary sources of potassium alkaline salts, namely fruit and vegetables, may be beneficial for postmenopausal women at risk for osteoporosis, particularly those consuming a diet generous in sodium chloride. (*J Clin Endocrinol Metab* 87: 2008–2012, 2002)

OSTEOPOROSIS AND ITS sequelae pose a substantial health burden for postmenopausal women. One white woman in six will suffer a hip fracture during her lifetime, and mortality after hip fracture ranges from 12–20% in the first year. Modifications in nutrition can help reduce the risk of bone loss and fracture; for example, the benefit of adequate calcium and vitamin D intake is well documented (1, 2). Conversely, dietary sodium chloride has been considered potentially detrimental because increasing dietary sodium chloride increases urinary calcium excretion (3–9).

Renal calcium reabsorption is directly proportional to sodium reabsorption. When dietary sodium chloride is increased, the fractional reabsorption of sodium is decreased, leading to a parallel reduction in calcium reabsorption. Approximately 1 mmol calcium is excreted for every 100 mmol of sodium excreted (10). A net deficit of only 1 mmol/d of calcium would result in losing one third of the calcium contained in the typical adult skeleton in just over two decades unless a compensatory increase in intestinal calcium absorption occurred.

The ability to compensate for increased dietary sodium chloride may be related to age and menopausal status. In young men and premenopausal women, increased dietary sodium and consequent hypercalciuria induced an increase in calcitriol (1,25-dihydroxyvitamin D) levels and intestinal calcium absorption (11). However, postmenopausal women

did not demonstrate increased calcitriol levels (12), suggesting that older women may be unable to compensate for urinary calcium losses induced by sodium.

Although the relationship between dietary sodium chloride and calcium excretion has been consistently demonstrated, studies have been mixed with regard to the effect of dietary salt on bone density and bone turnover markers, and potential interventions have not been examined. Therapies that decrease urinary calcium excretion could potentially prevent the calciuric effect of increased dietary sodium chloride. Alkaline salts of potassium (e.g. potassium bicarbonate, potassium citrate) significantly reduce urinary calcium excretion in young men and both premenopausal and postmenopausal women (13–15), even in the setting of a high sodium intake (16). Alkaline salts of potassium are a particularly attractive intervention as they occur naturally in fruits and vegetables. Amounts that decrease urinary calcium excretion could be consumed by increasing intake of fruits and vegetables.

In this study, we investigated the impact of increased dietary sodium chloride on urine calcium excretion, bone resorption, and bone formation in postmenopausal women and, further, whether oral potassium citrate attenuates the effects of increased dietary salt.

## Subjects and Methods

### Subjects

Sixty postmenopausal women were recruited through newspaper advertisements and direct mailings. Women were excluded if they were

Abbreviations: Cr, Creatinine; nMBCE, nanomoles bone collagen equivalent; NTX, N-telopeptide; UCSF, University of California San Francisco.

less than 2 yr past menopause, on medications known to affect bone metabolism, or had evidence of metabolic bone disease by history, physical, or laboratory exam. Women were also excluded if they had conditions potentially worsened by increased dietary sodium such as congestive heart failure, uncontrolled hypertension, myocardial infarction within the previous 12 months, or cardiac arrhythmias. Additionally, women were excluded if they had contraindications to potassium supplementation, namely active gastrointestinal disease, renal insufficiency [2 h creatinine (Cr) clearance <40 ml/min, or serum Cr >2.0 mg/dl], metabolic acidosis (serum bicarbonate <16 mmol/liter), uncontrolled diabetes mellitus (fasting glucose >200 mg/dl), or medications altering potassium metabolism (angiotensin-converting enzyme inhibitors,  $\beta$ -blockers, potassium-sparing diuretics). To minimize differences in calcium intake, participants discontinued all personal calcium supplements 1–2 wk before starting the low-salt diet and were placed on calcium carbonate 500 mg/d throughout the study. This study was approved by the University of California San Francisco (UCSF) Institutional Review Board, and all participants provided informed consent.

### Experimental protocol

After enrollment, a research dietitian instructed the study subjects in a diet containing 87 mmol/d (2000 mg) sodium. Participants were provided with information on reading food labels, lists of low-sodium foods, examples of low-sodium menus, and books listing the sodium content of common foods. Approximately 10 d into the low-salt diet, participants collected a 24-h urine sample to be assayed for sodium content to assess dietary compliance. Women whose urinary sodium was more than 10 mmol from the sodium intake goal were contacted, a 24-h dietary recall was performed to identify barriers to compliance, and the low-salt diet was reinforced. The subjects ingested the 87 mmol/d sodium diet for 3 wk. At the end of the third week, 24-h urine samples and fasting morning blood samples were collected in triplicate (three collections in the last 3–5 d of the low-salt diet). Triplicate collections were used to maximize precision, particularly in the 24-h urine collections because these samples were collected as outpatients.

After 3 wk on the low-salt diet, subjects began a high-sodium (225 mmol/d) diet: to the 87 mmol/d sodium diet, four oral sodium chloride pills, two weighed packets of salt to sprinkle on their food, and one cup of bouillon per day were added. Subjects were randomized to take either potassium citrate 90 mmol/d (UrocitK, Mission Pharmacal, San Antonio, TX) or indistinguishable placebo pills while on the high-salt diet. Approximately 12 d into the high-salt diet, participants collected a 24-h urine sample to assess dietary compliance. Again, women who were more than 10 mmol from the targeted sodium intake were contacted, barriers to compliance were rectified, and the high-salt regimen was reinforced. At the end of the fourth week on the high-salt diet, 24-h urine samples and fasting morning blood samples were collected in triplicate.

### Randomization and blinding

Study drug was provided to the UCSF investigational pharmacy by the manufacturer in sealed bottles labeled “potassium citrate or placebo” and one of two lot numbers. Only the manufacturer knew the lot number that was potassium citrate and the lot number that was placebo. Participants were randomized using a random numbers list provided by the investigational pharmacy. All study personnel and the participants were blinded to the treatment groups throughout the study.

### Biochemical assessment

Blood and urine samples were aliquoted and frozen at  $-70^{\circ}\text{C}$  until all samples from a single participant were collected. All samples from each participant were analyzed in a single batch to prevent interassay differences. Sodium, potassium, and calcium were measured using standard techniques in the UCSF clinical laboratory.

Net acid excretion was calculated as the sum of the excretion rates of titratable acid and ammonium minus that of bicarbonate. Titratable acid concentration was determined by titration, and urine ammonium concentration was determined by the phenol method. Urine total carbon dioxide content was determined by thermal conductivity ( $\text{CO}_2$  analyzer, CIBA Corning, Medfield, MA); the coefficient of variation of the method

is 0.7%. Urine bicarbonate concentration was calculated from the measured values of urine pH and carbon dioxide content using the Henderson-Hasselbach equation, where the solubility coefficient of  $\text{CO}_2$  is taken as 0.0309 and  $\text{pK}^1$  is corrected for ionic strength as follows:  $\text{pK}^1 = 6.33 - 0.5([\text{Na}^+] + [\text{K}^+])^{1/2}$ , where  $\text{Na}^+$  and  $\text{K}^+$  concentrations are expressed in eq/liter.

Urinary N-telopeptides (NTX) of type I collagen and osteocalcin were measured using the ELISA method (Metra Biosystems, Palo Alto, CA). Intact PTH was measured using a two-site immunoradiometric assay (Nichols Institute Diagnostics, San Juan Capistrano, CA). cAMP was measured using a RIA (INCSTAR Corp., Stillwater, MN).

### Statistical analysis

The triplicate values from each participant at the end of each diet interval were averaged to provide a single low-salt and high-salt value for each participant. Within group analyses (comparing low salt to high salt in each group) were performed using a paired *t* test on the mean values at the end of the low- and high-salt periods. Between group analyses (comparing placebo and potassium citrate groups) were performed using an unpaired *t* test. For the between-groups analyses, outcomes were assessed as the difference in values between the low-salt and high-salt diets. All analyses were performed using STATA software (version 5.0, 1997; Stata Corp., College Station, TX).

## Results

### Baseline characteristics and subject withdrawals

Fifty-two (87%) of the women completed the 7-wk study; 26 had been randomized to placebo, and 26 had been randomized to potassium citrate. Eight women dropped out due to nausea from the sodium chloride pills on the high-salt diet. Women who withdrew from the study did not differ with respect to age, years since menopause, baseline sodium intake, compliance with the low-salt diet, or randomization to treatment or placebo. The characteristics at study entry (*i.e.* before the low-salt diet) of the women who completed the study are shown in Table 1.

### Biochemical data

Urine sodium averaged  $96 \pm 5$  mmol/d (mean  $\pm$  SEM) on the low-salt diet and  $235 \pm 7$  mmol/d on the high-salt diet. Compliance to the low- and high-salt diets, as evidenced by 24-h urine sodium excretion, was not different between the women randomized to potassium citrate and the women randomized to placebo.

**TABLE 1.** Characteristics of subjects at study entry ( $n = 52$ )

	High salt + placebo ( $n = 26$ )	High salt + potassium citrate ( $n = 26$ )
Age (yr)	$63 \pm 8$	$65 \pm 8$
Age at menopause (yr)	$50 \pm 4$	$50 \pm 4$
Urine sodium (mmol/d)	$144 \pm 51$	$138 \pm 46$
Serum calcium (mg/dl)	$9.3 \pm 0.1$	$9.4 \pm 0.1$
Urine calcium (mg/d)	$215 \pm 79$	$201 \pm 101$
Osteocalcin (ng/ml)	$10.8 \pm 3.2$	$10.7 \pm 3.5$
Urine NTX (nM BCE/mmol Cr)	$47.0 \pm 25.5$	$44.6 \pm 28.6$
Fasting PTH (pg/ml)	$42.0 \pm 13.0$	$45.6 \pm 18.0$
cAMP (nmol/liter)	$2766 \pm 779$	$2957 \pm 829$

Values are expressed as the mean  $\pm$  SD. *P* value for comparison between women randomized to high salt plus placebo and women randomized to high salt plus potassium citrate by unpaired *t* test was not significant for all variables.

**Low-salt vs. high-salt plus placebo.** In the women randomized to placebo, urine calcium excretion increased significantly from  $200 \pm 14$  mg/d on the low-salt diet to  $242 \pm 16$  mg/d on the high-salt diet plus placebo ( $P = 0.002$ ). Urine NTX excretion increased significantly from  $38.7 \pm 4.2$  nanomoles bone collagen equivalents (nMBCE)/mmol Cr on the low-salt diet to  $45.1 \pm 3.8$  nMBCE/mmol Cr on the high-salt diet plus placebo ( $P = 0.001$ ). Osteocalcin decreased slightly, but significantly, from  $11.2 \pm 0.6$  ng/ml on the low-salt diet to  $10.6 \pm 0.6$  ng/ml on the high-salt diet plus placebo ( $P = 0.01$ ). There was no significant change in fasting PTH or cAMP between the low-salt and high-salt plus placebo diets. Additionally, urine potassium and net acid excretion were not different between the low-salt and high-salt diets in the women randomized to placebo (Table 2).

**Low salt vs. high salt plus potassium citrate.** In the women randomized to potassium citrate, urine calcium excretion decreased from  $200 \pm 14$  mg/d on the low-salt diet to  $192 \pm 19$  mg/d on the high-salt diet plus potassium citrate, a nonsignificant change ( $P = 0.5$ ). Urine NTX excretion was  $41.3 \pm 3.6$  nMBCE/mmol Cr on the low-salt diet and  $43.3 \pm 3.7$  nMBCE/mmol Cr on the high-salt plus potassium citrate diet, a nonsignificant change ( $P = 0.3$ ). There were no significant changes in osteocalcin, fasting PTH, or cAMP between the low-salt and high-salt plus potassium citrate diets. Urine potassium excretion increased from  $70 \pm 3$  mmol/d on the low-salt diet to  $141 \pm 6$  mmol/d on the high-salt diet plus potassium citrate ( $P < 0.001$ ). Urine net acid excretion de-

creased from  $32 \pm 3$  mmol/d on the low-salt diet to  $-28 \pm 4$  mmol/d on the high-salt plus potassium citrate diet ( $P < 0.001$ ) (Table 2).

**Potassium citrate vs. placebo.** Urinary calcium excretion increased by  $42 \pm 12$  mg/d (33%) between the low-salt and high-salt diet in the women randomized to high salt plus placebo, while decreasing by  $8 \pm 14$  mg/d (4%) in the women randomized to high salt plus potassium citrate. The difference between the two groups was significant ( $P = 0.008$ ). Urinary NTX excretion increased by  $6.4 \pm 1.4$  nMBCE/mmol Cr (23%) between the low-salt and high-salt diets in the women randomized to high salt plus placebo, while increasing by  $2.0 \pm 1.7$  nMBCE/mmol (7.5%) in the women randomized to high salt plus potassium citrate. The change in urine NTX excretion was significantly different between the two groups ( $P < 0.05$ ) (Table 3). There was no significant difference between the two groups in changes in osteocalcin, fasting PTH, or cAMP. Between the low- and high-salt diets, urine potassium increased by  $2 \pm 3$  mmol/d in the women randomized to high salt plus placebo, while increasing by  $72 \pm 5$  mmol/d in the women randomized to high salt plus potassium citrate, a significant difference ( $P < 0.001$ ). Urine net acid excretion decreased by  $3 \pm 3$  mmol/d between the low-salt and high-salt diets in the women randomized to high salt plus placebo, while decreasing by  $60 \pm 5$  mmol/d in the women randomized to high salt plus potassium citrate; the difference between the two groups was significant ( $P < 0.001$ ).

**TABLE 2.** Calcium, bone markers, and hormones: low salt (87 mmol/d sodium) vs. high salt (225 mmol/d sodium) plus placebo and low salt vs. high salt plus potassium citrate (90 mmol/d)

	Women randomized to placebo (n = 26)			Women randomized to potassium citrate (n = 26)		
	Low salt	High salt	P value vs. low salt	Low salt	High salt	P value vs. low salt
Urine calcium (mg/d)	$200 \pm 14$	$242 \pm 16$	0.002	$200 \pm 14$	$192 \pm 19$	0.5
Urine NTX (nMBCE/mmol Cr)	$38.7 \pm 4.2$	$45.1 \pm 3.8$	0.001	$41.3 \pm 3.6$	$43.3 \pm 3.7$	0.3
Serum calcium (mg/dl)	$9.4 \pm 0.1$	$9.4 \pm 0.1$	0.5	$9.4 \pm 0.1$	$9.4 \pm 0.1$	0.4
Osteocalcin (ng/ml)	$11.2 \pm 0.6$	$10.6 \pm 0.6$	0.01	$10.9 \pm 0.7$	$10.7 \pm 0.6$	0.3
Fasting PTH (pg/ml)	$41.7 \pm 2.7$	$42.7 \pm 2.9$	0.5	$49.2 \pm 3.8$	$48.5 \pm 3.4$	0.7
cAMP (nmol/liter)	$2630 \pm 132$	$2783 \pm 144$	0.2	$3013 \pm 185$	$3043 \pm 157$	0.8
Urine potassium (meq/d)	$75 \pm 4$	$77 \pm 4$	0.6	$70 \pm 3$	$141 \pm 6$	<0.001
Net acid excretion (meq/d)	$28 \pm 4$	$25 \pm 3$	0.3	$32 \pm 3$	$-28 \pm 4$	<0.001

All values are mean  $\pm$  SEM. The P value is for comparison between the low-salt and high-salt diets by paired *t* test within each group (placebo or potassium citrate).

**TABLE 3.** Change in calcium, bone turnover markers, and hormones from low salt (87 mmol/d sodium) to high salt (225 mmol/d sodium), placebo vs. potassium citrate (90 mmol/d)

	Change from low salt to high salt + placebo	Change from low salt to high salt + potassium citrate	P value placebo vs. potassium citrate
Urine calcium (mg/d)	$42 \pm 12$	$-8 \pm 14$	0.008
Urine NTX (nMBCE/mmol Cr)	$6.4 \pm 1.4$	$2.0 \pm 1.7$	0.049
Serum calcium (mg/dl)	$0.04 \pm 0.06$	$0.05 \pm 0.05$	0.89
Osteocalcin (ng/ml)	$-0.57 \pm 0.21$	$-0.22 \pm 0.23$	0.26
Fasting PTH (pg/ml)	$0.97 \pm 1.5$	$-0.74 \pm 1.8$	0.46
cAMP (nmol/liter)	$142.2 \pm 99.2$	$106.7 \pm 135.6$	0.83
Urine potassium (meq/d)	$2 \pm 3$	$72 \pm 5$	<0.001
Net acid excretion (meq/d)	$3 \pm 3$	$60 \pm 5$	<0.001

Values are expressed as the mean  $\pm$  SE. The P value is for comparison between women randomized to high salt plus placebo and women randomized to high salt plus potassium citrate by unpaired *t* test.



## Discussion

Although the effects of dietary sodium chloride on urine calcium excretion have been consistent, the effect on bone density, bone loss, and bone turnover is less clear. Dietary sodium has not been associated with bone mineral density in two cohorts in the United States (17, 18), and studies of the relationship between dietary sodium and bone loss have been mixed (19, 20). Cross-sectional studies have demonstrated a positive correlation between sodium and calcium excretion and urinary hydroxyproline (5, 6, 21), and hydroxyproline excretion appears to increase with salt administration (4, 22). However, of the two intervention studies using a more specific marker of bone resorption, deoxypyridinoline, one demonstrated an increase in bone resorption with increased dietary salt (23), but the other did not (24).

We found that increased dietary salt increased urinary calcium excretion in the women randomized to the high-salt diet plus placebo. Enhanced calcium excretion was associated with increases in a marker of bone resorption, NTX, suggesting that the increased calcium excretion due to increased dietary salt has skeletal effects. Additionally, the marker of bone formation, osteocalcin, was slightly, but significantly, reduced. The increased urinary calcium loss in the high salt plus placebo group did not result in changes in fasting PTH or cAMP, suggesting that, as seen in previous studies (12), postmenopausal women eating abundant salt do not show hormonal responses that could compensate for enhanced urinary calcium losses.

In contrast, urinary calcium and NTX excretion in the women randomized to high salt plus potassium citrate during the high-salt diet were not different from on the low-salt diet, despite increased urine sodium excretion. Additionally, the women taking potassium citrate during the high-salt diet were significantly different from the high salt plus placebo group with regard to changes in urine calcium and NTX, suggesting that potassium citrate prevented the increased calcium and bone resorption biomarker excretion caused by increased salt intake.

The ability of potassium citrate to reduce urine calcium excretion and markers of bone turnover may be multifactorial. Alkaline salts of potassium are both natriuretic and chloruretic, potentially reducing the extracellular volume expansion that occurs with increased salt intake. This effect may be partly responsible for the decreased calcium excretion that occurs when these agents are given with a high-salt diet (16). In our study, the addition of potassium citrate and sodium chloride to a low-salt diet resulted in no difference in urine calcium excretion compared with a low-salt diet.

In addition to effects on extracellular volume, alkaline salts of potassium reduce endogenous acid production and increase blood pH and plasma bicarbonate concentration (25, 26). Urinary calcium excretion varies directly with endogenous acid production as measured by urinary net acid excretion; increased net acid excretion is associated with decreased renal tubular reabsorption of calcium (27). In our study, net acid excretion decreased by 60 mmol/d in the women given potassium citrate; in fact, the average net acid excretion in this group was negative. This appears to be a significant factor; in our study, urine calcium and bone turn-

over markers were no different on a high-salt diet with potassium citrate than on a low-salt (87 mmol/d sodium) diet alone. Additionally, potassium itself may be a factor in calcium excretion. Potassium deficient diets are associated with increased urine calcium excretion, potassium supplementation reduces urinary calcium excretion (14, 25, 28), and potassium appears to have a direct effect on the kidney to promote calcium reclamation (29). Finally, the total sodium and chloride content of the diet may also be important. Sodium bicarbonate can improve calcium balance when substituted for sodium chloride in the diet (30). However, when dietary sodium chloride remains constant or is increased, potassium bicarbonate, but not sodium bicarbonate, improves calcium balance (13).

Our study examined the effects of dietary sodium chloride and oral potassium citrate over 1 month. Previous studies examining the effect of high-salt diets on bone turnover markers have been limited to 7–10 d. This study suggests that the effect of increased dietary sodium chloride could be ongoing. Larger studies with more long-term outcomes such as bone mineral density and fracture will be needed to define the role of dietary sodium chloride in postmenopausal bone loss and osteoporosis more completely.

However, the results of this randomized, double blind, placebo-controlled trial suggest that increased dietary salt not only enhances urinary calcium excretion, but also results in increased excretion of bone resorption biomarkers. The addition of oral potassium citrate to a high-salt diet prevented the increased calcium losses and rise in bone turnover markers caused by a high dietary salt intake. Prior investigations have also suggested that higher intake of dietary potassium and potassium-rich foods, namely fruit and vegetables, is associated with higher bone density in both pre- and postmenopausal women (31, 32). The amount of potassium that reduced urine calcium excretion and NTX in our study could be ingested by consuming 7–8 servings of potassium-rich fruit and vegetables each day.

Current dietary guidelines recommend moderate sodium intake (2400 mg/d), but research on the impact of dietary sodium has focused primarily on the cardiovascular system. This study provides further evidence that recommendations regarding moderate sodium intake and increased intake of dietary sources of potassium alkaline salts, namely fruit and vegetables, may also be beneficial to postmenopausal women at risk for osteoporosis.

## Acknowledgments

We acknowledge the generous donation of potassium citrate (UroclitK) and placebo tablets by Mission Pharmacal (San Antonio, TX).

Received July 30, 2001. Accepted January 31, 2002.

Address all correspondence and requests for reprints to: Dr. Deborah E. Sellmeyer, University of California, San Francisco, Prevention Sciences Group, 74 New Montgomery Street, Suite 600, San Francisco, California 94105. Email: dsellmeyer@psg.ucsf.edu.

This work was supported by NIH Grant RO-1 DK53172. These studies were performed in part in the General Clinical Research Center, Moffitt Hospital, University of California, San Francisco, with funds provided by USPHS National Center for Research Resources Grant 5 M01 RR-00079.

## References

1. Dawson-Hughes B, Dallal GE, Krall EA, Sadowski L, Sahyoun N, Tannenbaum S 1990 A controlled trial of the effect of calcium supplementation on bone density in postmenopausal women. *N Engl J Med* 323:878–883
2. Dawson-Hughes B, Harris SS, Krall EA, Dallal GE, Falconer G 1995 Rates of bone loss in postmenopausal women randomly assigned to one of two dosages of vitamin D. *Am J Clin Nutr* 61:1140–1145
3. Goulding A, McParland BE 1990 Fasting and 24-h urinary sodium/creatinine values in young and elderly women on low-salt and salt-supplemented regimens. *J Cardiovasc Pharmacol* 16(Suppl 7):S47–S49
4. Goulding A, Lim PE 1983 Effects of varying dietary salt intake on the fasting urinary excretion of sodium, calcium and hydroxyproline in young women. *N Z Med J* 96:853–854
5. Goulding A 1981 Fasting urinary sodium/creatinine in relation to calcium/creatinine and hydroxyproline/creatinine in a general population of women. *N Z Med J* 93:294–297
6. Chan ELP, Ho CS, MacDonald D, Ho SC, Chan TYK, Swaminathan R 1992 Interrelationships between urinary sodium, calcium, hydroxyproline and serum PTH in healthy subjects. *Acta Endocrinol (Copenh)* 127:242–245
7. Muldowney FP, Freaney R, Moloney MF 1982 Importance of dietary sodium in the hypercalciuria syndrome. *Kidney Int* 22:292–296
8. Shortt C, Madden A, Flynn A, Morrissey PA 1988 Influence of dietary sodium intake on urinary calcium excretion in selected Irish individuals. *Eur J Clin Nutr* 42:595–603
9. Need AG, Morris HA, Cleghorn DB, De Nichilo D, Horowitz M, Nordin BEC 1991 Effect of salt restriction on urine hydroxyproline excretion in postmenopausal women. *Arch Intern Med* 151:757–759
10. Nordin BEC, Need AG, Morris HA, Horowitz M 1993 The nature and significance of the relationship between urinary sodium and urinary calcium in women. *J Nutr* 123:1615–1622
11. Breslau NA, McGuire JL, Zerwekh JE, Pak CYC 1982 The role of dietary sodium on renal excretion and intestinal absorption of calcium and on vitamin D metabolism. *J Clin Endocrinol Metab* 55:369–373
12. Breslau NA, Sakhaee K, Pak CYC 1985 Impaired adaptation to salt-induced urinary calcium losses in postmenopausal osteoporosis. *Trans Assoc Am Physicians* 98:107–115
13. Lemann Jr J, Gray RW, Pleuss JA 1989 Potassium bicarbonate, but not sodium bicarbonate, reduces urinary calcium excretion and improves calcium balance in healthy men. *Kidney Int* 35:688–695
14. Lemann Jr J, Pleuss JA, Gray RW, Hoffmann RG 1991 Potassium administration reduces and potassium deprivation increases urinary calcium excretion in healthy adults [published erratum appears in *Kidney Int* 1991 Aug;40(2):388]. *Kidney Int* 39:973–983
15. Bell RR, Eldrid MM, Watson FR 1992 The influence of NaCl and KCl on urinary calcium excretion in healthy young women. *Nutr Res* 12:17–26
16. Morris RC, Sebastian A, Forman A, Tanaka M, Schmidlin O 1999 Normotensive salt-sensitivity: effects of race and dietary potassium. *Hypertension* 33:18–23
17. Greendale GA, Barrett-Connor E, Edelstein S, Ingles S, Haile R 1994 Dietary sodium and bone mineral density: results of a 16-year follow-up study. *J Am Geriatr Soc* 42:1050–1055
18. Dawson-Hughes B, Fowler SE, Dalsky G, Gallagher C 1996 Sodium excretion influences calcium homeostasis in elderly men and women. *J Nutr* 126:2107–2112
19. Reid IR, Ames RW, Evans MC, Sharpe SJ, Gamble GD 1994 Determinants of the rate of bone loss in normal postmenopausal women. *J Clin Endocrinol Metab* 79:950–954
20. Devine A, Criddle RA, Dick IM, Kerr DA, Prince RL 1995 A longitudinal study of the effect of sodium and calcium intakes on regional bone density in postmenopausal women. *Am J Clin Nutr* 62:740–745
21. Itoh R, Suyama Y 1996 Sodium excretion in relation to calcium and hydroxyproline excretion in a healthy Japanese population. *Am J Clin Nutr* 63:735–740
22. McParland BE, Goulding A, Campbell AJ 1989 Dietary salt affects biochemical markers of resorption and formation of bone in elderly women. *Br Med J* 299:834–835
23. Evans CE, Chugtai AY, Blumsohn A, Giles M, Eastell R 1997 The effect of dietary sodium on calcium metabolism in premenopausal and postmenopausal women. *Eur J Clin Nutr* 51:394–399
24. Lietz G, Avenell A, Robins SP 1997 Short-term effects of dietary sodium intake on bone metabolism in postmenopausal women measured using urinary deoxypyridinoline excretion. *Br J Nutr* 78:73–82
25. Sebastian A, Harris ST, Ottaway JH, Todd KM, Morris Jr RC 1994 Improved mineral balance and skeletal metabolism in postmenopausal women treated with potassium bicarbonate. *N Engl J Med* 330:1776–1781
26. Bushinsky DA 1998 Acid-base imbalance and the skeleton. In: Burckhardt P, Dawson-Hughes B, Heaney RP, eds. *Nutritional aspects of osteoporosis*. New York: Springer; 208–217
27. Lemann JJ 1999 Relationship between urinary calcium and net acid excretion as determined by dietary protein and potassium: a review. *Nephron* 81(Suppl 1):18–25
28. Lemann Jr J, Pleuss JA, Gray RW 1990 Dietary potassium deprivation increases daily and fasting urinary calcium excretion in healthy men. *J Bone Min Res* 5(Suppl 2):S203
29. Brunette MG, Mailloux J, Lajeunesse D 1992 Calcium transport through the luminal membrane of the distal tubule. I. Interrelationship with sodium. *Kidney Int* 41:281–288
30. Lutz J 1984 Calcium balance and acid-base status of women as affected by increased protein intake and by sodium bicarbonate ingestion. *Am J Clin Nutr* 39:281–288
31. New SA, Bolton-Smith C, Grubb DA, Reid DM 1997 Nutritional influences on bone mineral density: a cross-sectional study in premenopausal women. *Am J Clin Nutr* 65:1831–1839
32. New SA, Robins SP, Campbell MK, Martin JC, Garton MJ, Bolton-Smith C, Grubb DA, Lee SJ, Reid DM 2000 Dietary influences on bone mass and bone metabolism: further evidence of a positive link between fruit and vegetable consumption and bone health? *Am J Clin Nutr* 71:142–151