

Brief Review

Dietary Calcium and Blood Pressure in Experimental Models of Hypertension

A Review

Daniel C. Hatton, David A. McCarron

Restricted dietary calcium intake has been linked to elevated blood pressure in a number of epidemiologic studies. These findings, in conjunction with indications of calcium deficiency in clinical hypertension, suggest that repletion of dietary calcium may have a beneficial effect on blood pressure. This possibility has been explored extensively in both humans and animal models of hypertension. In this article, we review the animal work relating dietary calcium to blood pressure and examine some of the mechanisms that may be responsible. Much of the content is based on published experiments and earlier reviews by our laboratory on this subject.¹⁻³

Background

Calcium Metabolism in Experimental and Essential Hypertension

More than 20 epidemiologic investigations have reported an inverse relation between dietary calcium intake and blood pressure status.⁴⁻²⁸ The increased risk of elevated blood pressure with lower calcium intake appears to be related to an apparent deficit in calcium metabolism that occurs in a subset of hypertensive individuals. That deficit appears systemically as a small increase in circulating parathyroid hormone (PTH)^{29,30} levels, reduced serum phosphorus,³¹ and increased urinary calcium excretion.³²⁻³⁸ At the cellular level, the deficit in calcium metabolism is reflected in elevated intracellular free calcium levels, as indicated by studies examining platelets isolated from individuals with essential hypertension.³⁹⁻⁴⁷

The alterations in calcium metabolism observed in hypertensive humans are mirrored in experimental models of hypertension. Plasma ionized calcium has been reported to be decreased in the spontaneously hypertensive rat (SHR) relative to the normotensive Wistar-Kyoto (WKY) rat,⁴⁸⁻⁵¹ whereas PTH has been observed to be either elevated or unchanged.^{49,51} End-organ responsiveness to PTH appears to be abnormal based on findings of hypercalciuria,⁵¹⁻⁵⁴ decreased basal and stimulated 1,25-dihydroxyvitamin D₃ [1,25(OH)₂ vitamin D₃] production,⁵⁵⁻⁵⁸ and low to normal urinary nephrogenous cyclic

AMP in the presence of elevated PTH.^{52,59,60} At the cellular level, SHR exhibit depressed ATP-dependent calcium uptake capacity in crude microsomal fractions of aorta as well as in more purified sarcolemmal fractions of the mesenteric artery.⁶¹⁻⁶⁶ Therefore, it has been suggested that the vascular myocyte may not be able to maintain normal levels of intracellular calcium or properly remove calcium after activation.⁶⁷ It appears that the cell membrane of the SHR as exemplified by platelets and lymphocytes may be hyperpermeable to calcium because both basal and agonist-stimulated calcium uptake have been found to be elevated in the SHR.⁶⁸⁻⁷¹ Consequently, the SHR has abnormally high intracellular free calcium levels, as has been observed in essential hypertensive patients.

The inverse relation between dietary calcium and blood pressure identified through epidemiologic investigations suggests that supplemental dietary calcium may lower blood pressure. Numerous studies have tested this possibility. Of the published studies of calcium supplementation in humans, 24 reported a significant reduction in blood pressure⁷²⁻⁹⁵ in at least a segment of the study population, and 13 reported no difference in blood pressure.⁹⁶⁻¹⁰⁸ Recent reviews of these data^{109,110} indicated a modest, dose-dependent effect of calcium supplementation on blood pressure that may be most important in certain subsets of the population. These subsets include African Americans,¹¹¹⁻¹¹³ women with gestational hypertension,⁸³⁻⁸⁸ or salt-sensitive individuals.⁹⁰⁻⁹² Low calcium intake may be an especially important indicator because very low calcium diets (<400 to 600 mg/d) are most closely associated with elevated blood pressure,^{5,20-22,24,27,114,115} and work with experimental models indicates that low calcium diets elevate blood pressure (see below).

Dietary Calcium in Experimental Models of Hypertension

The outcome of calcium supplementation in experimental models of hypertension is much more consistent than it is in humans. As shown in Table 1, more than 80 studies have manipulated dietary calcium and measured blood pressure. Most have found an inverse relation between calcium intake and blood pressure. This relation has been found in a variety of genetic and experimental models of hypertension, including SHR; stroke-prone SHR; Dahl salt-sensitive (DS) rats; Lyon hypertensive rats; deoxycorticosterone-saline (DOC-saline) rats; one-kidney, one clip rats; and remnant kidney

Received May 25, 1993; accepted in revised form December 15, 1993.

From the Division of Nephrology and Hypertension, Oregon Health Sciences University, Portland.

Reprint requests to Daniel C. Hatton, PhD, Division of Nephrology and Hypertension, PP262, Oregon Health Sciences University, 3314 SW US Veterans Hospital Rd, Portland, OR 97201.

TABLE 1. Studies of Dietary Calcium and Blood Pressure in Experimental Animal Models

Study	Model	Calcium Diet	Age of Animal/ Treatment Length, wk*	Result in mm Hg SBP	Focus
Ambrozy et al 1991 ¹¹⁶	Zucker rat	0.5-1.5%	8/4	~ ↓ 30 in lean, ~ ↓ 20 in obese, $P < .05$	Vascular reactivity, VSM Ca^{2+} efflux
Anderson et al 1986 ¹¹⁷	Dog	CaCl_2 , 115 mEq	2	No effect	Diet-stress interaction
Anderson et al 1983 ¹¹⁸	SHR (n=7-8)	0.02-4.0%	16/2	↓ 19 MAP, $P < .005$	Calcium balance, PTH
Ando et al 1991 ¹¹⁹	SD/Ang II/salt	1.17-4.07%	12 days	↓ 35 MAP, $P < .01$	SNS
Ayachi 1979 ⁵⁴	SHR/WKY	1.2-2.5%	4/14	~ ↓ 21, $P < .05$	Metabolic study
Baksi 1988 ¹²⁰	SD	0.005/0.17/1.4/ 2.8%	4/8	~ ↓ 30, $P < .05$	Pressor response
Baksi 1989 ¹²¹	SD	0.005/0.17/1.4/ 2.8%	4/8	↓ 34 MAP, $P < .05$	CNS NE
Baksi 1988 ¹²²	SD/PTX	0.005/1.4/2.8%	4/8	↓ 43, $P < .05$	PTH
Baksi et al 1989 ¹²³	SD	0.005-1.4%	4/8	~ ↓ 100, $P < .01$	RAS
Belizán et al 1981 ¹²⁴	Pregnant SD (n=5,8)	0.0-0.6%	20/12	↓ 20, $P < .001$	Pregnancy-induced hypertension
Belizán et al 1984 ¹²⁵	PTX Wistar (n=9)	0.0-0.6%	8/10	↓ 19, $P < .005$	PTH
Blakeborough et al 1990 ¹²⁶	♀ SHR/WKY	0.06-0.6%	8/10	~ ↓ 16, $P < .05$	Intestine and kidney membrane enzyme
Bogden et al 1991 ¹²⁷	Wistar rat	0.02-4.0%	~4/31	~ ↑ 10 MAP, $P < .05$	Lead- Ca^{2+} interaction neoplasia
Bukoski et al 1989 ¹²⁸	SHR/WKY	0.5/1.0/2.0%	6/6 or 12	~ ↓ 12, $P < .05$	Vascular contractility
Bukoski and McCarron 1986 ¹²⁹	SHR/WKY	1.0-2.0%	6/8 or 15	↓ 19, $P < .05$	Vascular contractility
DiPette et al 1990 ¹³⁰	SD/DOC-salt	0.6-2.5%	~6/8	↓ 14 MAP, $P < .05$	Calcium-regulating hormones
Doris 1985 ¹³¹	♀ WKY/saline	1.0-2.5%	~8/26	↓ 10, $P < .01$	NaCl-Ca^{2+} interaction
Doris 1988 ¹³²	♀ WKY/saline	1.0-2.0%	12/26	~ ↓ 7, $P < .05$	NE and PTH
Evans et al 1990 ¹³³	SHR (n=9)	0.075/0.5/2.5%	8/23	~ ↓ 35, $P < .001$	$\text{Mg}^{2+}\text{-Ca}^{2+}$ interaction
Fujito et al 1991 ¹³⁴	SHR/WKY (n=5)	0.1/0.6/4.0%	6/14	~ ↓ 12, $P < .01$	Erythrocyte sodium transport
Furspan et al 1989 ¹³⁵	SHRSP/WKY (n=6)	1.0-2.5%	10/9	↓ 58, $P < .05$	Membrane stability
Ganguli et al 1986 ¹³⁶	DS rat (n=18)	0.5-3.0%	8/15	↓ 7 MAP, $P < .01$	CaCO_3 vs CaHPO_4
Geiger et al 1988 ¹³⁷	SHR/PTX	0.1-3.0%	4/4	↓ 26, $P < .05$	ANP
Hano et al 1991 ¹³⁸	SHR/WKY (n=10)	1.2% CaCl_2 in water	4/3	~ ↓ 25, $P < .01$	NE, platelet $[\text{Ca}^{2+}]$
Hatton et al 1986 ¹³⁹	SHR (n=12/21)	0.1-2.0%	Prenatal/4	↓ 30 MAP, $P < .01$	Early diet experience
Hatton et al 1987 ¹⁴⁰	SHR/WKY (n=8)	0.1/1.0/2.0%	6/13	~ ↓ 20, $P < .05$	Diet-stress interaction
Hatton et al 1988 ¹⁴¹	SHR (n=8)	0.1/1.0/2.0%	3/1	↓ 14 MAP, $P < .01$	Plasma volume, vascular contraction
Hatton et al 1989 ¹⁴²	SHR/WKY (n=8)	0.1/1.0/2.0%	4/12	~ ↓ 20 MAP, $P < .01$	SNS

SBP indicates systolic blood pressure; VSM, vascular smooth muscle; SHR, spontaneously hypertensive rats; MAP, mean arterial pressure; PTH, parathyroid hormone; SD, Sprague-Dawley rats; Ang II, angiotensin II; SNS, sympathetic nervous system; WKY, Wistar-Kyoto rats; CNS, central nervous system; NE, norepinephrine; PTX, parathyroidectomy; RAS, renin-angiotensin system; DOC, deoxycorticosterone; SHRSP, stroke-prone SHR; DS, Dahl salt-sensitive; ANP, atrial natriuretic peptide; ACI, American Cancer Institute; BP, blood pressure; LVH, left ventricular hypertrophy; PHF, parathyroid hypertensive factor; SHR-S, salt-sensitive SHR; SHR-R, salt resistant SHR; LHR, Lyon hypertensive rats; 2K1C, two-kidney, one clip hypertensive rats; ADX, adrenalectomized rats; and ald, aldosterone.

*Unless noted otherwise.

rats. Normotensive rats, such as WKY, Wistar, Fisher 344, and Sprague-Dawley rats, are also responsive to dietary calcium, but the changes in blood pressure are smaller and take prolonged periods of time to develop.

The most common method of manipulating dietary calcium has been by altering CaCO_3 in dry food. However, CaCl_2 in the drinking water has proved to be an effective means of altering blood pressure as well. The

TABLE 1. Continued

Study	Model	Calcium Diet	Age of Animal/ Treatment Length, wk*	Result in mm Hg SBP	Focus
Hatton et al 1991 ¹⁴³	SHR (n=8)	0.1-2.0%	3/1	~ ↓ 10 MAP, <i>P</i> < .01	Hematocrit
Hatton et al 1993 ¹⁴⁴	SHR (n=8/7)	0.1-2.0%	3/2	↓ 13 MAP, <i>P</i> < .005	Adrenergic receptors
Hattori et al 1991 ¹⁴⁵	Wistar/DOC-salt (n=9/12)	2.0% CaCl ₂ in water	5/4	↓ 56	Endogenous opioids
Hermesmeier et al 1990 ¹⁴⁶	SHRSP	0.2-2.0%	4/8	~ ↓ 25, <i>P</i> < .01	Calcium channels
Huie et al 1987 ¹⁴⁷	Fisher 344, Wistar-Furth, ACI	0.1-1.0%	4/25	↓ 15, <i>P</i> < .05, Fisher 344	Genetic variability
Huie et al 1987 ¹⁴⁸	SHR (n=8/10)	0.1/1.0/2.0%	6/13	~ ↓ 27	Stress-diet interaction
Jirakulsomchok et al 1990 ¹⁴⁹	SHR/WKY high salt	0.68-2.0%	7/2.5	↓ 42, <i>P</i> < .05	Natriuresis, diuresis
Jones et al 1986 ¹⁵⁰	Wistar (n=9)	1.0-2.5%	~4/6	↓ 27, <i>P</i> < .01	Electrolyte interaction
Jones and Huibonhoa 1988 ¹⁵¹	Wistar (n=8)	0.2/0.3/0.4/0.5%	~4/19	No effect	Marginal calcium diet
Kageyama et al 1986 ¹⁵²	SHR/WKY (n=10)	1.5% CaCl ₂ in water	6/3	↓ 26, <i>P</i> < .01	Vascular reactivity
Kageyama et al 1987 ¹⁵³	2K1C (n=30)	1.5% CaCl ₂ in water	~7/3	↓ 53, <i>P</i> < .005	RAS
Kageyama et al 1987 ¹⁵⁴	Wistar/DOC-saline (n=6/8)	1% CaCl ₂ in water	~8/3	~ ↓ 18, <i>P</i> < .01	Vascular reactivity
Kageyama and Bravo 1987 ¹⁵⁵	Dog/DOC-saline (n=7)	0.4-1.7%	Adult/15 days	↓ 24 MAP, <i>P</i> < .01	Peripheral resistance
Kang et al 1990 ¹⁵⁶	DS rat (n=10)	1.0-2.0%	11/30	↓ 17, <i>P</i> < .05	Sodium transport
Karanja et al 1987 ¹⁵⁷	SHR (n=7)	0.1/0.25/0.5/1.0/2.0%	6/14	↓ 13, <i>P</i> < .05	Calcium-BP dose response
Karanja et al 1989 ¹⁵⁸	SHR (n=10)	0.25-2.0%	3/26	↓ 16, <i>P</i> < .01	Fat/calcium
Kohno et al 1989 ¹⁵⁹	SHR/WKY (n=30)	0.4-2.9%	10/3	↓ 22, <i>P</i> < .05	ANP
Koide and Tuan 1989 ¹⁶⁰	Chick embryo (n=8-9)	0%	Prenatal	↓ 3 MAP, <i>P</i> < .05	Adrenergic regulation
Kurtz and Morris 1986 ¹⁶¹	DOC-saline (n=7-9)	0.5-4.0%	4/9	↓ 26 MAP	DOC-calcium interaction
Lau et al 1984 ¹⁶²	♀ SHR/WKY multiple groups	0.22/1.2/4.3%	Multiple experiments	~ ↓ 15, <i>P</i> < .05	PTX, PO ₄ , volume, hypercalcemia
Lau et al 1986 ¹⁶³	♀ SHR (n=10)	0.87-2.0%	3/12	↓ 16, <i>P</i> < .05	NaCl-Ca ²⁺ interaction, RAS
Lau et al 1986 ¹⁶⁴	SHR/WKY (n=11-46)	0.35/0.87/1.35%	3-15/5-11	No effect	Calcium balance
Lee et al 1984 ¹⁶⁵	Turkey (n=10)	0.98/1.96%	33/15	↓ 29 MAP, <i>P</i> < .005	LVH, stress
Lewanczuk et al 1990 ¹⁶⁶	SHR (n=10)	0.02/0.6/2.0%	4/4-8	↓ 33 MAP, <i>P</i> < .01	PHF
Luft et al 1988 ¹⁶⁷	SHRSP, WKY (n=20)	0.25-4.0%	6/16	↓ 24, <i>P</i> < .05, no effect of higher Mg	Mg homeostasis, catecholamines
Mangiarua et al 1990 ¹⁶⁸	SHR/WKY (n=12-60)	0.02/1.0/4.0%	5/8	↓ 24 MAP, <i>P</i> < .05	Hypertensive factor
McCarron et al 1981 ¹⁵¹	SHR/WKY (n=12)	0.25/0.5/4.0%	10/38	~ ↓ 30, <i>P</i> < .001	Calcium metabolism
McCarron 1982 ¹⁶⁹	WKY (n=12)	0.25/0.5/4.0%	8-10/24-26	↓ 14, <i>P</i> < .001	Calcium balance

amount of calcium has varied considerably, ranging from 5% to virtually no calcium in the diet. Most studies have used either 0.5% or 1.0% as the normal value for

dietary calcium. The smaller value corresponds to the recommended level of dietary calcium for rats based on the American Institute for Nutrition 1976 report.²⁰¹

TABLE 1. Continued

Study	Model	Calcium Diet	Age of Animal/ Treatment Length, wk*	Result in mm Hg SBP	Focus
McCarron et al 1985 ¹⁷⁰	SHR/WKY (n=7)	0.1/1.0/2.0%	6/14	↓ 56, $P<.01$	Na-Ca ²⁺ interaction, Ca ²⁺ flux
Muntzel et al 1989 ¹⁷¹	SHR (n=10-14)	0.1/1.0/2.0%	Prenatal/28 days postnatal	~ ↓ 14 MAP, $P<.01$	Early exposure
Oparil et al 1991 ¹⁷²	SHR-S, SHR-R, WKY (n=10-12)	0.68-2.0%	8/2	~ ↓ 14 MAP, $P<.01$	NaCl, SNS
Oshima et al 1992 ⁶⁸	SHRSP (n=15)	0.2-2.0%	3/6	~ ↓ 35, $P<.001$	Cellular calcium
Pamnani et al 1990 ¹⁷³	Wistar/reduced renal mass (n=21)	~1.0-3.0%	~10/5	↓ 40, $P<.05$	Electrolyte metabolism
Pang et al 1992 ¹⁷⁴	SHR (n=12)	0.2/0.4/0.8%	12/8	↓ 12 MAP, $P<.05$	Calcium antagonists
Pernot et al 1978 ¹⁷⁵	SD DOC-saline (n=7-12)	3.5% CaCl ₂ in water	~6/10	↓ 25, $P<.01$	PTH
Pernot et al 1985 ¹⁷⁶	SD DOC-saline SHR, ♀ LHR	DOC=1.3% CaCl ₂ in water, SHR=0.3-1.2%, LHR=0.6-2.5%	6/16 DOC, 6/44 SHR, 6/22 LHR	~ ↓ 30 DOC, ~ ↓ 18 SHR, ~ ↓ 16 LHR	Compare models
Pernot et al 1990 ¹⁷⁷	LHR (n=7-12)	0.03/0.6/2.5%	4/23	~ ↓ 22, $P<.01$	Vascular reactivity
Peuler et al 1987 ¹⁷⁸	DS rat (n=10-11)	0.4-4.0%	4/6-7	~ ↓ 30 MAP, $P<.05$	SNS
Peuler and Mark 1989 ¹⁷⁹	DS rat (n=10-13)	0.4-2.0%	6/6	~ ↑ 25, $P<.05$	Calcium BP, SNS
Peuler 1991 ¹⁸⁰	♀ SHRSP/DS rat (n=8-13)	0.4-2.0%	4-6/6-22	DS= ↑ 17 MAP, SHRSP= ↓ 38 MAP, $P<.05$	Divergent effects calcium on BP
Peuler and Schelper 1992 ¹⁸¹	♀ SHRSP (n=11-13)	0.4-2.0%	4 Weeks to death	~ ↓ 45 MAP, $P<.05$	Stroke
Porsti et al 1990 ¹⁸²	SHR DOC-saline (n=12)	1.5% CaCl ₂ in water	8/9	↓ 24 MAP, $P<.05$	Vascular reactivity, Ca ²⁺ -ATPase
Porsti et al 1992 ¹⁸³	SHR/WKY (n=16/9)	1.1-3.1%	8/12	↓ 20 MAP, $P<.0001$	Vascular relaxation, platelet [Ca ²⁺] _i , Na,K-ATPase
Portsi 1992 ¹⁸⁴	SHR/WKY (n=16/9)	1.1-2.1%	8/12	↓ 18 MAP, $P<.0001$	Vascular reactivity, platelet [Ca ²⁺] _i
Resnick et al 1986 ¹⁸⁵	Wistar DOC-saline/2K1C (n=10)	0.2-1.8%	~6/4	↓ 16 DOC, ↑ 21 2K1C, $P<.05$	RAS
Resnick et al 1986 ¹⁸⁶	Wistar DOC-saline/2K1C	1.2-1.8%	~6/4	↓ 23 DOC, ↑ 18 2K1C, $P<.05$	RAS
Saito et al 1991 ⁹¹	WKY DOC-saline (n=8-30)	1.0-4.0% CaCl ₂	6/4	↓ 55, $P<.01$	Electrolyte balance
Schleiffer et al 1984 ¹⁸⁷	SHR (n=7-9)	0.0/0.3/1.2%	5/39	~ ↓ 18, $P<.05$	BP
Scrogin et al 1991 ¹⁸⁸	SHR (n=8-9)	0.2-2.0%	4/8	~ ↓ 18, $P<.05$	NaCl-Ca ²⁺ , BP reactivity
Scrogin et al 1991 ¹⁸⁹	SHR (n=8-9)	0.2-2.0%	4/8	↓ 14, $P<.05$	Stress
Semafuko and Morris 1991 ¹⁹⁰	SHR/ADX-aldo (n=8-10/4-5)	0.5-2.5%	3/17 SHR, 6/2 SHR/ADX	↓ 20 SHR, ↓ 70 SHR/ADX, $P<.01$	Calcium/aldosterone
Stern et al 1984 ⁵⁰	SHR/WKY (n=10)	0.4-2.8%	6/4	No effect	Calcium metabolism
Stern et al 1987 ¹⁹¹	SHR (n=18)	0.4-2.5%	10/4	No effect	Vascular reactivity

TABLE 1. Continued

Study	Model	Calcium Diet	Age of Animal/ Treatment Length, wk*	Result in mm Hg SBP	Focus
Tamura 1987 ¹⁹²	SHR (n=6)	0.8-3.2%	5/4	↓ 19, <i>P</i> < .01	Mg ²⁺ , phosphorus
Tenner et al 1989 ¹⁹³	SHR/WKY (n=15-20)	0.02/0.1/0.5/ 2.5%	5/15	~ ↓ 45, <i>P</i> < .001	PTH, vascular reactivity
Togari et al 1990 ¹⁹⁴	Wistar (n=6)	0.01-0.3%	3/7	↓ 28	PTH-induced hypotension
Tresham et al 1988 ¹⁹⁵	Sheep (n=4)	50-200 mmol	Adult/6	No effect	BP
Wiecek et al 1989 ¹⁹⁶	PTX SHR/WKY (n=5-10)	0.95/1.6/2.4/ 3.0%	4/11	No effect in intact rats	PTH, vascular reactivity
Wuorela et al 1990 ¹⁹⁷	SHR DOC (n=12)	1.5% CaCl ₂ in water	9/4	~ ↓ 21, <i>P</i> < .01	Electrolyte balance, Ca ²⁺ -ATPase
Wuorela et al 1992 ¹⁹⁸	SHR DOC (n=12)	1.5% CaCl ₂ in water	8/9	↓ 26, <i>P</i> < .001	Ca ²⁺ -ATPase Na-K ratio
Wyss et al 1989 ¹⁹⁹	SHR-S (n=12-18)	0.68-2.0%	7/2	↓ 16 MAP, <i>P</i> < .05	Hypothalamic NE
Yang et al 1989 ²⁰⁰	SHR-S (n=6)	0.68-2.0%	7/2	↓ 16 MAP, <i>P</i> < .05	Hypothalamic α ₂ receptors

However, 1.0% is often considered the normal level because it is the amount contained in major brands of rat chow.

Based on the available literature, it appears that studies which have used fairly substantial differences in calcium intake (5- to 10-fold) between treatment groups have most consistently reported significant differences in blood pressure. Using five diets that ranged between 0.1% and 2.0% calcium, Karanja et al¹⁵⁷ reported a nonlinear inverse relation between the calcium content of the diet and blood pressure in SHR maintained on a concurrent high sodium chloride diet. When calcium was varied by a factor of two or less, differences in blood pressure were less likely.

The overall composition of the diet has varied considerably, reflecting the variety of dietary sources available to researchers around the world. The nutrients that have received the most attention in relation to calcium are other electrolytes. Sodium, chloride, potassium, magnesium, and phosphorus have frequently been modified along with calcium. Less consideration has been given to other factors in the diet, even though other nutrients may influence calcium metabolism. Fiber content is known to influence the absorption of calcium,²⁰² and the lipid profile of the diet has been shown to interact with calcium to alter blood pressure.¹⁵⁸

How rapidly blood pressure changes after dietary manipulation depends on the age of the animal as well as the composition of the diet. With normal levels of sodium chloride in the diet, 3-week-old animals show alterations in blood pressure to dietary calcium within 4 to 5 days.¹⁴¹ At 4 weeks of age, 2 weeks or longer may be required before a change in blood pressure is observed, whereas at 6 weeks of age, more than 2 months may pass without a change in blood pressure.⁵¹ In contrast, the effects of calcium may be apparent within 2 weeks in an 8-week-old salt-sensitive SHR if the animal is on a high sodium chloride diet.^{172,199,200}

The diet-related changes in blood pressure do not appear to be due to variations in body weight. Karanja

et al¹⁵⁷ reported that weight gain was greatest in SHR consuming 1.0% calcium diets, yet the rate of blood pressure rise was about half that of rats fed 0.25% calcium diets. It is of interest to note that animals on enriched calcium diets eat as much as 30% more but weigh less than animals on normal calcium diets.²⁰³

Proposed Mechanisms for the Antihypertensive Actions of Calcium

A multitude of potential mechanisms may explain the effect of dietary calcium on blood pressure. As dietary calcium changes, so do PTH, calcitriol, calcitonin, calcitonin gene-related peptide (CGRP), atrial natriuretic peptide (ANP), renin-angiotensin system activity, sympathetic nervous system activity, metabolism of other electrolytes, calcium binding proteins, intracellular free calcium levels, membrane-bound calcium, and electrolyte fluxes. Any one or combination of these changes could be responsible for altering blood pressure. In the following sections, several mechanisms that have been postulated to mediate the effects of dietary calcium on blood pressure will be explored. The proposed mechanisms are summarized in Table 2.

Vascular Smooth Muscle

Calcium plays a key role in vascular smooth muscle function. Calcium influx through receptor- and voltage-operated calcium channels is thought to initiate vascular contraction, and the fall in the intracellular free calcium concentration is thought to result in relaxation or vasodilation. Therefore, how the vascular smooth muscle cell (VSMC) handles calcium is critical to vascular tone and blood pressure. Anything that causes a perturbation in the processes that control cellular regulation of calcium would very likely change vascular tone.

In the following section, the effects of dietary calcium on VSMC regulation of calcium will be considered. In some instances, the effects on vascular smooth muscle are estimated from the effect of dietary calcium on platelets or other cell types. Although there are impor-

TABLE 2. Proposed Mechanisms of Action of Supplemental Dietary Calcium on Blood Pressure in Experimental Models of Hypertension

Site of Action	Mechanism
Cellular	↑ Membrane stabilization
	↑ Na,K-ATPase
	↑ Ca-ATPase
	↑ Calmodulin
	↓ Intracellular calcium
Vascular	↓ Contractility
	↑ Relaxation
Calcium-regulating hormones	↓ PTH
	↓ 1,25(OH) ₂ vitamin D ₃
	↑ Calcitonin
Calcium-sensitive hormones	↓ Parathyroid hypertensive factor
	↑ Atrial natriuretic peptide
	↓ Renin-angiotensin system
	↑ CGRP
Sympathetic nervous system	↓ CNS outflow
	↓ Circulating catecholamines
	↓ α_1 -Adrenergic receptor-mediated responses
Renal	↑ Natriuresis
	↑ Prostaglandins
	↑ Natriuresis

↑ indicates increase in; ↓, decrease in; PTH, parathyroid hormone; CGRP, calcitonin gene-related peptide; and CNS, central nervous system.

tant differences among cell types, it is assumed that the effect of dietary calcium on cellular calcium regulation is not restricted to VSMCs, and therefore, other cell types can be used as surrogates. This assumption is bolstered by the work of Oshima et al,⁶⁸ who observed a decline in platelet and lymphocyte intracellular free calcium in stroke-prone SHR maintained on high calcium/high sodium chloride diets. In their study, stroke-prone SHR were maintained on low or high calcium (0.2% versus 2.0%) and either low or high sodium (0.3% versus 3.1%). Animals on high sodium/low calcium diets had elevated blood pressure and high levels of basal and stimulated intracellular free calcium in both circulating platelets and intrathymic lymphocytes. Supplemental dietary calcium corrected the abnormalities in intracellular calcium. The similarity of findings in two different cell types indicates that the dietary calcium-induced correction of intracellular abnormalities of calcium metabolism may be a generalized phenomenon and not cell specific.

Membrane Stabilization

Based on numerous reports of abnormalities in cell membrane transport systems associated with hypertension, Bohr and colleagues²⁰⁴ have hypothesized that there is a defect in the lipid bilayer of vascular smooth muscle membrane in hypertensive individuals. This defect results in decreased membrane fluidity and enhanced excitability as a consequence of increased Ca²⁺ flux across the cell membrane. Bohr and coworkers^{135,204-209} have provided persuasive evidence that in-

creased extracellular calcium causes a reduction in vascular contractility through a process of membrane stabilization. Increased binding of calcium to the plasma membrane decreases membrane permeability to both monovalent and divalent cations, resulting in diminished vascular contractility.

Membrane stabilization is an attractive hypothesis for explaining the effects of dietary calcium, but most studies of membrane stabilization have been done in vitro using extracellular calcium levels that may not correspond to the magnitude of change that might be expected from manipulations of dietary calcium. To evaluate the effect of dietary calcium on cellular function, Furspan and Bohr²¹⁰ fed stroke-prone SHR and WKY rats either 1% or 2.5% calcium diets for 10 weeks. They then measured blood pressure, serum ionized calcium, and lymphocyte intracellular sodium, potassium, and calcium, as well as net passive fluxes of sodium and potassium. They found that supplemental calcium reduced blood pressure, increased serum ionized calcium, reduced net potassium efflux, and reduced free ionized calcium within the lymphocyte. The changes were attributed to increased extracellular calcium and resultant membrane stabilization in the calcium-supplemented stroke-prone SHR.

Na,K-ATPase

In a series of studies, Porsti et al¹⁸²⁻¹⁸⁴ reported that increasing dietary calcium lowers blood pressure and improves vascular relaxation in the SHR. They found that supplemental calcium normalized DOC-induced

elevations of blood pressure as well as the increased vascular contractility to norepinephrine that is characteristic of vessels from DOC-treated animals.¹⁸² Dietary calcium did not alter contractility to norepinephrine in SHR not treated with DOC, suggesting that the change in maximal contractility resulted from a calcium-induced modulation of the DOC treatment. On the other hand, supplemental calcium did enhance relaxation to sodium nitroprusside and acetylcholine in both the DOC-calcium group and the calcium-supplemented control SHR. The authors suggest that the improved relaxation may have been a consequence of increased Na,K-ATPase activity. In a previous report, they noted that potassium-induced relaxation of mesenteric arterial rings was augmented by calcium supplementation in SHR.¹⁸³ Because ouabain was able to prevent the relaxation, the difference between diet groups was attributed to increased Na,K-ATPase activity. This outcome is consistent with the notion of a reciprocal relation between intracellular calcium and Na,K-ATPase activity; as intracellular calcium declines, Na,K-ATPase activity increases.²¹¹

The results from Porsti et al¹⁸²⁻¹⁸⁴ are commensurate with other findings showing the maximal contractility to norepinephrine in intact vessels is not altered by supplemental dietary calcium. Aorta, mesenteric, and tail arteries from animals fed high calcium diets have not shown diminished reactivity to norepinephrine or potassium chloride (KCl).^{*} If anything, there may be increased contractility. Ambrozy et al¹¹⁶ found increased sensitivity to phenylephrine in Zucker obese and lean rats fed high calcium diets, whereas Pernot et al¹⁷⁷ found an increase in maximal contractility to norepinephrine in Lyon hypertensive rats on high calcium diets.

Despite the *in vitro* results, blood pressure reactivity *in vivo* has been reported to be reduced to norepinephrine in animals fed supplemental calcium. Several investigators have reported diminished pressor responses to norepinephrine in both hypertensive^{142,153,180} and normotensive^{120,132} rats fed high calcium diets. These results would not appear to be a consequence of diminished vascular contractility given the results from isolated vessels cited above. However, the *in vivo* results may be related to circulating factors that modify the response to norepinephrine or to difficulties related to vascular relaxation.

Ca²⁺-ATPase

Just as Furspan and Bohr²¹⁰ observed lower levels of intracellular free calcium in lymphocytes from animals on high calcium diets, Porsti et al¹⁸³ reported lower levels of intracellular free calcium in platelets isolated from animals maintained on supplemental calcium diets. These favorable changes in intracellular calcium regulation may be related to increased Ca²⁺-ATPase activity. Porsti et al¹⁸² and Wuorela et al¹⁹⁸ observed an increase in the maximal velocity of calcium transport by "inside-out" red blood cell vesicles from SHR on calcium-supplemented diets. The greater velocity of calcium transport is indicative of increased Ca²⁺-ATPase activity. Wuorela et al¹⁹⁸ also reported that nitroprusside-induced relaxation of norepinephrine-contracted mes-

enteric arterial rings was enhanced in calcium-supplemented animals, perhaps as a consequence of an increased ability to extrude calcium from the cell.

Calmodulin

Using duodenal enterocytes, Roullet et al²¹² have observed that calmodulin levels are lower in SHR than WKY rats and that increasing dietary calcium eliminates that difference. Subsequent work indicated that dietary calcium can actually upregulate calmodulin levels (C.M. Roullet, personal communication, November 1993). Correction of a defect in calmodulin activity by calcium could provide a mechanism whereby a multitude of molecular and cellular processes might be modified. Calmodulin plays a pivotal role in intracellular calcium regulation and could be responsible for diet-induced variations in Ca²⁺-ATPase.^{213,214}

Calcium-Regulating Hormones

Until recently, calcium-regulating hormones were rarely considered outside of their traditional role in calcium homeostasis. Calcium-regulating hormones exert tight control over serum free ionized calcium levels (for review, see Reference 215). Reductions in serum ionized calcium provoke release of PTH from the parathyroid gland. PTH in turn stimulates the production of calcitriol [1,25(OH)₂ vitamin D₃], the most active metabolite of vitamin D, in the proximal convoluted tubule cell. Calcitriol enters the circulation to reach its target tissues, the principal targets being the renal and intestinal epithelia where 1,25(OH)₂ vitamin D₃ increases the synthesis of the calcium-binding proteins calbindin and integral membrane calcium-binding protein. An increase in these calcium-binding proteins is reflected by greater intestinal absorption and renal reabsorption of calcium. When serum free ionized calcium levels are too high, calcitonin is released from the thyroid gland. This compound acts to lower serum free ionized calcium by inhibiting osteoclast activity and preventing mobilization of calcium from bone.

It has become increasingly clear that PTH, calcitriol, and perhaps calcitonin possess vasoactive properties and thus may play a role in blood pressure regulation. Furthermore, other vasoactive peptides and hormones have been identified that are responsive to dietary calcium such as CGRP and parathyroid hypertensive factor. Because these compounds are responsive to changes in calcium intake, it follows that variations in calcium-regulating hormones may account for calcium-induced alterations in blood pressure.

Parathyroid hormone. Hyperparathyroidism is associated with elevated blood pressure in humans,^{32,216,217} and PTH is elevated in many essential hypertensive patients.^{32,35,217-223} These observations suggest that elevated PTH may in some way be responsible for the high blood pressure. If so, dietary calcium may lower blood pressure by suppressing PTH. However, such an effect would be paradoxical because PTH is a potent vasodilator. Intravenous injections of PTH result in a prompt, dose-dependent fall in blood pressure.^{224,225} Nevertheless, parathyroidectomy has been consistently shown to lower blood pressure in animal models of hypertension.²²⁶⁻²²⁸ Furthermore, transplantation of parathyroid glands between WKY rats and SHR results in higher

*References 116, 128, 129, 141, 177, 183, 184.

blood pressure in the WKY rat and lower blood pressure in the SHR.^{226,229,230}

The resolution of the paradox may be related to a number of factors. First, it should be noted that at physiological concentrations in blood, PTH does not appear to be vasoactive.²³¹⁻²³³ Thus, the influence of PTH on blood pressure may be independent of its vasodilating effects. Second, prolonged exposure to low calcium diets, which elevates circulating PTH levels,⁵⁰ diminishes the magnitude of the blood pressure decline after an acute injection of PTH.¹¹⁸ The potential development of tolerance to the vasodilator effects of PTH may be associated with a simultaneous enhancement in response to other vasoactive drugs. This situation would be analogous to the effect of alcohol exposure on arterial pressure. Acutely, alcohol is a vasodilator, but with continued exposure it is associated with hypertension. Several studies have shown that as the vasodilator effects of alcohol decline, there is an increase in intracellular free calcium in VSMCs that may favor enhanced responses to other agonists (for review, see Reference 234).

PTH may act as a calcium ionophore to promote increased intracellular calcium levels in VSMCs and thereby increase vascular tone. The possibility of PTH as an ionophore in vascular smooth muscle is suggested by findings that low concentrations of PTH induce calcium influx into isolated cardiac cells²³⁵ as well as osteoblasts.²³⁶ The relation between PTH and platelet intracellular calcium has been examined in humans and in rats. Oshima et al²³⁷ could find no effect of parathyroidectomy on platelet free intracellular calcium in SHR. In humans, there is little intercorrelation between intracellular free calcium and PTH activity.²³⁸ Thus, the evidence does not favor the ionophoric effect of PTH in hypertension.

It is possible that PTH has central nervous system effects that promote an elevation of blood pressure. In a brief article, Delbarre et al²³⁹ reported that intracerebral ventricular administration of PTH resulted in significantly elevated blood pressure in the SHR. Although this study is apparently the only one of its kind, there are other indications that PTH has central nervous system effects. Geiger et al²⁴⁰ reported altered cyclic AMP in various brain regions in the rat that were attributable to variations in PTH. More recently, Islam et al²⁴¹ found evidence that PTH diminished synaptosomal phospholipid content in uremic rats. In a subsequent study, Smogorzewski et al²⁴² reported that PTH was responsible for a higher calcium content in rat brain synaptosomes as well as reduced Na,K-ATPase activity. PTH also modified epinephrine content, uptake, and release.

An additional explanation for the PTH paradox is the putative parathyroid hypertensive factor. Lewanczuk et al²⁴³ have reported the presence of a circulating hypertensive factor that is present in both SHR and human plasma.²⁴⁴ When injected into normotensive rats for bioassay, the hypertensive factor stimulates vascular smooth muscle calcium uptake and causes an elevation in blood pressure with a time lag of about 2 hours. Parathyroidectomy eliminates the circulating factor, suggesting that the parathyroid gland is the source.²²⁶ If so, this would shift focus away from PTH to the parathyroid gland itself and in so doing would reconcile the

disparate blood pressure results that have previously been ascribed to PTH. Pertinent to this review, the purported factor is responsive to variations in dietary calcium. Increased dietary calcium reduces circulating hypertensive factor, whereas restricted calcium increases it.¹⁶⁷

Calcitriol. As mentioned previously, low calcium diets provoke an increase in the synthesis and release of 1,25(OH)₂ vitamin D₃, (see Reference 215), which itself may mediate an increase in blood pressure. Receptors for 1,25(OH)₂ vitamin D₃ have now been demonstrated in vascular tissue.^{246,247} Calcitriol has been shown to stimulate Ca²⁺-ATPase in vascular smooth muscle, suggesting that 1,25(OH)₂ vitamin D₃ may play a role in regulating cellular calcium metabolism.^{248,249} A direct effect of calcitriol on VSMC intracellular free Ca²⁺ has been reported for calcitriol in intact mesenteric resistance arteries as well as isolated VSMCs. Bukoski et al²⁵⁰ and Xue et al²⁵¹ have shown that short-term incubation of intact mesenteric resistance vessels with 1,25(OH)₂ vitamin D₃ increases the Ca²⁺ transient induced by norepinephrine in the SHR. No effects were seen in the WKY rat. Similar effects are observed after systemic injections of calcitriol. Bukoski et al^{128,252} observed that both acute injections and 3-day administration of the vitamin at physiological levels significantly enhanced maximal force generation of isolated vessels from both hypertensive and normotensive animals. These findings suggest direct effects of 1,25(OH)₂ vitamin D₃ on Ca²⁺ metabolism of intact resistance vessels. They further suggest that the observed response to calcitriol differs depending on whether the vascular tissue is isolated from the normal or hypertensive animal.

Despite evidence of direct effects on vascular smooth muscle, observations of elevated blood pressure as a consequence of exposure to 1,25(OH)₂ vitamin D₃ have been inconsistent. The animals that Bukoski et al²⁵² examined for vascular contractility did not have elevated blood pressure as a consequence of exposure to calcitriol. Likewise, Hatton et al²⁵³ found evidence of enhanced vascular contractility in vessels from animals given 1,25(OH)₂ vitamin D₃ injections for 7 consecutive days, but there were no changes in blood pressure. Using a higher concentration of 1,25(OH)₂ vitamin D₃, Shimosawa et al²⁵⁴ did observe a potentiation of pressor responses to both norepinephrine and angiotensin II in animals treated with calcitriol or the noncalcemic analogue 22-oxacalcitriol for 14 days.

Most recently, Bukoski and Xue²⁵⁵ found evidence that daily injections of 1,25(OH)₂ vitamin D₃ over a 28-day period caused sustained elevations in blood pressure in normotensive Wistar rats. However, chronic administration of 1,25(OH)₂ vitamin D₃ in SHR produced only a transient elevation in blood pressure after 5 weeks of treatment, suggesting a modest effect of the hormone on blood pressure.²⁵⁶ As reported in previous studies, vascular contractility was significantly enhanced in the SHR.

Thus, considering the direct effects on the vasculature, the observation of potentiated pressor responses, and recent reports of elevated blood pressure, 1,25(OH)₂ vitamin D₃ should be considered as an agent that has the potential to alter blood pressure. Further

work is needed to understand the cellular events underlying the vascular smooth muscle response to calcitriol.

Calcitonin. In general, the effects of calcitonin on calcium regulation are opposite to those of PTH and $1,25(\text{OH})_2$ vitamin D_3 . Although supplemental dietary calcium can be expected to elevate calcitonin levels, calcitonin has not been routinely measured in studies of the effect of dietary calcium manipulations on blood pressure. Nevertheless, there is evidence that calcitonin may have vasoactive properties.

There have been conflicting reports on the relation of calcitonin to blood pressure. Bindels et al⁴⁹ found significantly elevated calcitonin in young SHR, along with reduced serum ionized calcium and elevated $1,25(\text{OH})_2$ vitamin D_3 . Presumably, the disparate results between elevated calcitonin in the presence of reduced serum ionized calcium were related to phosphate metabolism. These findings suggest that calcitonin is associated with elevated blood pressure. In fact, Clementi et al²⁵⁷ reported elevated blood pressure after intracerebroventricular administration of 0.4 IU salmon calcitonin. The increased blood pressure was accompanied by increased plasma renin activity. It is unlikely that the increased plasma renin activity contributed to the elevation of blood pressure, because intravenous and intramuscular administration of calcitonin elevated plasma renin activity without altering blood pressure.

Other investigators have found conflicting evidence. Delbarre et al²³⁹ observed a reduction in blood pressure with intracerebroventricular administration of calcitonin but an elevation of blood pressure with peripheral administration. Wegener and McCarron²⁵⁸ were unable to detect any effect of calcitonin on blood pressure through intravenous administration.

Additional research will be needed to assess the effects of calcitonin on blood pressure and its relation to dietary calcium. Calcitonin may well have an influence on blood pressure via the central nervous system. Although calcitonin is a thyroid hormone, a number of calcitonin receptors in the brain²⁵⁹ mediate a variety of effects, including anorexia,²⁶⁰ analgesia,²⁶¹ and altered pituitary release of hormones.²⁶² However, at the present time there is insufficient evidence to evaluate the possibility that calcitonin modulates blood pressure via the central nervous system.

Calcitonin gene-related peptide. In contrast to calcitonin, CGRP is known to be vasoactive. This 37-amino acid neuropeptide results from alternative processing of the calcitonin gene. It is widely distributed in the central and peripheral nervous systems²⁶³ and is one of the most potent vasodilators yet discovered. Intravenous infusion of CGRP produces marked hypotension through a reduction in total peripheral resistance.^{264,265} Central nervous system administration of CGRP produces variable effects depending on the dose and site of administration. For example, injection of 0.2 pmol CGRP into the nucleus tractus solitarius produced a depressor response, whereas 2 pmol resulted in a pressor response.²⁶⁶ Likewise, injections into the central nucleus of the amygdala resulted in a pressor response,²⁶⁷ but intrathecal injections caused a reduction in blood pressure.²⁶⁸

Of particular interest are observations that levels of CGRP in spinal cord vary with dietary calcium. DiPette et al²⁶⁹ demonstrated that low levels of dietary calcium

reduced dorsal horn CGRP content, and supplemental calcium increased CGRP levels. Although DiPette et al²⁶⁹ did not assess circulating levels of CGRP in their study, others²⁷⁰ have found it to be lower in SHR than WKY rats. Given the vasoactive properties of CGRP and its presence in central and peripheral nervous system sites involved in cardiovascular regulation, CGRP may be involved in the blood pressure alterations that result from dietary calcium manipulations.

Sympathetic Nervous System

There are consistent reports of altered sympathetic nervous system activity associated with variations in dietary calcium. Winternitz et al^{271,272} and Oparil et al²⁷³ have reported that increased sodium chloride causes an elevation of blood pressure in salt-sensitive SHR. The increased blood pressure is associated with elevated circulating norepinephrine and reduced hypothalamic norepinephrine.²⁷⁴ Provision of supplemental calcium reverses the increased blood pressure induced by sodium chloride while normalizing circulating and central nervous system changes in catecholamines.^{172,199,200} Peuler et al¹⁷⁸ reported that 4.8% calcium diets reduce blood pressure in DS rats by attenuating sympathetic nervous system outflow. Interestingly, 2% calcium diets aggravated hypertension in this strain¹⁷⁹ while reducing blood pressure in the stroke-prone SHR.¹⁸⁰

There are several possible mechanisms through which calcium may modify the sodium chloride effect on sympathetic nervous system outflow. One possibility is that the electrolytes may interact within the central nervous system. Infusion of sodium chloride into the cerebral ventricles²⁷⁵ or nucleus tractus solitarius²⁷⁶ has been shown to cause an elevation of blood pressure. In contrast, intracerebroventricular calcium causes a reduction in blood pressure.²⁷⁷ Therefore, it appears that both calcium and sodium chloride are reciprocally vasoactive in the central nervous system. There is evidence that manipulations of dietary calcium result in altered levels of central nervous system calcium. Long-term alterations in dietary calcium have been shown to change cerebrospinal fluid and brain levels of calcium in rats. Harris et al²⁷⁸ found the reduction in calcium levels to be greatest in the brain stem (24%) after 4 weeks of restricted dietary calcium (0.02%), with the soluble and microsomal fractions of calcium selectively reduced. Likewise, Tai et al²⁷⁹ found that calcium fluxes into brain and cerebrospinal fluid were linearly related to plasma ionized calcium. Thus, it appears that although the brain is protected from acute changes in serum ionized calcium by low cerebrovascular permeability to calcium, long-term changes in dietary calcium can alter brain levels of calcium, albeit to a lesser extent than the changes in serum calcium.²⁸⁰

The interaction between sodium and calcium may occur at the receptor level. Gavras²⁸¹ has hypothesized that sodium may directly influence neural activity and cause increased sympathetic nervous system outflow by attenuating the affinity of α_2 -adrenergic receptors for their agonists. Because activation of hypothalamic α_2 receptors causes inhibition of sympathetic outflow, reduced affinity would result in disinhibition of the sympathetic system. In contrast to sodium, divalent cations such as calcium increase the affinity of the receptors for the ligand.²⁸²

Whether changes in central nervous system calcium levels are responsible for alterations in central catecholamine levels remains to be demonstrated. Nevertheless, reductions in dietary calcium comparable to those used to modify central nervous system calcium levels are reported to alter central nervous system norepinephrine levels. Baksi¹²² reported that 8 weeks of a 0.005% calcium diet reduced hypothalamic norepinephrine and dopamine content in Sprague-Dawley rats. Restricted dietary calcium has also been reported to alter peripheral catecholamine levels. Baksi and Hughes²⁸³ and Hagi-hara et al²⁸⁴ reported altered adrenal catecholamine levels with reduced calcium diets, and Luft et al¹⁶⁷ observed reduced epinephrine levels and epinephrine responses to stress in stroke-prone SHR on high calcium diets. Scrogin et al¹⁸⁸ also observed diminished circulating epinephrine in calcium-supplemented SHR. Felicetta et al²⁸⁵ observed an increased urinary dopamine-norepinephrine excretion ratio during a high calcium/high sodium chloride diet. Other investigators have reported no difference in circulating catecholamines as a consequence of altered calcium diets.^{142,152,191}

Responses to norepinephrine have been reported to be potentiated by restricted calcium diets and dampened by high calcium diets. Hatton et al^{142,144} reported diminished pressor responses to exogenous norepinephrine in SHR on high calcium diets. The reduced pressor response did not occur in response to angiotensin II, suggesting that it was not due to a generalized change in vascular responsiveness. Doris,¹³² Kageyama et al,¹⁵²⁻¹⁵⁴ Baksi,¹²¹ and Peuler¹⁸⁰ have reported similar results with regard to norepinephrine.

The reduced responsiveness to norepinephrine may be related to altered adrenergic receptor activity. Hatton et al¹⁴⁴ demonstrated that dietary calcium specifically modulates the α_1 -adrenergic receptor. Blockade of α_1 -adrenergic receptors with phentolamine or prazosin eliminated the difference in blood pressure that prevailed in animals on high and low calcium diets. Blockade of α_2 -, β_1 -, or β_2 -adrenergic receptors had no such effect. Likewise, pharmacologic reduction of blood pressure with CGRP, sodium nitroprusside, or the converting enzyme inhibitor captopril had no differential effect on blood pressure. The results of this study strongly suggest that dietary calcium modifies the α_1 -adrenergic receptor. The nature of the effect remains to be determined, but preliminary data indicate that there may be a difference in receptor expression because binding to tritiated prazosin was greater in kidneys from animals fed low calcium diets.

Electrolyte Interactions

A change in the level of one dietary electrolyte can have extensive ramifications on other dietary electrolytes. Increased dietary calcium, for example, can increase sodium excretion, reduce magnesium absorption, and reduce circulating phosphorus levels. Likewise, increasing sodium chloride in the diet causes calcium wasting, but if sodium levels are altered independently of chloride, calcium is unaffected. Consequently, when manipulation of a dietary electrolyte is shown to have an effect on blood pressure, the possibility exists that the blood pressure effect is due to reciprocal changes in other electrolytes.

Phosphate depletion. Early in the study of the effect of dietary calcium on blood pressure, Lau and coworkers¹⁶³ proposed that calcium-induced phosphate depletion accounted for the blood pressure reduction. However, rather than a positive relation between phosphate and blood pressure as predicted by Lau et al,¹⁶² Bindels et al⁴⁹ demonstrated that supplementing the SHR diet with phosphate results in a blood pressure reduction during mid-adolescence. To specifically determine the role of phosphorus in calcium-induced reductions in blood pressure, Tamura¹⁹² varied both calcium and phosphate and found that calcium lowered blood pressure regardless of serum phosphate levels. Thus, it would appear that the effect of calcium on blood pressure is relatively independent of phosphate.

Magnesium. Dietary calcium intake can have a pronounced effect on magnesium metabolism.¹⁹² With increased dietary calcium intake, serum magnesium levels decline. Furthermore, observations of increased intracellular calcium with low calcium diets suggest that intracellular magnesium may be reduced.²⁴⁴ Because of the interactive effects of dietary calcium and magnesium, it has been suggested that some of the effect of calcium on blood pressure may be mediated through alterations in magnesium. Indeed, Tamura¹⁹² found a significant positive correlation of serum magnesium levels with systolic blood pressure across groups receiving different levels of dietary calcium. However, in a factorial study of calcium and magnesium, Evans et al¹³³ found no effect of magnesium on blood pressure and no interactive effects of the two cations on blood pressure.

Sodium chloride/calcium interactions. Numerous studies have found that calcium attenuates sodium chloride-dependent, low-renin hypertension. Calcium lowers blood pressure in several salt-sensitive models of hypertension, including DOC-saline rats,^{91,130,145,154,176,186} reduced renal mass/saline rats, DS rats,^{156,173,178,182} salt-sensitive SHR,^{149,172,199,200} salt-fed WKY rats,^{131,132} angiotensin II-salt rats,¹¹⁹ and adrenalectomy/aldosterone-induced hypertension.¹⁹⁰ The mechanisms through which calcium attenuates salt-induced forms of hypertension have not been identified, but numerous candidates have been suggested.

Calcium-induced natriuresis. Increasing dietary calcium facilitates natriuresis in a number of ways. Increased circulating ANP, reduced sympathetic nervous system outflow, reduced α_1 -adrenergic receptor activity, reduced angiotensin II receptor expression, reduced circulating PTH, and direct effects of calcium on sodium excretion at the proximal tubule may all play a role. The most familiar natriuretic effect of calcium is the interaction that occurs at the renal level. Increasing calcium facilitates natriuresis and diuresis in part by inhibiting sodium reabsorption in the proximal tubule.²⁸⁶ Conversely, reducing serum ionized calcium levels by acute volume expansion retards natriuresis and diuresis.²⁸⁷ This occurs, at least in part, as a consequence of increased circulating PTH acting on the tubules to alter production of prostaglandins, with a resultant reduction in sodium excretion²⁸⁸ as well as direct effects of calcium on prostaglandin production. Less well known are the effects of increasing dietary calcium on other systems that promote natriuresis. Geiger et al¹³⁷ and Kohno et al¹⁵⁹ have reported increased circulating ANP after supplemental calcium diets. Such an increase in ANP

would promote natriuresis as well as cause peripheral vasodilation and reduced blood pressure. Levi and Henrich²⁸⁹ recently reported that high calcium diets reduced angiotensin II binding in renal brush-border membranes and decreased brush-border membrane Na-H antiport activity in SHR. Angiotensin II increases proximal tubular sodium transport by increasing Na-H antiport activity. Decreased activity in this system would enhance sodium excretion.

The effect of calcium on sympathetic nervous system activity would also facilitate sodium excretion. The reduction in sodium chloride-induced increases in sympathetic nervous system outflow that have been reported¹⁷² can be expected to result in increased sodium excretion, as would the diminished α_1 -adren-ergic activity during high calcium diets, as reported by Hatton et al.¹⁴⁴

Given the multiple mechanisms through which increased dietary calcium facilitates sodium excretion, it is not surprising that increasing the proportion of calcium in the diet results in natriuresis.* In some models of hypertension, natriuresis may be an important component of the antihypertensive effect of dietary calcium. This would apply regardless of how sodium chloride might provoke hypertension, because removal of sodium chloride would remove the initial stimulus for elevated blood pressure.

Not all evidence favors a natriuretic explanation for the blood pressure effects of dietary calcium. In SHR that are not salt sensitive, increasing dietary sodium chloride does not alter blood pressure. However, increasing dietary calcium does lower blood pressure. Studies from our laboratory¹⁷⁰ and that of Hamet et al²⁹⁰ have demonstrated that the antihypertensive action of calcium may actually be enhanced by simultaneous sodium chloride supplementation. This interactive effect of these nutrients is not likely to be related to natriuresis because, as mentioned above, manipulation of sodium chloride alone does not alter blood pressure in these animals.

Thus, it may be necessary to distinguish between hypertensive models when assessing the interactive effects of dietary calcium and sodium chloride. For models in which calcium prevents a sodium chloride-induced elevation in blood pressure, such as in DOC-saline hypertension, natriuresis might be suspected. Otherwise, natriuresis seems less likely. Even when natriuresis is an unlikely cause for the blood pressure-lowering effect of dietary calcium, as in the SHR, sodium chloride and calcium may interact to potentiate the antihypertensive effects of supplemental calcium.

Sodium chloride-induced calciuresis. Just as calcium promotes natriuresis, sodium chloride promotes calciuresis. Consequently, as sodium chloride is increased in the diet, more calcium is excreted. Resnick and colleagues^{244,291,292} hypothesize that this dietary sodium chloride-induced calciuresis results in reduced serum ionized calcium and increased circulating levels of calcium-regulating hormones including calcitriol, PTH, and/or parathyroid hypertensive factor. In turn, these hormones stimulate cellular calcium uptake and disrupt cellular electrolyte balance, resulting in elevated blood

pressure. According to this hypothesis, calcium supplementation increases available calcium, prevents the fall in ionized calcium that provokes the increase in calcium-regulating hormones, and reverses the salt-induced hypertension.

Favoring this hypothesis are indications that increased dietary sodium chloride causes renal wasting of calcium^{293,294} and stimulates calcitriol and PTH production.²⁹³ Elevated calcitriol and PTH have been found in low-renin human hypertension²⁹² as well as in experimental forms of hypertension.⁹¹ Kotchen et al²⁹⁵ reported that after 5 days of a high salt diet, young DS rats exhibited increased calcium excretion in both urine and feces, whereas ionized calcium was reduced and calcitriol and PTH were elevated relative to Dahl salt-resistant rats. The differences in calcium excretion occurred with no changes in blood pressure. The excessive calcium excretion could be avoided by sodium loading with anions other than chloride, suggesting that the combination of sodium and chloride is critically important to the calciuresis that occurs. It is notable that high sodium diets that do not contain chloride also do not cause hypertension in the DS rat. Similarly, Kurtz and Morris¹⁶¹ found increased calcium excretion in DOC-saline hypertension when the animals were fed sodium chloride. However, if an equimolar amount of sodium bicarbonate was substituted for sodium chloride, the increases in both calcium excretion and blood pressure were reversed.

To test the hypothesis that sodium chloride causes hypertension as a consequence of alterations in calcium-regulating hormones, Resnick et al¹⁸⁶ fed supplemental dietary calcium to low-renin, DOC-salt hypertensive rats and to renin-dependent Goldblatt hypertensive rats. Calcium excretion was substantially greater in the DOC-salt rats relative to the two-kidney, one clip Goldblatt rats. Modest increases in dietary calcium (1.2% versus 1.8%) caused a reduction of blood pressure in the DOC-salt animal but resulted in an increase in blood pressure in the Goldblatt model of hypertension.

Although these results are dramatic, other investigators have found that supplemental dietary calcium reduces renovascular hypertension. Kageyama et al¹⁵³ reduced blood pressure in two-kidney, one clip Goldblatt hypertensive rats by providing supplemental calcium (1.5%) in the drinking water. The reduction in blood pressure was accompanied by a suppression of plasma renin and aldosterone as well as diminished pressor responses to norepinephrine, leading the authors to conclude that the reduction in blood pressure may have been a consequence of reduced renin-angiotensin system activity.

Nonvascular Mechanisms

Nonvascular mechanisms must also be considered in explaining the role of dietary calcium in blood pressure regulation. Tordoff and colleagues^{296,298} have provided important new insights as to potential nonvascular mechanisms. Their studies in three genetically distinct animal models indicate that dietary calcium regulates sodium chloride appetite or intake. Low dietary calcium stimulates sodium chloride intake, and higher intakes of calcium suppress sodium chloride appetite. Thus, while volume may not be changed at higher calcium intakes,

*References 54, 91, 119, 131, 137, 149, 156, 159, 173, 190, 191, 286-288.

exposure to sodium chloride is reduced with potentially important effects on sodium-regulating hormones such as renin, aldosterone, and angiotensin II.

Change in blood viscosity is another nonvascular mechanism potentially affected by dietary calcium. Hatton et al¹⁴³ have measured changes in hematocrit that occur with calcium supplementation as an indirect assessment of viscosity. They found that there is an inverse relation between calcium intake and hematocrit. However, the changes in blood pressure observed with the higher calcium intake were independent of changes in hematocrit.

Hamet and Tremblay²⁹⁹ have linked correction of the abnormal HSP70 gene expression of experimental hypertension to increased dietary calcium intake. The HSP70 gene, which exists in the major histocompatibility complex and segregates closely with hypertension, has its expression corrected (suppressed) by dietary calcium.²⁹⁹

Summary

More than 80 studies have reported lowered blood pressure after dietary calcium enrichment in experimental models of hypertension. The evidence presented here suggests that dietary calcium may act concurrently through a number of physiological mechanisms to influence blood pressure. The importance of any given mechanism may vary depending on the experimental model under consideration.

Supplemental dietary calcium is associated with reduced membrane permeability, increased Ca^{2+} -ATPase and Na,K-ATPase, and reduced intracellular calcium. These results suggest that supplemental calcium may limit calcium influx into the cell and improve the ability of the VSMC to extrude calcium. This could be a direct effect of calcium on the VSMC or an indirect effect mediated hormonally.

The calcium-regulating hormones have all been found to have vasoactive properties and therefore may influence blood pressure. Furthermore, CGRP and the proposed parathyroid hypertensive factor are both vasoactive substances that are responsive to dietary calcium. Therefore, diet-induced variations in calcium-regulating hormones may influence blood pressure.

Modulation of the sympathetic nervous system is another important way that dietary calcium can influence blood pressure. There is evidence of altered norepinephrine levels in the hypothalamus as a consequence of manipulations of dietary calcium as well as changes in central sympathetic nervous system outflow. Dietary calcium has also been shown to specifically modify α_1 -adrenergic receptor activity in the periphery.

In some experimental models of hypertension, dietary calcium may alter blood pressure by changing the metabolism of other electrolytes. For example, the ability of calcium to prevent sodium chloride-induced elevations in blood pressure may be attributed to natriuresis. However, natriuresis does not account for all of the interactive effects of calcium and sodium chloride on blood pressure. Sodium chloride-induced hypertension may be due in part to calcium wasting and subsequent elevation of calcium-regulating hormones. Chloride is an important mediator of this effect because it appears that sodium does not cause calcium wasting when it is not combined with chloride.

More attention to the central nervous system effects of dietary calcium is needed. Not only can calcium itself influence neural function, but many of the calcium-regulating hormones appear to affect the central nervous system. The influence of calcium and calcium-regulating hormones on central nervous system activity may have important implications for blood pressure regulation and also may extend to other aspects of physiology and behavior.

References

1. Young EW, Bukoski RD, McCarron DA. Calcium metabolism in essential hypertension. *Proc Soc Exp Biol Med*. 1988;187:123-141.
2. McCarron DA. Calcium metabolism and hypertension. *Kidney Int*. 1989;35:717-736.
3. Morris CD, Hatton DC, Young EW, Karanja N, McCarron DA. Calcium nutrition and metabolism and pressure control. In: Rapaport E, ed. *Cardiology Update: Reviews for Physicians*. New York, NY: Elsevier Science Publishing Co, Inc; 1990:175-211.
4. McCarron DA, Morris CD, Cole C. Dietary calcium in human hypertension. *Science*. 1982;217:267-269.
5. McCarron DA, Morris CD, Henry HJ, Stanton JL. Blood pressure and nutrient intake in the United States. *Science*. 1984;224:1392-1398.
6. Bei-fang Z, Xi-gui W, Shou-qi T. Dietary patterns in 10 groups and the relationship with blood pressure. *Chin Med J (Engl)*. 1989;102:257-261.
7. Ackley S, Barrett-Connor E, Suarez L. Dairy products, calcium, and blood pressure. *Am J Clin Nutr*. 1983;38:457-461.
8. Garcia-Palmieri MR, Costas R Jr, Cruz-Vidal M, Sorlie PD, Tillotson J, Havlik RJ. Milk consumption, calcium intake, and decreased hypertension in Puerto Rico: Puerto Rico Heart Health Program Study. *Hypertension*. 1984;6:322-328.
9. Harlan WR, Hull AL, Schmoeder RL, Landis JR, Thompson RE, Larkin FA. Blood pressure and nutrition in adults: The National Health and Nutrition Examination Survey. *Am J Epidemiol*. 1984;120:17-28.
10. Gruchow HW, Sobocinski KA, Barborki JJ. Alcohol, nutrient intake, and hypertension in US adults. *JAMA*. 1985;253:1567-1571.
11. Liebman M, Chopin LF, Carter E, Clark AJ, Disney GW, Hegsted M, Kenney MA, Kirmani AZ, Koonce KL, Korslund MK, Moak SW, McCoy H, Stallings SF, Wakefield T. Factors related to blood pressure in a biracial adolescent female population. *Hypertension*. 1986;8:843-850.
12. Gruchow HW, Sobocinski KA, Barborki JJ. Threshold effects of dietary calcium on blood pressure. *J Hypertens*. 1986;4 (suppl 5):S355-S357.
13. Harlan WR, Hull AL, Schmoeder RL, Landis JR, Larkin FA, Thompson FE. High blood pressure in older Americans: The First National Health and Nutrition Examination Survey. *Hypertension*. 1984;6:802-809.
14. Harlan WR, Landis JR, Schmoeder RL, Goldstein NG, Harlan LC. Blood lead and blood pressure: relationship in the adolescent and adult US population. *JAMA*. 1985;253:530-534.
15. Johnson NE, Smith EL, Freudenheim JL. Effects on blood pressure of calcium supplementation in women. *Am J Clin Nutr*. 1985;42:12-17.
16. Kromhout D, Bosschieter EB, Coulander CL. Potassium, calcium, alcohol intake and blood pressure: the Zutphen Study. *Am J Clin Nutr*. 1985;41:1299-1304.
17. Sempos C, Cooper R, Kovar MG, Johnson C, Drizd T, Yetley E. Dietary calcium and blood pressure in National Health and Nutrition Examination Surveys I and II. *Hypertension*. 1986;8:1067-1074.
18. Reed D, McGee D, Yano K, Hankin J. Diet, blood pressure, and multicollinearity. *Hypertension*. 1985;7:405-410.
19. Kok FJ, Vandenbroucke JP, Van Der Heide-Wessel C, Van Der Heide RM. Dietary sodium, calcium, and potassium, and blood pressure. *Am J Epidemiol*. 1986;123:1043-1048.
20. Joffres MR, Reed DM, Yano K. Relationship of magnesium intake and other dietary factors to blood pressure: the Honolulu Heart Study. *Am J Clin Nutr*. 1987;45:469-475.
21. Witteman JCM, Willett WC, Stampfer MJ, Colditz GA, Sacks FM, Speizer FE, Rosner B, Hennekens CH. A prospective study of nutritional factors and hypertension among U.S. women. *Circulation*. 1989;80:1320-1327.

22. McGarvey ST, Zinner SH, Willett WC, Rosner B. Maternal prenatal dietary potassium, calcium, magnesium, and infant blood pressure. *Hypertension*. 1991;17:218-224.
23. Hamet P, Dagnault-Gelinas M, Lambert J, Ledoux M, Whissell-Cambiotti L, Bellavance F, Mongeau E. Epidemiological evidence of an interaction between calcium and sodium intake impacting on blood pressure. *Am J Hypertens*. 1992;5:378-385.
24. Marcoux S, Brisson J, Fabia J. Calcium intake from dairy products and supplements and the risks of preeclampsia and gestational hypertension. *Am J Epidemiol*. 1991;133:1266-1272.
25. Resnick LM, Laragh JH. The hypotensive effect of short-term oral calcium loading in essential hypertension. *Clin Res*. 1983;31:334A. Abstract.
26. De Swiet M. The epidemiology of hypertension in children. *Br Med Bull*. 1986;42:172-175.
27. Criqui MH, Langer RD, Reed DM. Dietary alcohol, calcium, and potassium: independent and combined effects on blood pressure. *Circulation*. 1989;80:609-614.
28. Harlan WR, Harlan LC. Blood pressure and calcium and magnesium intake. In: Laragh JH, Brenner BM, eds. *Hypertension: Pathophysiology, Diagnosis, and Management*. New York, NY: Raven Press, Publishers; 1990:229-240.
29. Wessels F, Hoffmann D, Wagner H, Zunkley H. Influence of family history of hypertension on the relationships between blood pressure, body weight, electrolyte metabolism, renin, prolactin and parathormone. *Clin Sci*. 1981;61(suppl 7):359s-362s.
30. Rosenthal FD, Roy S. Hypertension and hyperparathyroidism. *Br Med J*. 1972;4:396-397.
31. McCarron DA. Low serum concentrations of ionized calcium in patients with hypertension. *N Engl J Med*. 1982;307:226-228.
32. Kesteloot H, Geboers J. Calcium and blood pressure. *Lancet*. 1982;1:813-815.
33. McCarron DA, Pingree PA, Rubin RJ, Gaucher SM, Molitch M, Krutzik S. Enhanced parathyroid function in essential hypertension: a homeostatic response to a urinary calcium leak. *Hypertension*. 1980;2:162-168.
34. Strazzullo P, Nunziata V, Cirillo M, Giannattasio R, Ferrara LA, Mattioli PL, Mancini M. Abnormalities of calcium metabolism in essential hypertension. *Clin Sci*. 1983;65:137-141.
35. Kesteloot H, Joossens JV. Relationship of dietary sodium, potassium, calcium, and magnesium, with blood pressure: Belgian interuniversity research on nutrition and health. *Hypertension*. 1988;12:594-599.
36. Hvarfner A, Bergstrom R, Morlin C, Wide L, Ljunghall S. Relationships between calcium metabolic indices and blood pressure in patients with essential hypertension as compared with a healthy population. *J Hypertens*. 1987;5:451-456.
37. Young EW, Morris CD, McCarron DA. Urinary excretion in essential hypertension. *J Lab Clin Med*. 1992;120:624-632.
38. Luft FC, Zemel MB, Sowers JA, Fineberg NS, Weinberger MH. Sodium bicarbonate and sodium chloride: effects on blood pressure and electrolyte homeostasis in normal and hypertensive man. *J Hypertens*. 1990;8:663-670.
39. Erne P, Bolli P, Burgisser E, Bühler FR. Correlation of platelet calcium with blood pressure: effect of antihypertensive therapy. *N Engl J Med*. 1984;310:1084-1088.
40. Lechi A, Lechi C, Bonadonna G, Sinigaglia D, Corradini P, Polignano R, Arosio E, Covi G, de Togni P. Increased basal and thrombin-induced free calcium in platelets of essential hypertensive patients. *Hypertension*. 1987;9:230-235.
41. Haller H, Lenz T, Thiede M, Lundersdorf M, Kribben A, Distler A, Philipp T. Platelet intracellular free calcium and hypertension. *J Clin Hypertens*. 1987;3:12-19.
42. Pritchard K, Raine AEG, Ashley CC, Castell LM, Somers V, Osborn C, Ledingham JGG, Conway J. Correlation of blood pressure in normotensive and hypertensive individuals with platelet but not lymphocyte intracellular free calcium concentrations. *Clin Sci*. 1989;76:631-635.
43. Cooper R, Lipowski J, Ford E, Shamsi N, Feinberg H, Le Breton G. Increased membrane-bound calcium in platelets of hypertensive patients. *Hypertension*. 1989;13:139-144.
44. Astarie C, Levenson J, Simon A, Meyer P, Devynck M-A. Platelet cytosolic proton and free calcium concentrations in essential hypertension. *J Hypertens*. 1989;7:485-491.
45. Touyz RM, Milne FJ, Reinach SG. Intracellular Mg^{2+} , Ca^{2+} , Na^{+} and K^{+} in platelets and erythrocytes of essential hypertension patients: relation to blood pressure. *Clin Exp Hypertens A*. 1992;14:1189-1209.
46. Le Quan Sang KH, Laude D, Devynck M-A. Platelet cytosolic free Ca^{2+} concentration and serotonin content in essential hypertension. *J Hypertens*. 1987;5:237-240.
47. Hvarfner A, Larsson R, Morlin C, Rastad J, Wide L, Akerstrom G, Ljunghall S. Cytosolic free calcium in platelets: relationships in blood pressure and indices of systemic calcium metabolism. *J Hypertens*. 1988;6:71-77.
48. Wright GL, Rankin GO. Concentrations of ionic and total calcium in plasma of four models of hypertension. *Am J Physiol*. 1982;243:H365-H370.
49. Bindels RJM, Van Den Broek LAM, Jongen MJM, Hackeng WHL, Lowik CWGM, Van Os CH. Increased plasma calcitonin levels in young spontaneously hypertensive rats: role in disturbed phosphate homeostasis. *Pflugers Arch*. 1987;408:395-400.
50. Stern N, Lee DBN, Silis V, Beck FWJ, Deftos L, Manolagas SC, Sowers JR. Effects of high calcium intake on blood pressure and calcium metabolism in young SHR. *Hypertension*. 1984;6:639-646.
51. McCarron DA, Yung NN, Ugoretz BA, Krutzik S. Disturbances of calcium metabolism in the spontaneously hypertensive rat. *Hypertension*. 1981;3(suppl 1):1-162-1-173.
52. Young EW, Hsu CH, Patel S, Simpson RU, Komanicky P. Metabolic degradation and synthesis of calcitriol in spontaneously hypertensive rat. *Am J Physiol*. 1987;252:E778-E782.
53. Hsu CH, Chen PS, Caldwell RM. Renal phosphate excretion in spontaneously hypertensive and normotensive Wistar Kyoto rats. *Kidney Int*. 1984;25:789-795.
54. Ayachi S. Increased dietary calcium lowers blood pressure in the spontaneously hypertensive rat. *Metabolism*. 1979;28:1234-1238.
55. Lucas PA, Brown RC, Drüeke T, Lacour B, Metz JA, McCarron DA. Abnormal vitamin D metabolism, intestinal calcium transport, and bone calcium status in the spontaneously hypertensive rat compared with its genetic control. *J Clin Invest*. 1986;78:221-227.
56. Schedl HP, Miller DL, Horst RL, Wilson HD, Natarajan K, Conway T. Intestinal calcium transport in the spontaneously hypertensive rat: response to calcium depletion. *Am J Physiol*. 1986;250:G412-G419.
57. Young EW, Patel SR, Hsu CH. Plasma $1,25(OH)_2D_3$ response to parathyroid hormone, cyclic adenosine monophosphate, and phosphorus depletion in the spontaneously hypertensive rat. *J Lab Clin Med*. 1986;108:562-566.
58. Kurtz TW, Portale AA, Morris RC Jr. Evidence for a difference in vitamin D metabolism between spontaneously hypertensive and Wistar-Kyoto rats. *Hypertension*. 1986;8:1015-1020.
59. Hsu CH, Chen P-S, Smith DE, Yang C-S. Pathogenesis of hypercalciuria in spontaneously hypertensive rats. *Miner Electrolyte Metab*. 1986;12:130-141.
60. McCarron DA, Shneidman RJ, Lee DM. Ca^{2+} defects in experimental hypertension SHR renal response to chronic infusion of human PTH (1-34). *Kidney Int*. 1987;32(suppl 22):s249-s253.
61. Bhalla RC, Webb RC, Singh D, Ashley T, Brock T. Calcium fluxes, calcium binding, and adenosine cyclic 3'5'-monophosphate-dependent protein kinase activity in the aorta of spontaneously hypertensive and Kyoto Wistar normotensive rats. *Mol Pharmacol*. 1978;14:468-477.
62. Kwan C-Y, Belbeck L, Daniel EE. Abnormal biochemistry of vascular smooth muscle plasma membrane isolated from hypertensive rats. *Mol Pharmacol*. 1980;17:137-140.
63. Aoki K, Yamashita K, Tomita N, Tazumi K, Hotta K. ATPase activity and Ca^{++} binding ability of subcellular membrane of arterial smooth muscle in spontaneously hypertensive rat. *Jpn Heart J*. 1974;15:180-181.
64. Shibata S, Kuchii M, Taniguchi T. Calcium flux and binding in the aortic smooth muscle from the spontaneously hypertensive rat. *Blood Vessels*. 1975;12:279-289.
65. Webb RC, Bhalla RC. Altered calcium sequestration by subcellular fractions of vascular smooth muscle from spontaneously hypertensive rats. *J Mol Cell Cardiol*. 1976;8:651-661.
66. Kwan CY, Daniel EE. Arterial muscle abnormalities of hydralazine treated spontaneously hypertensive rats. *Eur J Pharmacol*. 1982;82:1878-1890.
67. Young EW, Bukoski RD, McCarron DA. Calcium metabolism in experimental genetic hypertension. In: Laragh JH, Brenner BM, eds. *Hypertension: Pathophysiology, Diagnosis, and Management*. New York, NY: Raven Press, Publishers; 1990:977-987.
68. Oshima T, Young EW, Hermsmeyer K, McCarron DA. Modification of platelet and lymphocyte calcium handling and blood pressure by dietary sodium and calcium in genetically hypertensive rats. *J Lab Clin Med*. 1992;119:151-158.

69. Oshima T, Young EW, McCarron DA. Abnormal platelet and lymphocyte calcium handling in prehypertensive rats. *Hypertension*. 1991;18:111-115.
70. Bruschi G, Bruschi ME, Caroppo M, Orlando G, Pavarani C, Cavatorta A. Intracellular free $[Ca^{2+}]$ in circulating lymphocytes of spontaneously hypertensive rats. *Life Sci*. 1984;35:535-542.
71. Vasdev S, Thompson P, Trigg C, Fernandez P, Bolli P, Ananthanarayanan VS. Fura-2 used as a probe to show elevated intracellular free calcium in platelets of Dahl-sensitive rats fed a high salt diet. *Biochem Biophys Res Commun*. 1988;154:380-386.
72. Belizan JM, Villar J, Pineda O, Gonzalez AE, Sainz E, Garrera G, Sibrian R. Reduction of blood pressure with calcium supplementation in young adults. *JAMA*. 1983;249:1161-1166.
73. Luft FC, Aronoff GR, Sloan RS, Fineberg NS, Weinberger MH. Short-term augmented calcium intake has no effect on sodium homeostasis. *Clin Pharmacol Ther*. 1986;39:414-419.
74. Tabuchi Y, Ogihara T, Hashizume K, Saito H, Kumahara Y. Hypotensive effect of long-term oral calcium supplementation in elderly patients with essential hypertension. *J Clin Hypertens*. 1986;3:254-262.
75. Lyle RM, Melby CL, Hyner GC, Edmondson JW, Miller JZ, Weinberger MH. Blood pressure and metabolic effects of calcium supplementation in normotensive white and black men. *JAMA*. 1987;257:1772-1776.
76. Lasaridis AN, Kaisis CN, Zananiri KI, Syrganis CD, Tourkantonis AA. Oral calcium supplementation promotes renal sodium excretion in essential hypertension. *J Hypertens*. 1987;5:S307-S309.
77. Bierenbaum ML, Wolf E, Bisgeier G, Maginnis WP. Dietary calcium: a method of lowering blood pressure. *Am J Hypertens*. 1988;1:149s-152s.
78. Gilliland M, Zawada ET, McClung D, TerWee J. Preliminary report: natriuretic effect of calcium supplementation in hypertensive women over forty. *J Am Coll Nutr*. 1987;6:139-143.
79. Vinson JA, Mazur T, Bose P. Comparison of different forms of calcium on blood pressure of normotensive young males. *Nutr Rep Int*. 1987;36:497-505.
80. Takagi Y, Fukasa M, Takata B, Fujimi T, Fujita T. Calcium treatment of essential hypertension in elderly patients evaluated by 24 H monitoring. *Am J Hypertens*. 1991;4:836-839.
81. Van Beresteijn ECH, Van Schaalk M, Schaafsma G. Milk: does it affect blood pressure? A controlled intervention study. *J Intern Med*. 1990;228:477-480.
82. Buonopane GJ, Kilara A, Smith JS, McCarthy RD. Effect of skim milk supplementation on blood cholesterol concentration, blood pressure, and triglycerides in a free-living human population. *J Am Coll Nutr*. 1992;11:56-67.
83. Belizan JM, Villar J, Zalzar A, Rojas L, Chan D, Bryce GF. Preliminary evidence of the effect of calcium supplementation on blood pressure in normal pregnant women. *Am J Obstet Gynecol*. 1983;146:175-180.
84. Villar J, Repke J, Belizan JM, Pareja G. Calcium supplementation reduces blood pressure during pregnancy: results of a randomized clinical trial. *Obstet Gynecol*. 1987;70:317-322.
85. López-Jaramillo P, Narváez M, Weigel RM, Yépez R. Calcium supplementation reduces the risk of pregnancy-induced hypertension in an Andes population. *Br J Obstet Gynaecol*. 1989;96:648-655.
86. López-Jaramillo P, Narváez M, Yépez R. Effect of calcium supplementation on the vascular sensitivity to angiotensin II in pregnant women. *Am J Obstet Gynecol*. 1987;156:261-262. Letter.
87. Marya RK, Rathee S, Manrow M. Effect of calcium and vitamin supplementation on toxemia of pregnancy. *Gynecol Obstet Invest*. 1987;24:38-42.
88. Belizan JM, Villar J, Gonzalez L, Campodonica L, Berjel E. Calcium supplementation to prevent hypertensive disorders of pregnancy. *N Engl J Med*. 1991;325:1389-1405.
89. Knight KB, Keith RE. Calcium supplementation on normotensive and hypertensive pregnant women. *Am J Clin Nutr*. 1992;55:891-895.
90. Saito K, Sano H, Furuta Y, Fukuzaki H. Effect of oral calcium on blood pressure response in salt-loaded borderline hypertensive patients. *Hypertension*. 1989;13:219-226.
91. Saito K, Sano H, Furuta Y, Yamonishi J, Omatsu T, Ito Y, Fukujaki H. Calcium supplementation in salt-dependent hypertension. *Contrib Nephrol*. 1991;90:25-35.
92. Weinberger MH, Wagner UL, Fineberg NS. The blood pressure effects of calcium supplementation in humans of known sodium responsiveness. *Am J Hypertens*. 1993;6:799-805.
93. Lyle RM. Does baseline serum total calcium level influence the blood pressure response to calcium supplementation? A double-blind study. *Neth J Med*. 1992;41:48-55.
94. Meese RB, Gonzalez DG, Casparian JM, Ram CVS, Pak CYC, Kaplan NM. Failure of calcium supplements to relieve hypertension. *Am J Med Sci*. 1987;294:219-224.
95. Grobbee DE, Hofman A. Calcium supplementation lowers blood pressure in young mildly hypertensive subjects: a randomized, controlled trial. *Lancet*. 1986;2:703-707.
96. Bloomfield RL, Young LD, Zurek G, Felts JH, Straw MK. Effects of oral calcium carbonate on blood pressure in subjects with mildly elevated arterial pressure. *J Hypertens*. 1986;4(suppl 5):s351-s354.
97. Strazzullo P, Siani A, Guglielmi S, DiCarlo A, Galletti F, Cirillo M, Mancini M. Controlled trial of long-term oral calcium supplementation in essential hypertension. *Hypertension*. 1986;8:1084-1088.
98. Thomsen K, Nilas L, Christiansen C. Dietary calcium intake and blood pressure in normotensive subjects. *Acta Med Scand*. 1987;222:51-56.
99. Siani A, Strazzullo P, Guglielmi S, Pacioni D, Giacco A, Iacone R, Mancini M. Controlled trial of low calcium versus high calcium intake in mild hypertension. *J Hypertens*. 1988;6:253-256.
100. Cappuccio FP, Markandu ND, Singer DRJ, Smith SJ, Shore AC, MacGregor GA. Does oral calcium supplementation lower high blood pressure? A double blind study. *J Hypertens*. 1987;5:67-71.
101. Van Beresteijn ECH, Schaafsma G, De Waard H. Oral calcium and blood pressure: a controlled intervention trial. *Am J Clin Nutr*. 1986;44:883-888.
102. Nowson C, Morgan T. Effect of calcium carbonate on blood pressure in normotensive and hypertensive people. *Hypertension*. 1989;13:630-639.
103. Whelton PK, Appel L, Charleston J, Dalcin AT, Ewart C, Fried L, Kaidy D, Klag MJ, Kumanyika S, Steffen L, Walker WG, Oberman A, Counts K, Hataway H, Raczynski J, Rappaport N, Weinsier R, Borhani NO, Bernauer E, Borhani P, De La Cruz C, Ertl A, Heustis D, Lee M, Lovelace W, O'Connor E, Peel L, Sugars C, Taylor JO, Corkery BW, Evans DA, Keough ME, Morris MC, Pistorino E, Sacks F, Cameron M, Corrigan S, Wright NK, Applegate WB, Brewer A, Goodwin L, Miller S, Murphy J, Randle J, Sullivan J, Lasser NL, Batey DM, Dolan L, Hamill S, Kennedy P, Lasser VI, Kuller LH, Caggiula AW, Milas NC, Yamamoto ME, Vogt TM, Greenlick MR, Hollis J, Stevens V, Cohen JD, Mattfeldt-Beman M, Brinkmann C, Roth K, Shepek L, Hennekens CH, Buring J, Cook N, Danielson E, Eberlein K, Gordon D, Hebert P, Macfadyen J, Mayrent S, Rosner B, Satterfield S, Tosteson H, Van Denburgh M, Cutler JA, Brittain E, Farrand M, Kaufmann P, Lakatos E, Obarzanek E, Belcher J, Dommeyer A, Mills I, Neibling P, Woods M, Goldman BJK, Blethen E. The effects of non-pharmacologic interventions on blood pressure of persons with high normal levels. *JAMA*. 1992;267:1213-1220.
104. Kawasaki N, Matsui K, Ito M, Nakamura T, Yoshimura T, Ushijima H, Maeyama M. Effect of calcium supplementation on the vascular sensitivity to angiotensin II in pregnant women. *Am J Obstet Gynecol*. 1985;153:576-582.
105. Morris CD, McCarron DA. Effect of calcium supplementation in an older population with mildly increased blood pressure. *Am J Hypertens*. 1992;5:230-237.
106. Singer DRJ, Markandu ND, Cappuccio FP, Beynon GW, Shore AC, Smith SJ, MacGregor GA. Does oral calcium lower blood pressure: a double-blind study. *J Hypertens*. 1985;3:661-671.
107. Zoccali C, Mallamaci F, Delfino D, Ciccarelli M, Parlongo S, Iellamo D, Moscato D, Maggiore Q. Long-term oral calcium supplementation in essential hypertension: a double-blind, randomized, crossover study. *J Hypertens*. 1990;4(suppl 6):S676-S678.
108. Zoccali C, Mallamaci F, Delfino D, Ciccarelli M, Parlongo S, Iellamo D, Moscato D, Maggiore Q. Double-blind randomized, crossover trial of calcium supplementation in essential hypertension. *J Hypertens*. 1988;6:451-455.
109. Cutler JA, Brittain E. Calcium and blood pressure: an epidemiologic perspective. *Am J Hypertens*. 1990;3:137S-146S.
110. Mikami H, Ogihara T, Tabuchi Y. Blood pressure response to dietary calcium intervention in humans. *Am J Hypertens*. 1990;3:147S-151S.
111. Zemel MB, Gualdoni SM, Sowers JR. Reductions in total and extracellular water associated with calcium-induced natriuresis and the antihypertensive effect of calcium in blacks. *Am J Hypertens*. 1988;1:70-72.

112. Zemel MB, Kraniak J, Standley PR, Sowers JR. Erythrocyte cation metabolism in salt-sensitive hypertensive blacks as affected by dietary sodium and calcium. *Am J Hypertens*. 1988;1:386-392.
113. Ernst FA, Enwonwu CO, Francis RA. Calcium attenuates cardiovascular reactivity to sodium and stress in Blacks. *Am J Hypertens*. 1990;3:451-457.
114. Karanja N, Morris CD, Rufolo P, Singelar G, Illingworth DR, McCarron DA. The impact of increasing calcium through the diet on nutrient composition and plasma lipids in humans. 1994;59: 900-907.
115. Gillman MW, Oliveria SA, Moore LL, Ellison RC. Inverse association of dietary calcium with systolic blood pressure in young children. *JAMA*. 1992;267:2340-2343.
116. Ambroz SL, Shehin SE, Chiou C-Y, Sowers JR, Zemel MB. Effects of dietary calcium on blood pressure, vascular reactivity and vascular smooth muscle calcium efflux rate in Zucker rats. *Am J Hypertens*. 1991;4:592-596.
117. Anderson DE, Murphy P, Kearns P, Kearns W. High calcium intake does not prevent stress-salt hypertension in dogs. *Pav J Biol Sci*. 1986;21:25-31.
118. Anderson S, Grady JR, Ellison DH, McCarron DA. Calcium balance and parathyroid hormone mediated vasodilation in the spontaneously hypertensive rat. *Hypertension*. 1983;5(suppl 1): 1-59-1-63.
119. Ando K, Sato Y, Ono A, Takahashi K, Shimomura T, Ogata E, Fujita T. Antihypertensive effect of dietary calcium loading in angiotensin II-salt rats. *Am J Physiol*. 1991;261:R1070-R1074.
120. Baksi SN. Altered pressor response to norepinephrine in calcium and vitamin D deficient rats. *Clin Exp Hypertens A*. 1988;10: 811-832.
121. Baksi SN. Blood pressure and hypothalamic norepinephrine and dopamine in calcium-deficient rats. *Am J Hypertens*. 1989;2: 462-464.
122. Baksi SN. Hypotensive action of parathyroid hormone in hypoparathyroid and hyperparathyroid rats. *Hypertension*. 1988; 11:509-513.
123. Baksi SN, Abhold RH, Speth RC. Low-calcium diet increases blood pressure and alters peripheral but not central angiotensin II binding sites in rats. *J Hypertens*. 1989;7:423-427.
124. Belizán JM, Pineda O, Sainz E, Menéndez LA, Villar J. Rise of blood pressure in calcium-deprived pregnant rats. *Am J Obstet Gynecol*. 1981;141:163-169.
125. Belizán JM, Villar J, Self S, Pineda O, González I, Sainz E. The mediating role of the parathyroid gland in the effect of low calcium intake on blood pressure in the rat. *Arch Latinoam Nutr*. 1984;34:666-675.
126. Blakeborough P, Neville SG, Rolls BA. The effect of diets adequate and deficient in calcium on blood pressures and the activities of intestinal and kidney plasma membrane enzymes in normotensive and spontaneously hypertensive rats. *Br J Nutr*. 1990;63:65-78.
127. Bodgen JD, Gertner SB, Kemp FW, McLeod R, Bruening KS, Chung HR. Dietary lead and calcium: effects on blood pressure and renal neoplasia in Wistar rats. *J Nutr*. 1991;121:718-728.
128. Bukoski RD, DeWan P, McCarron DA. 1,25(OH)₂ vitamin D₃ modifies growth and contractile function of vascular smooth muscle of spontaneously hypertensive rats. *Am J Hypertens*. 1989; 2:553-556.
129. Bukoski RD, McCarron DA. Altered aortic reactivity and lowered blood pressure associated with high calcium intake. *Am J Physiol*. 1986;251:H976-H983.
130. DiPette DJ, Greulich PE, Nickols GA, Graham GA, Green A, Cooper CW, Holland OB. Effect of dietary calcium supplementation on blood pressure and calciotropic hormones in mineralocorticoid-salt hypertension. *J Hypertens*. 1990;8:515-520.
131. Doris PA. Sodium and hypertension: effect of dietary calcium supplementation on blood pressure. *Clin Exp Hypertens A*. 1985; 7:1441-1456.
132. Doris PA. Interactions between sodium and calcium intake and blood pressure responses to norepinephrine and parathyroid hormone. *Horm Metab Res*. 1988;20:163-167.
133. Evans GH, Weaver CM, Harrington DD, Babbs CF Jr. Association of magnesium deficiency with the blood pressure-lowering effects of calcium. *J Hypertens*. 1990;8:327-337.
134. Fujito K, Yokomatsu M, Ishiguro N, Numahata H, Tomino Y, Koide H. Effects of dietary Ca²⁺ on erythrocyte Na⁺-transport systems in spontaneously hypertensive rats. *Clin Sci*. 1991;81: 107-112.
135. Furspan PB, Rinaldi GJ, Hoffman K, Bohr DF. Dietary calcium and cell membrane abnormality in genetic hypertension. *Hypertension*. 1989;13:727-730.
136. Ganguli MC, Tobian L, Iwai J. Reduction of blood pressure in salt-fed Dahl salt-sensitive rats with diets rich in olive oil, safflower oil or calcium biphosphate but not with calcium carbonate. *J Hypertens*. 1986;4(suppl 5):S168-S169.
137. Geiger H, Bahner U, Palkovits M, Seewaldt B, Heidland A. Is the effect of calcium diet or parathyroidectomy on the development of hypertension in spontaneously hypertensive rats mediated by atrial natriuretic peptide? *Kidney Int*. 1988;34(suppl 25):S93-S97.
138. Hano T, Baba A, Takeda J, Nishio I, Masuyama Y. Antihypertensive effects of oral calcium supplementation in spontaneously hypertensive rats. *Contrib Nephrol*. 1991;90:36-41.
139. Hatton DC, Huie PS, Muntzel MS, Metz JA, Karanja N, McCarron DA. Blood pressure development and serum calcium in suckling spontaneously hypertensive rat pups: effects of maternal dietary calcium. *J Hypertens*. 1986;4(suppl 3):S465-S467.
140. Hatton DC, Huie PE, Muntzel MS, Metz JA, McCarron DA. Stress-induced blood pressure responses in SHR: effect of dietary calcium. *Am J Physiol*. 1987;252:R48-R54.
141. Hatton DC, Muntzel M, McCarron DA, Pressley M, Bukoski RD. Early effects of dietary calcium on blood pressure, plasma volume, and vascular reactivity. *Kidney Int*. 1988;34:16-18.
142. Hatton DC, Scrogin KE, Metz JA, McCarron DA. Dietary calcium alters blood pressure reactivity in spontaneously hypertensive rats. *Hypertension*. 1989;13:622-629.
143. Hatton D, Muntzel M, Absalon J, Lashley D, McCarron DA. Dietary calcium and iron: effects on blood pressure and hematocrit in young spontaneously hypertensive rats. *Am J Clin Nutr*. 1991;53:542-546.
144. Hatton DC, Scrogin KE, Levine D, Feller D, McCarron DA. Dietary calcium modulates blood pressure through an α -1 adrenergic mechanism. *Am J Physiol*. 1993;264:F234-F238.
145. Hattori K, Sano H, Kubota Y, Kawahara J, Miki T, Suzuki H, Fukuzaki H. The effect of Ca and Mg supplementation and the role of the opiodergic system on the development of DOCA-salt hypertension. *Am J Hypertens*. 1991;4:72-75.
146. Hermesmeier K, Hatton DA, Karanja N, McCarron D. Effects of dietary calcium on nimodipine-sensitive calcium channel function in stroke-prone spontaneously hypertensive rats. *Stroke*. 1990; 21(suppl IV):IV-98-IV-101.
147. Huie PE, Hatton DC, Muntzel MS, Metz JA, McCarron DA. Genetic variability in response to dietary calcium. *Life Sci*. 1987; 41:2185-2193.
148. Huie PE, Hatton DC, Muntzel MS, Metz JA, McCarron DA. Psychosocial stress, dietary calcium and hypertension in the spontaneously hypertensive rat. *Physiol Behav*. 1987;40:425-429.
149. Jirakulsomchok S, Mozaffari MS, Oparil S, Wyss JM. Dietary Ca²⁺ increases natriuretic and diuretic responses to volume loading in NaCl-sensitive spontaneously hypertensive rats. *J Hypertens*. 1990;8:947-951.
150. Jones MR, Ghaffari F, Tomerson BW, Clemens RA. Hypotensive effect of a high calcium diet in the Wistar rat. *Miner Electrolyte Metab*. 1986;12:85-91.
151. Jones MR, Huibonhoa A. Effect of marginal calcium intake on blood pressure in the Wistar rat. *Clin Exp Hypertens A*. 1988;10: 801-810.
152. Kageyama Y, Suzuki H, Hayashi K, Saurta T. Effects of calcium loading on blood pressure in spontaneously hypertensive rats: attenuation of the vascular reactivity. *Clin Exp Hypertens A*. 1986; 8:355-370.
153. Kageyama Y, Suzuki H, Arima K, Saruta T. Oral calcium treatment lowers blood pressure in renovascular hypertensive rats by suppressing the renin-angiotensin system. *Hypertension*. 1987; 10:375-382.
154. Kageyama Y, Suzuki H, Arima K, Kondo K, Saruta T. Effects of calcium loading in DOCA-salt hypertensive rats. *Jpn Circ J*. 1987; 51:1315-1324.
155. Kageyama Y, Bravo EL. Neurohumoral and hemodynamic responses to dietary calcium supplementation in deoxycorticosterone-salt hypertensive dogs. *Hypertension*. 1987;9(suppl III): III-166-III-170.
156. Kang JS, Cregor MD, Smith JB. Effect of calcium on blood pressure, platelet aggregation and erythrocyte sodium transport in Dahl salt-sensitive rats. *J Hypertens*. 1990;8:245-250.
157. Karanja N, Metz JA, Mercer LP, McCarron DA. Analysis of BP response and Ca²⁺ metabolism using the saturation kinetics model. *Am J Physiol*. 1987;253:R501-R508.

158. Karanja N, Phanouvong T, McCarron DA. Blood pressure in spontaneously hypertensive rats fed butterfat, corn oil, or fish oil. *Hypertension*. 1989;14:674-679.
159. Kohno M, Murakawa K-I, Yasunari K, Yokokawa K, Kurihara N, Takeda T. Possible involvement of atrial natriuretic factor in the antihypertensive action of a high-calcium diet in spontaneously hypertensive and Wistar-Kyoto rats. *Metabolism*. 1989;38:997-1004.
160. Koide M, Tuan RS. Adrenergic regulation of calcium-deficient hypertension in chick embryos. *Am J Physiol*. 1989;257:H1200-H1209.
161. Kurtz TW, Morris RC Jr. Attenuation of deoxycorticosterone-induced hypertension by supplemental dietary calcium. *J Hypertens*. 1986;4(suppl 5):S129-S131.
162. Lau K, Chen S, Eby B. Evidence for the role of PO_4 deficiency in antihypertensive action of a high-Ca diet. *Am J Physiol*. 1984;246:H324-H331.
163. Lau K, Thomas D, Salvi D, Martin C, Tan S. Effects of high-calcium and/or high-sodium diet on basal and angiotensin II-stimulated blood pressure in the spontaneously hypertensive rat. *J Hypertens*. 1986;4(suppl 5):S126-S128.
164. Lau K, Gaftan U, Rydell D, Eby B, Pesigan M, Tropp I, Garro J, Zikos D. Evidence against the role of calcium deficiency in genetic hypertension. *Hypertension*. 1986;8:45-49.
165. Lee JC, McGrath CJ, Leighton AT Jr, Gross WB. Effect of dietary calcium on blood pressure of turkeys. *Poult Sci*. 1984;63:993-996.
166. Lewanczuk RZ, Chen A, Pang PTK. The effect of dietary calcium on blood pressure in spontaneously hypertensive rats may be mediated by parathyroid hypertensive factor. *Am J Hypertens*. 1990;3:349-353.
167. Luft FC, Ganten U, Meyer D, Steinberg H, Gless KJ, Unger T, Ganten D. Effect of high calcium diet on magnesium, catecholamines, and blood pressure of stroke-prone spontaneously hypertensive rats. *Proc Soc Exp Biol Med*. 1988;187:474-481.
168. Mangiarua EI, Wright GL, McCumbee WD. Effect of dietary calcium on *in vitro* aortic tissue responsiveness to a hypertensive factor. *Clin Exp Hypertens A*. 1990;12:1255-1279.
169. McCarron DA. Blood pressure and calcium balance in the Wistar-Kyoto rat. *Life Sci*. 1982;30:683-689.
170. McCarron DA, Lucas PA, Shneidman RJ, Lacour B, Drücke T. Blood pressure development of the spontaneously hypertensive rat after concurrent manipulations of dietary Ca^{2+} and Na^+ : relation to intestinal Ca^{2+} fluxes. *J Clin Invest*. 1985;76:1147-1154.
171. Muntzel MS, Hatton DC, Metz JA, McCarron DA. Dietary calcium alters blood pressure in neonatal spontaneously hypertensive rats. *Am J Hypertens*. 1989;2:158-162.
172. Oparil S, Chen YF, Jin HK, Yang RH, Wyss JM. Dietary Ca^{2+} prevents NaCl -sensitive hypertension in spontaneously hypertensive rats via sympatholytic and renal effects. *Am J Clin Nutr*. 1991;54:227S-236S.
173. Pamnani MB, Chen S, Bryant HJ, Schooley JF, Haddy FJ. Effect of increased dietary calcium on the development of reduced renal mass saline hypertension in rats. *J Am Coll Nutr*. 1990;9:35-43.
174. Pang PKT, Benishin CG, Lewanczuk RZ. Combined effect of dietary calcium and calcium antagonists on blood pressure reduction in spontaneously hypertensive rats. *J Cardiovasc Pharmacol*. 1992;19:442-446.
175. Pernot F, Berthelot A, Gairard A. Pression artérielle et régime hypercalcique chez le Rat normal ou en cours d'hypertension minéralocorticoïde (DOCA + chlorure de sodium). *CR Seances Soc Biol Fil*. 1978;172:1214-1219.
176. Pernot F, Schleiffer R, Berthelot A, Vincent M, Sassard J, Gairard A. Calcium alimentaire et hypertension artérielle chez le rat. *Arch Mal Coeur Vaiss*. 1985;11:1725-1729.
177. Pernot F, Schleiffer R, Bergmann C, Vincent M, Sassard J, Gairard A. Dietary calcium, vascular reactivity, and genetic hypertension in the Lyon rat strain. *Am J Hypertens*. 1990;3:846-853.
178. Peuler JD, Morgan DA, Mark AL. High calcium diet reduces blood pressure in Dahl salt-sensitive rats by neural mechanisms. *Hypertension*. 1987;9(suppl III):III-159-III-165.
179. Peuler JD, Mark AL. Aggravation of salt-induced hypertension in Dahl rats by 2% supplemental dietary calcium. *Hypertension*. 1989;13:929-934.
180. Peuler JD. Contrasting hemodynamic effects of high oral calcium in genetic models of salt-sensitive hypertension. *Clin Exp Hypertens A*. 1991;13:709-717.
181. Peuler JD, Schelper RL. Partial protection from salt-induced stroke and mortality by high oral calcium in hypertensive rats. *Stroke*. 1992;23:532-538.
182. Porsti I, Arvola P, Wuorela H, Ilkka M, Saynvalammi P, Huhtala H, Metsa-Ketela T, Vapaatalo V. Effects of a high calcium diet and deoxycorticosterone on vascular smooth muscle responses in spontaneously hypertensive rats. *J Hypertens*. 1990;8:835-841.
183. Porsti I, Arvola P, Wuorela H, Vapaatalo H. High calcium diet augments vascular potassium relaxation in hypertensive rats. *Hypertension*. 1992;19:85-92.
184. Porsti I. Arterial smooth muscle contractions in spontaneously hypertensive rats on a high-calcium diet. *J Hypertens*. 1992;10:255-263.
185. Resnick LM, Gupta RK, Sosa RE, Corbett ML, Sealey JE, Laragh JH. Effects of altered dietary calcium intake in experimental hypertension: role of intracellular free magnesium. *J Hypertens*. 1986;4(suppl 5):S182-S185.
186. Resnick LM, Sosa RE, Corbett ML, Gertner JM, Sealey JE, Laragh JH. Effects of dietary calcium on sodium volume vs renin-dependent forms of experimental hypertension. *Trans Assoc Am Physicians*. 1986;99:172-179.
187. Schleiffer R, Pernot F, Berthelot A, Gairard A. Low calcium diet enhances development of hypertension in the spontaneously hypertensive rat. *Clin Exp Hypertens A*. 1984;6:783-793.
188. Scrogin KE, Hatton DC, McCarron DA. The interactive effects of dietary sodium chloride and calcium on cardiovascular stress responses. *Am J Physiol*. 1991;261:R945-R949.
189. Scrogin KE, Hatton DC, McCarron DA. Effects of dietary sodium and calcium on blood pressure reactivity in the SHR and WKY. *Clin Exp Hypertens A*. 1991;13:699-707.
190. Semafuko WEB, Morris DJ. Effect of high calcium diet on the development of high blood pressure in intact spontaneously hypertensive rats and in adrenalectomized spontaneously hypertensive rats treated with aldosterone. *Steroids*. 1991;56:131-135.
191. Stern N, Golub M, Nyby M, Berger M, Eggena P, Lee DBN, Tuck ML, Brickman AS. Effect of high calcium intake on pressor responsiveness in hypertensive rats. *Am J Physiol*. 1987;252:H1112-H1119.
192. Tamura H. Influence of a high calcium diet on the development of hypertension in the spontaneously hypertensive rat. *Jpn J Nephrol*. 1987;29:107-113.
193. Tenner TE, Buddingh F, Yang M-L, Yang C-M, Pang PKT. Dietary calcium and development of hypertension in spontaneously hypertensive rats. *Magnesium*. 1989;8:288-298.
194. Togari A, Shintani S, Arai M, Matsumoto S, Nagatsu T. Acute effect of parathyroid hormone-(1-34) fragment on blood pressure in rats fed a low calcium diet. *Gen Pharmacol*. 1990;21:547-549.
195. Tresham JJ, McGuire P, Coghlan JP, Whitworth JA, Scoggins BA. The effects of calcium and vitamin D on blood pressure in conscious sheep. *Clin Exp Hypertens A*. 1988;10:1085-1106.
196. Wiecek A, Kuczerka M, Ganten U, Ritz E, Mann JFE. Influence of parathyroidectomy on blood pressure and vascular reactivity in spontaneously hypertensive rats. *Clin Exp Hypertens A*. 1989;11:1515-1533.
197. Wuorela H, Porsti I, Siltaloppi E, Arvola Nurmi A-K, Saynvalammi P, Vapaatalo H. Deoxycorticosterone prevents the blood pressure-lowering effect of calcium in spontaneously hypertensive rats. *Methods Find Exp Clin Pharmacol*. 1990;12:37-42.
198. Wuorela H, Arvola P, Porsti I, Siltaloppi E, Saynvalammi P, Vapaatalo H. The effect of high calcium intake on Ca^{2+} ATPase and the tissue Na:K ratio in spontaneously hypertensive rats. *Naunyn Schmiedeberg Arch Pharmacol*. 1992;345:117-122.
199. Wyss JM, Chen Y-F, Meng QC, Jin HK, Jirakulsomchok S, Oparil S. Dietary Ca^{2+} prevents NaCl -induced exacerbation of hypertension and increases hypothalamic norepinephrine turnover in spontaneously hypertensive rats. *J Hypertens*. 1989;7:711-719.
200. Yang R-H, Jin H, Chen Y-F, Oparil S, Wyss JM. Dietary Ca^{2+} supplementation prevents the exaggerated responsiveness of anterior hypothalamic α_2 -adrenoceptors in NaCl -loaded spontaneously hypertensive rats. *J Cardiovasc Pharmacol*. 1989;13:162-167.
201. American Institute for Nutrition Ad Hoc Committee on Animal Nutrition. 1976 Report on Animal Nutrition. *J Nutr*. 1977;107:1340-1348.
202. Charles P. Calcium absorption and calcium bioavailability. *J Intern Med*. 1992;231:161-168.
203. Metz JA, Karanja N, Torok J, McCarron DA. Modification of total body fat in spontaneously hypertensive rats and Wistar-Kyoto rats by dietary calcium and sodium. *Am J Hypertens*. 1988;1:58-60.

204. Dominiczak AF, Lazar DF, Das AK, Bohr DF. Lipid bilayer in genetic hypertension. *Hypertension*. 1991;18:748-757.
205. Bohr DF. Vascular smooth muscle: dual effect of calcium. *Science*. 1963;139:597-599.
206. Bohr DF. What makes the pressure go up? A hypothesis. *Hypertension*. 1981;3(suppl II):11-160.
207. Rinaldi G, Bohr DF. Potassium-induced relaxation of the arteries in hypertension: modulation by extracellular calcium. *Am J Physiol*. 1989;256:H707-H712.
208. Bohr DF, Wu CC. Mechanisms of calcium relaxation of vascular smooth muscle. *Am J Physiol*. 1991;261:H1411-H1416.
209. Dominiczak AF, Bohr DF. Cell membrane abnormalities and the regulation of intracellular calcium concentration in hypertension. *Clin Sci*. 1990;79:415-423.
210. Furspan PB, Bohr DF. Calcium-related abnormalities in lymphocytes from genetically hypertensive rats. *Hypertension*. 1986;8(suppl II):11-123-II-126.
211. Shiffman FH, Bose R. A role of calcium in altered sodium ion transport of hypertensives? *Life Sci*. 1988;42:1573-1581.
212. Roulet CM, Roulet J-B, Duchambon P, Thomasset M, Lacour B, McCarron DA, Drieker T. Abnormal intestinal regulation of calbindin-D9K and calmodulin by dietary calcium in genetic hypertension. *Am J Physiol*. 1991;261:F474-F480.
213. Larsen FL, Vincenzi FF. Calcium transport across the plasma membrane: stimulation by calmodulin. *Science*. 1979;204:306-308.
214. Cox JA, Comte M, Stein EA. Activation of human erythrocyte Ca^{2+} -dependent Mg^{2+} -activated ATPase by calmoduline and calcium: quantitative analysis. *Proc Natl Acad Sci U S A*. 1982;79:4265-4269.
215. Drücke TB. Mechanisms of action of calcium absorption: factors that influence bioavailability. In: Langford H, Levine B, Ellenbogen L, eds. *Nutritional Factors in Hypertension*. New York, NY: Alan R Liss, Inc; 1990:155-173.
216. Hellstrom J, Birke G, Edvall CA. Hypertension in hyperparathyroidism. *Br J Urol*. 1958;30:13-24.
217. Leug MC. Hypertension and primary hyperparathyroidism: a five year case review. *South Med J*. 1982;75:1371-1374.
218. Grobbee DE, Hackeng WHL, Birkenhager JC, Hofman A. Intact parathyroid hormone (1-84) in primary hypertension. *Clin Exp Hypertens A*. 1986;8:299-308.
219. Grobbee DE, Hackeng WHL, Birkenhager JC, Hofman A. Raised plasma intact parathyroid hormone concentrations in young people with mildly raised blood pressure. *Br Med J*. 1988;296:814-816.
220. Zachariah PK, Schwartz GL, Strong CG, Ritter SG. Parathyroid hormone and calcium: a relationship in hypertension. *Am J Hypertens*. 1988;1:79S-82S.
221. Ellison DH, Shneidman R, Morris C, McCarron DA. Effects of calcium infusion on blood pressure in hypertensive and normotensive humans. *Hypertension*. 1986;8:497-505.
222. Young EW, McCarron DA, Morris CD. Calcium regulating hormones in essential hypertension. *Am J Hypertens*. 1990;3:161S-166S.
223. Van Hooft IMS, Grobbee DE, Frölich M, Pols HAP, Hofman A. Alterations in calcium metabolism in young people at risk for primary hypertension. *Hypertension*. 1993;21:267-272.
224. Crass MF III, Moore PL, Strickland ML, Pang PKT, Citak MS. Cardiovascular responses to parathyroid hormone. *Am J Physiol*. 1985;249:E187-E194.
225. DiPette DJ, Christenson W, Nickols MA, Nickols GA. Cardiovascular responsiveness to parathyroid hormone (PTH) and PTH-related protein in genetic hypertension. *Endocrinology*. 1992;130:2045-2051.
226. Pang PKT, Lewanczuk RZ. Parathyroid origin of a new circulating hypertensive factor in spontaneously hypertensive rats. *Am J Hypertens*. 1989;2:898-902.
227. Mann JFE, Wiecek A, Bommer J, Ganten U, Ritz E. Effects of parathyroidectomy on blood pressure in spontaneously hypertensive rats. *Nephron*. 1987;45:46-52.
228. Gairard A, Berthelot A, Schleiffer R, Pernot F. Parathyroidectomy significantly decreases hypertension in spontaneously hypertensive and deoxycorticosterone plus saline treated rats. *Can J Physiol Pharmacol*. 1982;60:208-212.
229. Neuser D, Schulte-Brinkmann R. Influence of transplantation of parathyroid glands on blood pressure development in stroke prone spontaneously hypertensive rats and in normotensive Wistar Kyoto rats. *Clin Exp Hypertens A*. 1991;13:325-338.
230. Neuser D, Schulte-Brinkmann R, Kazda S. Development of hypertension in WKY rats after transplantation of parathyroid glands from SHR/SP. *J Cardiovasc Pharmacol*. 1990;6:971-974.
231. Pang PKT, Tenner TE Jr, Yee JA, Yang M, Janssen HF. Hypotensive action of parathyroid hormone preparations on rats and dogs. *Proc Natl Acad Sci U S A*. 1980;77:675-678.
232. Nickols GA. Increased cAMP in cultured vascular smooth muscle cells and relaxation of aortic strips by parathyroid hormone. *Eur J Pharmacol*. 1985;116:137-144.
233. Zemel MB, Sowers JR. Salt sensitivity and systemic hypertension in the elderly. *Am J Cardiol*. 1988;61:7h-12h.
234. Altura BM, Altura BT. Peripheral and cerebrovascular actions of ethanol, acetaldehyde, and acetate: relationship to divalent cations in alcoholism. *Clin Exp Res Commun*. 1987;11:99-111.
235. Bogin E, Massry SG, Harary I. Effect of parathyroid hormone on rat heart cells. *J Clin Invest*. 1981;67:1215-1227.
236. Reid IR, Civitelli R, Halstead LR, Avioli LV, Hruska KA. Parathyroid hormone acutely elevates intracellular calcium in osteoblast-like cells. *Am J Physiol*. 1987;252:E45-E51.
237. Oshima T, Schleiffer R, Young EW, McCarron DA, Bukoski RD. Parathyroidectomy lowers blood pressure independently of changes in platelet free calcium. *J Hypertens*. 1991;9:155-158.
238. Brickman A, Nyby M, von Hungen K, Eggena P, Tuck M. Parathyroid hormone, platelet calcium, and blood pressure in normotensive subjects. *Hypertension*. 1991;18:176-182.
239. Delbarre B, Delbarre G, Caulin A. Opposite effects exist in cardiovascular control after central and peripheral administration of calcitonin (CT) and parathyroid hormone (PTH). *Clin Exp Hypertens A*. 1984;6:2079-2084.
240. Geiger H, Palkovits M, Schmid G, Bahner U, Heidland A. Altered cAMP content in specific brain areas of spontaneously hypertensive rats dependent on calcium status or parathyroidectomy. *Am J Nephrol*. 1986;6(suppl 1):139-144.
241. Islam A, Smogorzewski M, Massry SG. Effect of chronic renal failure and parathyroid hormone on phospholipid content of brain synaptosomes. *Am J Physiol*. 1989;256:F705-F710.
242. Smogorzewski M, Piskorska G, Borum PR, Massry SG. Chronic renal failure, parathyroid hormone and fatty acids oxidation in skeletal muscle. *Kidney Int*. 1988;33:555-560.
243. Lewanczuk RZ, Wang J, Zhang Z-R, Pang PKT. Effects of spontaneously hypertensive rat plasma on blood pressure and tail artery calcium uptake in normotensive rats. *Am J Hypertens*. 1989;2:26-31.
244. Resnick LM, Gupta RK, Lewanczuk RZ, Pang PKT, Laragh JH. Intracellular ions in salt-sensitive essential hypertension: possible role of calcium-regulating hormones. *Contrib Nephrol*. 1991;90:88-93.
245. Goff JP, Reinhardt TA, Beckman MJ, Horst RL. Contrasting effects of exogenous 1,25-dihydroxyvitamin D [$1,25\text{-(OH)}_2\text{D}$] versus endogenous $1,25\text{-(OH)}_2\text{D}$, induced by dietary calcium restriction, on vitamin D receptors. *Endocrinology*. 1990;126:1031-1035.
246. Koh E, Morimoto S, Fukuo K, Itoh K, Hironaka T, Shiraishi T, Onishi T, Kumahara Y. $1,25\text{-Dihydroxyvitamin D}_3$ binds specifically to rat vascular smooth muscle cells and stimulates their proliferation in vitro. *Life Sci*. 1988;42:215-223.
247. Mitsuhashi T, Morris RC, Ives HE. $1,25\text{-Dihydroxyvitamin D}_3$ modulates growth of vascular smooth muscle cells. *J Clin Invest*. 1991;87:1889-1895.
248. Kawashima H. $1,25\text{-Dihydroxyvitamin D}_3$ stimulates Ca-ATPase in a vascular smooth muscle cell line. *Biochem Biophys Res Commun*. 1988;150:1138-1143.
249. Inoue T, Kawashima H. $1,25\text{-Dihydroxyvitamin D}_3$ stimulates $^{45}\text{Ca}^{2+}$ uptake by cultured vascular smooth muscle cells derived from rat aorta. *Biochem Biophys Res Commun*. 1988;152:1388-1394.
250. Bukoski RD, Xue H, McCarron DA. Effect of $1,25\text{(OH)}_2$ vitamin D_3 and ionized Ca^{2+} on ^{45}Ca uptake by primary cultures of aortic myocytes of spontaneously hypertensive and Wistar Kyoto normotensive rats. *Biochem Biophys Res Commun*. 1987;146:1330-1335.
251. Xue H, McCarron DA, Bukoski RD. $1,25\text{(OH)}_2$ vitamin D_3 -induced ^{45}Ca uptake in vascular myocytes cultured from spontaneously hypertensive and normotensive rats. *Life Sci*. 1991;49:651-659.
252. Bukoski RD, Wang D, Wagman DW. Injection of $1,25\text{-(OH)}_2$ vitamin D_3 enhances resistance artery contractile properties. *Hypertension*. 1990;16:523-531.
253. Hatton DC, Xue H, DeMerritt JA, McCarron DA. $1,25\text{(OH)}_2$ vitamin D_3 -induced alterations in vascular reactivity in the SHR. *Am J Med Sci*. 1994;307:S154-S158.

254. Shimomura T, Ando K, Fujita T. Enhancement of vasoconstrictor response by a noncalcemic analogue of vitamin D₃. *Hypertension*. 1993;21:253-258.
255. Bukoski RD, Xue H. On the vascular inotropic action of calcitriol. *Am J Hypertens*. 1993;6:388-396.
256. Bukoski RD, Li J, Bo J. Effect of long-term administration of 1,25(OH)₂ vitamin D₃ on blood pressure and resistance artery contractility in the spontaneously hypertensive rat. *Am J Hypertens*. 1993;6:944-950.
257. Clementi G, Rapisarda E, Giore CE, Prato A, Amico Roxas M, Millia C, Bernardine R, Maugeri S, Scapagnini U. Effects of salmon calcitonin on plasma renin activity and systolic blood pressure in the rat. *Neurosci Lett*. 1986;66:351-355.
258. Wegener LL, McCarron DA. Acute intravenous calcitonin: failure to modify systemic blood pressure. *Peptides*. 1988;9:1191-1193.
259. Fisher JA, Sager SM, Martin JB. Characterization and regional distribution of calcitonin binding sites in the rat brain. *Life Sci*. 1981;29:663-671.
260. de Beaurepaire R, Freed WJ. Anatomical mapping of the rat hypothalamus for calcitonin-induced anorexia. *Pharmacol Biochem Behav*. 1987;27:177-182.
261. Martín MI, Goicoechea C, Colado MI, Alfaro MJ. Analgesic effect of salmon-calcitonin administered by two routes: effect on morphine analgesia. *Eur J Pharmacol*. 1992;224:77-82.
262. Chihara K, Iwasaki J, Iwasaki Y, Minamitani N, Kaji H, Fujita T. Central nervous system effect of calcitonin: stimulation of prolactin release in rats. *Brain Res*. 1982;248:331-339.
263. Okimura Y, Chihara K, Abe H, Kita T, Kashio Y, Sato M, Fujita T. Calcitonin gene-related peptide-like immunoreactivity in the central nervous system and peripheral organs of rats. *Regul Pept*. 1987;17:327-337.
264. Gardiner SM, Compton AM, Bennett T. Regional hemodynamic effects of calcitonin gene-related peptide. *Am J Physiol*. 1989;256:R332-R338.
265. DiPette DJ, Schwarzenberger K, Kerr N, Holland OB. Systemic and regional hemodynamic effects of calcitonin gene-related peptide. *Hypertension*. 1987;9(suppl III):III-142-III-146.
266. Vallejo M, Lightman S, Marshall I. Central cardiovascular effects of calcitonin gene-related peptide: interaction with noradrenaline in the nucleus tractus solitarius of rats. *Exp Brain Res*. 1988;70:221-224.
267. Brown MR, Gray TS. Peptide injections into the amygdala of conscious rats: effects on blood pressure, heart rate and plasma catecholamines. *Regul Pept*. 1988;21:95-106.
268. Rochford J, Yashpal K, Henry JL. Intrathecal administration of calcitonin gene-related peptide (CGRP) increases heart rate and decreases arterial pressure in the urethane anesthetized rat. *Brain Res Bull*. 1990;25:809-816.
269. DiPette DJ, Westlund KN, Holland OB. Dietary calcium modulates spinal cord content of calcitonin gene-related peptide in the rat. *Neurosci Lett*. 1988;95:335-340.
270. Xu D, Wang X, Wang J-P, Yuan Q-X, Fiscus RR, Chang J-K, Tang J. Calcitonin gene-related peptide (CGRP) in normotensive and spontaneously hypertensive rats. *Peptides*. 1989;10:309-312.
271. Winternitz SR, Wyss JM, Meadows JR, Oparil S. Increased noradrenaline content of hypothalamic nuclei in association with worsening of hypertension after high sodium intake in the young spontaneously hypertensive rat. *Clin Sci Mol Med*. 1982;63:339s-342s.
272. Winternitz SR, Oparil S. Sodium-neural interactions in the development of spontaneous hypertension. *Clin Exp Hypertens A*. 1982;4:751-760.
273. Oparil S, Meng QC, Chen Y-F, Yang R-H, Jin H, Wyss JM. Genetic basis of NaCl-sensitive hypertension. *J Cardiovasc Pharmacol*. 1988;12(suppl 3):S56-S69.
274. Chen Y-F, Meng Q, Wyss JM, Jin H, Oparil S. High NaCl diet reduces hypothalamic norepinephrine turnover in hypertensive rats. *Hypertension*. 1988;11:55-62.
275. Kawano Y, Ferrario CM. Neurohormonal characteristics of cardiovascular response due to intraventricular hypertonic NaCl. *Am J Physiol*. 1984;247:H422-H428.
276. Gavras H, Bain GT, Bland L, Vlahakos D, Gavras I. Hypertensive response to saline microinjection in the area of the nucleus tractus solitarius of the rat. *Brain Res*. 1985;343:113-199.
277. Borowitz JL, Stebbins G, Isom G. Studies of the hypotensive effect of calcium injected into lateral brain ventricles. *Res Commun Chem Pathol Pharmacol*. 1987;55:67-74.
278. Harris RA, Carnes DL, Forte LR. Reduction of brain calcium after consumption of diets deficient in calcium or vitamin D. *J Neurochem*. 1981;36:460-466.
279. Tai C-Y, Smith QR, Rapoport SI. Calcium influxes into brain and cerebrospinal fluid are linearly related to plasma ionized calcium concentration. *Brain Res*. 1986;385:227-236.
280. Murphy VA, Smith QR, Rapoport SI. Homeostasis of brain and cerebrospinal fluid calcium concentrations during chronic hypo- and hypercalcemia. *J Neurochem*. 1986;47:1735-1741.
281. Gavras H. How does salt raise blood pressure? A hypothesis. *Hypertension*. 1986;8:83-88.
282. Rouot BM, U'Prichard DC, Snyder SH. Multiple α_2 -noradrenergic receptor sites in rat brain: selective regulation of high-affinity [³H]clonidine binding by guanine nucleotides and divalent cations. *J Neurochem*. 1980;34:374-384.
283. Baksi SN, Hughes MJ. Alteration of adrenal catecholamine levels in the rat after dietary calcium and vitamin D deficiencies. *J Auton Nerv Syst*. 1984;11:393-396.
284. Hagihara M, Togari A, Matsumoto S, Nagatsu T. Dietary calcium deprivation increased the levels of plasma catecholamines and catecholamine-synthesizing enzymes of adrenal glands in rats. *Biochem Pharmacol*. 1990;39:1229-1231.
285. Felicetta JV, Beck FWJ, Metz JA, McCarron DA, Sowers JR. Diet and catecholamines in the spontaneously hypertensive rat. *J Hypertens*. 1986;4(suppl 5):S55-S57.
286. Friedman PA, Figueiredo JF, Maack T, Windhager EE. Sodium-calcium interactions in the renal proximal convoluted tubule of the rabbit. *Am J Physiol*. 1981;240:F558-F568.
287. Wesson DE. Hypocalcemia-associated modulation of renal response to acute volume expansion in rats. *Am J Physiol*. 1987;253:F726-F733.
288. Lahera V, Fiksen-Olsen MJ, Romero JC. Stimulation of renin release by intrarenal calcium infusion. *Hypertension*. 1990;15(suppl 1):I-149-I-152.
289. Levi M, Henrich WL. Dietary calcium modulates renal BBM angiotensin II binding and Na⁺-H⁺ antiporter activity in SHR. *Am J Physiol*. 1991;29:F657-F662.
290. Hamet P, Mongeau E, Lambert J, Bellavance F, Daignault-Gelinas M, Ledoux M, Whissell-Cambiotti L. Interactions among calcium, sodium, and alcohol intake as determinants of blood pressure. *Hypertension*. 1991;17(suppl 1):I-150-I-154.
291. Resnick LM. Calcium and hypertension: the emerging connection. *Ann Intern Med*. 1985;103:944-945.
292. Resnick LM, Nicholson JP, Laragh JH. Calcium metabolism and the renin-aldosterone system in essential hypertension. *J Cardiovasc Pharmacol*. 1985;7(suppl 6):S187-S193.
293. Gregor JL, Krashoc CL, Krzykowski CE. Calcium, sodium and chloride interactions in rats. *Nutr Res*. 1987;7:401-412.
294. Galletti F, Rutledge A, Triggie DJ. Dietary sodium intake: influence on calcium channels and urinary calcium excretion in spontaneously hypertensive rats. *Biochem Pharmacol*. 1991;41:893-896.
295. Kotchen TA, Ott CE, Whitescarver SA, Resnick LM, Gertner JM, Blehschmidt NG. Calcium and calcium regulating hormones in the 'prehypertensive' Dahl salt sensitive rat (calcium and salt sensitive hypertension). *Am J Hypertens*. 1989;2:747-753.
296. Tordoff MG, Ulrich PM, Schulkin J. Calcium deprivation increases salt intake. *Am J Physiol*. 1990;259:R411-R419.
297. Tordoff MG. Influence of dietary calcium on sodium and calcium intake of spontaneously hypertensive rats. *Am J Physiol*. 1992;262:R370-R381.
298. Tordoff MG. Calcium deprivation increases NaCl intake of Fischer-344 rats. *Physiol Behav*. 1991;49:113-155.
299. Hamet P, Tremblay J. Influence of dietary calcium on the expression of the HSP-70 gene and blood pressure. Presented at 14th Scientific Meeting of the International Society of Hypertension; June 1992; Madrid, Spain.

KEY WORDS • Ca²⁺-transporting ATPase • parathyroid hormones • calcitriol • sympathetic nervous system • natriuresis • muscle, smooth, vascular

Dietary calcium and blood pressure in experimental models of hypertension. A review.

D C Hatton and D A McCarron

Hypertension. 1994;23:513-530

doi: 10.1161/01.HYP.23.4.513

Hypertension is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 1994 American Heart Association, Inc. All rights reserved.

Print ISSN: 0194-911X. Online ISSN: 1524-4563

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://hyper.ahajournals.org/content/23/4/513>

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Hypertension* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

Reprints: Information about reprints can be found online at:
<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Hypertension* is online at:
<http://hyper.ahajournals.org/subscriptions/>